

TWO RESULTS ON THE 2-LOCAL EHP SPECTRAL SEQUENCE

M. G. BARRATT, F. COHEN, B. GRAY, M. MAHOWALD, AND W. RICHTER

(Communicated by Thomas Goodwillie)

ABSTRACT. The E_2 -term of the 2-local EHP spectral sequence is shown to be a $\mathbb{Z}/2$ module. 4 is the order of the identity map on the double loop space of the fiber $W(n)$ of the double suspension $E^2: S^{2n-1} \rightarrow \Omega^2 S^{2n+1}$.

1. INTRODUCTION

Restrict attention to the category of 2-local spaces. The EHP fibrations [6, 1]

$$\Omega^2 S^{2q+1} \xrightarrow{P} S^q \xrightarrow{E} \Omega S^{q+1} \xrightarrow{H} \Omega S^{2q+1}$$

give a tower of fibrations converging to $Q(S^0)$, whose homotopy spectral sequence is the EHP spectral sequence [9] $E_1^{p,q} = \pi_{p+q}(S^{2q-1}) \Rightarrow \pi_p^S$, with differentials $d_r: E_r^{p,q} \rightarrow E_r^{p-r, q-1}$. James [6] proved that $2^{2n}\pi_*(S^{2n+1}) = 0$, by showing that $2\pi_*(S^{2n+1}) \subset \text{Im}(E)$ and $E(2\pi_*(S^{2n})) \subset \text{Im}(E^2)$. Thus the E_∞ -term of the EHP spectral sequence is a $\mathbb{Z}/2$ module. James's work was translated to spaces [1, §5]:

Lemma 1.1. (1) $\Omega^2 S^{2n+1} \xrightarrow{\Omega H} \Omega^2 S^{4n+1} \xrightarrow{2} \Omega^2 S^{4n+1}$ is nullhomotopic.

(2) There exists a map ϕ making the following diagram homotopy commutative.

$$\begin{array}{ccccc} \Omega^2 S^{2n} & \xrightarrow{2} & \Omega^2 S^{2n} & \xrightarrow{\Omega^2 E} & \Omega^3 S^{2n+1} \\ & & & & \uparrow \Omega E^2 \\ & & & & \Omega S^{2n-1} \end{array}$$

ϕ (diagonal arrow from $\Omega^2 S^{2n}$ to ΩS^{2n-1})

Mahowald [8] made the following conjecture, which will follow from Lemma 1.1 and an extensive amount of diagram chasing.

Received by the editors June 8, 1993.

1991 Mathematics Subject Classification. Primary 55Q40, 55Q15, 55Q25.

Key words and phrases. Exponents, EHP fibrations and spectral sequence, $W(n)$, coliftings.

Research supported by the NSF.

Theorem 1.2. *The E_2 -term of the EHP spectral sequence is a $\mathbb{Z}/2$ module. The 4th power map of $\Omega^2 W(n)$ is nullhomotopic, and $\pi_*(W(n))$ has exponent 4.*

Let $d_1^+ : \Omega^3 S^{4n+1} \xrightarrow{\Omega P} \Omega S^{2n} \xrightarrow{H} \Omega S^{4n-1}$ and $d_1^- : \Omega^3 S^{4n+3} \xrightarrow{\Omega P} \Omega S^{2n+1} \xrightarrow{H} \Omega S^{4n+1}$ denote the composites which realize the first EHP differential. Selick [11] improved the James exponent to $2^{2n-[n/2]}$. Cohen [1, §6] reformulated this as a compression of the H -space squaring map on $\Omega^4 S^{4n+1}$ through $\Omega^2 S^{4n-1}$. Theorem 1.2 is implied by the following compression result, which extends their work.

Theorem 1.3. *There exist maps $\mathcal{F}^+ : \Omega^2 S^{4n-1} \rightarrow \Omega^4 S^{4n+1}$ and $\mathcal{F}^- : \Omega^2 S^{4n-3} \rightarrow \Omega^2 W(n)$ making the following diagrams homotopy commutative.*

$$\begin{array}{ccc}
 \Omega^4 S^{4n+1} & \xrightarrow{\Omega d_1^+} & \Omega^2 S^{4n-1} \\
 & \searrow 2 & \downarrow \mathcal{F}^+ \\
 & & \Omega^4 S^{4n+1}
 \end{array}
 \qquad
 \begin{array}{ccc}
 \Omega^4 S^{4n-1} & \xrightarrow{\Omega d_1^-} & \Omega^2 S^{4n-3} \\
 & \searrow 2 & \downarrow \mathcal{F}^- \\
 \Omega^4 S^{4n-1} & \xrightarrow{\Omega^2 \partial} & \Omega^2 W(n)
 \end{array}$$

2. PROOFS

For any space X , we will denote by $2 : \Omega X \rightarrow \Omega X$ the H -space squaring map. We will often use the following fact. For any map $f : \Omega X \rightarrow \Omega Y$, the composites $\Omega^2 X \xrightarrow{\Omega f} \Omega^2 Y \xrightarrow{2} \Omega^2 Y$ and $\Omega^2 X \xrightarrow{2} \Omega^2 X \xrightarrow{\Omega f} \Omega^2 Y$ are homotopic. We will use the following result about coliftings, which we state without proof.

Lemma 2.1. *Let $F \xrightarrow{i} E \xrightarrow{p} B$ be a fibration, and let $f : B \rightarrow X$ be a map such that $f \cdot p : E \rightarrow X$ is nullhomotopic. Then Ωf factors through ∂ , by a colifting $\mathcal{B} : F \rightarrow \Omega X$, which makes the following diagram commute up to homotopy.*

$$\begin{array}{ccc}
 \Omega B & \xrightarrow{\partial} & F \\
 & \searrow \Omega f & \swarrow \mathcal{B} \\
 & & \Omega X
 \end{array}$$

Proof of Theorem 1.3. The EHP fibration $\Omega S^{2n} \xrightarrow{\Omega E} \Omega^2 S^{2n+1} \xrightarrow{\Omega H} \Omega^2 S^{4n+1}$ and Lemmas 1.1(1) and 2.1 give a colifting $\mathcal{B} : \Omega S^{2n} \rightarrow \Omega^3 S^{4n+1}$ making the diagram

$$\begin{array}{ccc}
 \Omega^3 S^{4n+1} & \xrightarrow{\Omega P} & \Omega S^{2n} \\
 & \searrow 2 & \swarrow \mathcal{B} \\
 & & \Omega^3 S^{4n+1}
 \end{array}$$

homotopy commutative. But $\mathcal{B} \cdot E : S^{2n-1} \rightarrow \Omega^3 S^{4n+1}$ is nullhomotopic. By Lemma 2.1 and the EHP fibration $\Omega^2 S^{4n-1} \xrightarrow{P} S^{2n-1} \xrightarrow{E} \Omega S^{2n}$, there exists

a colifting $\mathcal{F}^+ : \Omega^2 S^{4n-1} \rightarrow \Omega^4 S^{4n+1}$ making the diagram

$$\begin{array}{ccc} \Omega^2 S^{2n} & \xrightarrow{\Omega H} & \Omega^2 S^{4n-1} \\ & \searrow \Omega \beta & \swarrow \mathcal{F}^+ \\ & \Omega^4 S^{4n+1} & \end{array}$$

homotopy commutative. This proves the first part of Theorem 1.3.

By Lemma 1.1(2), the composite $\Omega^2 S^{2n} \xrightarrow{2-\Omega E \cdot \phi} \Omega^2 S^{2n} \xrightarrow{\Omega^2 E} \Omega^3 S^{2n+1}$ is nullhomotopic. Hence there exists a map $\psi : \Omega^2 S^{2n} \rightarrow \Omega^4 S^{4n+1}$ making the diagram

$$\begin{array}{ccccc} & \Omega^2 S^{2n} & \xrightarrow{\Omega H} & \Omega^2 S^{4n-1} & \\ \psi \swarrow & & \searrow 2-\Omega E \cdot \phi & & \searrow 2 \\ \Omega^4 S^{4n+1} & \xrightarrow{\Omega^2 P} & \Omega^2 S^{2n} & \xrightarrow{\Omega H} & \Omega^2 S^{4n-1} \end{array}$$

commute up to homotopy, since (cf. [1, Proof of Lemma 4.1]) ΩH is linear. We have an induced map of fibers $\beta : \Omega S^{2n-1} \rightarrow \Omega W(n)$, obtained by pulling back the outer trapezoid to the left, making the following diagram homotopy commutative.

$$\begin{array}{ccccccc} \Omega^3 S^{4n-1} & \xrightarrow{\Omega P} & \Omega S^{2n-1} & \xrightarrow{\Omega E} & \Omega^2 S^{2n} & \xrightarrow{\Omega H} & \Omega^2 S^{4n-1} \\ \downarrow 2 & & \downarrow \beta & & \downarrow \psi & & \downarrow 2 \\ \Omega^3 S^{4n-1} & \xrightarrow{\Omega \partial} & \Omega W(n) & \xrightarrow{\Omega j} & \Omega^4 S^{4n+1} & \xrightarrow{\Omega d_1^+} & \Omega^2 S^{4n-1} \end{array}$$

But $\beta \cdot E : S^{2n-2} \rightarrow \Omega W(n)$ is nullhomotopic. The EHP fibration $\Omega^2 S^{4n-3} \xrightarrow{P} S^{2n-2} \xrightarrow{E} \Omega S^{2n-1}$ and Lemma 2.1 then yield the colifting $\mathcal{F}^- : \Omega^2 S^{4n-3} \rightarrow \Omega^2 W(n)$ making the following diagram homotopy commutative.

$$\begin{array}{ccc} \Omega^2 S^{2n-1} & \xrightarrow{\Omega H} & \Omega^2 S^{4n-3} \\ & \searrow \Omega \beta & \swarrow \mathcal{F}^- \\ & \Omega^2 W(n) & \square \end{array}$$

Proof of Theorem 1.2. By Theorem 1.3, $\text{Ker}(d_1^+)_* \subset \pi_*(S^{4n+1})$ is a $\mathbb{Z}/2$ module. Thus each $E_2^{*,2n+1}$ is a $\mathbb{Z}/2$ module. By Theorem 1.3, any cycle $\alpha \in \text{Ker}(d_1^-)_* \subset \pi_*(S^{4n-1})$ satisfies $2\alpha \in \text{Im}(d_1^+)_*$. Hence each $E_2^{*,2n}$ is a $\mathbb{Z}/2$ module.

We have the fibration sequence $\Omega^2 S^{4n-1} \xrightarrow{\partial} W(n) \xrightarrow{j} \Omega^3 S^{4n+1} \xrightarrow{d_1^+} \Omega S^{4n-1}$. By Theorem 1.3, the composite $\Omega W(n) \xrightarrow{\Omega j} \Omega^4 S^{4n+1} \xrightarrow{2} \Omega^4 S^{4n+1}$ is nullhomotopic. As indicated by the following homotopy commutative diagram, there exists a lifting $f : \Omega W(n) \rightarrow \Omega^3 S^{4n-1}$ of the H space squaring map of $\Omega W(n)$

through $\Omega\partial$.

$$\begin{array}{ccccc}
 \Omega^3 S^{4n-1} & \xrightarrow{\Omega\partial} & \Omega W(n) & \xrightarrow{\Omega j} & \Omega^4 S^{4n+1} \\
 \swarrow f & & \uparrow 2 & \nearrow * & \uparrow 2 \\
 & & \Omega W(n) & \xrightarrow{\Omega j} & \Omega^4 S^{4n+1} \\
 & & & & \searrow \mathcal{F}^+ \\
 & & & & \Omega^2 S^{4n-1} \\
 & & & & \uparrow \Omega d_1^+
 \end{array}$$

We have the following homotopy commutative diagrams.

$$\begin{array}{ccc}
 \Omega^2 S^{4n-1} & \xrightarrow{P} & S^{2n-1} \\
 \searrow \partial & & \nearrow i \\
 & & W(n)
 \end{array}$$

$$\begin{array}{ccccc}
 & & \Omega^3 S^{4n-1} & \xrightarrow{d_1^-} & \Omega S^{4n-3} \\
 & \nearrow f & \downarrow \Omega\partial & \searrow \Omega P & \nearrow H \\
 \Omega W(n) & \xrightarrow{2} & \Omega W(n) & \xrightarrow{\Omega i} & \Omega S^{2n-1}
 \end{array}$$

By looping the above parallelogram and applying Lemma 1.1(1), we see that the composite $\Omega^2 W(n) \xrightarrow{\Omega f} \Omega^4 S^{4n-1} \xrightarrow{\Omega d_1^-} \Omega^2 S^{4n-3}$ is nullhomotopic. The composite $\Omega^2 W(n) \xrightarrow{\Omega f} \Omega^4 S^{4n-1} \xrightarrow{2} \Omega^4 S^{4n-1} \xrightarrow{\Omega^2 \partial} \Omega^2 W(n)$ is nullhomotopic, by Theorem 1.3. Hence $4: \Omega^2 W(n) \rightarrow \Omega^2 W(n)$, the 4th power map, is nullhomotopic. \square

3. REMARKS

James [6] also showed that $2E(x) = 0$ for all $x \in \text{Ker}(E^2) \subset \pi_*(S^q)$. When $q = 2n - 1$, this gives evidence for Theorem 1.2, as it is implied by $4\pi_*(W(n)) = 0$. We used the case $q = 2n$ of James's result in an earlier version of our paper.

Richter [10] strengthened Theorem 1.3, showing that $2 \simeq -\Omega E^2 \cdot d_1^+$ on $\Omega^3 S^{4n+1}$ and $2 - \Omega^3(2i) \simeq -\Omega E^2 \cdot d_1^-$ on $\Omega^3 S^{4n-1}$, solving a conjecture of Gray [3] and Mahowald, which Harper [5] proved at odd primes. At an odd prime p , Cohen, Moore, and Neisendorfer [2] showed that the p^{th} power map on $\Omega W(n)$ is nullhomotopic. Gray [4] showed that $W(n)$ deloops, essentially by delooping the map d_1^+ . It was already known that $\pi_*(W(2))$ had exponent 4, by Cohen's [1, Theorem 19.1] splitting $\Omega^2 S^5\{2\} \simeq W(2) \times \Omega^2 S^3\{3\}$.

Mahowald [8] further conjectured that $(d_1^-)_* = 0: \pi_{*+2}(S^{4n-1}) \rightarrow \pi_*(S^{4n-3})$. Note that James shows that $\text{Ker}(E) \subset \pi_*(S^{2n+1})$ is a $\mathbb{Z}/4$ module.

The conjecture implies that $\text{Ker}(E)$ is a $\mathbb{Z}/2$ module. By [10], the conjecture also implies

Conjecture C2. For any element $\alpha \in \pi_*(S^{4n-1})$, $(2i) \cdot \alpha = 2\alpha \in \pi_*(S^{4n-1})$.

One might wonder whether $2 \simeq \Omega^k(2i)$ on $\Omega^k S^{4n-1}$ for some k . Note [1, §§11 and 12] that away from Arf invariant one or Hopf invariant one dimensions, k must be at least 3.

ACKNOWLEDGMENT

Thanks to Paul Burchard for the commutative diagram package, which uses $\mathcal{L}^{\mathcal{M}\mathcal{S}}$ -TEX fonts.

REFERENCES

1. F. Cohen, *A course in some aspects of classical homotopy theory*, Algebraic Topology, Proceedings Seattle 1985 (H. Miller and D. Ravenel, eds.), Lecture Notes in Math., vol. 1286, Springer, New York, 1987, pp. 1–92.
2. F. Cohen, J. C. Moore, and J. Neisendorfer, *Exponents in homotopy groups*, Ann. of Math. (2) **110** (1979), 549–565.
3. B. Gray, *Unstable families related to the image of J* , Math. Proc. Cambridge Philos. Soc. **96** (1984), 95–113.
4. ———, *On the iterated suspension*, Topology **27** (1988), 301–310.
5. J. Harper, *A proof of Gray's conjecture*, Proc. Internat. Conf. Algebraic Topology (M. Mahowald and S. Priddy, eds.), Contemp. Math., vol. 96, Amer. Math. Soc., Providence, RI, 1989, pp. 189–195.
6. I. M. James, *On the suspension sequence*, Ann. of Math. (2) **65** (1957), 74–107.
7. M. Mahowald, *The image of J in the EHP sequence*, Ann. of Math. (2) **116** (1982), 65–112.
8. M. Mahowald and D. Ravenel, *The root invariant in stable homotopy*, Topology **32** (1993), 865–898.
9. D. Ravenel, *Complex cobordism and stable homotopy groups of spheres*, Pure Appl. Math., vol. 121, Academic Press, Boston, MA, 1986.
10. W. Richter, *The H -space squaring map on $\Omega^3 S^{4n+1}$ factors through the double suspension*, Proc. Amer. Math. Soc. (submitted).
11. P. Selick, *2-primary exponents for the homotopy groups of spheres*, Topology **23** (1984), 97–99.

(M. G. Barratt, M. Mahowald, and W. Richter) DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON, ILLINOIS 60208-2730

Current address, W. Richter: Department of Mathematics, Purdue University, West Lafayette, Indiana 47907

E-mail address, M. Mahowald: mark@math.nwu.edu

E-mail address, W. Richter: richter@math.purdue.edu

(F. Cohen) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NEW YORK 14627-0001

(B. Gray) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT CHICAGO CIRCLE, CHICAGO, ILLINOIS 60680-4348

E-mail address: brayton@math.nwu.edu