

1112-35-667

Stan Alama* (alama@mcmaster.ca), Department of Mathematics & Statistics, McMaster University, Hamilton, ON L8S 4K1, Canada, **Lia Bronsard** (bronsard@mcmaster.ca), Department of Mathematics & Statistics, McMaster University, Hamilton, ON L8S 4K1, Canada, and **Xavier Lamy** (xavierlamy@gmail.com), Institut Camille Jordan, Université Claude Bernard Lyon 1, 69622 Villeurbanne cedex, France. *Minimizers of the Landau-de Gennes energy around a spherical colloid particle.*

We consider energy minimizing configurations of a nematic liquid crystal around a spherical colloid particle, in the context of the Landau-de Gennes model. The nematic is assumed to occupy the exterior of a ball B_{r_0} , and satisfy homeotropic weak anchoring at the surface of the colloid and approach a uniform uniaxial state as $|x| \rightarrow \infty$. We study the minimizers in two different limiting regimes: for balls which are small $r_0 \ll L^{\frac{1}{2}}$ compared to the characteristic length scale $L^{\frac{1}{2}}$, and for large balls, $r_0 \gg L^{\frac{1}{2}}$. The relationship between the radius and the anchoring strength W is also relevant. For small balls we obtain a limiting quadrupolar configuration, with a “Saturn ring” defect for relatively strong anchoring, corresponding to an exchange of eigenvalues of the Q -tensor. In the limit of very large balls we obtain an axisymmetric minimizer of the Oseen–Frank energy, and a dipole configuration with exactly one point defect is obtained. (Received August 11, 2015)