

1105-11-349

Patrick J Dynes (pdynes@clemons.edu), 110 Cherry Hill Avenue, Goose Creek, SC 29445,
Brian McDonald* (bmcdon11@u.rochester.edu), 12947 Wembly Court, Carmel, IN 46033, and
**Kimsey Tor, Andrew Best, Xixi Edelsbrunner, Madeleine Weinstein and Steven J
Miller.** *Benfordness of Zeckendorf Decompositions.*

We report on connections we have established between Zeckendorf decompositions and Benford's law. Zeckendorf showed that every positive integer can be decomposed uniquely into a sum of non-consecutive Fibonacci numbers; this result has been extended to decompositions arising from many other recurrence relations. Additionally, the Fibonacci numbers are known to satisfy Benford's law of digit bias, which means that the density of elements with first digit d is $\log_{10}(1 + \frac{1}{d})$. According to this law, the smaller the digit, the more likely it is to occur as a leading digit.

We prove that for a randomly selected integer between 1 and the n th Fibonacci number, as $n \rightarrow \infty$ the leading digits of the Fibonacci summands in its Zeckendorf decomposition are arbitrarily close to Benford almost surely. The proof proceeds by first analyzing random subsets of Fibonacci numbers for Benfordness. The main ingredient there is showing sets of density are preserved under this process. Using this, prove a correspondence between Zeckendorf decompositions and random subsets of Fibonacci numbers. In those sets the Fibonacci numbers are chosen with a probability $p = 1/\varphi^2$ (where φ is the golden mean) if the previous Fibonacci number wasn't chosen, and 0 otherwise. (Received September 23, 2014)