

1105-34-361

A Abebe* (abraham@temple.edu), **M Chhetri** and **R Shivaji**. *Positive solutions for a class of multiparameter elliptic systems.*

We consider an elliptic system of the form

$$\left. \begin{aligned} -\Delta_p u &= \lambda_1 f_1(u) + \mu_1 \frac{g_1(v)}{v^{\alpha_1}} & \text{in } \Omega; \\ -\Delta_q v &= \lambda_2 \frac{f_2(u)}{u^{\alpha_2}} + \mu_2 g_2(v) & \text{in } \Omega; \\ u = v &= 0 & \text{on } \partial\Omega, \end{aligned} \right\}$$

where $p, q > 1$, $\Delta_m w := |\nabla w|^{m-2} \nabla w$ is the m -Laplacian operator for $m > 1$, $\Omega \subset \mathbb{R}^N$ is a bounded domain with smooth boundary, $\lambda_i, \mu_i > 0$ are parameters and $0 \leq \alpha_i < 1$ are fixed constants for $i = 1, 2$. The nonlinearities $f_i, g_i : [0, \infty) \rightarrow \mathbb{R}$ are continuous functions satisfying certain p, q -sublinear or combined sublinear conditions at infinity. When $g_1(0) < 0$ and $f_2(0) < 0$, we discuss existence of a positive solution for $\lambda_i + \mu_i \gg 1$ for $i = 1, 2$. We also discuss a multiplicity result when $\alpha_1 = 0 = \alpha_2$. Method of sub- and supersolutions are employed to establish these results.

(Received September 24, 2014)