

1132-35-226

Ugur G. Abdulla*, 150 West University Blvd., Melbourne, FL 32901. *The Wiener Test on the Removability of the Logarithmic Singularity for the Elliptic PDEs with Measurable Coefficients and Its Consequences.*

This paper introduces the notion of *log*-regularity (or *log*-irregularity) of the boundary point ζ (possibly $\zeta = \infty$) of the arbitrary open subset Ω of the Greenian deleted neighborhood of ζ in \mathbb{R}^2 concerning second order uniformly elliptic equations with bounded and measurable coefficients, according as whether the *log*-harmonic measure of ζ is null (or positive). A necessary and sufficient condition for the removability of the logarithmic singularity, that is to say for the existence of a unique solution to the Dirichlet problem in Ω in a class $O(\log |\cdot - \zeta|)$ is established in terms of the Wiener test for the *log*-regularity of ζ . From a topological point of view, the Wiener test at ζ presents the minimal thinness criteria of sets near ζ in minimal fine topology. Precisely, the open set Ω is a deleted neighborhood of ζ in minimal fine topology if and only if ζ is *log*-irregular. From the probabilistic point of view, the Wiener test presents asymptotic law for the *log*-Brownian motion near ζ conditioned on the logarithmic kernel with pole at ζ . (Received July 23, 2017)