

1133-35-156

**Ugur G. Abdulla\***, 150 West University Blvd., Melbourne, FL 32901. *The Wiener Test on the Removability of the Logarithmic Singularity for the Elliptic PDEs with Measurable Coefficients and Its Consequences.*

This paper introduces the notion of *log*-regularity (or *log*-irregularity) of the boundary point  $\zeta$  (possibly  $\zeta = \infty$ ) of the arbitrary open subset  $\Omega$  of the Greenian deleted neighborhood of  $\zeta$  in  $\mathbb{R}^2$  concerning second order uniformly elliptic equations with bounded and measurable coefficients, according as whether the *log*-harmonic measure of  $\zeta$  is null (or positive). A necessary and sufficient condition for the removability of the logarithmic singularity, that is to say for the existence of a unique solution to the Dirichlet problem in  $\Omega$  in a class  $O(\log |\cdot - \zeta|)$  is established in terms of the Wiener test for the *log*-regularity of  $\zeta$ . From a topological point of view, the Wiener test at  $\zeta$  presents the minimal thinness criteria of sets near  $\zeta$  in minimal fine topology. Precisely, the open set  $\Omega$  is a deleted neighborhood of  $\zeta$  in minimal fine topology if and only if  $\zeta$  is *log*-irregular. From the probabilistic point of view, the Wiener test presents asymptotic law for the *log*-Brownian motion near  $\zeta$  conditioned on the logarithmic kernel with pole at  $\zeta$ . (Received July 23, 2017)