

Theta Constants, Riemann Surfaces and the Modular Group

Hershel M. Farkas
Irwin Kra

**Graduate Studies
in Mathematics**

Volume 37



American Mathematical Society

Selected Titles in This Series

- 37 **Hershel M. Farkas and Irwin Kra**, Theta constants, Riemann surfaces and the modular group, 2001
- 36 **Martin Schechter**, Principles of functional analysis, second edition, 2001
- 35 **James F. Davis and Paul Kirk**, Lecture notes in algebraic topology, 2001
- 34 **Sigurdur Helgason**, Differential geometry, Lie groups, and symmetric spaces, 2001
- 33 **Dmitri Burago, Yuri Burago, and Sergei Ivanov**, A course in metric geometry, 2001
- 32 **Robert G. Bartle**, A modern theory of integration, 2001
- 31 **Ralf Korn and Elke Korn**, Option pricing and portfolio optimization: Modern methods of financial mathematics, 2001
- 30 **J. C. McConnell and J. C. Robson**, Noncommutative Noetherian rings, 2001
- 29 **Javier Duoandikoetxea**, Fourier analysis, 2001
- 28 **Liviu I. Nicolaescu**, Notes on Seiberg-Witten theory, 2000
- 27 **Thierry Aubin**, A course in differential geometry, 2001
- 26 **Rolf Berndt**, An introduction to symplectic geometry, 2001
- 25 **Thomas Friedrich**, Dirac operators in Riemannian geometry, 2000
- 24 **Helmut Koch**, Number theory: Algebraic numbers and functions, 2000
- 23 **Alberto Candel and Lawrence Conlon**, Foliations I, 2000
- 22 **Günter R. Krause and Thomas H. Lenagan**, Growth of algebras and Gelfand-Kirillov dimension, 2000
- 21 **John B. Conway**, A course in operator theory, 2000
- 20 **Robert E. Gompf and András I. Stipsicz**, 4-manifolds and Kirby calculus, 1999
- 19 **Lawrence C. Evans**, Partial differential equations, 1998
- 18 **Winfried Just and Martin Weese**, Discovering modern set theory. II: Set-theoretic tools for every mathematician, 1997
- 17 **Henryk Iwaniec**, Topics in classical automorphic forms, 1997
- 16 **Richard V. Kadison and John R. Ringrose**, Fundamentals of the theory of operator algebras. Volume II: Advanced theory, 1997
- 15 **Richard V. Kadison and John R. Ringrose**, Fundamentals of the theory of operator algebras. Volume I: Elementary theory, 1997
- 14 **Elliott H. Lieb and Michael Loss**, Analysis, 1997
- 13 **Paul C. Shields**, The ergodic theory of discrete sample paths, 1996
- 12 **N. V. Krylov**, Lectures on elliptic and parabolic equations in Hölder spaces, 1996
- 11 **Jacques Dixmier**, Enveloping algebras, 1996 Printing
- 10 **Barry Simon**, Representations of finite and compact groups, 1996
- 9 **Dino Lorenzini**, An invitation to arithmetic geometry, 1996
- 8 **Winfried Just and Martin Weese**, Discovering modern set theory. I: The basics, 1996
- 7 **Gerald J. Janusz**, Algebraic number fields, second edition, 1996
- 6 **Jens Carsten Jantzen**, Lectures on quantum groups, 1996
- 5 **Rick Miranda**, Algebraic curves and Riemann surfaces, 1995
- 4 **Russell A. Gordon**, The integrals of Lebesgue, Denjoy, Perron, and Henstock, 1994
- 3 **William W. Adams and Philippe Loustau**, An introduction to Gröbner bases, 1994
- 2 **Jack Graver, Brigitte Servatius, and Herman Servatius**, Combinatorial rigidity, 1993
- 1 **Ethan Akin**, The general topology of dynamical systems, 1993

This page intentionally left blank

Theta Constants, Riemann Surfaces and the Modular Group

This page intentionally left blank

Theta Constants, Riemann Surfaces and the Modular Group

An Introduction with Applications to
Uniformization Theorems, Partition Identities
and Combinatorial Number Theory

Hershel M. Farkas
Irwin Kra

Graduate Studies
in Mathematics

Volume 37



American Mathematical Society
Providence, Rhode Island

Editorial Board

Steven G. Krantz
David Saltman (Chair)
David Sattinger
Ronald Stern

2000 *Mathematics Subject Classification*. Primary 30F35, 30F30, 11F20, 11F25, 11F30, 11P81, 11P82, 11P83, 14H42, 14H45, 14H55, 20H10.

ABSTRACT. The book is an introduction to the theory of Theta Functions and Theta Constants with rational characteristics showing their close connections to Riemann Surfaces and the Modular Group. Applications to uniformization theorems, partition identities and combinatorial number theory are developed. Projective embeddings of modular curves and Ramanujan partition congruences are explored. Theta identities and their combinatorial contents are studied. Many examples are provided, and directions for future studies are indicated.

Library of Congress Cataloging-in-Publication Data

Farkas, Hershel M.

Theta constants, Riemann surfaces, and the modular group : an introduction with applications to uniformization theorems, partition identities and combinatorial number theory / Hershel M. Farkas, Irwin Kra.

p. cm. — (Graduate studies in mathematics, ISSN 1065-7339 ; v. 37)

Includes bibliographical references and index.

ISBN 0-8218-1392-7 (acid-free paper)

1. Functions, Theta. 2. Riemann surfaces. 3. Modular groups. I. Kra, Irwin. II. Title. III. Series.

QA345.F37 2001
515'.984—dc21

2001035711

Copying and reprinting. Individual readers of this publication, and nonprofit libraries acting for them, are permitted to make fair use of the material, such as to copy a chapter for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgment of the source is given.

Republication, systematic copying, or multiple reproduction of any material in this publication is permitted only under license from the American Mathematical Society. Requests for such permission should be addressed to the Assistant to the Publisher, American Mathematical Society, P. O. Box 6248, Providence, Rhode Island 02940-6248. Requests can also be made by e-mail to reprint-permission@ams.org.

© 2001 by the American Mathematical Society. All rights reserved.

The American Mathematical Society retains all rights
except those granted to the United States Government.
Printed in the United States of America.

∞ The paper used in this book is acid-free and falls within the guidelines
established to ensure permanence and durability.

Visit the AMS home page at URL: <http://www.ams.org/>

10 9 8 7 6 5 4 3 2 1 06 05 04 03 02 01

Contents

Introduction	xv
Chapter 1. The modular group and elliptic function theory	1
§1. Möbius transformations	2
§2. Riemann surfaces	4
§3. Kleinian groups	5
3.1. Generalities	5
3.2. The situation of interest	8
§4. The elliptic paradise	9
4.1. The family of tori	9
4.2. The algebraic curve associated to a torus	14
4.3. Invariants for tori	23
4.4. Tori with symmetries	28
4.5. Congruent numbers	31
4.6. The plumbing construction	31
4.7. Teichmüller and moduli spaces for tori	33
4.8. Fiber spaces – the Teichmüller curve	33
§5. Hyperbolic version of elliptic function theory	37
5.1. Fuchsian representation	38
5.2. Symmetries of once punctured tori	41
5.3. The modular group	43

5.4.	Geometric interpretations	45
5.5.	The period of a punctured torus	47
5.6.	The function of degree two on the once punctured torus	48
5.7.	The quasi-Fuchsian representation	48
§6.	Subgroups of the modular group	49
6.1.	Basic properties	49
6.2.	Poincaré metric on simply connected domains	50
6.3.	Fundamental domains	52
6.4.	The principal congruence subgroups $\Gamma(k)$	54
6.5.	Adjoining translations: The subgroups $G(k)$	62
6.6.	The Hecke subgroups $\Gamma_o(k)$	63
6.7.	Structure of $\Gamma(k, k)$	65
6.8.	A two parameter family of groups	66
§7.	A geometric test for primality	68
Chapter 2.	Theta functions with characteristics	71
§1.	Theta functions and theta constants	72
1.1.	Definitions and basic properties	72
1.2.	The transformation formula	81
1.3.	More transformation formulae	87
§2.	Characteristics	89
2.1.	Classes of characteristics	89
2.2.	Integral classes of characteristics	93
2.3.	Rational classes of characteristics	93
2.4.	Invariant classes for $\Gamma(k)$	97
2.5.	Punctures on $\mathbb{H}^2/\Gamma(k)$ and the classes $X_o(k)$	98
2.6.	The classes in $X_o(k)$	100
2.7.	Invariant quadruples	105
2.8.	Towers	105
§3.	Punctures and characteristics	106
3.1.	A correspondence	106
3.2.	Branching	106
§4.	More invariant classes	107
4.1.	Invariant classes for $G(k)$	107

4.2.	Characterization of $G(k)$	110
4.3.	The surface $\mathbb{H}^2/G(k)$	112
4.4.	Invariant classes for $\Gamma_o(k)$	113
4.5.	More homomorphisms	116
§5.	Elliptic function theory revisited	117
5.1.	Function theory on a torus	117
5.2.	Projective embeddings of the family of tori	125
§6.	Conformal mappings of rectangles and Picard's theorem	126
6.1.	Reality conditions	127
6.2.	Hyperbolicity and Picard's theorem	128
§7.	Spaces of N -th order θ -functions	129
§8.	The Jacobi triple product identity	138
8.1.	The triple product identity	138
8.2.	The quintuple product identity	143
Chapter 3.	Function theory for the modular group Γ and its subgroups	147
§1.	Automorphic forms and functions	148
1.1.	Two Banach spaces	148
1.2.	Poincaré series	151
1.3.	Relative Poincaré series	152
1.4.	Projections to the surface	154
1.5.	Factors of automorphy	157
1.6.	Multiplicative q -forms	159
1.7.	Residues	161
1.8.	Weierstrass points for subspaces of $\mathbb{A}(\mathbb{H}^2, G, e)$	162
§2.	Automorphic forms constructed from theta constants	165
2.1.	The order of automorphic forms at cusps – Fourier series expansions at $i\infty$	165
2.2.	Automorphic forms for $\Gamma(k)$	172
2.3.	Meromorphic automorphic functions for $\Gamma(k)$	176
2.4.	Evaluation of automorphic functions at cusps	177
2.5.	Automorphic forms and functions for $G(k)$	178
2.6.	Automorphic forms and functions for $\Gamma_o(k)$	178
2.7.	The structure of $\bigoplus_{q=0}^{\infty} \mathbb{A}_q(\mathbb{H}^2, \Gamma)$ and $\bigoplus_{q=0}^{\infty} \mathbb{A}_q^+(\mathbb{H}^2, \Gamma)$	179

§3. Some special cases ($k' = 1$)	183
3.1. $k = 1$	183
3.2. $k = 2$	185
3.3. $k = 3$	190
3.4. $k = 4$	198
3.5. $k = 5$	201
3.6. $k = 6$	204
§4. Primitive invariant automorphic forms	213
4.1. An index 4 subgroup of $\Gamma(k)$ for even k	213
4.2. A Hilbert space of modified theta constants	215
4.3. Projective representation of $\text{Aut } \mathbb{H}^2/\Gamma(k)$	218
4.4. More Hilbert spaces of modified theta constants	223
§5. Orders of automorphic forms at cusps	225
5.1. Calculations via $\Gamma_o(k)$	225
5.2. The general case	227
§6. The field of meromorphic functions on $\overline{\mathbb{H}^2/\Gamma(k)}$	228
6.1. Functions of small degree	228
6.2. $G(k)$ -invariant functions	230
6.3. Generators for the function field $\mathcal{K}(\Gamma(k))$	231
§7. Projective representations	235
§8. Some special cases ($k' = k$)	239
8.1. $k = 3$	239
8.2. $k = 5$	242
8.3. The function field for $\overline{\mathbb{H}^2/\Gamma(7)}$	245
8.4. The projective embedding of $\overline{\mathbb{H}^2/\Gamma(7)}$	246
8.5. $k = 11$	248
8.6. $k = 13$	249
8.7. $k = 9$	250
§9. The function field of $\overline{\mathbb{H}^2/\Gamma(k)}$ over $\overline{\mathbb{H}^2/\Gamma}$	253
§10. Equations that are satisfied by the embedding	253
10.1. The residue theorem	253
10.2. The algorithm	254
10.3. Three term identities	255
10.4. Examples of equations	256

§11. Some special cases (restricted characteristics)	260
11.1. Characteristics with $m' = k$	260
11.2. Characteristics with $m = k$	260
11.3. Ratios	263
Chapter 4. Theta constant identities	265
§1. Dimension considerations	268
1.1. The septuple product identity	268
1.2. Further generalizations	272
§2. Uniformization considerations	274
§3. Elliptic functions as quotients of N -th order theta functions	274
3.1. The Jacobi quartic and derivative formula revisited	274
3.2. More identities - revisited	276
3.3. More identities - new results	277
3.4. More first order applications	279
3.5. Some modular equations	284
§4. Identities which arise from modular forms	291
4.1. Multiplicative meromorphic forms	292
4.2. Cusp forms for Γ	294
4.3. Some special results for the primes 5 and 7	297
§5. Ramanujan's τ -function	297
§6. Identities among infinite products	299
§7. Identities via logarithmic differentiation	301
§8. Averaging automorphic forms	308
§9. The groups $G(k)$	312
Chapter 5. Partition theory: Ramanujan congruences	325
§1. Calculations of $P_N(n)$	331
§2. Some preliminaries	333
2.1. $\Gamma(p, q)$ -invariant functions	333
2.2. Calculation of divisor of $\eta(N \cdot)$	342
2.3. Coset representatives	344
§3. Generalities on constructions of $\Gamma_o(k)$ -invariant functions	345
3.1. The basic problems	345

3.2. Some generalities	346
§4. Constructions of (group) $\Gamma_o(k)$ -invariant functions	347
4.1. The direct construction	347
4.2. Averaging $\Gamma(k^n, k)$ -invariant functions	349
4.3. Bases for $\mathcal{K}(\Gamma_o(k))_0$ and $\mathcal{K}(\Gamma_o(k))_\infty$	361
4.4. Precomposing with A_k	363
§5. Partition identities	369
§6. Production of constant functions	375
6.1. The Frobenius automorphism	375
6.2. Constant functions	378
6.3. Congruences	380
6.4. Functions $F_{k,n,N}$ for negative N	383
6.5. Functions $F_{k,-N}$ of small degree	386
§7. Averaging operators	388
7.1. Automorphisms of $\mathcal{K}(\Gamma_o(k))$	388
7.2. Other linear maps	391
§8. Modular equations	392
8.1. $k = 2$	393
8.2. $k = 3$	395
8.3. $k = 5$	396
8.4. $k = 7$	398
8.5. $k = 13$	399
§9. The ideal of partition identities	399
§10. Examples: Calculations for small k	405
10.1. $k = 2$	405
10.2. $k = 3$	408
10.3. $k = 5$	411
10.4. $k = 7$	413
10.5. $k = 11$	414
10.6. $k = 13$	418
10.7. $k = 4$	419
10.8. $k = 6$	423
§11. The higher level Ramanujan congruences	424
11.1. The level two and three congruences for small primes	424

11.2.	The level n congruences for the prime 2	426
11.3.	The level n congruences for the prime 5	428
11.4.	The level two congruences for the prime 11	430
§12.	Taylor series expansions for infinite products	430
Chapter 6.	Identities related to partition functions	439
§1.	Some more identities related to covering maps	439
1.1.	$k = 2$	440
1.2.	$k = 3$	440
1.3.	$k = 5$	442
1.4.	$k = 7$	443
1.5.	$k = 11$	444
§2.	The j -function and generalizations of the discriminant Δ	447
§3.	Congruences for the Laurent coefficients of the j -function	453
3.1.	Averaging f_k^i	457
3.2.	Completion of the proof of Theorem 3.6 for $k = 5$	458
3.3.	Proof of Theorem 3.6 for $k = 11$, $n = 1$	459
3.4.	A further analysis of the $k = 2$ case	461
Chapter 7.	Combinatorial and number theoretic applications	463
§1.	Generalities on partitions	464
1.1.	Euler series and some old identities	467
1.2.	Partitions and sums of divisors	472
1.3.	Lambert series	473
§2.	Identities among partitions	480
2.1.	A curious property of 8	481
2.2.	A curious property of 3	481
2.3.	A curious property of 7	482
§3.	Partitions, divisors, and sums of triangular numbers	482
3.1.	Sums of 4 squares	486
3.2.	A remarkable formula	489
3.3.	Weighted sums of triangular numbers	493
§4.	Counting points on conic sections	495
§5.	Continued fractions and partitions	499

§6. Return to theta functions	504
Bibliography	511
Bibliographical Notes	513
Index	527

Introduction

The theory of compact Riemann surfaces brings together diverse areas of mathematics. Its building blocks include vast areas of analysis (including Lie theory), geometry/topology and algebra. This was our point of view in our book on Riemann surfaces [6] and it dictated the material to be included in that volume. In particular, we presented a modern approach to the theory of compact Riemann surfaces based on classical methods that prepared the reader to study the modern theories of moduli of surfaces. In this book we head in a different direction and develop another classical connection: to combinatorial number theory. We do not neglect, however, the connections to the problem of uniformizing surfaces represented by very special Fuchsian groups. Problems in number theory can be reformulated as questions about Riemann surfaces, and many of the answers to some of these questions are obtained using function theory. Even though it is an old idea to use function theory (compact Riemann surfaces and automorphic forms) to study analytic and combinatorial number theory and there are many results in these fields, we found it hard to dig out the underlying function theory in the publications of number theorists. No doubt, this is our failing. But since others may also have a deficiency in this area, we decided to organize the material from this point of view. There is new material in this book that has not previously appeared in print, and part of our aim is to present this material to as wide an audience as possible. Our more important aim, however, is to expose the reader to a beautiful chapter in function theory and its applications.

The main actors in our presentation are genus one theta functions and theta constants¹ (including the classical η -function), the modular group $\Gamma = \mathrm{PSL}(2, \mathbb{Z})$, and some of the Riemann surfaces that arise as quotients of the action of finite index subgroups of Γ on \mathbb{H}^2 . We are particularly interested in the principal congruence subgroups $\Gamma(k)$ and the related subgroups $\Gamma_o(k)$ for (usually small) primes k . Some very interesting combinatorial identities follow from the function theory on these surfaces.²

Theta functions and theta constants with integral characteristics are classical objects intimately connected with the principal congruence subgroup of level 2, $\Gamma(2)$. This theory is well understood and has as one of its consequences the theorem of Picard: every entire function which omits two values is constant. As is well known, the basic ingredients in the proof of Picard's theorem are that the holomorphic universal covering of the sphere punctured at three points is the upper half plane and that its fundamental group is $\Gamma(2)$. We use theta constants with even integral characteristics to construct the universal covering map, and in this way obtain, without using the general uniformization theorem, the hyperbolicity of the three times punctured sphere. The universal covering map is constructed here as a quotient of fourth powers of any two of the three theta constants. We noticed that in this construction the three even characteristics correspond in a natural way to the three punctures on $\mathbb{H}^2/\Gamma(2)$, and we began to wonder about natural generalizations. In this book, we present the answer to these inquiries. We uniformize the Riemann surfaces $\mathbb{H}^2/\Gamma(k)$ using theta constants with special rational characteristics, and establish a one-to-one (almost canonical) correspondence between the punctures on $\mathbb{H}^2/\Gamma(k)$ and certain equivalence classes of characteristics. For example, the four punctures on $\mathbb{H}^2/\Gamma(3)$ correspond to the characteristics

$$\begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}, \begin{bmatrix} \frac{1}{3} \\ \frac{1}{3} \end{bmatrix}, \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix}, \begin{bmatrix} \frac{1}{3} \\ \frac{5}{3} \end{bmatrix}.$$

Furthermore, the Riemann surface is uniformized by a quotient of cubes of any two of the four corresponding theta constants. Similar, but obviously more complicated, expressions uniformize the surfaces represented by the higher level congruence groups. Multiple uniformizations of the same

¹Thus the θ -functions we study, $\theta[\chi](\zeta, \tau)$, depend on three variables: a characteristic $\chi \in \mathbb{R}^2$; a variable $\zeta \in \mathbb{C}$; and a parameter $\tau \in \mathbb{H}^2$, the upper half plane. Fixing the variable $\zeta = 0$ yields the family of theta constants, an abuse of notation since these are holomorphic functions on \mathbb{H}^2 ; as functions of the local coordinates $q = e^{2\pi i \tau}$ these are classically known as q -series. We will use the symbol x for the local variable, since tradition in (parts of complex analysis) reserves the letter q for the weight of an automorphic form.

²What is interesting is clearly in the eyes of the beholder. The identities we discuss are obviously interesting to us. The reader must decide whether or not to share our enthusiasm.

Riemann surface lead to theta identities. It is an open problem to determine uniformizations of all four punctured spheres by the methods described above.

The theta constants which appear in our constructions of the uniformizing functions for $\mathbb{H}^2/\Gamma(3)$ are closely related with the formulae used by Euler and Ramanujan in the theory of partitions. Specifically, we note that³

$$\theta \left[\begin{array}{c} \frac{1}{3} \\ 1 \end{array} \right] (0, \tau) = \exp\left(\frac{\pi\iota}{6}\right) x^{\frac{1}{24}} \sum_{n=-\infty}^{\infty} (-1)^n x^{\frac{3n^2+n}{2}} = \exp\left(\frac{\pi\iota}{6}\right) x^{\frac{1}{24}} \prod_{n=1}^{\infty} (1-x^n),$$

where $x = \exp\left(\frac{2\pi\iota\tau}{3}\right)$. Continuing in this direction, we discover that uniformizations of the Riemann surfaces $\mathbb{H}^2/\Gamma(k)$ involve functions which appear in the Jacobi triple product. We give a function theoretic proof of this famous formula and then generalize it to the quintuple and septuple product identities, explaining along the way why the formulae obtained are natural from the point of view of the theory of N -th order theta functions. The highlights of the book are systematic studies of theta constant identities, uniformizations of surfaces represented by subgroups of the modular group, partition identities and Fourier series coefficients of automorphic functions, and identities involving the σ -function and Fourier series coefficients of automorphic forms. More detailed information on the contents of each of the chapters follows.

In Chapter 1 we explain the genesis of the modular group in our theory. This group appears naturally when one classifies compact Riemann surfaces of genus one (elliptic curves) up to conformal equivalence. We discuss the generators of this group, find all the fixed points of elements of this group and describe some of the subgroups we shall need in the sequel. Almost everything we do in this chapter is well known and covered in a standard course on complex variables. We describe the structure of the Riemann surfaces \mathbb{H}^2/G for subgroups G of $\text{PSL}(2, \mathbb{Z})$. In order to show that this well known and elementary material has nontrivial consequences, we use this theory to show that factors of integers of the form $N^2 + 1$ are always the sums of two squares, and we give a geometric criterion for $N^2 + 1$ to be a prime number. The result is that $N^2 + 1$ is prime if and only if the portion in the upper half plane of the straight line joining the origin to the point $N + \iota$ in the complex plane intersects the orbit of ι under $\text{PSL}(2, \mathbb{Z})$ in exactly two points, namely $N + \iota$ and $\frac{N+\iota}{N^2+1}$. We have included some of the function theoretic prerequisites in this chapter. However, most of the prerequisites will be described when needed. In general, we provide full definitions of all concepts. We do not repeat proofs or arguments readily available in

³The reader may at this point conclude that the η -function is a disguised theta constant with a rational characteristic. It will also become obvious that the prime 3 plays a special role in our drama.

other books, but do reproduce, usually in modified form, proofs from many research papers. The bibliography of relevant books (after the last chapter) is followed by a set of bibliographical notes containing an (incomplete) list of research and expository notes on the material covered by this volume.

In Chapter 2 we define the theta functions and theta constants with characteristics and specialize to rational characteristics of the form $\begin{bmatrix} m \\ k \\ m' \\ k \end{bmatrix}$ with m , m' and k integers⁴ of the same parity in order to construct a correspondence between equivalence classes of sets of characteristics and the punctures on the surface $\mathbb{H}^2/\Gamma(k)$. In this chapter we derive a most important property of the theta functions and theta constants, the transformation formula (a significant generalization of known transformation rules) for the action of $\mathrm{PSL}(2, \mathbb{Z})$ on the upper half plane, and we give a function theoretic proof of the Jacobi triple product formula and some generalizations. The transformation formula allows us to use theta functions to construct modular and cusp forms for subgroups of $\mathrm{PSL}(2, \mathbb{Z})$. The function theoretic proof of the Jacobi triple product formula yields new proofs of important identities of Jacobi and Euler that are needed for our presentation of partition theory in Chapter 5. We construct theta constant identities which turn out to agree with discoveries of Ramanujan. Our derivations of these identities are on the one hand quite natural, and on the other hand lead to simpler expressions of the equivalent identities discovered by Ramanujan in the sense that they do not involve irrationalities (extracting roots of single valued functions) until they are artificially introduced. It appears that the theta constants which we use are a lot richer than the ones that Ramanujan had at his disposal.

Chapter 3, in a sense, contains the most important material of the book. In it we construct automorphic forms and functions for the principal congruence subgroups and some related groups. The theory we describe is particularly well suited for the study of $\mathbb{H}^2/\Gamma(k)$, and we obtain holomorphic mappings of these Riemann surfaces into projective spaces of rather low dimensions. Some interesting geometry and topology emerges as we observe connections of the principal congruence subgroups with the Platonic solids. This phenomenon first occurs for $k = 3, 4$ and 5 . In these cases, $\Gamma/\Gamma(k) \cong \mathrm{PSL}(2, \mathbb{Z}_k)$ are the symmetry groups of the regular tetrahedron, octahedron and icosahedron, respectively. This suggests a relation between the images of these curves in the projective space and the regular solids and leads to a generalization of the regular solids based on curves of (some) positive genera. While our development is most suited for the groups $\Gamma(k)$, for many of the most important applications we need to construct automorphic

⁴Assume, unless otherwise stated, for these introductory remarks that k is a (positive) prime.

forms for $\Gamma_o(k)$. Part of the extra difficulties involves the presence of torsion in these groups. We need more detailed analysis, in this and subsequent chapters, to handle these richer groups.

Chapter 4 is a systematic study of theta identities. Theta constant identities are interesting for several reasons. One reason is their inherent elegance and symmetry. There is something tantalizingly beautiful about the identities of Jacobi, for example,

$$\begin{aligned} \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, \tau) \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (z, \tau) \\ = \theta^2 \begin{bmatrix} 0 \\ 1 \end{bmatrix} (0, \tau) \theta^2 \begin{bmatrix} 0 \\ 1 \end{bmatrix} (z, \tau) + \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, \tau) \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (z, \tau) \end{aligned}$$

or its restriction to $z = 0$ (known as the Jacobi quartic identity)

$$\theta^4 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, \tau) = \theta^4 \begin{bmatrix} 0 \\ 1 \end{bmatrix} (0, \tau) + \theta^4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, \tau).$$

Aside from the inherent beauty of the form there is a combinatorial content to the identity. It relates the number of representations of an integer as a sum of four squares to its representation as a sum of four triangular numbers. This is of course just the beginning of a chapter. As one delves deeper into the theory, one finds more and more beautiful identities with more and more combinatorial content.

We present four distinct ways of constructing such identities. In two of these methods, we use the classical technique of constructing finite dimensional linear spaces of theta functions or modular forms or functions on certain Riemann surfaces and use linear algebra and the simple notions of independence and dependence. We present another newer technique which makes use of the fact that we can use theta functions to construct elliptic functions and the fact that the sum of the residues of an elliptic function in a period parallelogram vanishes. This technique is very powerful and succeeds in giving a rather large set of identities. The main idea here is to construct the correct elliptic function, which turns out to be more an art than a science, that leads to an interesting identity. The fourth method for constructing identities uses uniformizations of Riemann surfaces. In a very nontrivial sense the next two chapters are also studies of theta identities, this time of a very special sort with a very special purpose.

In Chapter 5 we turn to the congruences discovered by Ramanujan for the partition function and show how they follow in a rather simple way from function theory on the appropriate Riemann surfaces. The main ingredient is the construction of the same function in more than one way. Some of the constructions involve averaging operators. It turns out that the averaging processes produce in some cases constant functions. We study the

many implications of these constructions, especially of the appearance of constant functions. This material may not be new to the literature. We present it in a unified way based on function theoretic foundations that most of the time remove and in general isolate the mysteries in many of the research monographs on the subject. This chapter is based almost entirely on the properties of the classical η -function, a very special case of the theory described in the previous chapters. We need to know rather detailed information on its multiplier system. The needed number theoretic arguments are found in [16], for example. We would like to avoid dependence on these. We have only partially succeeded in doing so and have hence not included in this volume most of the results of this effort.

In Chapter 6 we begin by reviewing some concepts from covering space theory and show how these ideas lead to beautiful identities among theta constants and their interpretation as identities among infinite products. Here the main tools are the uniformizations of the Riemann surfaces in question. We then continue by showing how many of the ideas used in the congruences related with the partition function and its generalizations carry over to other modular forms. We treat in particular the j -function and the congruences satisfied by the coefficients of its Laurent series expansion.

In Chapter 7 we show how statements about partitions are related to other combinatorial quantities such as representations of positive integers as sums of squares or of triangular numbers, and most importantly to the divisors of an integer. In particular, we describe relations to the question of primality of integers depending on statements about partitions. This suggests that while primality is usually thought of as a subject in multiplicative number theory, it can also be viewed as a part of additive number theory.

We give some examples to show what type of results can be expected in this chapter in the expectation that these applications are the main interests of some readers of this text. We emphasize that these results were not the reason for writing this book. The list of examples is by no means exhaustive. Let $\sigma(n)$ denote the classical σ -function; that is, $\sigma(n)$ is the sum of the divisors of the positive integer n . We show that

$$\sum_{j=-\infty}^{\infty} \sigma\left(n - \frac{3j^2 + j}{2}\right) = 0$$

whenever n is not of the form $\frac{3m^2+m}{2}$ with $m \in \mathbb{Z}$. A companion related result is

$$\sum_{j=0}^{\infty} \sigma\left(n - \frac{j^2 + j}{2}\right) = 4 \sum_{j=0}^{\infty} \sigma\left(\frac{n}{2} + \frac{j^2 + j}{4}\right)$$

whenever n is not of the form $\frac{m^2+m}{2}$ with $m \in \mathbb{Z}$. We obtain a variant of Jacobi's result on the number of ways a positive integer can be written

as a sum of four squares by replacing squares with triangular numbers, obtaining, in our view, a cleaner result. A last example of the type of result we will discuss in this chapter is the following. Let S be a set consisting of the positive integers with an additional copy of those positive integers congruent to zero modulo 7. Decompose S into its even and odd parts, E and O respectively. Denote by $P_E(k)$ and $P_O(k)$ the number of partitions of k with parts taken from the sets E and O respectively. We prove that for all nonnegative integers k ,

$$P_E(2k) = P_O(2k + 1).$$

The prerequisites for this book are a thorough understanding of material traditionally covered in first year graduate courses – especially the contents of the complex analysis course. We review, however, the most salient points about elliptic function theory portions of this course. Although a knowledge of Riemann surfaces and Fuchsian groups is helpful, it is not needed by the reader who is willing to accept the summaries of the required material (with references to the literature). Although we do not, in general, reproduce material available in other textbooks, we make an exception for material on theta functions and theta constants despite the availability of excellent sources (for example, [23]). We do so, not only for the convenience of the reader, but also to emphasize our point of view. We have also ignored, to a great extent, the combinatorial and special functions connection. These are discussed in [2], [7] and [10], for example.

A road map

While we did not intend to write an encyclopedic text, the result has been quite a large book. We take the liberty of offering the readers our suggestions for possible ways of going through this text, which was written with several different types of readers in mind. These range from the beginning graduate mathematics student through the professional mathematician whose interests are either in combinatorial mathematics (partition theory, representation as sums of squares, counting points on conic sections) or function theory (Riemann surfaces, modular forms). Theoretical physicists might be interested in portions of the material we cover.

The reader is expected to have a reasonable knowledge of the theory of functions of a complex variable, through the Riemann mapping theorem, and enough mathematical maturity to follow an argument even though unfamiliar with the proofs of all the tools used. Thus, the book can be used as a text for a topics course in either analysis or analytic number theory, and as

such a reasonable approach would be to go through Chapter 1 sections 1-4.5 and sections 6 and 7. The above material is essentially background material to acquaint the reader with the domains on which we will be doing analysis to obtain the combinatorial results. This reader should then continue with Chapter 2, where the theory of the one dimensional theta function is presented. This chapter should be read in its entirety. In Chapter 3 the reader or instructor pressed for time could read quickly the first section for the definitions and then go on to a careful study of sections 2 through 8.4. Chapter 4 should be read in its entirety. The above could constitute a one semester topics course for beginning graduate students.

The above suggestion leaves out Chapters 5, 6 and 7, which occupy much of this book and are an important part of it, since they deal with the theory of congruences for the Ramanujan partition function, the congruences for the j -function, and the combinatorial interpretations of many of the identities derived in Chapter 4. In a one-year course the material studied could include Chapter 5 through section 10.8 and section 12,⁵ all of Chapters 6 and 7.

The professional mathematician who is interested in Riemann surface theory should study Chapter 1, including sections 4.6 through 5.7, in order to get a picture of where the theory can possibly go. If conversant with the theory of the Riemann theta function, the reader can skip section 1 of Chapter 2 and read the remainder of that chapter. The reader should then proceed to the beginning of Chapter 3. Some of the introductory material of this chapter can be skipped or read quickly to get acquainted with the notation used; the choice of which of the remaining sections to read should be guided by interests; we suggest that this include section 10. Chapter 4 should be read in its entirety and then Chapter 5 and Chapter 6, once again guided by the interests of the reader. Chapter 7 should also be read in its entirety.

The professional mathematician whose interests are in combinatorial mathematics may wish to begin by looking at Chapter 7 and then proceed backwards through the theory. Needless to say, Chapters 2, 3 and 4 will have to be read at some point, and if interested in Ramanujan congruences, Chapter 5 is a must. In any event section 12 of Chapter 5 should be reviewed.

There is lots of flexibility in the way the text can be studied and/or approached. We trust the various readers will find their way through the maze and enjoy the material they stop to study or, as we and others have said, will enjoy this tour of Ramanujan's garden and the flowers they pick there.

⁵In a course where all nonstandard material is included, the instructor might want to spend some time on the multiplicative properties of the η -function. These could be taken from Knopp's book [16].

We have included descriptions of many special cases and summarized the results of many calculations. Most of the nontrivial calculations used the symbolic manipulation programs MATHEMATICA and/or MAPLE. In order to excite the reader about the flowers in the beautiful garden we are cultivating here, we start each chapter with (what we regard as) a handsome example of what will follow. We have included a number of accessible exercises and research level problems. The latter may be quite challenging and are at times only conjectures. The reader should also approach the many special cases we have included as challenges to obtain independent solutions. They are presented in the spirit of exercises, with solutions supplied by the authors.

Numbering systems. The book consists of seven chapters and a set of bibliographical notes that will be maintained and updated on the web. Chapters are subdivided into sections; these into subsections. Definitions, lemmas, propositions, theorems, exercises, problems and remarks are labeled consecutively as a single group within each section. A typical item is Theorem section.number; number starts with 1 for the first item in the section. Thus, for example, in Chapter 2, Definition 2.32 (in section 2) in our numbering scheme is followed by Lemma 4.1 (in section 4). Equations that will be referenced subsequently in the text are labeled by a decimal: chapter.number; number starts with 1 for the first numbered equation in the chapter. Tables and figures are numbered consecutively in the book.

Acknowledgments. We thank

1. the mathematicians who have had the patience to listen to our lectures on the subject of this book: whether our students at the Hebrew University in Jerusalem or the State University of New York in Stony Brook, who at times have provided us with constructive criticism, or colleagues at various institutions who have invited us to speak about this subject;
2. our graduate students who read previous drafts of the book and pointed out errors and ambiguities (needless to say, the remaining errors and ambiguities are the authors' responsibilities);
3. our undergraduate students who helped with the MATHEMATICA calculations and the figures;
4. the editorial and technical staff of the American Mathematical Society, especially Deborah Smith and Janet Pecorelli who guided us through the final preparation of the manuscript; and most importantly,

5. our respective wives, who patiently suffered with us through this long project.

Finally we wish to acknowledge and thank the U.S. - Israel Binational Science Foundation, the U.S. National Science Foundation, The Hebrew University and the State University of New York at Stony Brook for their support over the years that helped produce the research for this book; for providing us with the needed time, facilities and equipment; and for making possible visits by each of us to the institution of the other, not only to work on this volume.

Hershel M. Farkas
Jerusalem, Israel

Irwin Kra
Stony Brook, NY, U.S.A.

This page intentionally left blank

Bibliography

- [1] L.V. Ahlfors. *Complex Analysis (third edition)*. McGraw-Hill, 1979.
- [2] G.E. Andrews. *The Theory of Partitions*, volume 2 of *Encyclopedia of Mathematics and its Applications*. Addison-Wesley, 1976.
- [3] J. Bak and D.J. Newman. *Complex Analysis*. Springer-Verlag, 1982.
- [4] A.F. Beardon. *The Geometry of Discrete Groups*, volume 91 of *Graduate Texts in Mathematics*. Springer-Verlag, 1983.
- [5] J.H. Conway. *The Atlas of Finite Simple Groups*. Oxford Clarendon Press, 1985.
- [6] H.M. Farkas and I. Kra. *Riemann Surfaces (second edition)*, volume 71 of *Graduate Texts in Mathematics*. Springer-Verlag, 1992.
- [7] N.J. Fine. *Basic Hypergeometric Series and Applications*, volume 27 of *Mathematical Surveys and Monographs*. American Mathematical Society, 1989.
- [8] L.R. Ford. *Automorphic Functions*. Chelsea, 1951.
- [9] J.B. Fraleigh. *A First Course in Abstract Algebra*. Addison-Wesley, 1969.
- [10] G. Gasper and M. Rahman. *Basic Hypergeometric Series*, volume 35 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, 1990.
- [11] P. Griffiths and J. Harris. *Principles of Algebraic Geometry*. John Wiley & Sons, 1978.
- [12] R.C. Gunning. *Lectures on Modular Forms*, volume 48 of *Annals of Mathematics Studies*. Princeton University Press, 1962.
- [13] G.H. Hardy and E.M. Wright. *An Introduction to the Theory of Numbers (fifth edition)*. Clarendon Press, 1995.
- [14] Y. Iwayoshi and M. Taniguchi. *An Introduction to Teichmüller Theory*. Springer-Verlag, 1992.
- [15] A.W. Knap. *Elliptic Curves*, volume 40 of *Mathematical Notes*. Princeton University Press, 1992.
- [16] M.I. Knopp. *Modular Functions in Analytic Number Theory*. Chelsea Publishing Company, 1993.
- [17] N. Koblitz. *Introduction to Elliptic Curves and Modular Forms (second edition)*, volume 97 of *Graduate Texts in Mathematics*. Springer-Verlag, 1993.

-
- [18] I. Kra. *Automorphic Forms and Kleinian Groups*. Benjamin, 1972.
 - [19] S. Lang. *Algebra*. Addison-Wesley, 1971.
 - [20] S. Lang. *Algebraic Number Theory (second edition)*. Springer-Verlag, 1994.
 - [21] B. Maskit. *Kleinian Groups*, volume 287 of *Grundlehren der mathematischen Wissenschaften*. Springer-Verlag, 1988.
 - [22] T. Miyake. *Modular Forms*. Springer-Verlag, 1989.
 - [23] D. Mumford. *Tata Lectures on Theta I*. Birkhäuser, 1983.
 - [24] H.E. Rauch and H.M. Farkas. *Theta Functions with Applications to Riemann Surfaces*. Williams and Wilkins, 1974.
 - [25] H.E. Rauch and A. Lebowitz. *Elliptic Functions, Theta Functions and Riemann Surfaces*. Williams and Wilkins, 1973.
 - [26] J.-P. Serre. *A Course in Arithmetic*, volume 7 of *Graduate Texts in Mathematics*. Springer-Verlag, 1973.
 - [27] G. Shimura. *Introduction to the Arithmetic Theory of Automorphic Functions*, volume 11 of *Publications of the Mathematical Society of Japan, Kanô memorial lectures I*. Iwanami Shoten and Princeton University Press, 1971.
 - [28] H. Weber. *Lehrbuch der Algebra (dritter band)*. Chelsea, 1961.

Bibliographical Notes

The bibliographical entries in the main body of the text contained only references to books that filled in gaps in our presentation or described prerequisites. The main purpose of these notes is to give the reader an idea of the many relevant books and scientific manuscripts in the literature on one variable theta functions as related to the topics of the book. It is incomplete and will be maintained and updated periodically on the web at

<http://www.math.sunysb.edu/~irwin/bkbiblio.dvi>

We (our respective e-mail addresses are

farkas@sunset.huji.ac.il

and

irwin@math.sunysb.edu)

would appreciate receiving corrections and additions from readers, especially from experts. The authors of the current volume are not experts in the field of combinatorial number theory and have attempted in their work to open this area to a wider audience. We have emphasized the function theoretic aspects and may have missed many points known to number theorists and combinatorialists. Edited communications to us will be posted on the web and acknowledged appropriately.

1. Books – background material

1.1. Analytic and other prerequisites. The basic facts on complex analysis come from [1]. We have assumed the reader is familiar with the material of a first year graduate course on the subject.

We refer the reader to [1, Ch. 7] for elementary facts concerning elliptic functions and curves. Many more complete treatments of function theory on tori are available in the literature including, for example, [60].

The necessary preliminaries about Riemann surfaces were taken from [35]. There are of course many other excellent books on this subject; among them we list only three: [100] (an English translation of an earlier classic), [76] and [45].

The prerequisites on automorphic forms and Fuchsian groups may be found in portions of [17], [74], [52] and [65]. The computations of the topological invariants associated to a subgroup of the modular group are based on [96].

In Chapter 3, we discussed automorphic forms. Although not extremely relevant for our work, we refer to two papers on the subject (in part because they deal with averaging processes, a topic central to this book): the first by one of the coauthors of this volume [66]; the second by McMullen [75]. The reader interested in number theory should consult [58].

1.2. Other approaches. There is some overlap between our treatment of theta functions and resulting identities (especially involving the partition function) and that of the number theoretic work [61].

[7] is a thorough treatment of the theory of partitions, particularly the interplay between combinatorial and analytic methods.

The book [48] on hypergeometric functions has many points of contact with Ramanujan identities and partition theory.

The encyclopedic book [9] contains chapters on q -series, partitions and the Rogers-Ramanujan identities.

Each of the books mentioned in this subsection contains an extensive bibliography.

1.3. Recent developments. Although it is not our purpose to record recent progress in Teichmüller theory or other fields encountered in our presentation but not directly related to the main topics under study, we mention that the problem of describing quasi-Fuchsian space for once punctured tori as a subset of \mathbb{C}^2 is treated in [95] in the smooth, not complex analytic, setting.

2. Papers by other authors

2.1. Curves represented by subgroups of Γ . In §8.7 of Chapter 3, we described the curve $\mathbb{H}^2/\Gamma(9)$. A refinement of the description can be found in [64].

2.2. The j -function. Divisibility properties of the Fourier coefficients of the j -function for the primes 5, 7 and 11 (the level 1 and 2 only for the prime 11) are found in [70]. The complete picture for the prime 11 is established in [11]. Multiplicative properties (see §2.4) of these Fourier coefficients are studied in [68].

2.3. The partition functions P_N . For $N = 1$, Ramanujan [86], [87] conjectured partition congruences for the primes 5, 7 and 11 and established the level 1 and 2 results. Watson [99] proved the conjecture for all levels n for the prime 5 and a suitably modified level n conjecture for the prime 7. More work is required for the prime 11 since the surface involved is no longer of genus 0. The level 1 congruence was proven by Winkler [101]; Atkin [11] established the general conjecture (see also the earlier manuscript [69]). The level 1 congruence for the prime 5 has probably received more attention than any other case; among the shortest proofs are [28] and [55]. Partition congruences (as well as properties of the Fourier coefficients of the j -function) are studied in [14].

Among the papers studying the function P_N for arbitrary N are [13], [50] and [56]; the computer connection is explored in [12].

For negative N , one obtains recursion identities rather than only congruences for the partition coefficients $P_N(n)$. This case has been thoroughly investigated by M. Newman, whose papers on this and related topics include [78], [79], [80], [82], [81], [83], [84] and [85].

2.4. The Hecke connection. A series $\sum_{n=0}^{\infty} a_n x^n$ that converges on the unit disc is *multiplicative* if $a_{nm} = a_n a_m$ whenever $(n, m) = 1$. The multiplicativity of Ramanujan's τ -function (η^{24}), a theorem first proved by Mordell [77], was established in Chapter 4. In [29], the authors determine all the multiplicative products of η -functions. Two further papers should be mentioned here which connect the η -function to the representation theory of Lie algebras. One is the 1972 paper of Macdonald [71]; the second is the 1978 paper of V. Kac [59]. See also [72]. Hecke operators are discussed in portions of [60, Chs. VIII, IX and X].

One of the motivations of the authors in writing this volume was to show that η is in fact just one example of a theta function with rational characteristics and that introduction of these more general functions can be extremely useful in the theory of uniformizations of the Riemann surfaces which arise from congruence subgroups of the modular group and in the study of theta constant identities. While this was presumably known to Klein, it is only with the development of the theory of theta characteristics that it becomes an efficient procedure.

2.5. The η -function. Papers dealing principally or exclusively with the eta-function and having at least tangential connection to our work include [51], [29], [73] and [72].

2.6. Theta identities. This rich subject is studied in many papers. Among the papers dealing with short and elementary proofs of the Jacobi triple product formula and the quintuple product identity are [3] and [27]. Proofs of the latter also appear in [98], [94] and [49]. [47] is a follow up of some of the work in our book.

2.7. Combinatorial proofs and interpretations of some identities. Among the many papers on the subject are [4] and [46].

3. Papers by the authors

Our interest in this subject may be traced to an early paper by one of the authors [31]. His interest in the subject reemerged as a consequence of the doctoral studies of one of his students resulting in [32] and [33].

The collaboration that led to this book started with [36] and [37], where the use of theta constants with rational characteristics was applied to the study of uniformization of surfaces represented by subgroups of the modular group. In [34] modified theta constants were introduced as a tool in the study of uniformization of surfaces represented by the prime level principal congruence subgroups; related topics are investigated in part of [43]. Our papers on partitions and the Ramanujan τ -function include [38], [67], [41], [39] and [42]. Our work that relates to Hecke operators includes [44]. The quintuple product identity is reproven in [40].

4. Original sources, conference proceedings

The primary sources for the mathematics of S. Ramanujan, the main subject of much of this book, are [54] and [89]; the Hardy lectures [53] are extremely useful in any attempt to evaluate Ramanujan's genius. So are his notebooks [88], [90] and [18]. The last of these contains extensive scholarly and bibliographic information. Among the many conference proceedings devoted to Ramanujan's legacy is [10]; [57] is among the latest. At least two journals, The Ramanujan Journal and The Hardy-Ramanujan Journal are named in his honor. Among the many books and papers that include the name Ramanujan in their title are [19], [20], [21], [22], [23], [24], [5], [6], [8], [25] and [26].

5. Related questions

Among the many issues related but not treated in the book are the higher level congruences for j and P_N and the Rogers-Ramanujan identities. These identities and their generalizations appear in many books and in the following papers: [91], [92], [98], [15], [16], [97] and [30], among others.

The Rogers-Ramanujan identities quoted in Chapter 7 read

$$\frac{1}{\prod_{n=0}^{\infty} (1 - x^{5n+1})(1 - x^{5n+4})} = 1 + \sum_{n=1}^{\infty} \frac{x^{n^2}}{\prod_{l=1}^n (1 - x^l)}$$

and

$$\frac{1}{\prod_{n=0}^{\infty} (1 - x^{5n+2})(1 - x^{5n+3})} = 1 + \sum_{n=1}^{\infty} \frac{x^{n^2+n}}{\prod_{l=1}^n (1 - x^l)}.$$

These identities are related via the Jacobi triple product formula to the theta constants which appeared in Chapter 3, namely

$$\theta \left[\begin{matrix} \frac{1}{5} \\ 1 \end{matrix} \right] (0, 5\tau) \text{ and } \theta \left[\begin{matrix} \frac{3}{5} \\ 1 \end{matrix} \right] (0, 5\tau).$$

The Jacobi triple product yields

$$\prod_{n=0}^{\infty} (1 - x^{5n+1})(1 - x^{5n+4})(1 - x^{5n+5}) = \sum_{n=-\infty}^{\infty} (-1)^n x^{\frac{5n^2+3n}{2}}$$

and

$$\prod_{n=0}^{\infty} (1 - x^{5n+2})(1 - x^{5n+3})(1 - x^{5n+5}) = \sum_{n=-\infty}^{\infty} (-1)^n x^{\frac{5n^2+n}{2}},$$

while straightforward calculation from the definitions shows that in terms of the local coordinate $x = \exp(2\pi i\tau)$,

$$\theta \left[\begin{matrix} \frac{1}{5} \\ 1 \end{matrix} \right] (0, 5\tau) = \exp\left(\frac{\pi i}{10}\right) x^{\frac{1}{40}} \sum_{n \in \mathbb{Z}} (-1)^n x^{\frac{n(5n+1)}{2}}$$

and

$$\theta \left[\begin{matrix} \frac{3}{5} \\ 1 \end{matrix} \right] (0, 5\tau) = \exp\left(\frac{3\pi i}{10}\right) x^{\frac{9}{40}} \sum_{n \in \mathbb{Z}} (-1)^n x^{\frac{n(5n+3)}{2}}.$$

Since

$$\eta(5\tau) = x^{\frac{5}{24}} \prod_{n=1}^{\infty} (1 - x^{5n}),$$

the left hand side (the infinite products) of the Rogers-Ramanujan identities can be interpreted as multiplicative automorphic forms for $\Gamma(5)$. Part of the fascination with these identities is that the right hand side (the infinite sum) does not seem to have such an interpretation.

The fact that the Rogers-Ramanujan identities are connected to the above theta constants was known to Schur. In [93], this connection is made and Schur derives the identities in two different ways. One way uses the theta constants but he also gives a second proof using combinatorial methods. The second proof uses an identity between the binomial coefficients extended to the q -binomial coefficients and some cosine identities.

There is also a generalization of the Rogers-Ramanujan identities known as the Gollnitz-Gordon identities where the modulus 8 replaces 5. While this topic is also not treated in the text, it seems interesting to point out that some of the techniques used to study these identities seem to be related to material in this text. It is pointed out in a paper by K. Alladi [2] that proofs of these identities can be based on splitting the respective functions (defined as infinite sums) into their even and odd parts and that these parts have interesting product representations. The variable x (in most of the literature q) is related to the parameter τ by $x = \exp(2\pi i\tau)$. Our proof of the quintuple product identity also used a decomposition of a function into even and odd parts in the variable $z = \exp(2\pi i\zeta)$. These ideas arise from certain continued fraction identities which were also considered by us in the last chapter of the book. The reader is encouraged to look at Alladi's paper and the references given there to the work of Andrews, Gordon and others.

Theta functions with characteristics enrich the set of tools available to investigate diverse questions. We believe that the material in the text opens areas for further investigations (in particular for doctoral work) and can be used to derive many identities not considered in the book. We illustrate both points by providing two different proofs of the Köhler-Macdonald (the name comes from [73]) identity quoted in Chapter 5. It reads

$$(1) \quad x \prod_{n=1}^{\infty} (1 - x^{6n})^5 (1 - x^{3n})^{-2} = \sum_{n=1}^{\infty} (-1)^{n-1} \left(\frac{n}{3}\right) n x^{n^2}.$$

The right hand side is easily seen to be the same as

$$x \sum_{m=0}^{\infty} (-1)^m (3m+1) x^{9m^2+6m} + x \sum_{m=0}^{\infty} (-1)^m (3m+2) x^{9m^2+12m+3}.$$

It now makes sense to replace x by $x^{\frac{1}{3}}$ and rewrite the identity as

$$\prod_{n=1}^{\infty} (1 - x^{2n})^5 (1 - x^n)^{-2} = \sum_{m=-\infty}^{\infty} (-1)^m (3m+1) x^{m(3m+2)}.$$

To translate this identity to the language of theta functions, we use the change of variable $x = \exp(2\pi i\tau)$. The left hand side of the Köhler-Macdonald

identity is thus

$$\exp\left(-\frac{2\pi i\tau}{3}\right)\eta(2\tau)^5\eta(\tau)^{-2}.$$

We have shown that

$$\theta' \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} (0, \tau) = \frac{2\pi i}{3} \exp\left\{\pi i \frac{1}{3}\right\} x^{\frac{1}{18}} \sum_{n \in \mathbb{Z}} (3n + 1)(-1)^n x^{\frac{n}{6}(3n+2)}.$$

Thus the right hand side of the Kähler-Macdonald identity is

$$\frac{3}{2\pi i} \exp\left\{-\pi i \frac{1}{3}\right\} \exp\left\{-2\pi i\tau \frac{1}{3}\right\} \theta' \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} (0, 6\tau).$$

We conclude that the function theoretic version of Kähler-Macdonald is

$$(2) \quad 2\pi i\eta(2\tau)^5\eta(\tau)^{-2} = 3 \exp\left\{-\pi i \frac{1}{3}\right\} \theta' \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} (0, 6\tau).$$

Our first proof of the above identity follows from the following product formula established by Y. Godin, which will appear as part of his doctoral thesis at the Hebrew University of Jerusalem. Godin proves

$$\begin{aligned} & \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau) \theta \begin{bmatrix} \delta \\ \delta' \end{bmatrix} (\zeta, 2\tau) \\ &= \sum_{\mu=0}^2 \theta \begin{bmatrix} \frac{2\mu+\epsilon+\delta}{3} \\ 2\epsilon' + \delta' \end{bmatrix} (3\zeta, 6\tau) \theta \begin{bmatrix} \frac{2\mu+\epsilon-2\delta}{3} \\ \epsilon' - \delta' \end{bmatrix} (0, 3\tau). \end{aligned}$$

It thus follows that

$$\begin{aligned} & \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (\zeta, \tau) \theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} (\zeta, \tau) \\ &= \theta \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} (0, 3\tau) \left(\theta \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} (3\zeta, 6\tau) - \theta \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} (-3\zeta, 6\tau) \right). \end{aligned}$$

Differentiation (with respect to ζ) and then evaluation at $\zeta = 0$ yields

$$\theta' \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} (0, 6\tau) = \frac{\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, \tau) \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} (0, 2\tau)}{\theta \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} (0, 3\tau)}.$$

An application of the Jacobi triple product formula completes the proof of (2).

It is appropriate to illustrate the combinatorial content of the Kähler-Macdonald identity. The identity may be rewritten as

$$\sum_{n=-\infty}^{\infty} (-1)^n (3n + 1) x^{3n^2+2n} = \prod_{n=1}^{\infty} (1 - x^{4n})^3 (1 - x^{4n-2})(1 + x^{2n-1})^2.$$

In this form the combinatorial content is as follows: Let S be the set containing 3 copies of the positive integers congruent to 0 modulo 4, 1 copy

of the positive integers congruent to 2 modulo 4, and 2 copies of the odd positive integers. Call such a partition *even* if it has an even number of even parts and call it *odd* otherwise. Denote by $E(N)$ and $O(N)$ the number of even and odd partitions of $N \in \mathbb{Z}^+$ using parts from the set S . The Köhler-Macdonald identity says that for all $N \in \mathbb{Z}^+$,

$$E(N) - O(N) = \begin{cases} 0 & \text{for } N \neq 3m^2 + 2m \text{ with } m \in \mathbb{Z} \\ (-1)^m(3m + 1) & \text{for } N = 3m^2 + 2m \text{ with } m \in \mathbb{Z} \end{cases} .$$

The second proof of the identity, more in the spirit of the book, is based on the connection of θ -constants to automorphic forms. It starts by “rewriting” (2) as

$$(3) \quad (2\pi)^{24} \eta(2\tau)^{120} \eta(\tau)^{-48} = 3^{24} \left(\theta' \left[\begin{matrix} \frac{2}{3} \\ 1 \end{matrix} \right] (0, 6\tau) \right)^{24} .$$

The equivalence class of the characteristic $\left[\begin{matrix} \frac{2}{3} \\ 1 \end{matrix} \right]$ is fixed by the group $\Gamma^o(6)$. Using this observation one shows that

$$f(\tau) = \left(\theta' \left[\begin{matrix} \frac{2}{3} \\ 1 \end{matrix} \right] (0, 6\tau) \right)^{24}$$

is a multiplicative 6-form for $\Gamma_o(6)$ (perhaps with a nontrivial multiplier system); that is,

$$\left(\theta' \left[\begin{matrix} \frac{2}{3} \\ 1 \end{matrix} \right] (0, 6\gamma(\tau)) \right)^{24} (\gamma'(\tau))^6 = c_\gamma \left(\theta' \left[\begin{matrix} \frac{2}{3} \\ 1 \end{matrix} \right] (0, 6\tau) \right)^{24} , \quad \tau \in \Gamma_o(6),$$

where c_γ is a constant of absolute value 1. The four punctures on $\mathbb{H}^2/\Gamma_o(6)$ are $P_\infty, P_{\frac{1}{2}}, P_0$ and $P_{\frac{1}{3}}$. The divisor (f) is computed to be $P_\infty^8 P_0^9 P_{\frac{1}{2}}^{18} P_{\frac{1}{3}}$.

The function $\frac{\eta(2\tau)^{120}}{\eta(\tau)^{48}}$ is a multiplicative 6-form for $\Gamma_o(2)$, hence certainly for $\Gamma_o(6)$. It has the same divisor as the automorphic form f . The formulae for the divisors show that these are ordinary cusp forms. Hence, except for the constants, (3) follows. If we extract a 24-th root and evaluate the one (so far undetermined) constant, we obtain (2).

The above two proofs are based on recognizing that both sides of (1) can be reinterpreted in terms of θ -constants. A proof of Köhler-Macdonald based on theta functions (in particular on [62]) can be found in [63]. It is different in flavor from the second of our proofs because it relies more directly and more exclusively on the properties of the η -function. Our second proof has an unexpected dividend. While it is more or less obvious that $\theta' \left[\begin{matrix} \frac{2}{3} \\ 1 \end{matrix} \right] (0, 6\tau)$ is a multiplicative form for $\Gamma_o(6)$, we have shown that it is a form for the bigger group $\Gamma_o(2)$.

As a special case of a more general formula Godin also proves

$$\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (0, \tau) \theta \begin{bmatrix} \delta \\ \delta' \end{bmatrix} (0, \tau) \\ = \sum_{n=0}^4 \theta \begin{bmatrix} \frac{2\mu+\epsilon+\delta}{5} \\ 4\epsilon' + \delta' \end{bmatrix} (0, 20\tau) \theta \begin{bmatrix} \frac{-2\mu-\epsilon+4\delta}{5} \\ -\epsilon' + \delta' \end{bmatrix} (0, 5\tau).$$

Choosing $\begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} \delta \\ \delta' \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ we obtain after some algebraic manipulation

$$\prod_{n=1}^{\infty} (1 + x^{2n-1})^2 \\ = \frac{1}{\prod_{n=0}^{\infty} (1 - x^{5n+1})(1 - x^{5n+4})(1 - x^{4(5n+1)})(1 - x^{4(5n+4)})} \\ + \frac{x}{\prod_{n=0}^{\infty} (1 - x^{5n+2})(1 - x^{5n+3})(1 - x^{4(5n+2)})(1 - x^{4(5n+3)})},$$

a formula, related to the Rogers-Ramanujan identities, which also has a clear combinatorial interpretation which we now give.

Let S be the set consisting of two copies of the odd positive integers. Denote the number of partitions of N from S by $S(N)$. Let T be the set consisting of the positive integers congruent to ± 1 modulo 5 with an extra copy of those which are also congruent to 0 modulo 4. Denote the number of partitions of N from T by $T(N)$. Finally, let U be the set of the positive integers congruent to ± 2 modulo 5 with an extra copy of those which are also congruent to 0 modulo 4. Denote the number of partitions of N from U by $U(N)$. The above identity says that for all $N \in \mathbb{Z}$, $N \geq 2$,

$$S(N) = T(N) + U(N - 1).$$

REFERENCES

- [1] L.V. Ahlfors. *Complex Analysis (third edition)*. McGraw-Hill, 1979.
- [2] K. Alladi. Some new observations on the Gollnitz-Gordon and Rogers-Ramanujan identities. *Trans. Amer. Math. Soc.*, 347:897–914, 1995.
- [3] G.E. Andrews. A simple proof of Jacobi’s triple product identity. *Proc. Amer. Math. Soc.*, 16:333–334, 1965.
- [4] G.E. Andrews. Enumerative proofs of certain q -identities. *Glasgow Math. J.*, 8:33–40, 1967.
- [5] G.E. Andrews. *On the general Rogers-Ramanujan theorem*, volume Number 152 of *Memoirs of the Amer. Math. Soc.* Amer. Math. Soc., 1974.
- [6] G.E. Andrews. On Rogers-Ramanujan type identities related to the modulus 11. *Proc. London Math. Soc.*, 30:330–346, 1975.
- [7] G.E. Andrews. *The Theory of Partitions*, volume 2 of *Encyclopedia of Mathematics and its Applications*. Addison-Wesley, 1976.
- [8] G.E. Andrews. An introduction to Ramanujan’s "lost" notebook. *Amer. Math. Monthly*, 86:89–108, 1979.

- [9] G.E. Andrews, R. Askey, and R. Roy. *Special Functions*, volume 71 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, 1999.
- [10] G.E. Andrews and R.A. Askey et al. (editors). *Ramanujan Revisited*. Academic Press, 1988.
- [11] A.O.L. Atkin. Proof of a conjecture of Ramanujan. *Glasgow Math. J.*, 8:14–32, 1967.
- [12] A.O.L. Atkin. Congruences and modular forms. In *Computers in Mathematical Research*, pages 8–19. North-Holland, 1968.
- [13] A.O.L. Atkin. Ramanujan congruences for $p_{-k}(n)$. *Can. J. Math.*, 20:67–78, 1968.
- [14] A.O.L. Atkin and J.N. O'Brien. Some properties of $p(n)$ and $c(n)$ modulo powers of 13. *Trans. Amer. Math. Soc.*, 126:442–459, 1967.
- [15] W.N. Bailey. Identities of the Rogers-Ramanujan type. *Proc. London Math. Soc.*, 50:1–10, 1949.
- [16] W.N. Bailey. On the simplification of some identities of the Rogers-Ramanujan type. *Proc. London Math. Soc.*, 1:217–221, 1951.
- [17] A.F. Beardon. *The Geometry of Discrete Groups*, volume 91 of *Graduate Texts in Mathematics*. Springer-Verlag, 1983.
- [18] B.C. Berndt. *Ramanujan's Notebooks Parts I-V*. Springer-Verlag, 1985–1998.
- [19] B.C. Berndt. Ramanujan's modular equations. In *Ramanujan Revisited*, pages 313–333. Academic Press, 1988.
- [20] B.C. Berndt and D.C. Bowman. Ramanujan's short unpublished manuscript on integrals and series related to euler's constant. In *Constructive, experimental, and nonlinear analysis (Limoges, 1999)*, pages 17–27. American Mathematical Society, 2000.
- [21] B.C. Berndt, H.H. Chan, and S.-S. Huang. Incomplete elliptic integrals in Ramanujan's lost notebook. *Contemporary Math.*, 254:79–126, 1998.
- [22] B.C. Berndt and P.T. Joshi. *Chapter 9 of Ramanujan's Second Notebook, Infinite series identities, transformations, and evaluations*, volume 23 of *Contemporary Mathematics*. American Mathematical Society, 1983.
- [23] B.C. Berndt, R.L. Lamphere, and B.M. Wilson. Chapter 12 of Ramanujan's second notebook: Continued fractions. *Rocky Mt. J. Math.*, 15:235–310, 1985.
- [24] B.C. Berndt and R.A. Rankin. *Ramanujan: Letters and Commentary*, volume 9 of *History of Mathematics*. American Mathematical Society, 1995.
- [25] S. Bhargava and C. Adiga. On some continued fraction identities of Srinivasa Ramanujan. *Proc. Amer. Math. Soc.*, 92:13–18, 1984.
- [26] S. Bhargava and C. Adiga. Two generalizations of Ramanujan's continued fraction. In *Number Theory, Lecture Notes in Mathematics 1122*, pages 56–62. Springer, Berlin, 1985.
- [27] L. Carlitz and M.V. Subbarao. A simple proof of the quintuple product identity. *Proc. Amer. Math. Soc.*, 32:42–44, 1972.
- [28] J.L. Drost. A shorter proof of the Ramanujan congruence modulo 5. *Amer. Math. Monthly*, 104:963–964, 1997.
- [29] D. Dummit, H. Kisilevsky, and J. McKay. Multiplicative products of η -functions. *Contemporary Math.*, 45:89–98, 1985.
- [30] L. Ehrenpreis. Function theory for Rogers-Ramanujan-like partition identities. *Contemporary Math.*, 143:259–320, 1993.
- [31] H.M. Farkas. Elliptic functions and modular forms. In *Contributions to Analysis*, pages 133–145. Academic Press, 1974.
- [32] H.M. Farkas and Y. Kopeliovich. New theta constant identities. *Israel J. Math.*, 82:133–140, 1993.
- [33] H.M. Farkas and Y. Kopeliovich. New theta constant identities II. *Proc. Amer. Math. Soc.*, 123:1009–1020, 1995.

- [34] H.M. Farkas, Y. Kopeliovich, and I. Kra. Uniformization of modular curves. *Comm. Anal. Geom.*, 4:207–259 and 681, 1996.
- [35] H.M. Farkas and I. Kra. *Riemann Surfaces (second edition)*, volume 71 of *Graduate Texts in Mathematics*. Springer-Verlag, 1992.
- [36] H.M. Farkas and I. Kra. Automorphic forms for subgroups of the modular group. *Israel J. Math.*, 82:87–131, 1993.
- [37] H.M. Farkas and I. Kra. Automorphic forms for subgroups of the modular group. II: Groups containing congruence subgroups. *J. d'Analyse Math.*, 70:91–156, 1996.
- [38] H.M. Farkas and I. Kra. Theta constant identities with applications to combinatorial number theory. *Contemporary Math.*, 211:227–252, 1997.
- [39] H.M. Farkas and I. Kra. A function theoretic approach to the Ramanujan partition identities with applications to combinatorial number theory. *Contemporary Math.*, 240:131–157, 1999.
- [40] H.M. Farkas and I. Kra. On the quintuple product identity. *Proc. Amer. Math. Soc.*, 127:771–778, 1999.
- [41] H.M. Farkas and I. Kra. Ramanujan partition identities. *Contemporary Math.*, 240:111–130, 1999.
- [42] H.M. Farkas and I. Kra. Partitions and theta constant identities. *Contemporary Math.*, 251:197–203, 2000.
- [43] H.M. Farkas and I. Kra. Special sets of points on compact Riemann surfaces. *Contemporary Math.*, 256:75–94, 2000.
- [44] H.M. Farkas and I. Kra. Three term theta identities. *Contemporary Math.*, 256:95–101, 2000.
- [45] O. Forster. *Lectures on Riemann Surfaces*, volume 81 of *Graduate Texts in Mathematics*. Springer-Verlag, 1981.
- [46] F.G. Garvan. Combinatorial interpretations of Ramanujan's partition identities. In *Ramanujan Revisited*, pages 29–45. Academic Press, 1988.
- [47] F.G. Garvan. A combinatorial proof of the Farkas-Kra theta function identities and their generalizations. *J. Math. Anal. Appl.*, 195:354–375, 1995.
- [48] G. Gasper and M. Rahman. *Basic Hypergeometric Series*, volume 35 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, 1990.
- [49] B. Gordon. Some identities in combinatorial analysis. *Quart. J. Math. Oxford*, 12:285–290, 1961.
- [50] B. Gordon. Ramanujan congruences for $p_{-k}(n) \equiv 11^r$. *Glasgow Math. J.*, 24:107–123, 1983.
- [51] B. Gordon and D. Sinor. Multiplicative properties of η -products. In *Number Theory, Madras*, pages 173–200. Springer Lecture Notes 1395, 1987.
- [52] R.C. Gunning. *Lectures on Modular Forms*, volume 48 of *Annals of Mathematics Studies*. Princeton University Press, 1962.
- [53] G.H. Hardy. *Ramanujan, Twelve Lectures on Subjects Suggested by His Life and Work*. Chelsea Publishing Company, 1959.
- [54] G.H. Hardy, P.V. Seshu Aiyar, and B.M. Wilson (editors). *Collected Papers of Srinivasa Ramanujan*. Chelsea Publishing Company, 1927.
- [55] M.D. Hirschhorn. Another short proof of Ramanujan's mod 5 partition congruence, and more. *Amer. Math. Monthly*, 106:580–583, 1999.
- [56] K. Hughes. Ramanujan congruences for $p_{-k}(n)$ modulo powers of 17. *Can. J. Math.*, 43:506–525, 1991.
- [57] M.E.H. Ismail and D.W. Stanton (editors). *q-Series from a Contemporary Perspective*, volume 254 of *Contemporary Mathematics*. American Mathematical Society, 2000.
- [58] H. Iwaniec. *Topics in Classical Automorphic Forms*, volume 17 of *Graduate Studies in Mathematics*. American Mathematical Society, 1997.

- [59] V.G. Kac. Infinite dimensional algebras, Dedakind's η -function, classical Möbius function and the very strange formula. *Adv. in Math.*, 30:85–136, 1978.
- [60] A.W. Knap. *Elliptic Curves*, volume 40 of *Mathematical Notes*. Princeton University Press, 1992.
- [61] M.I. Knopp. *Modular Functions in Analytic Number Theory*. Chelsea Publishing Company, 1993.
- [62] G. Köhler. Theta series on the theta group. *Abh. Math. Seminar Univ. Hamburg*, 58:15–45, 1988.
- [63] G. Köhler. Some eta-identities arising from theta series. *Math. Scand.*, 66:147–154, 1990.
- [64] Y. Kopeliovich and J.R. Quine. On the curve $X(9)$. *The Ramanujan J.*, 2:371–378, 1998.
- [65] I. Kra. *Automorphic Forms and Kleinian Groups*. Benjamin, 1972.
- [66] I. Kra. On the vanishing of and spanning sets for Poincaré series for cusp forms. *Acta Math.*, 153:47–116, 1984.
- [67] I. Kra. On a Ramanujan partition identity. In *Proceedings of the Ashkelon Conference on Complex Function Theory (1996)*, pages 167–172. Israel Conference Math. Proceedings, volume 11, Bar-Ilan University, 1997.
- [68] D.H. Lehmer. Properties of the coefficients of modular invariant $j(\tau)$. *Amer. J. Math.*, 64:488–502, 1942.
- [69] J. Lehner. Ramanujan identities involving the partition function for the moduli 11^α . *Amer. J. Math.*, 65:495–520, 1943.
- [70] J. Lehner. Divisibility properties of the Fourier coefficients of the modular invariant $j(\tau)$. *Amer. J. Math.*, 71:136–148, 1949.
- [71] I.G. Macdonald. Affine root systems and Dedakind's η -function. *Invent. Math.*, 15:91–143, 1972.
- [72] Y. Martin. Multiplicative η -quotients. *Trans. Amer. Math. Soc.*, 348:4825–4856, 1996.
- [73] Y. Martin and K. Ono. η -quotients and elliptic curves. *Proc. Amer. Math. Soc.*, 125:3169–3176, 1997.
- [74] B. Maskit. *Kleinian Groups*, volume 287 of *Grundlehren der mathematischen Wissenschaften*. Springer-Verlag, 1988.
- [75] C. McMullen. Amenability, Poincaré series and quasiconformal maps. *Invent. Math.*, 97:95–127, 1989.
- [76] R. Miranda. *Algebraic Curves and Riemann Surfaces*, volume 5 of *Graduate Studies in Mathematics*. American Mathematical Society, 1995.
- [77] L.J. Mordell. Note on certain modular relations considered by Messrs. Ramanujan, Darling, and Rogers. *Proc. London Math. Soc.*, 20:408–416, 1922.
- [78] M. Newman. Remarks on some modular identities. *Trans. Amer. Math. Soc.*, 73:313–320, 1952.
- [79] M. Newman. An identity for the coefficients of certain modular forms. *J. London Math. Soc.*, 30:488–493, 1955.
- [80] M. Newman. On the existence of identities for the coefficients of certain modular forms. *J. London Math. Soc.*, 31:350–359, 1956.
- [81] M. Newman. Congruences for the coefficients of modular forms and some new congruences for the partition function. *Can. J. Math.*, 9:549–552, 1957.
- [82] M. Newman. Some theorems about $p_r(n)$. *Can. J. Math.*, 9:68–70, 1957.
- [83] M. Newman. Further identities and congruences for the coefficients of modular forms. *Can. J. Math.*, 10:577–586, 1958.
- [84] M. Newman. Construction and application of a class of modular functions II. *Proc. London Math. Soc.*, 9:373–387, 1959.

-
- [85] M. Newman. Modular functions revisited. In *Analytic Number Theory*, pages 396–421. Springer Lecture Notes 899, 1980.
- [86] S. Ramanujan. Some properties of $p(n)$, the number of partitions of n . *Proc. Camb. Phil. Soc.*, 19:207–210, 1919.
- [87] S. Ramanujan. Congruence properties of partitions. *Math. Z.*, 9:147–153, 1921.
- [88] S. Ramanujan. *Notebooks (2 volumes)*. Tata Institute of Fundamental Research, 1957.
- [89] S. Ramanujan. *Collected Papers*. Chelsea, 1962.
- [90] S. Ramanujan. *The Lost Notebooks and other Unpublished Papers*. Narosa, 1988.
- [91] L.J. Rogers. Second memoir on the expansion of certain infinite products. *Proc. London Math. Soc.*, 25:318–343, 1894.
- [92] L.J. Rogers. On two theorems of combinatory analysis and allied identities. *Proc. London Math. Soc.*, 16:315–336, 1917.
- [93] I. Schur. Ein Beitrag zur additiven Zahlentheorie und zur Theorie der Kettenbrueche. *S.-B. Preuss. Akad. Wiss. Phys.-Math. Kl.*, pages 302–321, 1917.
- [94] D.B. Sears. Two identities of Bailey. *J. London Math. Soc.*, 27:510–511, 1952.
- [95] C. Series. Lectures on pleating coordinates on once punctured tori. *RIMS (Kyoto Univ.) Kokyuroku*, 1104:31–90, 1999.
- [96] G. Shimura. *Introduction to the Arithmetic Theory of Automorphic Functions*, volume 11 of *Publications of the Mathematical Society of Japan, Kanô memorial lectures 1*. Iwanami Shoten and Princeton University Press, 1971.
- [97] L.J. Slater. Further identities of the Rogers-Ramanujan type. *Proc. London Math. Soc.*, 54:147–167, 1952.
- [98] G.N. Watson. A new proof of the Rogers-Ramanujan identities. *J. London Math. Soc.*, 4:4–9, 1929.
- [99] G.N. Watson. Ramanujans Vermutung über Zerfallungszahlen. *J. Reine Angew. Math.*, 179:97–118, 1938.
- [100] H. Weyl. *The Concept of a Riemann Surface*. Addison-Wesley, 1955.
- [101] L. Winquist. An elementary proof of $p(11m+6) \equiv 0 \pmod{11}$. *J. Comb. Th.*, 6:56–59, 1969.

This page intentionally left blank

Index

- A_k , 345
- $D(n)$, 482
- $F_{k,N}$, 350
 - constant functions, 375
 - Laurent series expansion for, 352
- $F_{k,n,N}$, 350
 - Laurent series expansion for, 354
- $G(a, b)$, 40
 - fundamental domain, 40
- $G(k)$
 - invariant classes, 107
 - invariant functions, 230
- G_{2k+2} , 22
- $G_{2k+2}(\tau)$, 22
- $G_{k,n,N}(\tau)$, 363
- $LM(q)$, 180
- $N(k, m)$, 108
- P_N , 325
- $S_4(n)$, 486, 488
 - formula, 305
- $S_m(k)$, 476
- $T_4(n)$
 - formula, 305
- $T_m(k)$, 476
- $T_\nu(n)$, 448
- $U_{k,n}(f(\tau))$, 358
- $V(k)$, 215
 - dimension, 215
 - orthogonal basis, 221
 - quotients of functions, 228
 - Weierstrass points, 238
- $V'(k)$, 223
 - dimension, 215
 - Weierstrass points, 238
- $V_{k,M}(f)(\tau)$, 359
- $W_{k;m,n}(f)$, 391
- $X(G)$, 97
- $X(k)$, 94
- $X_0(k)$, 100
- $X_o(k)$, 175
- $Y(k)$, 94
- $Y_{k,N}(\tau)$, 357
- $Z(k)$, 93
- \mathbb{D} , 49
- Δ , 439
- $\Delta(\tau)$, 183
- $\Delta_\nu(\tau)$, 448
- $\Gamma(2)$, 23
 - fundamental domain for, 23
 - generators for, 23
- $\Gamma(k, k)$, 65
 - structure of, 65
- $\Gamma(p, q)$, 66
- $\Gamma/\Gamma(k)$
 - permutation action, 295
- $\Gamma_o(k)$, 113
- $\alpha(k)$, 328
- $\beta(\Gamma(k, k))$, 349
- $\beta(\Gamma_o(k))$, 349
- $\beta(k)$, 328
- $\chi(G)$, 9
- χ_0 , 94
- η -function, 241
- $\eta(\tau)$, 241, 289
- j -function, 27, 183, 439, 447
 - congruences, 457
 - level n for prime 5, 458
 - level 1 for prime 2, 457, 462
 - level 1 for prime 3, 457
 - level 1 for prime 5, 457

- level 1 for prime 7, 457
- level 1 for prime 11, 460
- Laurent coefficients, 451
- j -invariant, 27
- κ , 81
 - properties of, 87
- κ_J , 366
- $\lambda(\tau)$, 24
 - via theta functions, 124
- $[\cdot]$, 350
- $[\cdot]_*$, 109, 350
- $\oplus_{q=0}^{\infty} \mathbb{A}_q(\mathbb{H}^2, \Gamma)$, 179
 - dimension, 179
 - structure, 179
- $\oplus_{q=0}^{\infty} \mathbb{A}_q^+(\mathbb{H}^2, \Gamma)$, 179
 - dimension, 179
 - structure, 179
- $\pi_k(n)$, 393
- $\sigma(n)$, 301, 472, 483
 - recursion formulas, 490
- $\sigma_e(n)$, 492
- $\sigma_o(n)$, 492
- $d(n)$, 473
- eta-function, 289
- f_k , 347
 - Taylor series expansion for, 346
- $f_k(\tau)$, 346
- $f_{q,p}(\tau)$, 341
- $f_{r,s}^*$, 148
- h_k , 392
- $j(k, m)$, 109
- $m(\chi)$, 96
- $m'(\chi)$, 96
- q -form, 15
 - automorphic, 149
 - bounded, 149
 - integrable, 149
- $r(k, N, n)$, 384
- $\mathbb{A}_q(\Delta)$, 149
- $\mathbb{B}_q(\Delta)$, 149
- \mathbb{H}^2 , 49
- $\mathbb{H}^2/\Gamma(k)$, 98
 - automorphisms, 218
 - punctures on, 98
 - distinguished, 226
- $\mathbb{H}^2/G(k)$, 241, 289
- $\mathbb{H}^2/\Gamma(4)$, 419
 - punctures, 419
- $\overline{\mathbb{H}^2/\Gamma(7)}$
 - Weierstrass points, 245
- $\overline{\mathbb{H}^2/\Gamma(9)}$
 - equations, 257
- $\mathbb{H}^2/\Gamma(11)$, 248
- \mathcal{F}_N , 131, 137, 268
 - even functions in, 137
 - odd functions in, 137
- $\mathcal{G}(k)$, 111
- \mathcal{K} , 157
- $\mathcal{K}(\Gamma(k))$, 231
- $\mathcal{K}(\Gamma_o(k))_0$, 345
- $\mathcal{K}(\Gamma_o(k))_{\infty}$, 345
- $\mathcal{S}_4(n)$, 477
- $\mathcal{T}_4(n)$, 477
- $\tilde{T}_m(k)$, 476
- $\tilde{U}_k(j)$, 457
- $\tilde{U}_{k,n}$, 389
- $\tilde{\mathcal{T}}_4(n)$, 477
- \mathfrak{M}^* , 231
- $R\left(\frac{a}{b}\right)$, 350
- Abel's theorem, 18
- Automorphic forms, 165
 - for $\Gamma(k)$, 172
 - for $\Gamma_o(k)$, 178
 - for $G(k)$, 178
 - order, 165
 - theta constant, 168
 - divisor, 168
- Automorphy
 - canonical factor, 157
 - factor, 157
- Beltrami coefficients, 150
- Branch point, 8
- Branch value, 8
- Ceiling, 350
- Characteristics
 - adherent quadruple, 105
 - equivalence classes, 89
 - fundamental set, 89
 - integral, 93
 - rational, 93
 - parity of, 94
 - primitive k , 94
 - towers, 105
- Classes
 - invariants of group, 97
- Conformal map, 5
 - rectangle onto disc, 126
- Congruence subgroups, 54
 - $\Gamma(k)$, 54
 - adjoining translations, 62
- Congruences, 484
 - for $\sigma(n)$, 484
 - for $\mathcal{T}_4(n)$, 484
- Congruent numbers, 31
- Conic sections, 495
 - counting points, 495
- Constant functions, 375
- Conventions used, 232, 236, 337
- Cubic

- discriminant of, 289
- Cusp, 8, 156
 - distinguished, 226
 - forms, 156
 - width of, 162
- $D(k, N)$, 345
- Discontinuous action, 5
- Discriminant function, 439
- Divisor
 - q -canonical ramification, 155
 - theta constant, 168
- Divisors, 18, 483
 - group of, 18
 - sums of, 483
 - congruences, 484
 - sums of even, 492
 - sums of odd, 492
 - of theta functions, 79
- Elliptic curve, 9
- Elliptic function, 14
 - divisor of, 117
 - residues of, 15
 - theta function
 - N -th order, 274
- Equivalence modulo a group action, 6
- Eta-function, 241, 289
- Euler, 465, 467
 - φ function, 110
 - identity, 465
 - series, 467, 469
 - identity, 467
 - theorem of, 465
- Euler characteristic, 9
- Euler's identity, 72, 465
- Fibonacci sequence, 508
- Finite type, 8
- Fixed point
 - attractive, 4
 - repulsive, 4
- Floor, 109, 350
- Frobenius automorphism, 375
- Fuchsian group, 8
 - first kind, 8
 - second kind, 8
- Fundamental domain
 - for Γ , 12
 - for $\Gamma(2)$, 23
 - for $\Gamma(3)$, 60
 - for $\Gamma(4)$, 60
 - for $\Gamma(k)$, 55
 - for $G(a, b)$, 40
- Generating function, 369
- Hecke, 63, 294, 448
 - groups, 63, 349
 - operator, 294, 448
- Holomorphic map
 - branch number, 16
 - degree of, 16
- Horocycles, 51
- Horodisc, 8
- Hyperbolic
 - surface, 362
- Icosahedron, 242
 - group of symmetries, 242
- Identity, 480
 - among infinite products, 299
 - among partitions, 480
 - continued fraction, 507
 - cubic, 193
 - curious properties
 - of 3, 481
 - of 7, 482
 - of 8, 481
 - Euler, 72, 141, 431, 465
 - from logarithmic differentiation, 301
 - Jacobi, 405, 431
 - Jacobi triple product, 138, 436
 - for the prime 2, 440
 - for the prime 3, 440
 - for the prime 5, 442
 - for the prime 7, 443
 - for the prime 11, 444
 - products for 11, 446
 - quintuple product, 143, 436
 - Ramanujan, 313
 - Rogers-Ramanujan, 505
 - septuple product, 268, 271, 437
 - consequence of, 497
 - via covering maps, 439
- Invariants
 - classes of characteristic, 97
 - quadruples, 105
- Jacobi, 467
 - derivative formula, 122, 275, 292
 - generalization, 293
 - identity, 72, 288, 292, 467
 - quartic, 120
 - quartic, 186, 193, 275, 393, 407
- Jacobi triple product, 138, 436
- Jacobi variety, 17
- Klein, 32
- Klein 4-group, 32
- Kleinian group, 5
 - b -group, 7
 - conjugate, 7
 - definition, 6

- elementary, 6
 - Fuchsian group, 8
 - function group, 7
 - fundamental domain, 7
 - fundamental set, 7
 - limit set, 6
 - ordinary set, 5
 - stabilizer of a point, 6
- Lagrange, 482
- Lambert series, 473
- Legendre symbol, 314, 339
- Lehmer, 377
- Meromorphic function, 5
- Möbius transformation, 2
 - elliptic, 2
 - order, 3
 - hyperbolic, 2
 - loxodromic, 2
 - multiplier, 3
 - parabolic, 2
 - translation length, 3
- Modular equation, 284, 392
 - $k = 2$, 393
 - $k = 3$, 395
 - $k = 5$, 396
 - cubic, 285
 - quintic, 285
 - septic, 285
- Modular forms, 156
- Moduli space, 14
- Mordell's theorem, 297, 448
- Normal form, 3
- Operator
 - averaging, 358
- Partition, 464
 - $\sigma(n)$, 489
 - congruence, 370, 373, 374, 417, 426, 428, 430
 - continued fractions, 499
 - even, 464
 - graphical representation, 464
 - odd, 464
 - parity, 331
 - parts of, 464
- Partition coefficients, 325
 - three term recursions, 388, 447
- Partition congruences
 - level 1 for 2, 370
 - level 1 for 5, 300
 - level 1 for 7, 300
 - for P_4
 - level 1 for 7, 413
 - level 2 for 7, 413
- Partition functions, 300, 325
- Period parallelogram, 79
- Petersson pairing, 217
- Petersson scalar product, 150
- Picard
 - group, 6
 - theorem of, 126
- Plumbing construction, 31
- Poincaré, 50
 - density, 149
 - metric, 50
 - series
 - operator, 151, 344
 - relative, 152
- Precisely invariant set, 5
- Primality, 68
 - geometric test, 68
- Puncture, 226
 - distinguished, 226
- Quintuple product identity, 143, 436
- Ramanujan, 297
 - τ -function, 297
 - multiplicative properties, 298
 - congruence
 - level n , 327
 - level n for 2, 426
 - level n for 5, 428
 - level 1, 382
 - level 1 for 1, 405
 - level 1 for 3, 370, 373
 - level 1 for 5, 329, 370, 374, 411, 413
 - level 1 for 11, 417
 - level 1 for 7, 413, 414
 - level 2 for 2, 405
 - level 2 for 3, 405
 - level 2 for 5, 329, 405, 424, 426
 - level 3 for 5, 426
 - congruence for P_2
 - level n for 5, 430
 - level 2 for 5, 411
 - partition congruences, 297, 370
 - partition function, 300
 - polynomials, 332
- Ramification point, 8
- Riemann ζ -function, 22
- Riemann space, 31
 - tori, 33
- Riemann surface, 4
 - closed, 5
 - canonical homology basis, 17
 - degree of divisor, 18
 - divisor, 18
 - genus, 15
 - Jacobi variety, 17

- period matrix, 17
 - principal divisor, 18
- double, 8
- holomorphic equivalence, 5
- holomorphic map
 - branch number, 16
- hyperelliptic, 34
- open, 5
- orbifold, 14
- torus, 9
 - invariants, 23
- Riemann-Hurwitz, 16

- Septuple product identity, 271, 437
- Signature, 9
- Squares, 486
 - sums of, 476, 482, 486
 - sums of 4, 305
- Subgroups of Γ
 - Hecke groups, 63
 - principal congruence, 54

- Teichmüller curve, 33
- Teichmüller space, 23, 31
 - tori, 33, 41
 - punctured, 41
- Theta constants, 75
 - modified, 215
- Theta functions, 72
 - addition formula, 78
 - characteristics, 72, 89
 - classes, 89
 - equivalence, 89
 - functional equations, 74
 - heat equation, 73
 - N -th order, 129
 - spaces of, 132
 - transformation formula, 81
 - zeros of, 79, 132, 137
- Torus, 9
 - symmetries, 28
- Towers, 105
- Triangular numbers, 476
 - primality, 483
 - sums of, 476, 483
 - sums of 4, 305
 - weighted sums of, 493
- Trigonometric sums
 - cosines, 223, 494
 - sines, 319
 - tangents, 172, 319
- Triple product identity, 138

- Weierstrass
 - points, 162
 - for $\mathbb{H}^2/\Gamma(k, k)$, 341
 - weight, 163
 - Weierstrass \wp -function, 19
 - transformation formula, 22
 - via theta functions, 124
 - Wronskian, 163, 270

ISBN 0-8218-1392-7



9 780821 813928

GSM/37

AMS *on the Web*
www.ams.org