

# Lectures on Elliptic and Parabolic Equations in Sobolev Spaces

**N. V. Krylov**

**Graduate Studies  
in Mathematics**

**Volume 96**



**American Mathematical Society**

# Lectures on Elliptic and Parabolic Equations in Sobolev Spaces

*This page intentionally left blank*

# Lectures on Elliptic and Parabolic Equations in Sobolev Spaces

N. V. Krylov

in Mathematics

Volume 96



American Mathematical Society  
Providence, Rhode Island

## Editorial Board

David Cox (Chair)

Steven G. Krantz

Rafe Mazzeo

Martin Scharlemann

2000 *Mathematics Subject Classification*. Primary 35–01; Secondary 35J15, 35K10.

---

For additional information and updates on this book, visit  
**[www.ams.org/bookpages/gsm-96](http://www.ams.org/bookpages/gsm-96)**

---

### Library of Congress Cataloging-in-Publication Data

Krylov, N. V. (Nikolai Vladimirovich)

Lectures on elliptic and parabolic equations in Sobolev spaces / N. V. Krylov.

p. cm. — (Graduate studies in mathematics, ISSN 1065-7339 ; v. 96)

Includes bibliographical references and index.

ISBN 978-0-8218-4684-1 (alk. paper)

1. Differential equations, Elliptic. 2. Differential equations, Parabolic. 3. Sobolev spaces.

I. Title.

QA377.K7582 2008

515'.3533—dc22

2008016051

---

**Copying and reprinting.** Individual readers of this publication, and nonprofit libraries acting for them, are permitted to make fair use of the material, such as to copy a chapter for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgment of the source is given.

Republication, systematic copying, or multiple reproduction of any material in this publication is permitted only under license from the American Mathematical Society. Requests for such permission should be addressed to the Acquisitions Department, American Mathematical Society, 201 Charles Street, Providence, Rhode Island 02904-2294, USA. Requests can also be made by e-mail to [reprint-permission@ams.org](mailto:reprint-permission@ams.org).

© 2008 by the American Mathematical Society. All rights reserved.

The American Mathematical Society retains all rights  
except those granted to the United States Government.

Printed in the United States of America.

⊗ The paper used in this book is acid-free and falls within the guidelines  
established to ensure permanence and durability.

Visit the AMS home page at <http://www.ams.org/>

10 9 8 7 6 5 4 3 2 1 13 12 11 10 09 08

---

# Contents

Preface	xi
Chapter 1. Second-order elliptic equations in $W_2^2(\mathbb{R}^d)$	1
§1. The simplest equation $\lambda u - \Delta u = f$	2
§2. Integrating the determinants of Hessians (optional)	7
§3. Sobolev spaces $W_p^k(\Omega)$	8
§4. Second-order elliptic differential operators	14
§5. Multiplicative inequalities	17
§6. Solvability of elliptic equations with continuous coefficients	20
§7. Higher regularity of solutions	25
§8. Sobolev mollifiers	32
§9. Singular-integral representation of $u_{xx}$	38
§10. Hints to exercises	42
Chapter 2. Second-order parabolic equations in $W_2^{1,k}(\mathbb{R}^{d+1})$	45
§1. The simplest equation $u_t + a^{ij}(t)D_{ij}u - \lambda u = f$	45
§2. Sobolev spaces $W_p^{r,k+2r}(\Omega)$	51
§3. Parabolic equations with continuous coefficients	56
§4. Local or interior estimates	60
§5. The Cauchy problem	67

§6. Hints to exercises	70
Chapter 3. Some tools from real analysis	73
§1. Partitions and stopping times	73
§2. Maximal and sharp functions	78
2:1. The Fefferman-Stein theorem	78
2:2. Exercises (optional)	83
§3. Comparing maximal and sharp functions in the Euclidean space	87
§4. Hints to exercises	91
Chapter 4. Basic $\mathcal{L}_p$ estimates for parabolic and elliptic equations	93
§1. An approach to elliptic equations	94
§2. Preliminary estimates of $L$ -caloric functions	98
§3. Solvability of model equations	104
§4. Divergence form of the right-hand side for the Laplacian	113
§5. Hints to exercises	115
Chapter 5. Parabolic and elliptic equations in $W_p^{1,k}$ and $W_p^k$	117
§1. Better regularity for equations with coefficients independent of $x$	117
§2. Equations with continuous coefficients. The Cauchy problem	119
Chapter 6. Equations with VMO coefficients	125
§1. Estimating $\mathcal{L}_q$ oscillations of $u_{xx}$	125
§2. Estimating sharp functions of $u_{xx}$	130
§3. A priori estimates for parabolic and elliptic equations with VMO coefficients	134
§4. Solvability of parabolic and elliptic equations with VMO coefficients. The Cauchy problem	139
§5. Hints to exercises	143
Chapter 7. Parabolic equations with VMO coefficients in spaces with mixed norms	145
§1. Estimating sharp functions of $\ u_{xx}(t, \cdot)\ _{\mathcal{L}_q}$	146

---

§2. Existence and uniqueness results	149
§3. Hints to exercises	155
Chapter 8. Second-order elliptic equations in $W_p^2(\Omega)$	157
§1. Spaces of functions vanishing on the boundary	158
§2. Equations in half spaces	160
§3. Domains of class $C^k$ . Equations near the boundary	165
§4. Partitions of unity and the regularizer	171
§5. Solvability of equations in domains for large $\lambda$	174
§6. Hints to exercises	179
Chapter 9. Second-order elliptic equations in $W_p^k(\Omega)$	181
§1. Finite differences. Better regularity of solutions in $\mathbb{R}_+^d$ for model equations	181
§2. Equations in domains	187
§3. The oblique derivative problem in $\mathbb{R}_+^d$	190
§4. Local regularity of solutions	196
§5. Hints to exercises	199
Chapter 10. Sobolev embedding theorems for $W_p^k(\Omega)$	201
§1. Embedding for Campanato and Slobodetskii spaces	203
1:1. Embeddings in $C^\alpha$	203
1:2. Exercises (optional)	207
§2. Embedding $W_p^1(\Omega) \subset C^{1-d/p}(\Omega)$ . Morrey's theorem	209
§3. The Gagliardo-Nirenberg theorem	215
§4. General embedding theorems	216
§5. Compactness of embeddings. Kondrashov's theorem	223
§6. An application of Riesz's theory of compact operators	227
§7. Hints to exercises	229
Chapter 11. Second-order elliptic equations $Lu - \lambda u = f$ with $\lambda$ small	231
§1. Maximum principle for smooth functions	232
1:1. Maximum principle	232

1:2. Exercises (optional)	234
§2. Resolvent operator for $\lambda$ large	236
§3. Solvability of equations in smooth domains	241
§4. The way we proceed further	245
§5. Decay at infinity of solutions of $Lu = f$ in $\mathbb{R}^d$	246
§6. Equations in $\mathbb{R}^d$ with $\lambda$ small	249
§7. Traces of $W_p^k(\Omega)$ functions on $\partial\Omega$	255
§8. The maximum principle in $W_p^2$ . Green's functions	262
§9. Hints to exercises	263
Chapter 12. Fourier transform and elliptic operators	267
§1. The space $\mathcal{S}$	268
§2. The notion of elliptic differential operator	269
§3. Comments on the oblique derivative and other boundary-value problems. Instances of pseudo-differential operators	272
§4. Pseudo-differential operators	274
§5. Green's functions	280
§6. Existence of Green's functions	285
§7. Estimating $G$ and its derivatives	288
7:1. Differentiability of $G$ and estimates of its derivatives	288
7:2. Exercises (optional)	292
§8. Boundedness of the zeroth-order pseudo-differential operators in $\mathcal{L}_p$	293
§9. Operators related to the Laplacian	297
9:1. The operators $(1 - \Delta)^{\gamma/2}$ , Cauchy's operator, the Riesz and Hilbert transforms, the Cauchy-Riemann operator	297
9:2. Exercises (optional)	302
§10. An embedding lemma	304
§11. Hints to exercises	308
Chapter 13. Elliptic operators and the spaces $H_p^\gamma$	311
§1. The space $\mathcal{H}$	311

---

§2. Some properties of the space $\mathcal{H}$	315
§3. The spaces $H_p^\gamma$	317
3:1. Definition, solvability of elliptic equations in $H_p^\gamma$ , the equality $H_p^\gamma = W_p^\gamma$	317
3:2. Exercises (optional)	323
§4. Higher-order elliptic differential equations with continuous coefficients in $H_p^\gamma$	328
§5. Second-order parabolic equations. Semigroups (optional)	333
§6. Second-order divergence type elliptic equations with continuous coefficients	335
§7. Nonzero Dirichlet condition and traces	339
§8. Sobolev embedding theorems for $H_p^\gamma$ spaces	342
§9. Sobolev mollifiers	346
§10. Hints to exercises	350
Bibliography	353
Index	355

*This page intentionally left blank*

---

# Preface

These lectures concentrate on some basic facts and ideas of the modern theory of linear elliptic and parabolic partial differential equations (PDEs) in Sobolev spaces. We hope to show that this theory is based on some general and extremely powerful ideas and some *simple* computations. The main objects of study are the Cauchy problem for parabolic equations and the first boundary-value problem for elliptic equations, with some guidelines concerning other boundary-value problems such as the Neumann or oblique derivative problems or problems involving higher order elliptic operators acting on the boundary. The presentation has been chosen in such a way that after having followed the book the reader should acquire a good understanding of a wide variety of results and techniques.

These lecture notes appeared as the result of a two-quarter or a one-semester graduate course I gave at Moscow State University and the School of Mathematics, University of Minnesota, over a number of years and differ significantly from previous drafts. This book also includes some parts of the initial draft and as a whole is most appropriate for a two-quarter or a one-year course. Naturally, one cannot expect that in such a short course all important issues of the theory of elliptic and parabolic equations can be covered. Actually, even the area of second-order *elliptic* partial differential equations is so wide that one cannot imagine a book, let alone a textbook, of reasonable size covering all bases. Restricting further only to the theory of solvability in the *Sobolev* function spaces and *linear* equations still does not make the task realistic. Because of that we will only be concerned with some basic facts and ideas of the modern theory of linear elliptic and parabolic equations in Sobolev spaces. We refer the interested reader to the books [7], [9], and [15], which are classical texts and reference books in elliptic and

parabolic PDEs, and the literature therein for additional information on the subject.

I have been educated as a probabilist who in his early stages of research came across the necessity of using some PDE results and realized, with some deep disappointment mixed with astonishment, that at that time there were no simple *introductory* books about the modern theory available to a wide audience. The situation is slightly better now, forty-five years later, but yet by now, as has been pointed out above, the theory became so wide that it is impossible to have just one simple introductory book available to a wide audience. Indeed, one does have several introductory books in different areas of the theory including the book I wrote on Hölder space theory (see [12]) and the current textbook on Sobolev space theory.

As with almost any graduate textbook, this one is written for myself and my graduate students who, as I think, should know at least that much of the theory in order to be able to work on problems related to my own interests. That is why the choice of these “basic facts and ideas” and the exposition is by no means exhaustive but rather reflects the author’s taste and, in part, his view on what he should have known to be able to work in some areas of mathematics such as the theory of random diffusion processes. I also hope that the contents of the book will be useful to other graduate students and scientists in mathematics, physics, and engineering interested in the theory of partial differential equations.

### Comments on the structure of the book

We start with the  $\mathcal{L}_2$  theory of elliptic second-order equations in the whole space, first developing it for the Laplacian on the basis of the Fourier transform. Then we go to the  $\mathcal{L}_2$  theory for equations with variable coefficients by using partitions of unity, the method of “freezing the coefficients”, the method of a priori estimates, and the method of continuity. This is done in Chapter 1.

In Chapter 2 we deal with the  $\mathcal{L}_2$  theory of second-order parabolic equations along similar lines. As far as parabolic equations are concerned, in these notes we only concentrate on equations in the whole space and the Cauchy problem.

After that, in Chapter 3, we present some tools from real analysis, helping to pass from  $\mathcal{L}_2$  theory to  $\mathcal{L}_p$  theory with  $p \neq 2$ .

In Chapter 4 we derive basic  $\mathcal{L}_p$  estimates first for parabolic and then for elliptic equations. The estimates for the elliptic case turn out to follow immediately from the estimates for parabolic equations. On the other hand, for elliptic equations such estimates can be derived directly and we outline

how to do this in Section 4.1 and in a few exercises (see Exercises 1.1.5, 1.3.23, and 4.3.2 and the proof of Theorem 4.3.7).

Chapter 5 is devoted to the  $\mathcal{L}_p$  theory of elliptic and parabolic equations with continuous coefficients in the whole space. Chapter 6 deals with the same issues but for equations with VMO coefficients, which is quite a new development (only about 16 years old; compare it with the fundamental papers [1] of 1959–64).

Chapter 7 is the last one where we systematically consider parabolic equations. There the solvability of parabolic equations with VMO coefficients is proved in Sobolev spaces with mixed norms. Again as in Chapter 2, everything is done only for the equations in the whole space or for the Cauchy problem for equations whose coefficients are only measurable in the time variable. We return to this problem only briefly in Section 13.5 for equations with coefficients independent of time.

Starting from Chapter 8, we only concentrate on elliptic equations in

$$\mathbb{R}^d = \{x = (x^1, \dots, x^d) : x^i \in (-\infty, \infty)\}$$

or in domains  $\Omega \subset \mathbb{R}^d$ . It is worth noticing, however, that almost everything proved for the elliptic equations in Chapter 8 is easily shown to have a natural version valid for parabolic equations in  $\mathbb{R} \times \Omega$ . Also the Cauchy problem in  $(0, \infty) \times \Omega$  can be treated very similarly to what is done in Section 5.2. We do not show how to do that. Here we again run into choosing between what basic facts your graduate students should know and what are other very interesting topics in PDEs. Anyway, the interested reader can find further information about parabolic equations in [14], [15], and [18].

The reader can consult the table of contents as to what issues are investigated in the remaining chapters. We will only give a few more comments.

Chapters 12 and 13 can be studied almost independently of all previous with the exception of Chapter 3. There are many reasons to include their contents in a textbook, although it could be that this is the first time this is done. I wanted my graduate students to be exposed to equations in the function spaces of Bessel potentials since the modern theory of stochastic partial differential equations is using them quite extensively.

Some important topics are scattered throughout the book, the most notable are:

- Equations in divergence form; see Sections 4.4, 8.2, and 13.6.
- Boundary-value problems involving boundary differential operators; see Exercises 1.1.11 and 13.3.15 and Sections 9.3 and 12.3.

- Elliptic equations with measurable coefficients in two dimensions including the Neumann problem; see Exercises 1.4.7, 1.4.8, 1.4.9, 1.6.7, 8.2.6, 8.2.11, 11.5.5, and 11.6.5.

These notes are designed as a textbook and contain about 271 exercises, a few of which (almost all of the 63 exercises marked with an \*) are used in the main text. These are the simplest ones. However, many other exercises are quite difficult, despite the fact that their solutions are almost always short. Therefore, the reader should not feel upset if he/she cannot do them even after a good deal of thinking. Perhaps, hints for them should have been provided right after each exercise. We do give the hints to the exercises but only at the end of each chapter just to give the readers an opportunity to test themselves.

Some exercises which are not used in the main text are put in the main text where the reader has enough knowledge to solve them and thus learn more. Some other exercises less directly connected with the text are collected in optional subsections.

The theorems, lemmas, remarks, and such which are part of the main units of the text are numbered serially in a single system that proceeds by section. Exercise 1.7.6 is the sixth numbered unit in the seventh section in the first chapter. In the course of Chapter 1, this exercise is referred to as Exercise 7.6, and in the course of the seventh section of chapter 1 it is referred to as Exercise 6. Similarly and independently of these units formulas are numbered and cross-referenced.

### Basic notation

A complete reference list of notation can be found in the index at the end of the book. We always use the summation convention and allow constants denoted by  $N$ , usually without indices, to vary from one appearance to another even in the same proof. If we write  $N = N(\dots)$ , this means that  $N$  depends only on what is inside the parentheses. Usually, in the parentheses we list the objects that are fixed. In this situation one says that the constant  $N$  is *under control*. By domains we mean general open sets. On some occasions, we allow ourselves to use different symbols for the same objects, for example,

$$u_{x^i} = \frac{\partial u}{\partial x^i} = D_i u, \quad u_x = \text{grad } u = \nabla u, \quad u_{xx} = (u_{x^i x^j}).$$

Any  $d$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_d)$  of integers  $\alpha_k \in \{0, 1, 2, \dots\}$  is called a multi-index. For a multi-index  $\alpha$ ,  $k, j \in \{1, \dots, d\}$ , and  $\xi = (\xi^1, \dots, \xi^d) \in \mathbb{R}^d$  we denote

$$D_{k^j} u = D_j D_k u = u_{x^k x^j}, \quad |\alpha| = \alpha_1 + \dots + \alpha_d,$$

$$D^\alpha = D_1^{\alpha_1} \cdot \dots \cdot D_d^{\alpha_d}, \quad \xi^\alpha = (\xi^1)^{\alpha_1} \cdot \dots \cdot (\xi^d)^{\alpha_d}.$$

We also use the notation  $Du = u_x$  for the gradient of  $u$ ,  $D^2u = u_{xx}$  for the matrix of second-order derivatives of  $u$ , and  $D^n u$  for the set of all  $n$ th order derivatives of  $u$ . These  $D^n u(x)$  for each  $x$  can be considered as elements of a Euclidean space of appropriate dimension. By  $|D^n u(x)|$  we mean any fixed norm of  $D^n u(x)$  in this space.

In the case of parabolic equations we work with

$$\mathbb{R}^{d+1} = \{(t, x) : t \in \mathbb{R}, x \in \mathbb{R}^d\}.$$

For functions  $u(t, x)$  given on subdomains of  $\mathbb{R}^{d+1}$  we use the above notation only for the derivatives in  $x$  and denote

$$\partial_t u = \frac{\partial u}{\partial t} = u_t, \quad \partial_t D_k u = u_{tx^k} = u_{x^k t},$$

and so on.

If  $\Omega$  is a domain in  $\mathbb{R}^d$  and  $p \in [1, \infty]$ , by  $\mathcal{L}_p(\Omega)$  we mean the set of all Lebesgue measurable complex-valued functions  $f$  for which

$$\|f\|_{\mathcal{L}_p(\Omega)} := \left( \int_{\Omega} |f(x)|^p dx \right)^{1/p} < \infty$$

with the standard extension of this formula if  $p = \infty$ . We also define

$$\mathcal{L}_p = \mathcal{L}_p(\mathbb{R}^d).$$

One knows that  $\mathcal{L}_p(\Omega)$  is a Banach space. In the cases when  $f$  and  $g$  are measurable functions defined in the same domain  $D \subset \mathbb{R}^d$ , we write “ $f = g$  in  $D$ ” to mean that  $f$  equals  $g$  *almost everywhere* in  $D$  with respect to Lebesgue measure.

By  $C_0^\infty(\Omega)$  we mean the set of all infinitely differentiable functions on  $\Omega$  with compact support (contained) in  $\Omega$ . By the *support* of a function we mean the closure of the set where the function is different from zero. We call a subset of  $\mathbb{R}^d$  *compact* if it is closed and bounded. Of course, saying “compact support” is the same as saying “bounded support” and we keep “compact” just to remind us that we are talking about *closed* sets. We set

$$C_0^\infty = C_0^\infty(\mathbb{R}^d).$$

For  $k \in \{0, 1, 2, \dots\}$ , by  $C^k(\Omega)$  we denote the set of all  $k$  times continuously differentiable functions on  $\Omega$  with finite norm

$$\|u\|_{C^k(\Omega)} = \sum_{|\alpha| \leq k} \|D^\alpha u\|_{C(\Omega)},$$

where

$$\|u\|_{C(\Omega)} = \sup_{x \in \Omega} |u(x)|.$$

As usual,  $C(\Omega) = C^0(\Omega)$  and we drop  $\Omega$  in  $C^k(\Omega)$  and  $\mathcal{L}_p(\Omega)$  if  $\Omega = \mathbb{R}^d$ . The subset of  $C^k$  consisting of functions on  $\mathbb{R}^d$  with compact support is denoted  $C_0^k$ . In particular,  $C_0$  is the set of continuous functions with compact support.

By  $C^k(\bar{\Omega})$  we denote the subset of  $C^k(\Omega)$  consisting of all functions  $u$  such that  $u$  and  $D^\alpha u$  extend to functions continuous in  $\bar{\Omega}$  (the closure of  $\Omega$ ) whenever  $|\alpha| \leq k$ . For these extensions we keep the same notation  $u$  and  $D^\alpha u$ , respectively.

If  $\Omega$  is an unbounded domain, by  $C_0^k(\bar{\Omega})$  we mean the subset of  $C^k(\bar{\Omega})$  consisting of functions vanishing for  $|x|$  sufficiently large. Mainly, the notation  $C_0^k(\Omega)$  will be used for  $\Omega = \mathbb{R}^d$  and  $\Omega = \mathbb{R}_+^d$ , where

$$\mathbb{R}_+^d = \{(x^1, x') : x^1 > 0, x' = (x^2, \dots, x^d) \in \mathbb{R}^{d-1}\}.$$

Generally, speaking about functions on  $\mathbb{R}^d$ , we mean Lebesgue measurable functions.

### To the instructor

We begin with an  $\mathcal{L}_2$  theory of second-order elliptic equations of the type

$$a^{ij}(x)u_{x^i x^j}(x) + b^i(x)u_{x^i}(x) + c(x)u(x) = f(x), \quad x \in \mathbb{R}^d,$$

where the coefficients are assumed to be bounded and the matrix  $a = (a^{ij})$  symmetric and uniformly positive definite. If the matrix  $a$  is uniformly continuous and  $c$  is sufficiently large negative, we show that the equation is solvable for  $f \in \mathcal{L}_2$ . Much later in Section 11.6 on the basis of  $L_p$  theory, the restriction on  $c$  is replaced with  $c \leq -\delta$ , where  $\delta > 0$  is any constant.

Then the general scheme set out in Chapter 1 is repeated several times in the succeeding text in various settings without going into all the minor details each time.

Chapter 3 contains all the tools from real analysis that we use. Here is a major difference between these notes and one of the previous drafts. At

some stage, I was presenting the  $\mathcal{L}_p$  theory on the basis of the Calderón-Zygmund and Marzinkiewicz theorems interpolating between  $p = 1$  and  $p = 2$ . The present course is based on using the Fefferman-Stein theorem on sharp functions. One could use this theorem along with Stampacchia's interpolation theorem (and Marzinkiewicz's interpolation theorem) to interpolate between  $p = 2$  and  $p = \infty$ , which is done in some texts and in one of the drafts of these lectures. However, there is a shorter way to achieve the goal. The point is that it is possible to obtain *pointwise* estimates of the sharp function  $u_{xx}^\sharp$  of the second-order derivatives  $u_{xx}$  of the unknown solution  $u$  through the maximal function of the right-hand side of the equation. This fact has actually been very well known for quite a long time and allows one to get the  $\mathcal{L}_p$  estimates of  $u_{xx}$  for  $p > 2$  just by referring to the Fefferman-Stein and Hardy-Littlewood maximal function theorems.

A disadvantage of this approach is that the students are not exposed to such very powerful and beautiful tools as the Calderón-Zygmund, Stampacchia, and Marzinkiewicz theorems. On the other hand, there are two advantages. First, the Fefferman-Stein theorem is much more elementary than the Calderón-Zygmund theorem (see Chapter 3). Second, the approach based on the pointwise estimates allows us to prove existence theorems for equations with VMO coefficients with almost the same effort as in the case of equations with continuous coefficients.

Despite the fact that, as has been mentioned before, the  $\mathcal{L}_p$  theory for elliptic equations can be developed independently of the theory of parabolic equations, in the main text we prefer to derive elliptic estimates from parabolic ones for the following reasons. In the first place, we want the reader to have some insight into the theory of parabolic equations. Secondly, to estimate the  $\mathcal{L}_2$  oscillation of  $u_{xx}$  over the unit ball  $B_1$  centered at the origin for a  $C_0^\infty$  function  $u$ , through the maximal function of  $\Delta u$  we split  $f := \Delta u$  into two parts:  $f = h + g$  where  $h \in C_0^\infty$  and  $h = f$  in the ball of radius 2 centered at the origin. Then we define  $v$  and  $w$  as solutions of the equations  $\Delta v = h$  and  $\Delta w = g$ . The trouble is that to find appropriate  $v$  and  $w$  is not so easy. Say, if  $d = 1$  and we want the equations  $\Delta v = h$  and  $\Delta w = g$  to be satisfied in the whole space, quite often  $v$  and  $w$  will be unbounded. In the parabolic case this difficulty does not appear because we can find  $v$  and  $w$  as solutions of the Cauchy problem with zero initial condition, when the initial condition is given for  $t$  lying outside the domain where we are estimating the  $\mathcal{L}_2$  oscillation of the solution.

One more point worth noting is that one can have a short course on elliptic equations, and after going through Chapter 1, go directly to Chapters 8–10 if one is only interested in the case  $p = 2$ . If  $d = 2$ , one can also include Chapter 11. Adding to this list Chapter 3 and Section 4.1 would allow the

reader to follow all the material in full generality apart from what concerns parabolic equations and equations with VMO coefficients. In this case one also has to follow the proof of Lemma 6.3.8 in order to get control on the  $\mathcal{L}_p$  norm of solutions. Finally, doing Exercise 4.3.2 allows one to include the results of Chapter 6 related to the elliptic equations with VMO coefficients.

If one wants to give a course containing both Hölder space and Sobolev space theories, then one can start with part of the present notes, use the possibility of obtaining basic  $C^{2+\alpha}$  estimates by doing Exercises 4.3.2, 4.3.3, and 10.1.8, and then continue lecturing on Hölder space theory following one's favorite texts. By the way, this switch to Hölder space theory can be done right after Chapter 1 if only elliptic equations are to be treated. For parabolic equations this switch is possible after going through Chapters 1 and 2 and doing Exercises 4.3.5, 4.3.6, 10.1.9, and 10.1.10.

### Acknowledgements

I worked on these lecture notes and taught courses in Moscow and Minneapolis for a number of years. Many graduate students were exposed to constantly evolving versions of the notes and contributed to correcting a few errors and sometimes wrong or misleading hints to exercises. Their impact is greatly appreciated. Several of my colleagues have influenced the contents of the notes. One of them is my friend Eugene Fabes, who died prematurely in 1997. I learned a very substantial part of real analysis attending his brilliant and enthusiastic lectures and having many discussions.

My deepest thanks are due to N.N. Ural'tseva who heroically undertook the task of reading through the final draft of the notes. I have greatly profited from her comments.

Also, in the final stages of the work I was partially supported by the NSF grants DMS-0140405 and DMS-0653121.

Nicolai Krylov, Minneapolis, August 2007

# Bibliography

- [1] S. Agmon, A. Douglis, and L. Nirenberg, *Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions, I*, Comm. Pure Appl. Math., Vol. 12, 1959, 623–727; *II*, *ibid.*, Vol. 17 (1964), 35–92.
- [2] H. Amann, *Maximal regularity for nonautonomous evolution equations*, Adv. Non-linear Stud., Vol. 4 (2004), No. 4, 417–430.
- [3] L. Bers, F. John, and M. Schechter, “Partial differential equations”, Interscience Publishers—John Wiley & Sons, New York—London—Sydney, 1964.
- [4] N. Dunford and J.T. Schwartz, “Linear operators, Part I, General theory”, Interscience Publishers, New York, 1958.
- [5] N. Dunford and J.T. Schwartz, “Linear operators, Part III. Spectral operators”, Wiley-Interscience, New York, 1971.
- [6] E. DiBenedetto, “Real analysis”, Birkhäuser Advanced Texts, Basel Textbooks, Birkhäuser Boston, Inc., Boston, MA, 2002.
- [7] L.C. Evans, “Partial differential equations”, Graduate Studies in Mathematics, 19, Amer. Math Soc., Providence, RI, 1998.
- [8] M. Giaquinta, “Introduction to regularity theory for nonlinear elliptic systems”, Birkhäuser, Basel–Boston–Berlin, 1993.
- [9] D. Gilbarg and N.S. Trudinger, “Elliptic Partial Differential Equations of Second Order”, Series: Classics in Mathematics, Springer, Berlin, Heidelberg, New York, 2001.
- [10] R. Haller-Dintelmann, H. Heck, and M. Hieber,  *$L^p - L^q$  estimates for parabolic systems in non-divergence form with VMO coefficients*, J. London Math Soc., Vol. 74 (2006), No. 3, 717–736.
- [11] Doyoon Kim, *Parabolic equations with measurable coefficients in  $L_p$ -spaces with mixed norms*, <http://aps.arxiv.org/abs/0705.3808>
- [12] N.V. Krylov, “Lectures on elliptic and parabolic equations in Hölder spaces”, Amer. Math. Soc., Providence, RI, 1996.
- [13] O.A. Ladyzhenskaya and N.N. Ural'tseva, “Linear and quasi-linear elliptic equations”, Nauka, Moscow, 1964, in Russian; English translation: Academic Press, New York, 1968; 2nd Russian ed. 1973.

- 
- [14] O.A. Ladyzhenskaya, V.A. Solonnikov, and N.N. Ural'tseva, "Linear and quasi-linear parabolic equations", Nauka, Moscow, 1967, in Russian; English translation: Amer. Math. Soc., Providence, RI, 1968.
  - [15] G.M. Lieberman, "Second order parabolic differential equations", World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
  - [16] A. Lunardi, "Analytic semigroups and optimal regularity in parabolic problems", Progr. Nonlinear Differential Equations Appl., Vol. 16, Birkhäuser Verlag, Basel, 1995.
  - [17] A. Maugeri, D. K. Palagachev, and L. G. Softova, "Elliptic and Parabolic Equations with Discontinuous Coefficients", Mathematical Research, Vol. 109, Wiley, Berlin, etc., 2000.
  - [18] V.A. Solonnikov, *On boundary value problems for parabolic systems of general form*, Proceedings of the Steklov Institute of Math. in the Academy of Sci. of the USSR, Vol. 83, 1965, 1-184, Amer. Math. Soc., Library of Congress Card 66-26640, 1967.
  - [19] E.M. Stein, "Harmonic analysis: Real-variable methods, orthogonality and oscillatory integrals", Princeton Univ. Press, Princeton, NJ, 1993.
  - [20] H. Triebel, "Theory of function spaces", Birkhäuser Verlag, Basel–Boston–Stuttgart, 1983.
  - [21] P. Weidemaier, *On the sharp initial trace of functions with derivatives in  $L_p(0, T; L_p(\Omega))$* , Bollettino U.M.I. (7), 9-B (1995), 321-338.
  - [22] P. Weidemaier, *On  $L_p$ -estimates of optimal type for the parabolic oblique derivative problem with VMO-coefficients—a refined version*, pp. 529–536 in Nonlinear Elliptic and Parabolic Problems, Progr. Nonlinear Differential Equations Appl., Vol. 64, Birkhäuser, Basel, 2005.

# Index

## Miscellanea

$(f, g)$ , 267  
 $\langle f, \phi \rangle$ , 281  
 $a_R^\#(x)$ , 130  
 $a_R^\#$ , 130  
 $a_\pm$ , 232  
 $\mathbb{B}$ , 87, 93  
 $c_d$ , 2, 267  
 $\mathcal{F}^0$ , 73  
 $f^\#$ , 80  
 $f_p^\#$ , 82  
 $f|_n$ , 75  
 $f|_\tau$ , 75  
 $g^\#$ , 87  
 $g_\Omega(x, y)$ , 262  
 $\kappa(L)$ , 277  
 $\kappa(\sigma)$ , 274  
 $N^\alpha(L, \mu)$ , 277  
 $N^\alpha(\sigma, \mu)$ , 274  
 $N_n(L, \mu)$ , 277  
 $N_n(\sigma, \mu)$ , 276  
 $\text{osc}_x(a, Q_r(t, x))$ , 130  
 $\mathbb{Q}$ , 87, 93  
 $\rho(\Omega)$ , 203  
 $\sigma(\xi)$ , 269  
 $\sigma_L$ , 277  
 $\sigma_L(\xi)$ , 269  
 $\sigma_{\partial\Omega}$ , 260  
 $x'$ , xvi, 11, 160  
 $\xi^\alpha$ , xiv

## Domains and sets

$B_1$ , 7  
 $B_r$ , 23, 60, 93  
 $B_r(x)$ , 60, 93, 165, 203

$C_n(i_1, \dots, i_d)$ , 74  
 $C_n(x)$ , 73  
 $Q_r$ , 60, 93  
 $Q_r(t, x)$ , 60, 93  
 $\mathbb{R}_T^{d+1, c}$ , 68  
 $\mathbb{R}^{d+1}$ , xv  
 $\mathbb{R}_{T, S}^{d+1}$ , 149  
 $\mathbb{R}_T^{d+1}$ , 46  
 $\mathbb{R}^d$ , xiii  
 $\mathbb{R}_+^d$ , xvi, 11, 160  
 $\mathbb{Z}$ , 73  
 $\Omega \in C^k$ , 165  
 $\Omega_T$ , 67  
 $\bar{\Omega}$ , xvi, 8  
 $\partial B$ , 3  
 $\partial\Omega$ , 158

## Function spaces

$BMO$ , 83  
 $BMO$ , 135  
 $\text{Camp}_\alpha(u)$ , 208  
 $C(\Omega)$ , xvi  
 $C^k(\bar{\Omega})$ , xvi, 8  
 $C_0^k$ , xvi  
 $C_0^k(\bar{\Omega})$ , xvi, 11  
 $C^n$ , 346  
 $C^n(\Omega)$ , xvi  
 $C^{r, k+2r}(\Omega)$ , 46  
 $C^{r, k+2r}(\bar{\Omega})$ , 46  
 $C_0^{r, k+2r}(\bar{\Omega})$ , 46  
 $C_0^{r, k+2r}(\Omega)$ , 46  
 $C^\gamma$ , 203  
 $C_0^\infty$ , xv, 267  
 $C_0^\infty(\mathbb{R}^d)$ , 267  
 $C_0^\infty(\Omega)$ , xv, 9

$C_b^\infty$ , 34  
 $C_{loc}^\infty(\Omega)$ , 10  
 $C_0$ , xvi  
 $H_p^\gamma$ , 318  
 $\mathcal{H}$ , 313  
 $L_p(\Omega)$ , xv  
 $\mathbb{L}$ , 73  
 $\mathcal{L}(1, \infty)$ , 312  
 $\mathcal{L}_{1,loc}$ , 73  
 $\mathcal{L}_p$ , xv, 267  
 $\mathcal{L}_p(\partial\Omega)$ , 260  
 $\mathcal{L}_p\text{-}\mathcal{L}_q(\mathbb{R}_{T,S}^{d+1})$ , 150  
 $\mathcal{L}^{[1,\infty]}$ , 312  
 $S$ , 268  
 $VMO$ , 134  
 $W_{p,q}^{1,2}$ , 153  
 $W_{p,q}^{1,2}(\mathbb{R}_{T,S}^{d+1})$ , 153  
 $W_{p,q}^{1,2}(\mathbb{R}_T^{d+1})$ , 153  
 $W_{p,loc}^k(\Omega)$ , 10  
 $W_p^k$ , 8  
 $W_p^k(\Omega)$ , 8  
 $W_p^{r,k+2r}$ , 51  
 $W_p^{r,k+2r}(\Omega)$ , 51  
 $\overset{0}{W}_{p,q}^{1,2}(\Omega)$ , 155  
 $\overset{0}{W}_p^{1,2}$ , 68  
 $\overset{0}{W}_p^k(\Omega)$ , 158  
**Operators**  
 $(1 - \Delta)^{\gamma/2}$ , 277, 297  
 $\Delta$ , 1, 270  
 $Du$ , xv  
 $D^\alpha$ , xiv  
 $D^\alpha u(t, x')$ , 258  
 $\partial_t u$ , xv, 45  
 $\delta_{kh} f$ , 181  
 $D_k$ , xiv  
 $f_A$ , 73  
 $F$ , 2, 267  
 $F^{-1}$ , 268  
 $\mathcal{G}_\lambda^0$ , 54  
 $\mathcal{G}_\lambda$ , 48  
 $L_\psi$ , 278  
 $\mathbb{M}g$ , 87  
 $\mathcal{M}f$ , 78  
 $u(\varepsilon)$ , 34  
 $u(\ell)$ , 190  
 $u_B$ , 94, 208  
 $u_B(t)$ , 94  
 $u_Q$ , 93  
 $u_t$ , xv  
 $u_{xx}$ , xiv, xv  
 $u_x$ , xiv  
 $T_\Gamma$ , 256  
 $f$ , 73, 87, 93

$\frac{\partial}{\partial \bar{z}}$ , 272  
 $\bar{\partial}$ , 272  
 $\nabla u$ , xiv

### Norms

$[u]_{C^\alpha}$ , 203  
 $[u]_{C^\alpha(\Omega)}$ , 207  
 $[u]_{W_p^k(\Omega)}$ , 10, 182  
 $[u]_{BMO(\mathbb{R}^d)}$ , 214  
 $[\cdot]_{BMO}$ , 83  
 $\|f\|_{\mathcal{L}_p(\Omega)}$ , xv  
 $\|g\|_{H_p^\gamma}$ , 318  
 $\|u\|_{C^k(\Omega)}$ , xvi  
 $\|u\|_{C^\alpha}$ , 203  
 $\|u\|_{C^\gamma}$ , 203  
 $\|u\|_C$ , 203  
 $\|u\|_{W_p^k(\Omega)}$ , 8, 10  
 $\|u\|_{W_p^{r,k+2r}(\Omega)}$ , 51

### Sizes

$|A|$ , 73  
 $|B|$ , 87, 93  
 $|Q|$ , 93  
 $|\alpha|$ , xiv  
 $|\Gamma|$ , 207

a priori estimate, 16  
 adapted sequences, 84  
 adjoint symbol, 326  
 Agmon's inequality, 326  
 analytic semigroup, 333

bounded mean oscillation, 135  
 Brouwer's fixed point theorem, 8

caloric function, 67  
 $L$ -caloric function, 102  
 Campanato space, 207  
 Cauchy's operator, 297  
 Cauchy-Riemann operator, 272  
 characteristic polynomial, 43, 270  
 classical maximal function, 87  
 classical sharp function, 87  
 compact set, xv  
 constant of ellipticity, 14, 271, 274, 277  
 constant under control, xiv

Davis's inequality, 84, 85  
 defining sequence, 8, 51  
 derivative in the sense of distributions, 9  
 domain, xiv, 8  
 dyadic cubes, 74  
 dyadic maximal function, 91  
 dyadic sharp function, 87

eigenfunctions, 228

- eigenvalues, 228  
elliptic symbol, 274
- Fan Ky minimax theorem, 8  
Fefferman-Stein theorem, 81  
filtration of partitions, 74  
first boundary-value problem, 158  
formally adjoint operator, 270, 277  
Fourier transform, 2, 267  
freezing the coefficients, 22  
functions of bounded mean oscillation, 83
- generalized derivative, 8, 9  
global barriers, 232  
global regularity, 29, 57, 120, 189  
grad, xiv  
Green's function, 25, 254, 262  
Green's measure, 245
- Hardy's inequality, 160  
Hardy-Littlewood theorem, 88  
Hardy-Littlewood-Sobolev inequality, 343  
harmonic function, 67  
Hilbert transform, 299  
Hilbert's identity, 334  
Hölder space, 203  
homogeneous Cauchy's operator, 299  
homogeneous elliptic operator, 271  
Hopf's lemma, 5
- interior cone condition, 212  
interior diameter, 203
- John-Nirenberg class, 83  
John-Nirenberg space, 135  
John-Nirenberg theorem, 83
- kernel, 301  
Kondrashov's theorem, 223
- Laplace's operator, 1, 270  
Laplacian, 1  
Lebesgue differentiating theorem, 80  
Lipschitz continuous function, 38  
local regularity, 30, 60, 122, 196, 237
- $m$ th order operator, 269  
maximal function, 78  
maximal inequality, 78  
maximum principle, 4, 233, 243, 254, 262  
method of continuity, 15  
Minkowski's inequality, 33  
modification, 32, 207, 342  
multi-index, xiv  
multiplicative inequalities, 17, 326
- Neumann boundary-value condition, 190  
Neumann problem, 191  
Newtonian potential, 39
- oblique derivative problem, 190  
one-dimensional case, 50, 56
- parabolic dilation, 48  
parabolic dyadic cubes, 74  
parabolic Sobolev spaces, 51  
partition of unity, 23, 172  
Poincaré's inequality, 212, 226  
principal part, 270  
pseudo-differential operator, 277
- quadratic variation, 84
- resolvent operator, 236  
Riesz transforms, 299  
Riesz-Calderón-Zygmund decomposition, 77
- Sarason's class, 134  
second-order elliptic differential operator, 14  
sharp function, 80  
singular-integral operator, 41  
Slobodetskii space, 207  
Sobolev derivative, 9  
Sobolev mollifier, 32, 35, 348  
Sobolev spaces, 8, 158  
Sobolev-Poincaré inequality, 226  
spaces of Bessel potentials, 318  
spectrum, 228  
stopping time, 75  
strongly elliptic operator, 269, 277  
support, xv  
symbol, 274
- test function, 9  
two-dimensional case, 17
- uniformly elliptic operator, 328
- vanishing mean oscillation, 134  
von Neumann series, 241
- Weierstrass function, 307
- Young's inequality, 19, 33
- Zygmund space, 307

This book concentrates on the basic facts and ideas of the modern theory of linear elliptic and parabolic equations in Sobolev spaces.

The main areas covered in this book are the first boundary-value problem for elliptic equations and the Cauchy problem for parabolic equations. In addition, other boundary-value problems such as the Neumann or oblique derivative problems are briefly covered. As is natural for a textbook, the main emphasis is on organizing well-known ideas in a self-contained exposition. Among the topics included that are not usually covered in a textbook are a relatively recent development concerning equations with VMO coefficients and the study of parabolic equations with coefficients measurable only with respect to the time variable. There are numerous exercises which help the reader better understand the material.

After going through the book, the reader will have a good understanding of results available in the modern theory of partial differential equations and the technique used to obtain them. Prerequisites are basics of measure theory, the theory of  $L_p$  spaces, and the Fourier transform.

ISBN 978-0-8218-4684-1



9 780821 846841

GSM/96



For additional information  
and updates on this book, visit

[www.ams.org/bookpages/gsm-96](http://www.ams.org/bookpages/gsm-96)

AMS on the Web  
[www.ams.org](http://www.ams.org)