

# Dynamical Systems and Population Persistence

Hal L. Smith Horst R. Thieme

Graduate Studies in Mathematics Volume 118



**American Mathematical Society** 

# Dynamical Systems and Population Persistence

Hal L. Smith Horst R. Thieme

Graduate Studies in Mathematics

Volume 118



American Mathematical Society Providence, Rhode Island

#### **Editorial Board**

David Cox (Chair) Rafe Mazzeo Martin Scharlemann Gigliola Staffilani

2010 Mathematics Subject Classification. Primary 37N25, 92D25, 92D30; Secondary 37B25, 37Lxx.

> For additional information and updates on this book, visit www.ams.org/bookpages/gsm-118

#### Library of Congress Cataloging-in-Publication Data

Smith, Hal L.

Dynamical systems and population persistence / Hal L. Smith, Horst R. Thieme. p. cm. - (Graduate studies in mathematics ; v. 118)

Includes bibliographical references and index.

ISBN 978-0-8218-4945-3 (alk. paper)

1. Biology–Mathematical models. 2. Population biology. I. Thieme, Horst R., 1948-II. Title.

QH323.5.S58 2011 577.8'80151539-dc22

2010033476

Copying and reprinting. Individual readers of this publication, and nonprofit libraries acting for them, are permitted to make fair use of the material, such as to copy a chapter for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgment of the source is given.

Republication, systematic copying, or multiple reproduction of any material in this publication is permitted only under license from the American Mathematical Society. Requests for such permission should be addressed to the Acquisitions Department, American Mathematical Society, 201 Charles Street, Providence, Rhode Island 02904-2294 USA. Requests can also be made by e-mail to reprint-permission@ams.org.

> (c) 2011 by the American Mathematical Society. All rights reserved. The American Mathematical Society retains all rights except those granted to the United States Government.

Printed in the United States of America.

 $\infty$  The paper used in this book is acid-free and falls within the guidelines established to ensure permanence and durability. Visit the AMS home page at http://www.ams.org/

> 10987654321 $16\ 15\ 14\ 13\ 12\ 11$

To our doctoral advisors Willi Jäger (HRT) and Paul Waltman (HLS)

# Contents

Preface	ix
Introduction	1
Chapter 1. Semiflows on Metric Spaces	9
§1.1. Metric spaces	9
$\S1.2.$ Semiflows	17
$\S1.3.$ Invariant sets	19
$\S1.4.$ Exercises	25
Chapter 2. Compact Attractors	29
$\S2.1.$ Compact attractors of individual sets	30
§2.2. Compact attractors of classes of sets	36
$\S2.3.$ A sufficient condition for asymptotic smoothness	51
$\S2.4.$ $\alpha$ -limit sets of total trajectories	52
$\S2.5.$ Invariant sets identified through Lyapunov functions	52
$\S2.6.$ Discrete semiflows induced by weak contractions	54
$\S2.7.$ Exercises	57
Chapter 3. Uniform Weak Persistence	61
§3.1. Persistence definitions	61
§3.2. An SEIRS epidemic model in patchy host populations	64
$\S3.3.$ Nonlinear matrix models: Prolog	71
§3.4. The May-Leonard example of cyclic competition	78
§3.5. Exercises	84

v

Chapter	4. Uniform Persistence	87
$\S4.1.$	From uniform weak to uniform persistence	87
$\S4.2.$	From uniform weak to uniform persistence: Discrete case	91
$\S4.3.$	Application to a metered endemic model of SIR type	94
$\S4.4.$	From uniform weak to uniform persistence for time-set $\mathbb{R}_+$	97
$\S 4.5.$	Persistence à la Baron von Münchhausen	99
$\S4.6.$	Navigating between alternative persistence functions	107
$\S4.7.$	A fertility reducing endemic with two stages of infection	110
$\S4.8.$	Exercises	123
Chapter	5. The Interplay of Attractors, Repellers, and Persistence	125
$\S{5.1.}$	An attractor of points facilitates persistence	125
$\S{5.2.}$	Partition of the global attractor under uniform persistence	127
$\S{5.3.}$	Repellers and dual attractors	135
$\S{5.4.}$	The cyclic competition model of May and Leonard revisited	139
$\S{5.5.}$	Attractors at the brink of extinction	140
$\S{5.6.}$	An attractor under two persistence functions	141
$\S{5.7.}$	Persistence of bacteria and phages in a chemostat	142
$\S 5.8.$	Exercises	155
Chapter	6. Existence of Nontrivial Fixed Points via Persistence	157
$\S6.1.$	Nontrivial fixed points in the global compact attractor	158
$\S6.2.$	Periodic solutions of the Lotka-Volterra predator-prey model	160
$\S6.3.$	Exercises	162
Chapter	7. Nonlinear Matrix Models: Main Act	163
§7.1.	Forward invariant balls and compact attractors of bounded	
	sets	163
§7.2.	Existence of nontrivial fixed points	165
§7.3.	Uniform persistence and persistence attractors	167
§7.4.	Stage persistence	171
§7.5.	Exercises	175
Chapter	8. Topological Approaches to Persistence	177
$\S{8.1.}$	Attractors and repellers	177
$\S{8.2.}$	Chain transitivity and the Butler-McGehee lemma	180
§8.3.	Acyclicity implies uniform weak persistence	185
§8.4.	Uniform persistence in a food chain	191

§8.5.	The metered endemic model revisited	196
§8.6.	Nonlinear matrix models (epilog): Biennials	199
§8.7.	An endemic with vaccination and temporary immunity	209
§8.8.	Lyapunov exponents and persistence for ODEs and maps	215
§8.9.	Exercises	229
Chapter	9. An SI Endemic Model with Variable Infectivity	231
$\S{9.1.}$	The model	231
$\S{9.2.}$	Host persistence and disease extinction	236
$\S{9.3.}$	Uniform weak disease persistence	237
$\S{9.4.}$	The semiflow	239
$\S{9.5.}$	Existence of a global compact attractor	240
$\S{9.6.}$	Uniform disease persistence	245
$\S{9.7}.$	Disease extinction and the disease-free equilibrium	247
$\S{9.8.}$	The endemic equilibrium	249
$\S{9.9.}$	Persistence as a crossroad to global stability	250
$\S{9.10}.$	Measure-valued distributions of infection-age	254
Chapter	10. Semiflows Induced by Semilinear Cauchy Problems	261
$\S{10.1}.$	Classical, integral, and mild solutions	261
$\S{10.2}.$	Semiflow via Lipschitz condition and contraction principle	265
$\S{10.3.}$	Compactness all the way	266
$\S{10.4.}$	Total trajectories	271
$\S{10.5}.$	Positive solutions: The low road	273
$\S{10.6}.$	Heterogeneous time-autonomous boundary conditions	279
Chapter	11. Microbial Growth in a Tubular Bioreactor	283
$\S{11.1.}$	Model description	283
$\S{11.2.}$	The no-bacteria invariant set	287
$\S{11.3.}$	The solution semiflow	291
$\S{11.4.}$	Bounds on solutions and the global attractor	292
$\S{11.5.}$	Stability of the washout equilibrium	296
$\S{11.6}.$	Persistence of the microbial population	301
$\S{11.7.}$	Exercises	304
Chapter	12. Dividing Cells in a Chemostat	307
$\S{12.1.}$	An integral equation	309
$\S{12.2.}$	A $C_0$ -semigroup	314

$\S{12.3.}$	A semilinear Cauchy problem	318
$\S{12.4.}$	Extinction and weak persistence via Laplace transform	320
$\S{12.5.}$	Exercises	325
Chapter	13. Persistence for Nonautonomous Dynamical Systems	327
$\S{13.1.}$	The simple chemostat with time-dependent washout rate	327
$\S{13.2.}$	General time-heterogeneity	332
$\S{13.3.}$	Periodic and asymptotically periodic semiflows	335
$\S{13.4.}$	Uniform persistence of the cell population	336
$\S{13.5.}$	Exercises	339
Chapter	14. Forced Persistence in Linear Cauchy Problems	341
$\S{14.1.}$	Uniform weak persistence and asymptotic Abel-averages	342
§14.2.	A compact attracting set	343
$\S{14.3.}$	Uniform persistence in ordered Banach space	344
Chapter	15. Persistence via Average Lyapunov Functions	349
$\S{15.1.}$	Weak average Lyapunov functions	350
$\S{15.2.}$	Strong average Lyapunov functions	354
$\S{15.3.}$	The time-heterogeneous hypercycle equation	355
$\S{15.4.}$	Exercises	361
Appendiz	x A. Tools from Analysis and Differential Equations	363
§A.1.	Lower one-sided derivatives	363
§A.2.	Absolutely continuous functions	364
§A.3.	The method of fluctuation	365
§A.4.	Differential inequalities and positivity of solutions	367
§A.5.	Perron-Frobenius theory	372
§A.6.	Exercises	375
Appendiz	x B. Tools from Functional Analysis and Integral Equations	377
§B.1.	Compact sets in $L^p(\mathbb{R}_+)$	377
§B.2.	Volterra integral equations	378
§B.3.	Fourier transform methods for integro-differential equations	380
§B.4.	Closed linear operators	385
§B.5.	Exercises	390
Bibliogra	phy	391
Index		403

### Preface

The primary focus of this book is the mathematical theory of persistence. The theory is designed to provide an answer to such questions as which species, in a mathematical model of interacting species, will survive over the long term. In a mathematical model of an epidemic, will the disease drive a host population to extinction or will the host persist? Can a disease remain endemic in a population? Persistence theory can give a mathematically rigorous answer to the question of persistence by establishing an initial-condition-independent positive lower bound for the long-term value of a component of a dynamical system such as population size or disease prevalence.

Mathematically speaking, in its simplest formulation for systems of ordinary or delay differential equations, and for a suitably prescribed subset I of components of the system, persistence ensures the existence of  $\epsilon > 0$ such that  $\liminf_{t\to\infty} x_i(t) > \epsilon$ ,  $i \in I$  provided  $x_i(0) > 0$ ,  $i \in I$ . We say that these components persist uniformly strongly, or, more precisely, that the system is uniformly strongly  $\rho$ -persistent for the persistence function  $\rho(x) = \min_{i \in I} x_i$ . This persistence function  $\rho(x)$  may be viewed as the distance of state x to a portion of the boundary of the state-space  $\mathbb{R}^n_+$ , namely the states where one or more of species  $i \in I$  are extinct.

The adjective "strong" is often omitted; uniform weak  $\rho$ -persistence is defined similarly but with limit superior in place of limit inferior. The adjective "uniform" emphasizes that the lower bound  $\epsilon$  is independent of initial data satisfying the restriction  $x_i(0) > 0$ ,  $i \in I$ . Similarly, as in the definition of Lyapunov stability, the precise value of  $\epsilon$  is unspecified and usually difficult to estimate. Uniform persistence is a qualitative notion, not a quantitative one. However, in rare cases,  $\epsilon$  can be related to system parameters; this ideal situation is referred to as "practical persistence" [24, 25, 26, 28, 27, 35].

Weaker notions of weak and strong persistence drop the "uniformity with respect to initial data" (i.e.,  $\epsilon$ ), requiring merely positivity of the limit superior, respectively, the limit inferior.

The definition of persistence and the related concept of permanence (uniform persistence plus an upper bound on limit superior of all components) evolved in the late 1970s from the work of Freedman and Waltman [75], Gard [80, 81] Gard and Hallam [82], Hallam [96], and Schuster, Sigmund, Wolff [196]. Most of these early papers show weak persistence, but Schuster, Sigmund, and Wolff [196] prove uniform strong  $\rho$ -persistence for the hypercycle equation in the *n*-simplex with  $\rho(x) = x_1 \cdots x_n$  as persistence function.

The notion of a persistence function seems to have been introduced by Gard and Hallam [82, 80], though with a more technical intention than here. It was later superseded by a more general concept which combines the usual Lyapunov function methods with time averages [104] and became known as average Lyapunov function [109]. "Persistence function" (together with the  $\rho$ -symbol [80, 82]) is revived here as a means to make precise which parts of a system persist; in applications  $\rho$  has a very concrete and intuitive interpretation like the number of infected individuals to describe disease persistence in an epidemic model. Such a "hands on" interpretation would be lacking for a typical average Lyapunov function like  $x_1^{p_1} \cdots x_n^{p_n}$  [81, 82]. Zhao [238] uses the notion generalized distance function to stress the idea that  $\rho$  measures the distance to the brink of extinction.

Persistence theory developed rapidly in the 1980s because the necessary machinery from dynamical systems, a theory of attractors and repellers, was already in place. Early work focused on persistence of components of systems of ordinary differential equations. Later, this was extended to discrete time or difference equations, and then to infinite dimensional dynamical systems generated by delay differential equations and partial differential equations. Application of the theory was initially slow to catch on in the applied literature on population biology and epidemiology, but it has more recently become an accepted tool in theoretical population dynamics.

Although the theory is now quite "user friendly" in the sense that a user does not need to be an expert to use it, it is a mathematically sophisticated theory. Our motivation for writing this monograph grew out of the problem of teaching the theory to our graduate students. There are very few sources where one can find self-contained treatments that are accessible to graduate students. The survey articles by Waltman [231] and by Hutson and Schmitt [110] remain useful, although they do not contain more recent refinements of the theory which are scattered in the literature. Recent monographs by Cantrell and Cosner [25], Hofbauer and Sigmund [106], Thieme [217], and Zhao [238] are good sources, but their focus is broader than persistence theory.

This monograph began as a set of lecture notes for graduate students in a team-taught course on Dynamical Systems in Biology offered by the authors in fall 2005, spring 2007, and spring 2009 at Arizona State University. It contains a large number of homework exercises. A description of the contents of the chapters follows.

Chapters 1 through 8 contain our main results on persistence theory for autonomous dynamical systems.

Chapter 1 begins with a review of metric spaces, the natural abstract setting or "state space" for finite and infinite dimensional dynamics. The notion of a semiflow on a metric space is developed; it gives the dynamics. We distinguish discrete and continuous time semiflows simply by the time set: nonnegative integers in the former case, the nonnegative reals in the latter case. The basic properties of a semiflow are independent of the time set. This unified treatment of discrete time and continuous time semiflows allows us to unify the later treatment of persistence theory for discrete time and continuous time dynamics. In the literature, the theory developed separately for discrete and continuous time systems, but we have been largely successful in our attempt to present a unified treatment, avoiding as much as possible separate approaches. Persistence theory often requires that the underlying dynamics are dissipative in some sense. The strongest sense is that there is a compact attractor of bounded subsets. Although the trend of recent work, and one of our goals here, is to weaken these compactness requirements where possible, we present the theory of attractors in Chapter 2.

Chapter 3 begins with the definitions of persistence, both uniform weak and uniform strong persistence relative to a persistence function. However, its main focus is on uniform weak persistence and on elementary methods for establishing it. Several examples illustrating such methods are introduced. These include the continuous time model of an SEIRS infectious disease in a meta-population with host travel between patches, the classical May-Leonard system of three competing populations, and discrete time nonlinear matrix models of population dynamics. The latter include the LPA model of flour beetle dynamics and nonlinear versions of the well-studied Leslie-type demographic models.

Uniform strong persistence is the desired conclusion; uniform weak persistence is more easily obtainable. It has long been known that uniform weak persistence plus suitable compactness properties of the dynamical system give uniform strong persistence. See Freedman and Moson [72] for flows on locally compact metric spaces, Freedman, Ruan, and Tang [73] for flows, and Thieme [215] for semiflows on general metric spaces. In Chapter 4, we present a number of such results. Some of these have relatively weak compactness assumptions at the expense of lengthy and seemingly technical hypotheses. Others require more compactness assumptions but are more easily and concisely formulated. A number of applications are treated in detail, including those introduced in the previous chapter.

The choice of a persistence function may be not obvious; several different choices may be appropriate. The question then naturally arises as to whether, and how to prove that, persistence with respect to one such function implies persistence with respect to another persistence function. This issue is treated in Chapter 4.6.

For semiflows that are dissipative in a suitable strong sense and that are uniformly  $\rho$ -persistent, there is an elegant decomposition of the attractor into an extinction attractor, a persistence attractor, and a family of total trajectories whose  $\alpha$  limit sets are contained in the extinction attractor and whose  $\omega$  limit sets are contained in the persistence attractor. This result, versions of which were first proved by Hale and Waltman [95] and later by Zhao [238] and Magal and Zhao [158], is proved in Chapter 5.

The brief Chapter 6 explores various scenarios whereby one may establish that persistence implies the existence of a "persistence equilibrium", that is, an equilibrium  $x^*$  for which  $\rho(x^*) > 0$  where  $\rho$  denotes the persistence function. This provides an extra incentive for taking the trouble to establish persistence. The monograph by X.-Q. Zhao [238] contains a nice summary of the history of results in this direction. See the notes to Chapter 1 of [238]. Newer results also appear in the recent paper by Magal and Zhao [158].

Nonlinear matrix models, such as those introduced in Chapter 3, are increasingly being used in population modeling as indicated by the recent monographs [29, 44, 39]. Therefore, we devote Chapter 7 to applying the results of the previous chapters to them.

Chapter 8 treats the mathematically more sophisticated topological approach to persistence and its consequences. As Josef Hofbauer [**79**, **104**] has repeatedly pointed out, the theory of attractors and repellers, formulated by several mathematicians including Zubov, Ura, Kimura, and Conley, lead directly to proofs of many of the results of persistence theory. See [**79**] for many historical references to the work of Zubov and Ura and Kimura; the monograph of Bhatia and Szegő [**16**] contains some of this work. The notion of chain recurrence and chain transitivity of Conley [**33**] has also proved to

be very useful. These notions were originally established for flows on locally compact spaces but are needed for semiflows on potentially infinite dimensional Banach spaces for persistence theory. We give here a self-contained treatment of these ideas and how they are used in the theory. Most notable among the results implied by these ideas are the Butler-McGehee Theorem and the acyclicity theorem establishing uniform weak persistence.

These were first formulated by Butler, Freedman and Waltman [22, 23] for flows on locally compact spaces, extended to discrete time systems by Freedman and So [74] and Hofbauer and So [107]. They were later generalized to semiflows on infinite dimensional spaces by Hale and Waltman [95] assuming the existence of a compact attracting set, an assumption that was relaxed in [215].

Several more or less straightforward applications of the acyclicity approach to persistence are included. These include the classical three-level (ODE) food chain model considered by Hastings and collaborators, nonlinear matrix models for biennial species, and a metered epidemic model — a hybrid of both discrete and continuous time.

Finally, we show how classical Lyapunov exponents may be used to establish one of the key hypotheses in the acyclicity theorem, namely, that a compact invariant set, belonging to the "extinction set", is uniformly weakly repelling in directions normal to the extinction set. The use of Lyapunov exponents in the study of biological models was pioneered by Metz et al. [168], who proposed that the dominant Lyapunov exponent gives the best measure of invasion fitness, and by Rand et al. [180] who used it to characterize the invasion "speed" of a rare species. Roughly, a positive dominant Lyapunov exponent corresponding to a potential invading species in the environment set by a resident species attractor implies that the invader can successfully invade. Our treatment is patterned after the approach taken by Paul Salceanu in his thesis [186] and [187, 188, 189, 190].

Chapter 9 focuses on an SI epidemic model where infectives are structured by age since infection and where the force of infection depends on an age-since-infection weighted average of current infectives. The model can be reduced to a system of integral equations; existence and uniqueness of solutions, and boundedness of solutions are proved. The host is shown to (uniformly) persist, the basic replacement number  $\mathcal{R}_0$  is identified, disease extinction is shown to occur if  $\mathcal{R}_0 < 1$ , and uniform weak persistence of the disease is shown if  $\mathcal{R}_0 > 1$ . In order to obtain uniform persistence of the disease, it is useful to reformulate the dynamics as a semiflow on a suitable Banach space. This is done by showing that solutions satisfy a weakly formulated semilinear Cauchy problem. One can then show the existence of a compact attractor of bounded sets under suitable restrictions. This in turn facilitates the argument for uniform persistence of the disease when  $\mathcal{R}_0 > 1$ . The existence of an endemic equilibrium is also established, and rather unrestrictive conditions for its global stability are derived. It should be noted that persistence is indispensable for doing the latter because the Lyapunov function that is used is not defined on the whole state space but only on the persistence attractor (see also [155]).

Chapter 10 is devoted to a brief treatment of the semilinear Cauchy problem u' = Au + F(u),  $u(0) = u_0$  in a Banach space setting. Here Ais a closed linear operator and F is a nonlinear map. Notions of classical, integral, and mild solutions are defined, and the equivalence of mild and integral formulation is shown. Globally defined integral solutions are shown to define a semiflow, and local existence is established by the contraction mapping principle when F satisfies a Lipschitz condition. If F is suitably bounded, global in time existence is also shown. Conditions for the induced semiflow to be asymptotically smooth, a key requirement for showing the existence of a compact global attractor, are identified. As we have biological examples in mind, positivity of solutions must be satisfied. Conditions which ensure positivity of solutions are formulated.

Chapter 11 treats microbial growth on a growth-limiting nutrient in a tubular bioreactor. Fresh nutrient enters the left side of the tube, and unused nutrient and microbes leak out the right side of the tube in proportion to their concentration. Both nutrient and microbes are assumed to diffuse throughout the tube. The issue is whether or not the influx of nutrient is sufficient to allow the microbes to persist in the bioreactor. Relying heavily on the machinery of Chapter 10, we show that the system of reaction-diffusion equations generates a dissipative semiflow. Linearized stability analysis of the so-called washout equilibrium solution (no microbes) leads to a basic reproduction number  $\mathcal{R}_0$ . If  $\mathcal{R}_0 < 1$ , the microbes are "washed out" of the bioreactor, and, if  $\mathcal{R}_0 > 1$ , they uniformly persist and there is a unique colonization equilibrium.

Chapter 12 considers a model of microbial growth in a chemostat where microbial cells of different age take up nutrient at differing rates and divide at an age-dependent rate. Ignoring growth and uptake, focusing only on demographics of cell division, we begin by obtaining a renewal equation for cell population division rate and showing that it has a unique solution. This leads to the definition of a semigroup of operators and ultimately to a formulation of the full model, including growth and uptake, as an abstract ODE in a Banach space setting. Its mild solutions are shown to generate a semiflow. Consideration of the "washout state", absent microbes, allows identification of the basic "biomass production number" for the model. When it is less than one, and an additional condition satisfied, the microbes are washed out; when it exceeds one, the cell population persists uniformly weakly. Proofs of these results make use of the Laplace transform. In fact, uniform persistence of the cell population holds when the basic production number exceeds one under additional assumptions, but the proof is deferred to a later chapter. The problem is in establishing sufficient compactness of the semiflow. A different approach provides another route from uniform weak to uniform strong persistence, which succeeds for this model.

Chapter 13 is devoted to persistence for nonautonomous systems. Practical persistence is established, under suitable conditions, for a population of micro-organisms growing in a chemostat with time-dependent dilution rate using elementary arguments. It is also established that all positive solutions are asymptotic to each other. The abstract notion of a nonautonomous semiflow is introduced, corresponding definitions of persistence are given, and several results giving conditions under which uniform weak persistence implies uniform strong persistence are proved. Special attention is devoted to the case of periodic nonautonomous semiflows and nonautonomous semiflows that are asymptotic to such semiflows. The implication that uniform weak persistence implies uniform persistence for these cases is specialized. Finally, uniform persistence is established for the (autonomous) cell division model treated in Chapter 12 by using the methods developed for nonautonomous semiflows.

As noted in our description of Chapter 3 above, persistence functions were introduced early on in the history of persistence theory as a means to obtain uniform persistence (permanence) in much the same way that they are used in Lyapunov stability theory to obtain stability results for equilibria of dynamical systems [82], and there is now a well-developed approach to establishing persistence using so-called average Lyapunov functions (a generalization of persistence functions in the sense of Gard and Hallam [82]). The works of Fonda [71], Gard [80], Hofbauer [103], Hutson [109], and Schuster, Sigmund and Wolff [196] have been very influential. Some of their ideas, as well as later work, have been reviewed in the paper of Hutson and Schmitt [110] and the monograph of Sigmund and Hofbauer [106]. However, so far in this work, we have used persistence functions primarily as a means to precisely define what is meant by persistence, not as a tool with which to establish it.

In Chapter 15, taking inspiration from this large literature, we formulate some general results which yield persistence using the average Lyapunov function approach. The adjective "average" in the terminology signifies that a time-average of the function over a sufficiently large interval should be positive. We formulate an approach which works for nonautonomous semiflows and, as usual, seeks to minimize compactness requirements. These goals force rather technical statements, but the main results are simple: The existence of a weak average Lyapunov function  $\rho$  implies weak  $\rho$ -persistence; the existence of a strong average Lyapunov function implies uniform  $\rho$ -persistence. As an application, the hypercycle equation, treated in Chapter 12 of [106] in the autonomous case, is extended to the case where replication rates may be time-dependent.

The book ends with two appendices. The first, Appendix A, covers some useful techniques in differential equations which are not usually covered in a basic course. Chief among these are differential inequalities, a key tool in applied dynamics. Here, we mean Kamke's comparison theorem for ODEs and the strong maximum principle for PDEs. The former result is proved, the latter is merely stated and references are given. Dynamical systems in biology typically deal with nonnegative quantities, and therefore one needs to establish that solutions that begin nonnegative, remain so in the future. Another essential tool for dealing with positivity and stability is the Perron-Frobenius theory which we state but do not prove. Finally, an elementary but powerful method which can sometimes establish persistence is the method of fluctuation. It provides the means to explicitly estimate the limit inferior and limit superior of bounded components of solutions of systems of ordinary and delay differential equations.

Appendix B introduces selected useful tools from functional analysis. Among them are compactness criteria in  $L^p$  spaces, inequalities for Volterra integral equations, proof of the equivalence of integral and mild solutions of linear differential equations in Banach spaces, and Fourier transform methods for integro-differential equations. The latter leads to conditions implying that any bounded solution of a class of integro-differential inequalities or equations vanishes identically, and this result may be used to establish global stability results. These tools are used in Chapter 9 and Chapter 12.

One should also disclose what is not in this book that a reader might expect given the title. One such omission is the notion of robust persistence, more precisely, the reasonable expectation that the notion of uniform persistence should be structurally stable to small changes in system dynamics in some topology. For example, if the dissipative system takes the Kolmogorov form  $x'_i = x_i f_i(x)$  on  $\mathbb{R}^n_+$ , then small perturbations should mean small changes in the per capita growth rates  $f_i$ , say in the  $C^r$ -topology. Robust  $(C^r)$   $\rho$ -persistence with  $\rho(x) = \min_i x_i$  for this system would mean the existence of  $\epsilon, \delta > 0$  such that  $\liminf_{t\to\infty} x_i(t) > \epsilon$ ,  $\forall i$  provided x(t)satisfies  $x'_i = x_i g_i(x)$  where  $||f - g||_{C^r} < \delta$  and  $x_i(0) > 0$ ,  $\forall i$ . Such results were first established by Schreiber [193]. See also Hirsch et al. [101]. We do not include these results since they are partly covered in the monograph of Zhao [238]. Finally, we have not included recent work on persistence for stochastic systems [11, 105, 194] or for skew-product semiflows [169, 238].

There is a huge body of literature on persistence theory, and this book does not span nearly all of it. We ask the forgiveness of our valued fellow scholars whose works we have failed to reference.

We would like to acknowledge the many students, especially Thanate Dhirasakdanon, who have contributed to this work through their questions, suggestions, and their homework solutions.

We thank our wives, Kathryn Smith and Adelheid Thieme, for their unwavering support, though this endeavor must have been shrouded in mystery for them.

As much as any other science, mathematics takes place in a tapestry of teachers, peers, and students; we gratefully dedicate this monograph to our Ph.D. advisors Willi Jäger (HRT) and Paul Waltman (HLS).

Hal Smith was supported in part by NSF Grant DMS-0918440. Horst Thieme was supported in part by NSF Grant DMS-0715451.

## Bibliography

- R. Abraham, and J.E. Marsden, *Foundations of Mechanics*, 2nd edition, The Benjamin/Cummings Publishing Co. Inc., 1978.
- [2] A.S. Ackleh, B.G. Fitzpatrick, H.R. Thieme, Rate distributions and survival of the fittest: a formulation on the space of measures, Disc. Cont. Dyn. Sys. B 5 (2005), 917–928.
- [3] A.S. Ackleh, K. Ito, Measure-valued solutions for a hierarchically size-structured population, J. Differential Equations 217 (2005), 431–455.
- [4] R.M. Anderson, R.M. May, Infectious Diseases of Humans. Dynamics and Control, Oxford University Press, 1991.
- [5] W. Arendt, C.J.K. Batty, M. Hieber, F. Neubrander, Vector-valued Laplace Transforms and Cauchy Problems, Birkhäuser, 2001.
- [6] J. Arino, C.C. McCluskey, P. van den Driessche, Global results for an epidemic model with vaccination that exhibits backward bifurcation, SIAM J. Appl. Math. 64 (2003), 260–276.
- [7] L. Arnold, Random Dynamical Systems, Springer-Verlag, 1998.
- [8] P. Ashwin, J. Buescu, I. Stewart, From attractor to chaotic saddle: a tale of transverse instability, Nonlinearity 9 (1996), 703–737.
- [9] I. Barbalat, Systèmes d'équations différentielles d'oscillations nonlinéaires, Rev. Roumaine Math. Pures Appl. 4 (1959), 267–270.
- [10] M. Barnsley, *Fractals Everywhere*, 2nd edition, Academic Press, 1993.
- [11] M. Benaim, J. Hofbauer, W.H. Sandholm, Robust permanence and impermanence for stochastic replicator dynamics, J. Biol. Dynamics 2 (2008), 180–195.
- [12] E. Beretta, T. Hara, W. Ma, Y. Takeuchi, Global asymptotic stability of an SIR epidemic model with distributed time delay, Nonlinear Analysis 47 (2001), 4107-4115.
- [13] E. Beretta, F. Solimano, Y. Tang, Analysis of a chemostat model for bacteria and virulent bacteriophage, Discrete and Continuous Dynamical Systems-B, 2 (2002), 495–520.
- [14] H.C. Berg, Random Walks in Biology, Princeton Univ. Press, 1993.

- [15] A. Berman and R. Plemmons, Nonnegative Matrices in the Mathematical Sciences, Academic Press, 1979.
- [16] N.P. Bhatia and G.P. Szegő, Stability Theory of Dynamical Systems, Springer, 2002.
- [17] G.A. Bürger, Wunderbare Reisen zu Wasser und Lande, Feldzüge und lustige Abenteuer des Freiherrn von Münchhausen nach der Ausgabe von 1788 mit einem Anhang älterer Lügengeschichten - (I. Ruttmann, ed.) Philipp Reclam jun., 1969.
- [18] T.A. Burton, V. Hutson, Permanence for nonautonomous predator-prey systems, Differential Integral Equations 4 (1991), 1269–1280.
- [19] S.N. Busenberg and K.L. Cooke, The effect of integral conditions in certain equations modelling epidemics and population growth, J. Math. Biol. 10 (1980), 13–32.
- [20] S.N. Busenberg and K.L. Cooke, Vertically Transmitted Diseases, Springer, 1993.
- [21] S.N. Busenberg, K.L. Cooke, M.A. Pozio, Analysis of a model of a vertically transmitted disease, J. Math. Biol. 17 (1983), 305–329.
- [22] G. Butler, H.I. Freedman, P. Waltman, Uniformly persistent systems, Proc. Amer. Math. Soc. 96 (1986), 425–430.
- [23] G. Butler and P. Waltman, Persistence in dynamical systems, J. Diff. Eqns. 63 (1986), 255–263.
- [24] R.S. Cantrell and C. Cosner, Practical persistence in ecological models via comparison methods, Proc. Royal Soc. Edinburgh A 126 (1996), 247–272.
- [25] R.S. Cantrell and C. Cosner, Spatial Ecology via Reaction-Diffusion Equations, Wiley, 2003.
- [26] Y. Cao and T.C. Gard, Uniform persistence for population models with time delay using multiple Lyapunov functions, Diff. Int. Equations 6 (1993), 883-898.
- [27] Y. Cao and T.C. Gard, Practical persistence for differential delay models of population interactions, Differential Equations and Computational Simulations III (J. Graef, R. Shivaji, B. Soni, J. Zhu, eds.), Electronic J. Differential Equations, Conference 01, 1997, pp. 41–53.
- [28] Y. Cao, J. Fan, T.C. Gard, Persistence in Kolmogorov population models with time delay, Proc. First World Congress of Nonlinear Analysis (V. Lakshmikantham, ed.), 3741-3751, de Gruyter, 1996
- [29] H. Caswell, Matrix Population Models, Sinauer Assoc. Inc., 1989.
- [30] C.-H. Chiu and S.-B. Hsu, Extinction of the top predator in a three-level food chain, J. Math. Biol. 37 (1998), 372–380.
- [31] S.-N. Chow and J.K. Hale, Methods of Bifurcation Theory, Springer, 1982.
- [32] C.W. Clarke, Mathematical Bioeconomics: The Optimal Management of Renewable Resources, Wiley, 1976.
- [33] C. Conley, Isolated Invariant Sets and the Morse Index, CBMS 38, Amer. Math. Soc., 1978.
- [34] W. Coppel, Stability and Asymptotic Behavior of Differential Equations, Heath, 1965.
- [35] C. Cosner, Persistence (permanence), compressibility, and practical persistence in some reaction-diffusion models from ecology, Comparison Methods and Stability Theory (X. Liu, D. Siegel, eds.), Lecture Notes in Pure and Applied Mathematics, 162, Marcel Dekker, Inc., 1994.

- [36] R.F. Costantino, J.M. Cushing, B. Dennis, R.A. Desharnais, Experimentally induced transitions in the dynamic behavior of insect populations, Nature 375 (1995), 227–230.
- [37] P. Cull, Local and global stability for population models, Biol. Cybernetics 54 (1986), 141–149.
- [38] J.M. Cushing, Nonlinear Matrix Models and Population Dynamics, Natural Resource Modeling 2 (1988), 539–580.
- [39] J.M. Cushing, An Introduction to Structured Population Dynamics, CBMS-NSF Regional Conf. Series in Applied Math. vol. 77, 1998.
- [40] J.M. Cushing, Cycle chains and the LPA model, J. Difference Equations Appl. 9 (2003), 655–670.
- [41] J.M. Cushing, Nonlinear semelparous Leslie models, Math. Biosci. Eng. 3 (2006), 17–36.
- [42] J.M. Cushing, Three stage semelparous Leslie models, J. Math. Biol. 59 (2009), 75–104.
- [43] J.M. Cushing, R.F. Costantino, B. Dennis, R.A. Desharnais, S.M.Henson, Nonlinear population dynamics: Models, experiments and data, J. Theor. Biology 194 (1998), 1–9.
- [44] J.M. Cushing, R.F. Costantino, B. Dennis, R.A. Desharnais, S.M.Henson, Chaos in Ecology, experimental nonlinear dynamics, Academic Press, 2003.
- [45] J.M. Cushing and J. Li, On Ebenman's model for the dynamics of population with competing juveniles and adults, Bull. Math. Biol. 51 (1989), 687–713.
- [46] E.M.C. D'Agata, P. Magal, S. Ruan, G.F. Webb, Asymptotic behavior in nosocomial epidemic models with antibiotic resistance, Diff. Integral Equations 19 (2006), 573–600.
- [47] N. Davydova, Old and Young. Can they coexist? Thesis, University of Utrecht, 2004.
- [48] N.V. Davydova, O. Diekmann, S.A. van Gils, Year class coexistence or competitive exclusion for strict biennials?, J. Math. Biol. 46 (2003), 95-131.
- [49] K. Deimling, Nonlinear Functional Analysis, Springer, 1985.
- [50] W. Desch and W. Schappacher, Linearized stability for nonlinear semigroups, Differential Equations in Banach Spaces (A. Favini, E. Obrecht, eds.), 61-67, Lecture Notes in Mathematics 1223, Springer, 1986.
- [51] T. Dhirasakdanon, A Model of Infectious Diseases in Amphibian Populations with Ephemeral Habitat, Ph.D. Thesis, Arizona State University, 2010.
- [52] T. Dhirasakdanon and H.R. Thieme, Persistence of vertically transmitted parasite strains which protect against more virulent horizontally transmitted strains, Modeling and Dynamics of Infectious Diseases (Z. Ma, Y. Zhou, J. Wu, eds.), 187–215, World Scientific, 2009.
- [53] T. Dhirasakdanon and H.R. Thieme, Stability of the endemic coexistence equilibrium for one host and two parasites, Mathematical Modelling of Natural Phenomena 5 (2010), 109-138.
- [54] T. Dhirasakdanon, H.R. Thieme, P. van den Driessche, A sharp threshold for disease persistence in host metapopulations, J. Biol. Dynamics 1 (2007), 363-378
- [55] O. Diekmann and Ph. Getto, Boundedness, global existence and continuous dependence for nonlinear dynamical systems describing physiologically structured populations, J. Diff. Equations 215 (2005), 268–319.

- [56] O. Diekmann, M. Gyllenberg, H. Huang, M. Kirkilionis, J.A.J. Metz, H.R. Thieme, On the formulation and analysis of general deterministic structured population models. II. Nonlinear theory, J. Math. Biol. 43 (2001), 157–189.
- [57] O. Diekmann, M. Gyllenberg, J.A.J. Metz, Steady-state analysis of structured population models, Theor. Pop. Biol. 63 (2003), 309–338.
- [58] O. Diekmann, M. Gyllenberg, J.A.J. Metz, H.R. Thieme, On the formulation and analysis of general deterministic structured population models. I. Linear theory, J. Math. Biol. 36 (1998), 349–388.
- [59] O. Diekmann and S.A. van Gils, On the cyclic replicator equation and the dynamics of semelparous populations, SIAM J. Appl. Dyn. Sys. 8 (2009), 1160–1189.
- [60] O. Diekmann, S.A. van Gils, S.M. Verduyn Lunel, H.-O. Walter, *Delay Equations*. Functional, Complex, and Nonlinear Analysis, Springer, 1995.
- [61] J. Dugundji, *Topology*, Allyn and Bacon, Boston 1966, Prenctice Hall of India, New Delhi 1975.
- [62] N. Dunford and J.T. Schwartz, *Linear Operators, Part I: General Theory*, John Wiley & Sons, 1988.
- [63] B. Ebenman, Niche difference between age classes and intraspecific competition in age-structured populations, J. Theor. Biol. 124 (1987), 25–33.
- [64] B. Ebenman, Competition between age-classes and population dynamics, J. Theor. Biol. 131 (1988), 389–400.
- [65] G. Edgar, Measure, Topology, and Fractal Geometry, Springer-Verlag, 1990.
- [66] S.F. Ellermeyer, S.S. Pilyugin, R. Redheffer, Persistence criteria for a chemostat with variable nutrient input, J. Differential Equations 171 (2001), 132–147.
- [67] K.-J. Engel and R. Nagel, One-Parameter Semigroups for Linear Evolution Equations, Springer, 2000.
- [68] A. Fall, A. Iggidr, G. Sallet, J.J. Tewa, *Epidemiological models and Lyapunov functions*, Math. Model. Nat. Phenom. 2 (2007), 62-83.
- [69] Z. Feng and H.R. Thieme, Endemic models with arbitrarily distributed periods of infection. I. General theory, SIAM J. Appl. Math. 61 (2000), 803–833.
- [70] R. Ferriere and M. Gatto, Lyapunov exponents and the mathematics of invasion in oscillatory or chaotic populations, Theor. Population Biol. 48 (1995), 126–171.
- [71] A. Fonda, Uniformly persistent semi-dynamical systems, Proc. Amer. Math. Soc. 104 (1988), 111–116.
- [72] H. Freedman and P. Moson, Persistence definitions and their connections, Proc. Amer. Math. Soc. 109 (1990), 1025–1033.
- [73] H. Freedman, S. Ruan, M. Tang, Uniform persistence and flows near a closed positively invariant set, J. Dyn. Diff. Eqn. 6 (1994), 583-600
- [74] H. Freedman and J.W.-H. So, Persistence in discrete semi-dynamical systems, SIAM J. Math. Anal. 20 (1989), 930–938.
- [75] H. Freedman and P. Waltman, Mathematical analysis of some three species foodchain models, Math. Biosci. 33 (1977), 257–276.
- [76] H. Freedman and P. Waltman, Persistence in models of three interacting predatorprey populations, Math. Biosci. 68 (1984), 213–231.
- [77] L.Q. Gao, J. Mena-Lorca, H.W. Hethcote, Variations on a theme of SEI endemic models, Differential Equations and Applications to Biology and Industry (M. Martelli, C.L. Cooke, E. Cumberbatch, B. Tang, H.R. Thieme, eds.), 191-207, World Scientific, 1996.

- [78] B.M. Garay, Uniform persistence and chain recurrence, J. Math. Anal. Appl. 139 (1989), 372–381.
- [79] B.M. Garay and J. Hofbauer, Robust persistence for ecological differential equations, minimax, and discretizations, SIAM J. Math. Anal. 34 (2003), 1007–1039.
- [80] T.C. Gard, Persistence in food webs: Holling-type food chains, Math. Biosci. 49 (1980), 61–67.
- [81] T.C. Gard, Persistence in food chains with general interactions, Math. Biosci. 51 (1980), 165-174
- [82] T.C. Gard and T.G. Hallam, Persistence in food webs. I. Lotka-Volterra food chains, Bull. Math. Biol. 41 (1979), 877–891.
- [83] B.S. Goh, Global stability in many species systems, Amer. Nat. 111 (1977), 135– 142.
- [84] M. Gobbino and M. Sardella, On the connectedness of attractors for dynamical systems, J. Diff. Eqn. 133, 1-14.
- [85] G. Gripenberg, S.-O. Londen, O. Staffans, Volterra Integral Functional Equations, Cambridge University Press, 1990.
- [86] R.B. Guenther and J.W. Lee, Partial Differential Equations of Mathematical Physics and Integral Equations, Dover Publications Inc., 1988.
- [87] H. Guo and M.Y. Li, Global dynamics of a staged progression model for infectious diseases, Math. Biosci. Engin. 3 (2006), 513–525.
- [88] M. Gyllenberg, F. Jacobs, J. Metz, On the concept of attractor for communitydynamical processes II: the case of structured populations, J. Math. Biol. 47 (2003), 235–248.
- [89] K.P. Hadeler, Selection in a cyclically changing environment, Population Biology and Evolution (K. Wöhrmann, V. Loeschcke, eds.), 102-114, Springer, 1984.
- [90] J.K. Hale, Ordinary Differential Equations, Krieger, 1980.
- [91] J.K. Hale, Asymptotic Behavior of Dissipative Systems, American Mathematical Society, 1988.
- [92] J.K. Hale, *Dissipation and attractors*, Proceedings of the International Conference on Differential Equations (B. Fiedler, K. Groeger, and J. Sprekels, eds.), 622-637, World Scientific, 2000.
- [93] J.K. Hale, Dissipation and compact attractors, J. Dyn. Diff. Eqn. 18 (2006), 485-523.
- [94] J.K. Hale and S.M. Verduyn Lunel, Introduction to Functional Differential Equations, Springer, 1993.
- [95] J.K. Hale and P. Waltman, Persistence in infinite dimensional systems, SIAM J. Math. Anal. 20 (1989), 388–395.
- [96] T.G. Hallam, Structural sensitivity of grazing formulations in nutrient controlled plankton models J. Math. Biol. 5 (1978), 269–280.
- [97] A. Hastings and T. Powell, Chaos in a three-species food chain, Ecology 72(1991), 896–903.
- [98] H.W. Hethcote, The mathematics of infectious diseases, SIAM Review 42 (2000), 599–653.
- [99] E. Hewitt and K. Stromberg, Real and Abstract Analysis, Springer, 1965.
- [100] E. Hille and R.S. Phillips, Functional Analysis and Operator Semigroups, American Mathematical Society, 1957.

- [101] M.W. Hirsch, H.L. Smith, and X.-Q. Zhao, Chain transitivity, attractivity and strong repellors for semidynamical systems, J. Dynamics and Diff. Eqns. 13 (2001), 107–131.
- [102] W.M. Hirsch, H. Hanisch, J.-P. Gabriel, Differential equation models for some parasitic infections: methods for the study of asymptotic behavior, Comm. Pure Appl. Math. 38 (1985), 733–753.
- [103] J. Hofbauer, A general cooperation theorem for hypercycles, Monatshefte f
  ür Math. 91 (1981), 233–240.
- [104] J. Hofbauer, A unified approach to persistence, Acta Applicandae Math. 14 (1989), 11–22.
- [105] J. Hofbauer and S. Schreiber, Robust permanence for structured populations, J. Diff. Eqns. 248 (2010), 1955–1971.
- [106] J. Hofbauer and K. Sigmund, Evolutionary Games and Population Dynamics, Cambridge Univ. Press, 1998.
- [107] J. Hofbauer and J.W.-H. So, Uniform persistence and repellors for maps, Proceedings Amer. Math. Soc. 107 (1989), 1137–1142.
- [108] J. Hofbauer, J. Mallet-Paret, H.L. Smith, Stable periodic solutions for the hypercycle system, Journal of Dynamics and Differential Equations 3 (1991), 423–436.
- [109] V. Hutson, A theorem on average Liapunov functions, Monatshefte f
  ür Math. 98 (1984), 267–275.
- [110] V. Hutson and K. Schmitt, Permanence and the dynamics of biological systems, Math. Biosciences 111 (1992), 293–326.
- [111] M. Iannelli, M. Martcheva, X.-Z. Li, Strain replacement in an epidemic model with super-infection and perfect vaccination, Math. Biosci. 195 (2005), 23–46.
- [112] A. Iggidr, J. Mbang, G. Sallet, Stability analysis of within-host parasite models with delays, Math. Biosci. 209 (2007), 51–75.
- [113] F.J.A. Jacobs and J.A.J. Metz, On the concept of attractor for communitydynamical processes I: the case of unstructured populations, J.Math. Biol. 47 (2003), 222–234.
- [114] A. Katok and B. Hasselblatt, Introduction to the Modern Theory of Dynamical Systems, Cambridge University Press, 1995.
- [115] W.O. Kermack and A.G. McKendrick, A contribution to the mathematical theory of epidemics, Proc. Roy. Soc., A, 115 (1927), 700–721, reprinted in Bull. Math Biol. 53 (1991), 33–55.
- [116] W.O. Kermack and A.G. McKendrick, Contribution to the mathematical theory of epidemics. II. The problem of endemicity, Proc. Roy. Soc., A, 138 (1932), 55–85, Reprinted in Bull. Math Biol. 53 (1991), 57–87.
- [117] W.O. Kermack and A.G. McKendrick, Contribution to the mathematical theory of epidemics. III. Further studies on the problem of endemicity, Proc. Roy. Soc., A, 141 (1933), 94–122, reprinted in Bull. Math Biol. 53 (1991), 89–118.
- [118] J.R. Kirkwood, An Introduction to Analysis, PWS Publishing Company 1995.
- [119] A. Klebanoff and A. Hastings, Chaos in three species food chains, J. Math. Biol. 32 (1994), 427–451.
- [120] R. Kon, Multiple attractors in host-parasitoid interactions: Coexistence and extinction, Math. Biosci. 201 (2006), 172–183.
- [121] R. Kon, Nonexistence of synchronous orbits and class coexistence in matrix population models, SIAM J. Applied Math. 66 (2006), 616–626.

- [122] R. Kon and Y. Iwasa, Single-class orbits in nonlinear Leslie matrix models for semelparous populations, J. Math. Biol. 55 (2007), 781-802.
- [123] R. Kon, Y. Saito, Y. Takeuchi, Permanence of single-species stage structured models, J. Math. Biol. 48 (2004), 515–528.
- [124] R. Kon and Y. Takeuchi, *Permanence of host-parasitoid systems*, Nonlinear Anal., 47 (2001), 1383–1393.
- [125] A. Korobeinikov, Lyapunov functions and global properties for SEIR and SEIS epidemic models, Mathematical Medicine and Biology 21 (2004), 75–83.
- [126] A. Korobeinikov, Lyapunov functions and global stability for SIR and SIRS epidemiological models with non-linear transmission, Bull. Math. Biol. 30 (2006), 615–626.
- [127] A. Korobeinikov and G.C. Wake, Lyapunov functions and global stability for SIR, SIRS and SIS epidemiological models, Appl. Math. Letters 15 (2002), 955–961.
- [128] M.A. Krasnosel'skii, Positive Solutions of Operator Equations, Noordhoff, 1964.
- [129] U. Krause and R.D. Nussbaum, A limit set trichotomy for self-mappings of normal cones in Banach spaces, Nonlinear Analysis, TMA 20 (1993), 855–1993.
- [130] C. Kribs-Zaleta and J. Velasco-Hernández, A simple vaccination model with multiple endemic states, Math. Biosci. 164 (2000), 183–201.
- [131] Y. Kuznetsov, O. Feo, S. Rinaldi, Belyakov Homoclinic bifurcations in a tri-trophic food chain model, SIAM J. Appl. Math. 62 (2001), 462–487.
- [132] Y. Kuang, Global stability of Gause-type predator-prey systems, J. Math. Biol. 28 (1990), 463–474.
- [133] Y. Kuang and H. Freedman, Uniqueness of limit cycles in Gause-type predator-prey systems, Math. Biosci. 88 (1988), 67–84.
- [134] O. Ladyzhenskaya, Attractors for Semigroups and Evolution Equations, Cambridge University Press, 1991.
- [135] V. Lakshmikantham and S. Leela, Nonlinear Differential Equations in Abstract Spaces, Pergamon Press, 1981.
- [136] P. Lancaster and M. Tismenetsky, *The Theory of Matrices*, Academic Press, 1985.
- [137] J.P. LaSalle, The Stability of Dynamical Systems, SIAM, 1976.
- [138] D. Le and H.L. Smith, A parabolic system modeling microbial competition in an unmixed bio-reactor, Journal of Differential Equations 130 (1996), 59–91.
- [139] P. De Leenheer and H.L. Smith, Virus Dynamics: a global analysis, SIAM J. Appl. Math. 63(2003), 1313–1327.
- [140] R. Lenski and B. Levin, Constraints on the evolution of bacteria and virulent phage: a model, some experiments, and predictions for natural communities, Amer. Naturalist 125 (1985), 585–602.
- [141] P.H. Leslie, The use of matrices in certain population mathematics, Biometrika, 33 (1945), 183-212.
- [142] C.-K. Li and H. Schneider, Applications of Perron-Frobenius theory to population dynamics, J. Math. Biol. 44 (2002), 450–462.
- [143] J. Li, Simple stage-structured models for wild and transgenic mosquito populations, J. Difference Eqn. Appl. 15 (2009), 327-347.
- [144] M.Y. Li and J.S. Muldowney, Global stability for the SEIR model in epidemiology, Math. Biosci. 125 (1995), 155–164.

- [145] M.Y. Li, Z. Shuai, Global stability problem for coupled systems of differential equations on networks, J. Diff. Eqn. 248 (2010), 1-20.
- [146] M.Y. Li, Z. Shuai, C. Wang, Global stability of multi-group epidemic models with distibuted delays, J. Math. Anal. Appl. 361 (2010), 38-47.
- [147] G.M. Lieberman, Second Order Parabolic Differential Equations, World Scientific, 1996.
- [148] L.-P. Liou and K.-S. Cheng, On the uniqueness of a limit cycle for a predator-prey system, SIAM J. Math. Anal. 19 (1988), 867–878.
- [149] S. Liu, L. Wang, Global stability of an HIV-1 model with distributed intracellular delays and a combination therapy, Math. Biosci. Eng. 7 (2010), 675-685.
- [150] S.-O. Londen, On the variation of the solution of a nonlinear integral equation, J. Math. Anal. Appl. 52 (1975), 430–449.
- [151] S.-O. Londen, On some integral equations with locally finite measures and L<sup>∞</sup>perturbations, SIAM J. Math. Anal. 14 (1983), 1187–1203.
- [152] A.L. Lotka, Relation between birth rates and death rates, Science 26 (1907), 21–22.
- [153] A. Lunardi, Analytic Semigroups and Optimal Regularity in Parabolic Problems, Progress in Nonlinear Differential Equations and Their Applications 16, Birkhäuser, 1995.
- [154] P. Magal, Perturbation of a globally stable steady state and uniform persistence, J. Dyn. Diff. Eqn. 21 (2009), 1-20
- [155] P. Magal, C.C. McCluskey, G.F. Webb, Liapunov functional and global asymptotic stability for an infection-age model, Applicable Analysis, 89 (2010), 1109–1140.
- [156] P. Magal and S. Ruan, On integrated semigroups and age structured models in L<sup>p</sup> spaces, Diff. Integr. Eqn. 20 (2007), 197–239.
- [157] P. Magal and S. Ruan, Center manifolds for semilinear equations with non-dense domain and applications to Hopf bifurcation in age structured models, Memoirs of the AMS vol. 202, (2009), no. 951.
- [158] P. Magal and X.-Q. Zhao, Global attractors and steady states for uniformly persistent dynamical systems, SIAM J. Math. Anal. 37 (2005), 251–275.
- [159] R.H. Martin, Nonlinear Operators and Differential Equations in Banach Spaces, Wiley and Sons, 1976.
- [160] R.H. Martin Jr. and H.L. Smith, Abstract functional-differential equations and reaction-diffusion systems, Trans. Amer. Math. Soc. 321 (1990), 1–44.
- [161] M. Martcheva and H.R. Thieme, Progression age enhanced backward bifurcation in an epidemic model with super-infection, J. Math. Biol. 46 (2003), 385–424.
- [162] R.M. May and W.J. Leonard, Nonlinear aspects of competition between three species, SIAM J. Appl. Math. 29 (1975), 243–253.
- [163] C.C. McCluskey, Global stability for an SEIR epidemiological model with varying infectivity and infinite delay, Math. Biosci. and Eng. 6(2009) 603–610.
- [164] J.N. McDonald and N.A. Weiss, A Course in Real Analysis, Academic Press, 1999.
- [165] A.G. McKendrick, Applications of mathematics to medical problems, Proc. Edinb. Math. Soc. 14 (1926), 98–130.
- [166] R.C. McOwen, Partial Differential Equations, methods and applications, Prentice-Hall, 2003.
- [167] J.A.J. Metz, (2008) Fitness, Evol. Ecol. 2 (2008), 1599–1612.

- [168] J.A.J. Metz, R.M. Nisbet, S.A.H. Geritz, How should we define "fitness" for general ecological scenarios?, Tree 7 (1992), 198–202.
- [169] J. Mierczynski, W. Shen, X.-Q. Zhao, Uniform persistence for nonautonomous and random parabolic Kolmogorov systems, J. Diff. Equations, 204 (2004), 471–510.
- [170] K. Mischaikow, H.L. Smith, and H.R. Thieme, Asymptotically autonomous semiflows: Chain recurrence and Lyapunov functions, Trans. Amer. Math. Soc. 347 (1995), 1669–1685.
- [171] S. Muratori and S. Rinaldi, Low- and high-frequency oscillations in threedimensional food chains, SIAM J. Appl. Math. 52 (1992) 1688–1706.
- [172] M.A. Nowak and R.M. May, Virus Dynamics, Oxford Univ. Press, 2000.
- [173] L. Oksanen, S.D. Fretwell, J. Arruda, P. Niemela, Exploitation ecosystems in gradients of primary productivity, American Naturalist 118 (1981), 240–261.
- [174] C. V. Pao, Nonlinear Parabolic and Elliptic Equations, Plenum Press, 1992.
- [175] A. Pazy, Semigroups of Linear Operators and Applications to Partial Differential Equation, Springer, 1983.
- [176] A.S. Perelson and P.W. Nelson, Mathematical analysis of HIV-1 dynamics in vivo, SIAM Review 41 (1999), 3–44.
- [177] L. Perko, Differential Equations and Dynamical Systems, Springer-Verlag, (1991).
- [178] M. Protter and H.F. Weinberger, Maximum Principles in Differential Equations, Prentice Hall, 1967.
- [179] J. Prüss, L. Pujo-Menjouet, G.F. Webb, Analysis of a model for the dynamics of prions, Discrete Contin. Dyn. Syst. 6 (2006), 225–235.
- [180] D.A. Rand, H.B. Wilson, J.M. McGlade, Dynamics and evolution: evolutionarily stable attractors, invasion exponents and phenotype dynamics, Philosophical Transactions: Biological Sciences 343 (1994), 261–283.
- [181] G. Raugel, Global attractors in partial differential equations, Handbook of Dynamical Systems 2, 885–982, North-Holland, 2002.
- [182] M.G. Roberts and J.A.P. Heesterbeek, A simple parasite model with complicated dynamics, J. Math. Biol. 37 (1998), 272–290.
- [183] C. Robinson, Dynamical Systems, Stability, Symbolic Dynamics, and Chaos, 2nd ed., CRC Press, 1999.
- [184] G. Röst and J. Wu, SEIR epidemiological model with varying infectivity and infinite delay, Math. Biosci. Engin. 5 (2008), 389–401.
- [185] M. Salmani and P. van den Driessche, A model for disease transmission in a patchy environment, Disc. Cont. Dyn. Sys. -B 6 (2006), 185–202.
- [186] P.L. Salceanu, Lyapunov Exponents and Persistence in Dynamical Systems, Ph.D. Thesis, Arizona State University, 2009.
- [187] P.L. Salceanu and H.L. Smith, Persistence in a discrete-time, stage-structured epidemic model, J. Difference Equ. & Appl. 16 (2010), 73–103.
- [188] P.L. Salceanu and H.L. Smith, Persistence in a discrete-time stage-structured fungal disease model, J. Biol. Dynamics 3 (2009), 271–285.
- [189] P.L. Salceanu and H.L. Smith, Lyapunov exponents and persistence in discrete dynamical systems, Discrete & Continuous Dynamical Systems-B 12 (2009), 187– 203.
- [190] P.L. Salceanu and H.L. Smith, Lyapunov exponents and uniformly weak normally repelling invariant sets, Positive Systems (R. Bru and S. Romero-Vivo, eds.), Lecture Notes in Control and Information Sciences 389, Springer, 2009.

- [191] S.J. Schreiber, Persistence despite perturbations for interacting populations, J. of Theor. Biol. 242 (2006), 844–852.
- [192] S.J. Schreiber, On growth rates of subadditive functions for semiflows, J. Differential Equations 148 (1998), 334–350.
- [193] S.J. Schreiber, Criteria for C<sup>r</sup> robust permanence, J. Differential Equations 162 (2000), 400–426.
- [194] S.J. Schreiber, On persistence and extinction for randomly perturbed dynamical systems, Discrete Contin. Dyn. Syst. Ser. B 7 (2007), 457–463.
- [195] J. Schröder, Anwendung von Fixpunktsätzen bei der numerischen Behandlung nichtlinearer Gleichungen in halbgeordneten Räumen, Arch. Rat. Mech. Analysis 4 (1959/60), 177–192.
- [196] P. Schuster, K. Sigmund, R. Wolff, Dynamical systems under constant organization. III. Cooperative and competitive behavior of hypercycles, J. Diff. Eqns. 32 (1979), 357–368.
- [197] G.R. Sell and Y. You, Dynamics of Evolutionary Equations, Springer, 2002.
- [198] L.A. Segel, Modeling dynamic phenomena in molecular and cellular biology, Cambridge Univ. Press, 1987.
- [199] A. Singh and R.M. Nisbet, Semi-discrete host-parasitoid models, J. Theor. Biol. 247 (2007), 733–742.
- [200] H.L. Smith, Monotone Dynamical Systems: an introduction to the theory of competitive and cooperative systems, Amer. Math. Soc. Surveys and Monographs, vol. 41, 1995.
- [201] H.L. Smith, The discrete dynamics of monotonically decomposable maps, J. Math. Biol. 53 (2006), 747–758.
- [202] H.L Smith, Models of virulent phage growth with application to phage therapy, SIAM J. Appl. Math. 68 (2008), 1717–1737.
- [203] H.L. Smith and P. Waltman, The Theory of the Chemostat, Cambridge Studies in Mathematical Biology 13, Cambridge University Press 1995.
- [204] H.L. Smith and P. Waltman, Perturbation of a globally stable steady state, Proc. Amer. Math. Soc. 127 (1999), 447–453.
- [205] H.L. Smith and H. R. Thieme, *Persistence of bacteria and phages in a chemostat*, preprint.
- [206] H.L. Smith and X.-Q. Zhao, Robust persistence for semidynamical systems, Nonlinear Analysis 47 (2001), 6169–6179.
- [207] J. Smoller, Shock Waves and Reaction Diffusion Equations, Springer-Verlag, 1983.
- [208] H.W. Stech and M. Williams, Stability in a class of cyclic epidemic models with delay, J. Math. Biol. 11 (1981), 95–103.
- [209] P. Takáč, Convergence in the part metric for discrete dynamical systems in ordered topological cones, Nonlinear Analysis, TMA, 26 (1996), 1753–1777.
- [210] R. Temam, Infinite Dimensional Dynamical Systems in Mechanics and Physics, Springer, 1988
- [211] H.R. Thieme, Density-dependent regulation of spatially distributed populations and their asymptotic speed of spread, J. Math. Biology 8 (1979), 173–187.
- [212] H.R. Thieme, On a class of Hammerstein integral equations, Manuscripta mathematica 29 (1979), 49–84.

- [213] H.R. Thieme, Well-posedness of physiologically structured population models for Daphnia magna (How biological concepts can benefit by abstract mathematical analysis), J. Math. Biology 26 (1988), 299–317.
- [214] H.R. Thieme, Semiflows generated by Lipschitz perturbations of non-densely defined operators, Differential and Integral Equations 3 (1990), 1035–1066.
- [215] H.R. Thieme, Persistence under relaxed point-dissipativity (with applications to an endemic model), SIAM J. Math. Anal. 24 (1993), 407–435.
- [216] H.R. Thieme, Uniform persistence and permanence for non-autonomous semiflows in population biology, Math. Biosci. 166 (2000), 173–201.
- [217] H.R. Thieme, *Mathematics in Population Biology*, Princeton University Press, 2003.
- [218] H.R. Thieme, Pathogen competition and coexistence and the evolution of virulence, Mathematics for Life Sciences and Medicine, (Y. Takeuchi, Y. Iwasa, K. Sato, eds.), 123-153, Springer, 2007.
- [219] H.R. Thieme, Differentiability of convolutions, integrated semigroups of bounded semi-variation, and the non-homogeneous Cauchy problem, J. Evolution Equations 8 (2008), 283–305.
- [220] H.R. Thieme, Spectral bound and reproduction number for infinite dimensional population structure and time-heterogeneity, SIAM J. Appl. Math. 70 (2009), 188– 211.
- [221] H.R. Thieme and C. Castillo-Chavez, How may infection-age-dependent infectivity affect the dynamics of HIV/AIDS?, SIAM J. Appl. Math. 53 (1993), 1447–1479.
- [222] H.R. Thieme, A. Tridane, Y. Kuang, An epidemic model with post-contact prophylaxis of distributed length. I. Thresholds for disease persistence and extinction, J. Biological Dynamics 2 (2008), 221–239.
- [223] H.R. Thieme, A. Tridane, Y. Kuang, An epidemic model with post-contact prophylaxis of distributed length. II. Stability and oscillations if treatment is fully effective, Mathematical Modelling of Natural Phenomena, 3 (2008), 267-293
- [224] H.R. Thieme and P. van den Driessche, Global stability in cyclic epidemic models with disease fatalities (Proceedings of the conference on Differential Equations with Applications to Biology), Fields Inst. Comm. 21 (1999), 459–472.
- [225] H. R. Thieme and H. Voßeler, Semilinear perturbations of Hille Yosida operators. Mathematical Modelling of Population Dynamics (R. Rudnicki, ed.), 87-122. Banach Center Publ. 63, Polish Academy of Sciences, 2004.
- [226] H.R. Thieme and I.I. Vrabie, Relatively compact orbits and compact attractors for a class of nonlinear evolution equations, J. Dyn. Diff. Equ. 15 (2003), 731–750.
- [227] H.R. Thieme and J. Yang, An endemic model with variable re-infection rate and application to influenza, Math. Biosci. 180 (2002), 207–235.
- [228] A.C. Thompson, On certain contraction mappings in a partially ordered space, Proc. Amer. Math. Soc. 14 (1963), 438–443.
- [229] H. Thunberg, Periodicity versus chaos in one-dimensional dynamics, SIAM Review 43 (2001), 3–30.
- [230] P. Turchin, Complex Population Dynamics, Monographs in Population Biology, Princeton University Press, 2003.
- [231] P. Waltman, A brief survey of persistence in dynamical systems, Delay Differential Equations and Dynamical Systems (S. Busenberg, M. Martelli, eds.), 31–40, Lecture Notes in Math., 1475, Springer, 1991.

- [232] P. Waltman, A Second Course in Elementary Differential Equations, Dover Publications, 2004.
- [233] G.F. Webb, Theory of Nonlinear Age-dependent Population Dynamics, Marcel Dekker, 1985.
- [234] L. Xu, B. Chen, Y. Zhao, and Y. Cao, Normal Lyapunov exponents and asymptotically stable attractors, Dynamical Systems 23 (2008), 207–218.
- [235] F. Yang and S. Ruan, A generalization of the Butler-McGehee lemma and its applications in persistence theory, Diff. Int. Eqn. 9 (1996), 1321-1330
- [236] K. Yosida, Functional Analysis, second ed., Springer, 1968.
- [237] M.L. Zeeman, Hopf bifurcations in competitive three-dimensional Lotka-Volterra systems, Dynamics and Stability of Systems 8 (1993), 189–217.
- [238] X.-Q. Zhao, Dynamical Systems in Population Biology, CMS Books in Mathematics, Springer, 2003.
- [239] Y. Zhou and J.M. Cushing, The net reproductive value and stability in matrix population models, Nat. Res. Mod. 8 (1994), 297–333.

### Index

absolutely continuous, 54, 233, 251, 256, 298, 317 definition, 364 accumulation point, 12acyclic covering, 187 attract a neighborhood of itself, 42, 177 attractor, 30 average Lyapunov function strong, 354 weak, 350 Banach space, 16 bounded set, 13 Butler-McGehee lemma, 182 Cauchy sequence, 16 chain transitive set, 181 chemostat, 142 closed operator, 385 closed set, 12, 25 closure, 12  $\operatorname{compact}$ map, 58set, 11 compact attractor, 30 connected, 38 dual attractor, 137 stable, 137 extinction attractor, 129 global, 37 local, of compact sets, 38 stable, 47 of a class of sets, 36 of a set, 32, 39 of an open set, 40

of bounded sets, 37, 43, 147 of compact sets, 37, 38 of neighborhoods of compact sets, 37, 41 of points, 37, 40 persistence attractor, 129 upper semicontinuity of, 184 completeness, 16 condensing map, 158 cone, 273 connected set, 33 connected space, 33 convergence, 10 convergence of sets, 29 cooperative system, 370 decreasing, 53 diameter, 13 dissipative  $\rho$ -dissipative, 61 weakly  $\rho$ -dissipative, 62 distance from point to set, 11 from set to set, 14 dual attractor, 137 dual operator, 386 dual relation, 386 equilibrium, 23, 58 disease-free, 68 endemic, 45 fixed point, 23 flow, 18 flow-connected, 34 fluctuation method, 149, 366

increasing, 53 interior of a set, 13 interior point, 13 invariant, 19, 20, 30 backward, 19 forward, 19 isolated, 136 isolating neighborhood, 136 Laplace transform, 150, 311 truncated, 153 Leslie model, 72 limit, 10 limit inferior, 365 limit point, 12 limit set  $\alpha$ , 52  $\omega$ , 30 limit superior, 365 Lipschitz, 263 locally attracting, 177 LPA model, 72 Lyapunov exponent, 219 Lyapunov function, 53 matrix irreducible, 372 positive, 372 primitive, 172, 372 quasipositive, 65, 372 spectral bound of, 372 matrix cocycle, 219 matrix model, 71 metric, 9 Hausdorff, 15, 183 semimetric, 9 space, 9 minimal set, 19 Morse decomposition, 190 neighborhood, 13  $\epsilon$ -neighborhood of a set, 14 norm, 13 seminorm, 13 open ball, 12 open set, 13, 25 operator, 385 bounded linear, 385 closed, 385 linear, 385 permanent  $\rho$ -permanent, 62 persistence practical, x, 327 strong  $\rho$ -persistence, 61 uniform  $\rho$ -persistence, 61, 333

uniform weak  $\rho$ -persistence, 61, 333 weak  $\rho$ -persistence, 61 precompact set, 12 quasimonotone condition, 368 repeller associated with locally attracting set, 178 uniform, 136 uniform weak, 136 repeller neighborhood, 136 repelling uniformly weakly, 187, 189 weakly, 187, 188 reproduction number critical, 144 of bacteria in tubular bioreactor, 299 of infectious disease, 211, 237 of Leslie model, 74 of LPA model, 73 of phage, 144 semiflow, 17 asymptotically compact, 31 asymptotically periodic, 335 asymptotically smooth, 39 continuous, 17 discrete, 18 eventually bounded on a set M, 39 eventually uniformly  $\rho$ -positive on a set, 127 injective, 18 limit semiflow, 336 Lipschitz semiflow, 48 nonautonomous, 332 periodic, 335 point dissipative, 39 state-continuous, 17 state-continuous, uniformly in finite time, 17 time-continuous, 17 semigroup  $C_0$ -semigroup, 262 generator of, 262 solution classical, 261, 387 integral, 261, 387 mild, 263, 388 weak, 387 stable, 47 attractor, 4 dual attractor, 137 equilibrium, 152 forward invariant set, 47 heteroclinic orbit, 140 locally asymptotically, 4, 47 definition, 47 locally asymptotically relative to a set, 189 persistence attractor, 129

stable relative to a set, 189 subtangential condition, 20, 274 synchronous orbit, 171

total orbit, 20 periodic, 23 total trajectory, 20, 107, 272 periodic, 23 totally bounded set, 16 triangle inequality, 9

uniformly  $\rho\text{-positive set, }127$ 

The mathematical theory of persistence answers questions such as which species, in a mathematical model of interacting species, will survive over the long term. It applies to infinite-dimensional as well as to finite-dimensional dynamical systems, and to discrete-time as well as to continuous-time semiflows.

This monograph provides a self-contained treatment of persistence theory that is acces-

sible to graduate students. The key results for deterministic autonomous systems are proved in full detail such as the acyclicity theorem and the tripartition of a global compact attractor. Suitable conditions are given for persistence to imply strong persistence even for nonautonomous semiflows, and time-heterogeneous persistence results are developed using so-called "average Lyapunov functions".

Applications play a large role in the monograph from the beginning. These include ODE models such as an SEIRS infectious disease in a meta-population and discrete-time nonlinear matrix models of demographic dynamics. Entire chapters are devoted to infinite-dimensional examples including an SI epidemic model with variable infectivity, microbial growth in a tubular bioreactor, and an age-structured model of cells growing in a chemostat.





For additional information and updates on this book, visit www.ams.org/bookpages/gsm-118

