


# An Introduction to Extremal Kähler Metrics

**Gábor Székelyhidi**

**Graduate Studies  
in Mathematics**

**Volume 152**



**American Mathematical Society**

# An Introduction to Extremal Kähler Metrics



# An Introduction to Extremal Kähler Metrics

Gábor Székelyhidi

Graduate Studies  
in Mathematics

Volume 152



American Mathematical Society  
Providence, Rhode Island

## EDITORIAL COMMITTEE

Dan Abramovich  
Daniel S. Freed  
Rafe Mazzeo (Chair)  
Gigliola Staffilani

2010 *Mathematics Subject Classification*. Primary 53C25, 53C55, 14L24.

---

For additional information and updates on this book, visit  
[www.ams.org/bookpages/gsm-152](http://www.ams.org/bookpages/gsm-152)

---

### Library of Congress Cataloging-in-Publication Data

Székelyhidi, Gábor, 1981–

An introduction to extremal Kähler metrics / Gábor Székelyhidi.  
pages cm. — (Graduate studies in mathematics ; volume 152)

Includes bibliographical references and index.

ISBN 978-1-4704-1047-6 (alk. paper)

1. Geometry, Differential. I. Title.

QA641.S94 2014  
516.3'73—dc23

2014006619

---

**Copying and reprinting.** Individual readers of this publication, and nonprofit libraries acting for them, are permitted to make fair use of the material, such as to copy a chapter for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgment of the source is given.

Republication, systematic copying, or multiple reproduction of any material in this publication is permitted only under license from the American Mathematical Society. Requests for such permission should be addressed to the Acquisitions Department, American Mathematical Society, 201 Charles Street, Providence, Rhode Island 02904-2294 USA. Requests can also be made by e-mail to [reprint-permission@ams.org](mailto:reprint-permission@ams.org).

© 2014 by the American Mathematical Society. All rights reserved.  
The American Mathematical Society retains all rights  
except those granted to the United States Government.  
Printed in the United States of America.

∞ The paper used in this book is acid-free and falls within the guidelines  
established to ensure permanence and durability.  
Visit the AMS home page at <http://www.ams.org/>

10 9 8 7 6 5 4 3 2 1      19 18 17 16 15 14

To Sonja and Nóra



---

# Contents

Preface	xi
Introduction	xiii
Chapter 1. Kähler Geometry	1
§1.1. Complex manifolds	1
§1.2. Almost complex structures	4
§1.3. Hermitian and Kähler metrics	6
§1.4. Covariant derivatives and curvature	10
§1.5. Vector bundles	13
§1.6. Connections and curvature of line bundles	16
§1.7. Line bundles and projective embeddings	19
Chapter 2. Analytic Preliminaries	23
§2.1. Harmonic functions on $\mathbf{R}^n$	23
§2.2. Elliptic differential operators	25
§2.3. Schauder estimates	26
§2.4. The Laplace operator on Kähler manifolds	32
Chapter 3. Kähler-Einstein Metrics	35
§3.1. The strategy	36
§3.2. The $C^0$ - and $C^2$ -estimates	40
§3.3. The $C^3$ - and higher-order estimates	44
§3.4. The case $c_1(M) = 0$	47



---

§3.5. The case $c_1(M) > 0$	51
§3.6. Futher reading	53
Chapter 4. Extremal Metrics	57
§4.1. The Calabi functional	57
§4.2. Holomorphic vector fields and the Futaki invariant	62
§4.3. The Mabuchi functional and geodesics	67
§4.4. Extremal metrics on a ruled surface	71
§4.5. Toric manifolds	76
Chapter 5. Moment Maps and Geometric Invariant Theory	85
§5.1. Moment maps	85
§5.2. Geometric invariant theory (GIT)	90
§5.3. The Hilbert-Mumford criterion	94
§5.4. The Kempf-Ness theorem	97
§5.5. Relative stability	101
Chapter 6. K-stability	105
§6.1. The scalar curvature as a moment map	105
§6.2. The Hilbert polynomial and flat limits	109
§6.3. Test-configurations and K-stability	111
§6.4. Automorphisms and relative K-stability	115
§6.5. Relative K-stability of a ruled surface	116
§6.6. Filtrations	120
§6.7. Toric varieties	124
Chapter 7. The Bergman Kernel	129
§7.1. The Bergman kernel	129
§7.2. Proof of the asymptotic expansion	132
§7.3. The equivariant Bergman kernel	138
§7.4. The algebraic and geometric Futaki invariants	140
§7.5. Lower bounds on the Calabi functional	142
§7.6. The partial $C^0$ -estimate	150
Chapter 8. CscK Metrics on Blow-ups	153
§8.1. The basic strategy	153
§8.2. Analysis in weighted spaces	159
§8.3. Solving the non-linear equation when $n > 2$	172

---

§8.4. The case when $n = 2$	174
§8.5. The case when $M$ admits holomorphic vector fields	178
§8.6. K-stability of cscK manifolds	181
Bibliography	185
Index	191



---

# Preface

This book grew out of lecture notes written for a graduate topics course taught at the University of Notre Dame in the spring of 2012. The goal is to quickly introduce graduate students to ideas surrounding recent developments on extremal Kähler metrics. We make an effort to introduce the main ideas from Kähler geometry and analysis that are required, but the parts of the book on geometric invariant theory and K-stability would be difficult to follow without more background in complex algebraic geometry. A reader with a background in Riemannian geometry and graduate level analysis should be able to follow the rest of the book.

I would like to thank Tamás Darvas, Yueh-Ju Lin, and Valentino Tosatti for helpful comments about the manuscript.

Gábor Székelyhidi  
Notre Dame, 2014



---

# Introduction

A basic problem in differential geometry is to find canonical, or best, metrics on a given manifold. There are many different incarnations of this, perhaps the most well known being the classical uniformization theorem for Riemann surfaces. The study of extremal metrics is an attempt at finding a higher-dimensional generalization of this result in the setting of Kähler geometry. Extremal metrics were introduced by Calabi in the 1980s as an attempt to find canonical Kähler metrics on Kähler manifolds as critical points of a natural energy functional. The energy functional is simply the  $L^2$ -norm of the curvature of a metric. The most important examples of extremal metrics are Kähler-Einstein metrics and constant scalar curvature Kähler (or cscK) metrics.

It turns out that extremal metrics do not always exist, and the question of their existence is particularly interesting on projective manifolds. In this case, by works of Yau, Tian, and Donaldson, it was realized that the existence of extremal metrics is related to the stability of the manifold in an algebro-geometric sense, and obtaining a necessary and sufficient condition of this form for existence is the central problem in the field. Our goal in this book is to introduce the reader to some of the basic ideas on both the analytic and the algebraic sides of this problem. One concrete goal is to give a fairly complete proof of the following result.

**Theorem.** *If  $M$  admits a cscK metric in  $c_1(L)$  for an ample line bundle  $L \rightarrow M$  and if  $M$  has no non-trivial holomorphic vector fields, then the pair  $(M, L)$  is K-stable.*

The converse of this result, i.e. the existence of cscK metrics on K-stable manifolds, is the central conjecture in the field.

We will now give a brief description of the contents of the book. The first two chapters give a quick review of some of the background material that is needed. The first chapter contains the basic definitions in Kähler geometry, with a focus on calculations in local coordinates. The second chapter focuses on some of the analytic background required, in particular the Schauder estimates for elliptic operators, which we prove using a blow-up argument due to L. Simon.

The topic of Chapter 3 is Kähler-Einstein metrics, which are a special case of extremal metrics. We give a proof of Yau's celebrated theorem on the solution of the complex Monge-Ampère equation, leading to existence results for Kähler-Einstein metrics with zero or negative Ricci curvature. The case of positive Ricci curvature has only been understood very recently through the work of Chen, Donaldson, and Sun. The details of this are beyond the scope of this book, and we only give a very brief discussion in Section 3.5.

The study of general extremal metrics begins in Chapter 4. Following Calabi, we introduce extremal metrics as critical points of the Calabi functional, which is the  $L^2$ -norm of the scalar curvature:

$$\omega \mapsto \int_M S(\omega)^2 \omega^n,$$

defined for metrics  $\omega$  in a fixed Kähler class. An important discovery is that extremal metrics have an alternative variational characterization, as critical points of the (modified) Mabuchi functional. This is convex along geodesics in the space of Kähler metrics with respect to a natural, infinite-dimensional, Riemannian structure. Moreover the variation of the Mabuchi functional is closely related to the Futaki invariant, which plays a prominent role in the definition of K-stability. After giving the basic definitions, we construct an explicit family of extremal metrics on a ruled surface due to Tønnesen-Friedman in Section 4.4. This example illustrates how a sequence of extremal metrics can degenerate, and we return to it again in Section 6.5. In Section 4.5 we give an introduction to the study of extremal metrics on toric manifolds. Toric manifolds provide a very useful setting in which to study extremal metrics and stability, and while in the two-dimensional case the basic existence question is understood through the works of Donaldson and of Chen, Li, and Sheng, the higher-dimensional case remains an important problem to study.

In Chapter 5 we give an introduction to the relation between symplectic and algebraic quotients—the Kempf-Ness theorem—which, at least on a heuristic level, underpins many of the ideas that have to do with extremal metrics. The general setting is a compact group  $K$  acting by Hamiltonian isometries on a Kähler manifold  $M$ , with a moment map  $\mu : M \rightarrow \mathfrak{k}^*$ . The

Kempf-Ness theorem characterizes those orbits of the complexified group  $K^c$  which contain zeros of the moment map. The reason why this is relevant is that the scalar curvature of a Kähler metric, or rather the map  $\omega \mapsto S(\omega) - \hat{S}$  where  $\hat{S}$  is the average scalar curvature, can be realized as a moment map for a suitable infinite-dimensional Hamiltonian action. At the same time, orbits of  $K^c$  can be thought of as metrics in a given Kähler class, so an infinite-dimensional analog of the Kempf-Ness theorem would describe Kähler classes that contain cscK metrics. In Section 5.5 we will describe a suitable extension of the Kempf-Ness theorem dealing with critical points of the norm squared of a moment map, which in the infinite-dimensional setting are simply extremal metrics.

The notion of K-stability is studied in Chapter 6. It is defined in analogy with the Hilbert-Mumford criterion in geometric invariant theory by requiring that a certain weight—the Donaldson-Futaki invariant—is positive for all  $\mathbf{C}^*$ -equivariant degenerations of the manifold. These degenerations are called test-configurations. In analogy with the finite-dimensional setting of the Kempf-Ness theorem, the Donaldson-Futaki invariant of a test-configuration can be seen as an attempt at encoding the asymptotics of the Mabuchi functional “at infinity”, with the positivity of the weights ensuring that the functional is proper. In Section 6.6 we will describe test-configurations from the point of view of filtrations of the homogeneous coordinate ring of the manifold. It is likely that the notion of K-stability needs to be strengthened to ensure the existence of a cscK metric, and filtrations allow for a natural way to enlarge the class of degenerations that we consider. In the case of toric varieties, passing from test-configurations to filtrations amounts to passing from rational piecewise linear convex functions to all continuous convex functions, as we will discuss in Section 6.7.

The basic tool in relating the differential geometric and algebraic aspects of the problem is the Bergman kernel, which we discuss in Chapter 7. We first give a proof of a simple version of the asymptotic expansion of the Bergman kernel going back to Tian, based on the idea of constructing peaked sections of a sufficiently high power of a positive line bundle. Then, following Donaldson, we use this to show that a projective manifold which admits a cscK metric must be K-semistable. This is a weaker statement than the theorem stated above. The Bergman kernel also plays a key role in the recent developments on Kähler-Einstein metrics, through the partial  $C^0$ -estimate conjectured by Tian. We will discuss this briefly in Section 7.6.

In the final chapter, Chapter 8, the main result is a perturbative existence result for cscK metrics due to Arezzo and Pacard. Starting with a cscK metric  $\omega$  on  $M$  and assuming that  $M$  has no non-zero holomorphic vector fields, we show that the blow-up of  $M$  at any point admits cscK metrics in



suitable Kähler classes. The gluing technique used together with analysis in weighted Hölder spaces has many applications in geometric analysis. Apart from giving many new examples of cscK manifolds, this existence result is crucial in the final step of proving the theorem stated above, namely to improve the conclusion from K-semistability (obtained in Chapter 7) to K-stability. The idea due to Stoppa is to show that if  $M$  admits a cscK metric and is not K-stable, then a suitable blow-up of  $M$  is not even K-semistable. Since the blow-up admits a cscK metric, this is a contradiction.

There are several important topics that are missing from this book. We make almost no mention of parabolic equations such as the Calabi flow and the Kähler-Ricci flow. We also do not discuss in detail the existence theory for constant scalar curvature metrics on toric surfaces and for Kähler-Einstein metrics on Fano manifolds since each of these topics could take up an entire book. It is our hope that after studying this book the reader will be eager and ready to tackle these more advanced topics.

---

# Bibliography

1. M. Abreu, *Kähler geometry of toric varieties and extremal metrics*, Internat. J. Math. **9** (1998), 641–651.
2. V. Apostolov, D. M. J. Calderbank, P. Gauduchon, and C. W. Tønnesen-Friedman, *Hamiltonian 2-forms in Kähler geometry III, extremal metrics and stability*, Invent. Math. **173** (2008), no. 3, 547–601.
3. C. Arezzo and F. Pacard, *Blowing up and desingularizing constant scalar curvature Kähler manifolds*, Acta Math. **196** (2006), no. 2, 179–228.
4. ———, *Blowing up Kähler manifolds with constant scalar curvature II*, Ann. of Math. (2) **170** (2009), no. 2, 685–738.
5. C. Arezzo, F. Pacard, and M. A. Singer, *Extremal metrics on blow ups*, Duke Math. J. **157** (2011), no. 1, 1–51.
6. M. F. Atiyah, *Convexity and commuting Hamiltonians*, Bull. London Math. Soc. **14** (1982), no. 1, 1–15.
7. T. Aubin, *Métriques riemanniennes et courbure*, J. Differential Geom. **4** (1970), 383–424.
8. ———, *Équations du type Monge-Ampère sur les variétés kählériennes compactes*, Bull. Sci. Math. (2) **102** (1978), no. 1, 63–95.
9. ———, *Réduction de cas positif de l'équation de Monge-Ampère sur les variétés kählériennes compactes à la démonstration d'une inégalité*, J. Funct. Anal. **57** (1984), no. 2, 143–153.
10. ———, *Some nonlinear problems in Riemannian geometry*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 1998.
11. S. Bando and T. Mabuchi, *Uniqueness of Einstein Kähler metrics modulo connected group actions*, Algebraic geometry, Sendai, Adv. Stud. Pure Math., vol. 10, 1985, pp. 11–40.
12. R. Bartnik, *The mass of an asymptotically flat manifold*, Comm. Pure Appl. Math. **39** (1986), no. 5, 661–693.
13. E. Bedford and B. A. Taylor, *The Dirichlet problem for a complex Monge-Ampère equation*, Invent. Math. **37** (1976), no. 1, 1–44.

14. R. Berman, *K-polystability of Q-Fano varieties admitting Kähler-Einstein metrics*, arXiv:1205.6214v2.
15. B. Berndtsson, *A Brunn-Minkowski type inequality for Fano manifolds and some uniqueness theorems in Kähler geometry*, arXiv:1303.4975.
16. O. Biquard and Y. Rollin, *Smoothing singular extremal Kähler surfaces and minimal lagrangians*, arXiv:1211.6957.
17. Z. Błocki, *The Calabi-Yau theorem*, Complex Monge-Ampère equations and geodesics in the space of Kähler metrics, Lecture Notes in Math., vol. 2038, Springer, Heidelberg, 2012, pp. 201–227.
18. S. Boucksom and H. Chen, *Okounkov bodies of filtered linear series*, Compos. Math. **147** (2011), 1205–1229.
19. S. Boucksom, P. Eyssidieux, V. Guedj, and A. Zeriahi, *Monge-Ampère equations in big cohomology classes*, Acta Math. **205** (2010), no. 2, 199–262.
20. L. Caffarelli, J. J. Kohn, L. Nirenberg, and J. Spruck, *The Dirichlet problem for nonlinear second-order elliptic equations II. Complex Monge-Ampère and uniformly elliptic equations*, Comm. Pure Appl. Math. **38** (1985), 209–252.
21. E. Calabi, *Extremal Kähler metrics*, Seminar on Differential Geometry (S. T. Yau, ed.), Princeton, 1982.
22. ———, *Extremal Kähler metrics II*, Differential geometry and complex analysis, Springer, 1985, pp. 95–114.
23. H. D. Cao, *Deformation of Kähler metrics to Kähler-Einstein metrics on compact Kähler manifolds*, Invent. Math. **81** (1985), no. 2, 359–372.
24. D. Catlin, *The Bergman kernel and a theorem of Tian*, Analysis and geometry in several complex variables (Katata, 1997), Trends Math., Birkhäuser Boston Inc., 1999, pp. 1–23.
25. J. Cheeger and T. Colding, *On the structure of spaces with Ricci curvature bounded below. I*, J. Differential Geom. **46** (1997), no. 3, 406–480.
26. ———, *On the structure of spaces with Ricci curvature bounded below. II*, J. Differential Geom. **54** (2000), no. 1, 13–35.
27. ———, *On the structure of spaces with Ricci curvature bounded below. III*, J. Differential Geom. **54** (2000), no. 1, 37–74.
28. B. Chen, An-Min Li, and L. Sheng, *Extremal metrics on toric surfaces*, arXiv:1008.2607.
29. X. X. Chen, *The space of Kähler metrics*, J. Differential Geom. **56** (2000), no. 2, 189–234.
30. X. X. Chen, *Space of Kähler metrics III. On the lower bound of the Calabi energy and geodesic distance*, Invent. Math. **175** (2009), no. 3, 453–503.
31. X. X. Chen, S. K. Donaldson, and S. Sun, *Kähler-Einstein metrics and stability*, arXiv:1210.7494.
32. ———, *Kähler-Einstein metrics on Fano manifolds, I: Approximation of metrics with cone singularities*, arXiv:1211.4566.
33. ———, *Kähler-Einstein metrics on Fano manifolds, II: Limits with cone angle less than  $2\pi$* , arXiv:1212.4714.
34. ———, *Kähler-Einstein metrics on Fano manifolds, III: Limits as cone angle approaches  $2\pi$  and completion of the main proof*, arXiv:1302.0282.
35. X. X. Chen, C. LeBrun, and B. Weber, *On conformally Kähler, Einstein manifolds*, J. Amer. Math. Soc. **21** (2008), no. 4, 1137–1168.

36. X. X. Chen and G. Tian, *Geometry of Kähler metrics and foliations by holomorphic discs*, Publ. Math. Inst. Hautes Études Sci. (2008), no. 107, 1–107.
37. P. Cherrier, *Équations de Monge-Ampère sur les variétés Hermitiennes compactes*, Bull. Sci. Math. (2) **111** (1987), no. 4, 343–385.
38. T. Darvas, *Morse theory and geodesics in the space of Kähler metrics*, arXiv:1207.4465.
39. J.-P. Demailly, *Complex analytic and differential geometry*, <http://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>.
40. W. Y. Ding, *Remarks on the existence problem of positive Kähler-Einstein metrics*, Math. Ann. **282** (1988), no. 3, 463–471.
41. S. K. Donaldson, *Lecture notes for TCC course “Geometric Analysis”*, <http://www2.imperial.ac.uk/~skdona/GEOMETRICANALYSIS.PDF>.
42. ———, *Remarks on gauge theory, complex geometry and four-manifold topology*, Fields Medallists’ Lectures (Atiyah and Iagolnitzer, eds.), World Scientific, 1997, pp. 384–403.
43. ———, *Symmetric spaces, Kähler geometry and Hamiltonian dynamics*, Northern California Symplectic Geometry Seminar, Amer. Math. Soc. Transl. Ser. 2, vol. 196, Amer. Math. Soc., Providence, RI, 1999, pp. 13–33.
44. ———, *Scalar curvature and stability of toric varieties*, J. Differential Geom. **62** (2002), 289–349.
45. ———, *Interior estimates for solutions of Abreu’s equation*, Collect. Math. **56** (2005), no. 2, 103–142.
46. ———, *Lower bounds on the Calabi functional*, J. Differential Geom. **70** (2005), no. 3, 453–472.
47. ———, *Two-forms on four-manifolds and elliptic equations*, inspired by S. S. Chern, Nankai Tracts Math., vol. 11, World Sci. Publ., Hackensack, NJ, 2006, pp. 153–172.
48. ———, *Extremal metrics on toric surfaces: A continuity method*, J. Differential Geom. **79** (2008), no. 3, 389–432.
49. ———, *Constant scalar curvature metrics on toric surfaces*, Geom. Funct. Anal. **19** (2009), no. 1, 83–136.
50. ———, *Discussion of the Kähler-Einstein problem*, <http://www2.imperial.ac.uk/~skdona/KENOTES.PDF> (2009).
51. S. K. Donaldson and S. Sun, *Gromov-Hausdorff limits of Kähler manifolds and algebraic geometry*, arXiv:1206.2609.
52. P. Eyssidieux, V. Guedj, and A. Zeriahi, *Singular Kähler-Einstein metrics*, J. Amer. Math. Soc. **22** (2009), no. 3, 607–639.
53. A. Fujiki, *Moduli space of polarized algebraic manifolds and Kähler metrics [translation of Sūgaku **42** (1990), no. 3, 231–243;]*, Sugaku Expositions **5** (1992), no. 2, 173–191.
54. W. Fulton, *Introduction to toric varieties*, Ann. of Math. Stud., vol. 131, Princeton University Press, Princeton, N.J., 1993.
55. A. Futaki, *An obstruction to the existence of Einstein-Kähler metrics*, Invent. Math. **73** (1983), 437–443.
56. A. Futaki and T. Mabuchi, *Bilinear forms and extremal Kähler vector fields associated with Kähler classes*, Math. Ann. **301** (1995), 199–210.
57. P. Gauduchon, *Invariant scalar-flat Kähler metrics on  $\mathcal{O}(-1)$* , preprint, 2012.

58. D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, reprint of the 1998 ed., Classics in Mathematics, Springer-Verlag, Berlin, 2001.
59. P. Griffiths and J. Harris, *Principles of algebraic geometry*, Pure and Applied Mathematics, Wiley-Interscience, New York, 1978.
60. D. Guan, *On modified Mabuchi functional and Mabuchi moduli space of Kähler metrics on toric bundles*, Math. Res. Lett. **6** (1999), no. 5-6, 547–555.
61. V. Guillemin, *Kähler structures on toric varieties*, J. Differential Geom. **40** (1994), 285–309.
62. V. Guillemin and S. Sternberg, *Convexity properties of the moment mapping*, Invent. Math. **67** (1982), no. 3, 491–513.
63. ———, *Riemann sums over polytopes*, Ann. Inst. Fourier (Grenoble) **57** (2007), no. 7, 2183–2195.
64. R. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Differential Geom. **17** (1982), no. 2, 255–306.
65. A. Hwang and M. A. Singer, *A momentum construction for circle-invariant Kähler metrics*, Trans. Amer. Math. Soc. **354** (2002), no. 6, 2285–2325.
66. W. Jiang, *Bergman kernel along the Kähler Ricci flow and Tian’s conjecture*, arXiv:1311.0428.
67. F. C. Kirwan, *Cohomology of quotients in symplectic and algebraic geometry*, Princeton University Press, 1984.
68. S. Kołodziej, *The complex Monge-Ampère equation*, Acta Math. **180** (1998), no. 1, 69–117.
69. ———, *Hölder continuity of solutions to the complex Monge-Ampère equation with the right-hand side in  $L^p$ : The case of compact Kähler manifolds*, Math. Ann. **342** (2008), no. 2, 379–386.
70. C. LeBrun, *Counter-examples to the generalized positive action conjecture*, Comm. Math. Phys. **118** (1988), no. 4, 591–596.
71. C. LeBrun and S. R. Simanca, *On the Kähler classes of extremal metrics*, Geometry and global analysis (Sendai, 1993), Tohoku Univ., 1993, pp. 255–271.
72. ———, *Extremal Kähler metrics and complex deformation theory*, Geom. and Func. Anal. **4** (1994), no. 3, 298–336.
73. L. Lempert and L. Vivas, *Geodesics in the space of Kähler metrics*, Duke Math. J. **162** (2013), no. 7, 1369–1381.
74. M. Levine, *A remark on extremal Kähler metrics*, J. Differential Geom. **21** (1985), no. 1, 73–77.
75. C. Li and C. Xu, *Special test configurations and K-stability of Fano varieties*, arXiv:1111.5398.
76. A. Lichnerowicz, *Sur les transformations analytiques des variétés kählériennes compactes*, C. R. Acad. Sci. Paris **244** (1957), 3011–3013.
77. R. B. Lockhart and R. C. McOwen, *Elliptic differential operators on noncompact manifolds*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (4) **1** (1985), no. 3, 409–447.
78. Z. Lu, *On the lower order terms of the asymptotic expansion of Tian-Yau-Zelditch*, Amer. J. Math. **122** (1998), no. 2, 235–273.
79. T. Mabuchi, *K-stability of constant scalar curvature polarization*, arXiv:0812.4093.
80. ———, *K-energy maps integrating Futaki invariants*, Tohoku Math. J. **38** (1986), no. 4, 575–593.

81. ———, *Some symplectic geometry on compact Kähler manifolds*, Osaka J. Math. **24** (1987), 227–252.
82. M. Maruyama, *On automorphism groups of ruled surfaces*, J. Math. Kyoto University **11-1** (1971), 89–112.
83. Y. Matsushima, *Sur la structure du groupe d'homéomorphismes analytiques d'une certaine variété kählérienne*, Nagoya Math. J. **11** (1957), 145–150.
84. D. McDuff and D. Salamon, *Introduction to symplectic topology*, OUP, 1998.
85. D. Mumford, J. Fogarty, and F. Kirwan, *Geometric invariant theory*, third ed., Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)], vol. 34, Springer-Verlag, Berlin, 1994. MR1304906 (95m:14012)
86. F. Pacard, *Lectures on connected sum constructions in geometry and non-linear analysis*, <http://www.math.polytechnique.fr/~pacard/Publications/Lecture-Part-I.pdf>.
87. S. T. Paul, *Stable pairs and coercive estimates for the Mabuchi functional*, arXiv:1308.4377.
88. S. T. Paul and G. Tian, *CM stability and the generalised Futaki invariant II*, Astérisque **328** (2009), 339–354.
89. G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, math.DG/0211159.
90. D. H. Phong, N. Sesum, and J. Sturm, *Multiplier ideal sheaves and the Kähler-Ricci flow*, Comm. Anal. Geom. **15** (2007), no. 3, 613–632.
91. D. H. Phong, J. Song, and J. Sturm, *Complex Monge-Ampère equations*, Surveys in differential geometry, vol. 17, Int. Press, Boston, MA, 2012, pp. 327–410.
92. J. Ross and R. P. Thomas, *A study of the Hilbert-Mumford criterion for the stability of projective varieties*, J. Algebraic Geom. **16** (2007), no. 2, 201–255.
93. W.-D. Ruan, *Canonical coordinates and Bergman metrics*, Comm. Anal. Geom. **6** (1998), no. 3, 589–631.
94. S. Semmes, *Complex Monge-Ampère equations and symplectic manifolds*, Amer. J. Math. **114** (1992), 495–550.
95. S. R. Simanca, *Kähler metrics of constant scalar curvature on bundles over  $CP_{n-1}$* , Math. Ann. **291** (1991), no. 2, 239–246.
96. L. Simon, *Theorems on regularity and singularity of energy minimizing maps*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 1996.
97. Y. T. Siu, *Lectures on Hermitian-Einstein metrics for stable bundles and Kähler-Einstein metrics*, DMV Seminar 8, Birkhäuser Verlag, Basel, 1987.
98. J. Song and G. Tian, *The Kähler-Ricci flow on surfaces of positive Kodaira dimension*, Invent. Math. **170** (2007), no. 3, 609–653.
99. ———, *Canonical measures and Kähler-Ricci flow*, J. Amer. Math. Soc. **25** (2012), no. 2, 303–353.
100. J. Song and B. Weinkove, *Lecture notes on the Kähler-Ricci flow*, arXiv:1212.3653.
101. J. Stoppa, *K-stability of constant scalar curvature Kähler manifolds*, Adv. Math. **221** (2009), no. 4, 1397–1408.
102. J. Stoppa and G. Székelyhidi, *Relative K-stability of extremal metrics*, J. Eur. Math. Soc. **13** (2011), no. 4, 899–909.
103. G. Székelyhidi, *Blowing up extremal Kähler manifolds II*, arXiv:1302.0760.

104. ———, *Filtrations and test-configurations, with an appendix by S. Boucksom*, arXiv:1111.4986.
105. ———, *The partial  $C^0$ -estimate along the continuity method*, arXiv:1310.8471.
106. ———, *Extremal metrics and K-stability*, Ph.D. thesis, Imperial College, London, 2006.
107. ———, *Extremal metrics and K-stability*, Bull. Lond. Math. Soc. **39** (2007), no. 1, 76–84.
108. ———, *On blowing up extremal Kähler manifolds*, Duke Math. J. **161** (2012), no. 8, 1411–1453.
109. R. P. Thomas, *Notes on GIT and symplectic reduction for bundles and varieties*, Surveys in Differential Geometry **10** (2006): A Tribute to Professor S.-S. Chern.
110. G. Tian, *On a set of polarized Kähler metrics on algebraic manifolds*, J. Differential Geom. **32** (1990), no. 1, 99–130.
111. ———, *On Calabi's conjecture for complex surfaces with positive first Chern class*, Invent. Math. **101** (1990), no. 1, 101–172.
112. ———, *Kähler-Einstein metrics with positive scalar curvature*, Invent. Math. **137** (1997), 1–37.
113. ———, *Canonical metrics in Kähler geometry*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 2000.
114. ———, *Existence of Einstein metrics on Fano manifolds*, Metric and Differential Geometry, The Jeff Cheeger Anniversary Volume (X. Dai and X. Rong, eds.), Progress in Mathematics, vol. 297, Springer, Basel, 2012, pp. 119–159.
115. G. Tian and Z. Zhang, *Regularity of Kähler-Ricci flows on Fano manifolds*, arXiv:1310.5897.
116. C. W. Tønnesen-Friedman, *Extremal Kähler metrics on minimal ruled surfaces*, J. Reine Angew. Math. **502** (1998), 175–197.
117. V. Tosatti and B. Weinkove, *The complex Monge-Ampère equation on compact hermitian manifolds*, J. Amer. Math. Soc. **23** (2010), no. 4, 1187–1195.
118. ———, *The Calabi-Yau equation, symplectic forms and almost complex structures*, Geometry and analysis. No. 1, Adv. Lect. Math. (ALM), vol. 17, Int. Press, Somerville, MA, 2011, pp. 475–493.
119. X. Wang, *Moment map, Futaki invariant and stability of projective manifolds*, Comm. Anal. Geom. **12** (2004), no. 5, 1009–1037.
120. X.-J. Wang and B. Zhou, *On the existence and nonexistence of extremal metrics on toric Kähler surfaces*, Adv. Math. **226** (2011), no. 5, 4429–4455.
121. D. Witt Nyström, *Test configurations and Okounkov bodies*, Compos. Math. **148** (2012), no. 6, 1736–1756.
122. S.-T. Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation I*, Comm. Pure Appl. Math. **31** (1978), 339–411.
123. ———, *Open problems in geometry*, Proc. Symposia Pure Math. **54** (1993), 1–28.
124. S. Zelditch, *Szegő kernel and a theorem of Tian*, Int. Math. Res. Notices **6** (1998), 317–331.
125. ———, *Bernstein polynomials, Bergman kernels and toric Kähler varieties*, J. Symplectic Geom. **7** (2009), no. 2, 51–76.

---

# Index

- $\partial\bar{\partial}$ -lemma, 8
- Abreu's formula, 77
- almost complex structure, 4
  - integrable, 4
- Arezzo-Pacard theorem, 155
- Aubin-Yau theorem, 35
  
- Bergman kernel, 129
  - asymptotics, 130
  - equivariant, 138
- blow-up, 154
- bundle
  - curvature of, 16
  - first Chern class, 18
  - line, 15
  - positive, 18
  - vector, 13
- Burns-Simanca metric, 155
  
- Calabi functional, 57
  - lower bound, 143
- Chern connection, 16
- Christoffel symbols, 11
- continuity method, 37
- covariant derivatives, 10
- curvature
  - Ricci, 12
  - Riemannian, 11
  - scalar, 12
  
- Ding functional, 71
- Donaldson-Futaki invariant, 112
  - toric, 125
  
- elliptic operator, 25
  - regularity, 26
  
- Fano manifolds, 51
- flat limit, 109
- Futaki invariant, 65, *see also*
  - Donaldson-Futaki invariant
  
- geodesic equation, 68
- geometric invariant theory, 90
  - quotient, 93
  
- Hamiltonian
  - action, 86
  - construction, 85
- harmonic function, 23
- Hilbert polynomial, 109
- Hilbert-Mumford criterion, 95
- Hirzebruch-Riemann-Roch theorem, 131
- Hölder spaces, 27
  - weighted, 160
- holomorphy potential, 63
- Hörmander technique, 133
  
- indicial roots, 162
  
- K-energy, 68
- K-stability, 113
  - toric, 127
- $\bar{K}$ -stability, 123
- $\hat{K}$ -stability, 123
- Kähler-Ricci flow, 55
- Kempf-Ness theorem, 98
- Kodaira embedding theorem, 20



- Laplace operator, 32–33
  - weighted, 160
- Levine's example, 65
- Lichnerowicz operator, 59
- Lichnerowicz's theorem, 64
- linearization of action, 100
- Liouville's theorem, 25
  
- Mabuchi functional, 68
- Mabuchi metric, 68
- manifold
  - complex, 1
  - projective, 3
  - toric, 76, 124
- mean value theorem, 23
- metric
  - extremal, 57
  - Fubini-Study, 7
  - Hermitian, 6
  - Kähler, 6
  - Kähler-Einstein, 35
- moment map, 85
  - projective space, 88
- momentum profile, 72
- Monge-Ampère equation, 47
  
- normal coordinates, 9
  
- partial  $C^0$ -estimate, 150
- polystable point, 95
- projective space, 2
  
- reductive group, 92
- relative K-stability, 116
- relatively stable point, 103
  
- Schauder estimates
  - global, 32
  - local, 27, 30
- schemes, 92
- semistable point, 93
- stable point, 94
- symplectic potential, 79
  
- test-configuration, 111
  
- Yau's theorem, 13, 47
- Yau-Tian-Donaldson conjecture, 113
  - extremal metrics, 116
  - filtrations, 123

A basic problem in differential geometry is to find canonical metrics on manifolds. The best known example of this is the classical uniformization theorem for Riemann surfaces. Extremal metrics were introduced by Calabi as an attempt at finding a higher-dimensional generalization of this result, in the setting of Kähler geometry.

This book gives an introduction to the study of extremal Kähler metrics and in particular to the conjectural picture relating the existence of extremal metrics on projective manifolds to the stability of the underlying manifold in the sense of algebraic geometry. The book addresses some of the basic ideas on both the analytic and the algebraic sides of this picture. An overview is given of much of the necessary background material, such as basic Kähler geometry, moment maps, and geometric invariant theory. Beyond the basic definitions and properties of extremal metrics, several highlights of the theory are discussed at a level accessible to graduate students: Yau's theorem on the existence of Kähler-Einstein metrics, the Bergman kernel expansion due to Tian, Donaldson's lower bound for the Calabi energy, and Arezzo-Pacard's existence theorem for constant scalar curvature Kähler metrics on blow-ups.

ISBN: 978-1-4704-1047-6



9 781470 410476

GSM/I52



For additional information  
and updates on this book, visit

[www.ams.org/bookpages/gsm-I52](http://www.ams.org/bookpages/gsm-I52)

AMS *on the Web*  
[www.ams.org](http://www.ams.org)