



# **HARMONIC ANALYSIS IN EUCLIDEAN SPACES**

## **Part 1**

**PROCEEDINGS OF SYMPOSIA  
IN PURE MATHEMATICS  
Volume XXXV, Part 1**

**HARMONIC ANALYSIS  
IN EUCLIDEAN SPACES**

**AMERICAN MATHEMATICAL SOCIETY  
PROVIDENCE, RHODE ISLAND  
1979**

PROCEEDINGS OF THE SYMPOSIUM IN PURE MATHEMATICS  
OF THE AMERICAN MATHEMATICAL SOCIETY

HELD AT WILLIAMS COLLEGE  
WILLIAMSTOWN, MASSACHUSETTS  
JULY 10–28, 1978

EDITED BY  
GUIDO WEISS  
STEPHEN WAINGER

Prepared by the American Mathematical Society  
with partial support from National Science Foundation grant MCS 77-23480

**Library of Congress Cataloging in Publication Data**

Symposium in Pure Mathematics, Williams College, 1978.

Harmonic analysis in Euclidean spaces.

(Proceedings of symposia in pure mathematics; v. 35)

Includes bibliographies.

1. Harmonic analysis—Congresses. 2. Spaces, Generalized—Congresses. I. Wainger, Stephen,  
1936— II. Weiss, Guido L., 1928— III. American Mathematical Society. IV. Title.  
V. Series.

QA403.S9 1978

515'.2433

79-12726

ISBN 0-8218-1436-2 (v.1)

AMS (MOS) subject classifications (1970). Primary 22Exx, 26A51, 28A70,  
30A78, 30A86, 31Bxx, 31C05, 31C99, 32A30, 32C20, 35–XX,  
42–XX, 43–XX, 44–XX, 47A35, 47B35

Copyright © 1979 by the American Mathematical Society

Printed in the United States of America

All rights reserved except those granted to the United States Government.

This book may not be reproduced in any form without the permission of the publishers.


## CONTENTS OF VOLUME

### PART 1

Dedication to Nestor Rivière.....	vi
Contents of Part 1.....	xix
Photographs.....	xxiii
Preface.....	xxv
Chapter 1. Real harmonic analysis.....	3
Chapter 2. Hardy spaces and BMO.....	189
Chapter 3. Harmonic functions, potential theory and theory of functions of one complex variable.....	313

### PART 2

Contents of Part 2.....	v
Chapter 4. Several complex variables.....	3
Chapter 5. Pseudo differential operators and partial differential equations.....	105
Chapter 6. Harmonic analysis in other settings: Probability, martingales local fields, Lie groups and functions analysis.....	233



**These Proceedings are dedicated to**  
**NESTOR M. RIVIÈRE**  
**1940–1978**

NESTOR M. RIVIÈRE  
JUNE 10, 1940  
JANUARY 3, 1978

It is striking to contemplate the influence of Nestor Rivière upon so many areas in analysis, and even more striking to think of his influence upon so many people. His graciousness was reflected in his mathematical work. He loved to work with people and to share his ideas. Many of us attending this conference have benefited in our own work from these ideas and from the breadth of his mathematical knowledge. His collaborations with others were always marked by a brilliance, a willingness to listen, and an optimism that created an unending flow of ideas.

Born and raised in Buenos Aires, Argentina, Nestor entered the University at the age of 16. He received his Licenciatura in mathematics in 1960, married Marisa Renda in 1961 and taught in Buenos Aires and Bariloche until December 1962. At that time, with the help of A.P. Calderón, he came to the University of Chicago to pursue his mathematical studies. Nestor received his Ph.D. degree in 1966 and in the Fall of that year became a member of the faculty at the University of Minnesota. In April 1974 Marisa and Nestor's daughter, Melisa, was born.

Nestor was naturally influenced by his education at Chicago. Real and harmonic analysis always remained his primary mathematical interest. At Minnesota the environment was perfect for the development of this interest and for the application of his knowledge to problems in other areas, especially to certain areas of partial differential equations. From 1966 Nestor's work in real and harmonic analysis went along hand-in-hand with his work in P.D.E. . Below we review some of Nestor's work in four major areas: Singular Integrals, Multiplier Theory, Interpolation Theory, and Partial Differential Equations.

Nestor's love for mathematics and his desire to share ideas made him an exceptional teacher. During his years at Minnesota he supervised the theses of a number of students, among them were Eleonor Harboure de Aquilera, Nestor Aquilera, Norberto Fava, Robert Hanks, Wally Madych, and Felipe Zo.

### Singular Integrals

At the time Nestor was a student in Chicago the Calderón-Zygmund theory

of singular integral operators of elliptic type had already arrived to a well understood stage. The techniques of the 1952 paper, "On Certain Singular Integrals", were being used by B.F. Jones to study the  $L^p$ -continuity of singular integrals arising from parabolic equations. Rivièrè realized that the entire theory could be placed under one general setting dependent on a fixed notion of dilation, namely

$$\lambda^\alpha x = (\lambda^{\alpha_1} x_1, \dots, \lambda^{\alpha_n} x_n), \lambda > 0$$

where  $\alpha_1, \dots, \alpha_n$  are given positive numbers.

Associated with the above (nonisotropic) dilation is the metric,  $r(x)$ , defined for  $x \neq 0$  as the unique positive number satisfying

$$\sum_{j=1}^n \frac{x_j^2}{r^{2\alpha_j}} = 1.$$

$r(x)$  has the homogeneity property,  $r(\lambda^\alpha x) = \lambda r(x)$  and there is a polar decomposition of  $\mathbb{R}^n$  relative to  $r$ , i.e. each  $x \neq 0$  can be written as

$$x = r^\alpha \sigma, \quad |\sigma| = 1$$

and

$$dx = r^{(\sum \alpha_i) - 1} J(\sigma) dr d\sigma$$

with  $0 < J(\sigma) \in C^\infty(\Sigma)$ ,  $\Sigma = \{\sigma : |\sigma| = 1\}$ .

In this setting one can mimic the techniques of the 1952 paper of Calderón-Zygmund and prove the  $L^p$  continuity,  $1 < p < \infty$ , of convolution singular integrals of the form

$$(*) \quad \lim_{\epsilon \rightarrow 0} \int_{r(x-y) > \epsilon} k(x-y) f(y) dy$$

where

- i)  $k(x) \in C^1(\mathbb{R}^n \setminus \{0\})$ ,
- ii)  $k(\lambda^\alpha x) = \lambda^{-\sum \alpha_i} k(x)$ ,  $\lambda > 0$ ,  $x \neq 0$ ,
- iii)  $\int_{\Sigma} k(\sigma) J(\sigma) d\sigma = 0$  where  $d\sigma$  is area measure on  $\Sigma$ .

The proofs of the above results appeared in article [1] and the extensions of the results to certain nonconvolution type operators were given in [3].

Nestor went on to considerably generalize the setting in which one could consider convolution singular integral operators. In [13] he attaches the notion of a singular kernel with a one parameter family,  $\{U_r : r > 0\}$ , of open bounded neighborhoods of the origin satisfying the conditions:

$$i) \quad U_r \subset U_s, \quad r < s, \quad \bigcap_{r > 0} \bar{U}_r = \{0\},$$

- ii) The algebraic difference  $U_r - U_r \subset U_{\phi(r)}$ , with  $\phi: (0, \infty) \rightarrow (0, \infty)$  nondecreasing, continuous,  $\phi(r) \uparrow \infty$  as  $r \uparrow \infty$ .
- iii) The Lebesgue measure of  $U_{\phi(r)}$ , denoted by  $m(U_{\phi(r)})$ , is  $\leq Am(U_r)$ ,  $A$  independent of  $r$ .

Associated to such a family Rivière defined the notion of a singular kernel as a function,  $k(x) \in L^1_{loc}(\mathbb{R}^n \setminus \{0\})$  with the properties,

- i)  $\int_{U_s \setminus U_r} k(x) dx$  is bounded independently of  $s$  and  $r$  and  $\lim_{r \rightarrow 0^+} \int_{U_s \setminus U_r} k(x) dx$  exists for each fixed  $s > 0$ ,
- ii)  $\int_{U_{\phi(r)} \setminus U_r} |k(x)| dx \leq A$ , independent of  $r > 0$
- iii) There exists  $A > 0$  such that  $\int_{\mathbb{R}^n \setminus U_{\phi(r)}} |k(x-y) - k(x)| dx \leq A$  for all  $y \in U_r$  and for all  $r$ .

He then naturally defined the doubly truncated singular integral operator

$$K_{r,s}(f)(x) = \int_{U_s \setminus U_r} k(y)f(x-y)dy$$

and proved the following theorem, which was new even for the elliptic case, i.e.  $r(x) = |x|$  and  $\phi(r) = 2r$ .

Theorem. The operator  $K_x f(x) = \sup_{r,s} |K_{r,s}(f)(x)|$  is bounded from  $L^p \rightarrow L^p$ ,  $1 < p < \infty$ , and from  $L^1 \rightarrow$  weak  $L^1$ .

In particular for  $f \in L^p$ ,  $1 \leq p < \infty$   $\lim_{\substack{s \rightarrow \infty \\ r \rightarrow 0^+}} K_{r,s}(f)(x)$  exist pointwise for almost every  $x \in \mathbb{R}^n$

In 1973 Nestor, together with Steve Wainger and Alex Nagel, returned to a problem in singular integrals which was first formulated in the study of the singular integral operators involving "mixed homogeneous" kernels defined earlier. The problem was to find a "method of rotation" for these operators corresponding to that developed by Calderón and Zygmund. In the latter case the  $L^p$ -continuity of a singular integral of elliptic type arising from an odd kernel was reduced to the continuity of the one dimensional Hilbert transform. The problem was to find the appropriate one-dimensional operator for the mixed homogeneous operators coming from an odd kernel. A candidate was formulated as early as 1966, namely for  $x \in \mathbb{R}^n$

$$T_{\gamma} f(x) = \lim_{\epsilon \rightarrow 0} \int_{|t| > \epsilon} f(x_1 - \operatorname{sgn} t |t|^{\alpha_1}, \dots, x_n - \operatorname{sgn} t |t|^{\alpha_n}) \frac{dt}{t},$$

$$\alpha_i > 0, \quad i = 1, \dots, n.$$

This operator was called by Nagel, Rivière, and Wainger, the Hilbert transform of  $f$  along the curve,  $\gamma(t) = (\operatorname{sgn} t |t|^{\alpha_1}, \dots, \operatorname{sgn} t |t|^{\alpha_n})$ . In [24] they prove the continuity of  $T_{\gamma}$  on  $L^p(\mathbb{R}^n)$ ,  $1 < p < \infty$ , and as a consequence, the continuity on  $L^p(\mathbb{R}^n)$  of the mixed homogeneous operators in the form (\*) where the smoothness of  $k(x)$  is replaced by the condition

$$\int_{\Sigma} |k(\sigma)| \log^+ |k(\sigma)| d\sigma < \infty.$$

### Multiplier Theory

Rivière's interest in the theory of Fourier multipliers began as a graduate student in Chicago. In [1] there appears the extension of the Hörmander multiplier theorem to the case of multipliers,  $m(x)$ , behaving like smooth functions with mixed homogeneity zero. More precisely if  $r(x)$  denotes the metric, discussed in the previous part, corresponding to the given dilation

$$\lambda^{\alpha} x = (\lambda^{\alpha_1} x_1, \dots, \lambda^{\alpha_n} x_n), \quad \alpha_i > 0,$$

then  $m(x)$  is a Fourier multiplier on all  $L^p$ ,  $1 < p < \infty$  provided  $m$  is bounded and

$$R^{2(\alpha \cdot \beta) - |\alpha|} \int_{R/2 \leq r(x) \leq 2R} |D^{\beta} m(x)|^2 dx \leq C, \quad \text{independent of } R,$$

for all  $\beta$ ,  $|\beta| < N$  with  $N > \frac{|\alpha|}{2}$  ( $|\alpha| = \sum_i \alpha_i$ ).

When Nestor joined the faculty at Minnesota in 1966 he immediately began working with Walter Littman and Charles McCarthy on refinements of the Marcinkiewicz multiplier theorem in  $\mathbb{R}^n$  ([6]). At this same time he started studying a problem posed to him earlier by A.P. Calderón who asked if a bounded rational function on  $\mathbb{R}^d$ ,  $d > 1$ , was a Fourier multiplier on  $L^p$  for some interval of  $p$ 's around 2. Already Littman, McCarthy, and Rivière had given in [7] an example of a bounded rational function on  $\mathbb{R}^2$  which was not a multiplier on  $L^p(\mathbb{R}^2)$ ,  $1 < p < \frac{4}{3}$ . The example was  $\frac{1}{x^2 - y + i}$ . In [13]

Nestor extended the Marcinkiewicz multiplier result of [6] to operator valued multipliers and proved that any bounded rational function on  $\mathbb{R}^d$  is a multiplier on the space of functions

$$L^p(\mathbb{R}^{d-1}, L^2(\mathbb{R})) = \{f(t, x), t \in \mathbb{R}, x \in \mathbb{R}^{d-1} \text{ such that } (\int_{\mathbb{R}^{d-1}} (\int |f(t, x)|^2 dt)^{p/2} dx)^{1/p} < \infty\} \quad (1 < p < \infty).$$

Finally in [13] Nestor proved a version of the Hörmander multiplier theorem that not only considerably generalized the setting of the theorem but added an original twist which even for the elliptic case gave a very interesting result. It is in this setting that we would like to state the result.

Theorem. Let  $\beta_j, j=1, \dots, d$ , be positive integers such that  $\sum_{j=1}^d \beta_j^{-1} < 2$ . Assume  $m \in L^\infty$ , and

$$\sup_{\substack{n=0, \pm 1, \pm 2, \\ j=1, \dots, d}} 2^{(2\beta_j - n)d} \int_{2^{n-1} < |x| < 2^n} |D_{x_j}^{\beta_j} m(x)|^2 dx < \infty .$$

Then  $m$  is a Fourier multiplier on  $L^p(\mathbb{R}^d)$ ,  $1 < p < \infty$ .

The novelty of the above result is the "trade-off" of smoothness of the individual variables. One may assume a weak smoothness in one or several of the variables by requiring sufficient smoothness in the remaining ones.

Interpolation

Nestor began his studies at the University of Chicago in the area of interpolation. His unpublished thesis extended the Riesz-Thorin or Complex method of interpolation from Banach spaces to topological vector spaces,  $B$ , with a metric topology defined through an  $s$ -norm,  $0 < s \leq 1$ , i.e. a function  $\|\cdot\|_s: B \rightarrow [0, \infty)$  such that

- i)  $\|x\|_s = 0 \iff x = 0$
- ii)  $\|x+y\|_s \leq \|x\|_s + \|y\|_s$
- iii)  $\|\lambda x\|_s = |\lambda|^s \|x\|_s$ .

The metric is of course defined as  $d(x,y) = \|x-y\|_s$ . These spaces are called  $s$ -Banach spaces and prime examples are the Lebesgue and Hardy spaces,  $L^s(X, d\mu)$  and  $H^s(\mathbb{R}^n)$ ,  $0 < s < 1$ . In the thesis Nestor identifies, via the complex method, the intermediate spaces of various  $s$ -Banach spaces of functions and in particular shows that

$$[L^{p_1}(X, d\mu), L^{p_2}(X, d\mu)]_\alpha = L^p(X, d\mu)$$

where  $\frac{1}{p} = \frac{\alpha}{p_1} + \frac{(1-\alpha)}{p_2}$ ,  $0 < \alpha < 1$ ,  $0 < p_1, p_2 \leq \infty$ .

In [14] Nestor extends the techniques of the Marcinkiewicz interpolation theorem and as a consequence proves that any sublinear operator mapping boundedly

$$L^\infty(\mathbb{R}^n) \rightarrow BMO(\mathbb{R}^n)$$

and

$$L^1(\mathbb{R}^n) \rightarrow L(1, \infty)$$

must also map boundedly  $L^p \rightarrow L^p$  for  $1 < p < \infty$ . Here BMO denotes the space of functions with bounded mean oscillation, as defined by F. John and L. Nirenberg, and  $L(1, \infty)$  is the Lorentz space of functions commonly called "weak  $L^1$ ". This work of Nestor's, published in 1971, was his one mathematical paper written in Spanish.

Together with Yoram Sagher in [17], Nestor calculated the intermediate spaces,  $(H^1, C_\omega)_{\theta, p}$ , for the Lions-Peetre or real method of interpolation. Here  $H^1$  denotes the classical Hardy space of functions defined on  $\mathbb{R}^n$  and  $C_\omega$  denotes the class of continuous functions on  $\mathbb{R}^n$  vanishing at  $\infty$ . They proved

$$(H^1, C_\omega)_{\theta, p} = L^p \text{ for } \frac{1}{p} = (1 - \theta), \quad 0 < \theta < 1.$$

As a consequence, if  $M$  = space of finite measures then

$$(BMO, M)_{\theta, q} = L(p', q), \quad \frac{1}{p} = 1 - \theta, \quad \frac{1}{p} + \frac{1}{p'} = 1.$$

( $L(p', q)$  denotes the usual Lorentz space.)

The above results were extended in [19] where the equalities

$$(H^{p_0}, L^\infty)_{\theta, p} = H^p \text{ and } (H^{p_0}, H^{p_1})_{\theta, p} = H^p$$

were proved respectively for  $\frac{1}{p} = \frac{1 - \theta}{p_0}$  and  $\frac{1}{p} = \frac{1 - \theta}{p_0} + \frac{\theta}{p_1}$ ,  $0 < p_0, p_1 < \infty$ ,  $0 < \theta < 1$ .

Nestor's interest in interpolation remained throughout his career. In 1976 his student, Robert Hanks, identified in his thesis the intermediate space

$$(H^p, BMO)_{\theta, p} \text{ as } H^p \text{ for } p = \frac{p_0}{1 - \theta}, \quad 0 < p_0 < \infty.$$

As a consequence Nestor's result on sublinear operators described above was extended to the case

$$T: L^\infty \rightarrow BMO$$

$$T: H^1 \rightarrow L(1, \infty).$$

### Partial Differential Equations

As a graduate student Nestor was very interested in the use of singular integral operators as a general tool to study existence, uniqueness, and regularity for a large class of equations modeled mostly from elliptic operators. The use of the symbolic calculus, developed by A.P. Calderón and A. Zygmund in the elliptic case was adapted to parabolic operators in [4]. Any such operator, say  $L = \sum_{|\alpha| = 2b} A_\alpha(x, t) D_x^\alpha - D_t$ , was decomposed on smooth functions with support in  $\mathbb{R}_+^{n+1}$  as

$$L = S((-1)^{b+1} \Delta^b - D_t)$$

with  $S$  a parabolic singular integral operator. Assuming boundedness and uniform continuity on the coefficients the operator  $S$  was shown to be invertible on  $L^p(S_T)$  with  $S_T = \mathbb{R}^n \times (0, T)$  and  $1 < p < \infty$ . From this followed easily the existence and uniqueness for the problem

$$Lu = f \text{ in } S_T, u(x, 0) = 0$$

with  $f \in L^p(S_T)$  and  $u$  in the class of functions having spatial derivatives of orders  $\leq 2b$  and one time derivative in  $L^p(S_T)$ . Also in papers [28] and [29] one can again find the development of a symbolic calculus designed to give general algebraic conditions for the solvability of initial boundary value problems associated with the Navier-Stokes equations.

The final three years of Nestor's life were dedicated to problems in partial differential equations and some of his best work was done at this time. Together with Luis Caffarelli very precise regularity results in two dimensions were obtained for the free boundaries arising from the solution of the minimal energy problem above a given obstacle and from the solution of the minimal surface problem staying above an obstacle.

In the above situations we are given a bounded, connected domain  $D \subset \mathbb{R}^n$  and a function  $\varphi$ , the obstacle, defined on  $\bar{D}$ , satisfying:

- a)  $\varphi < 0$  on  $\partial D$
- b)  $\Delta\varphi$  and  $\nabla(\Delta\varphi)$  do not vanish simultaneously.

We let  $v(x)$ ,  $x \in D$ , be the solution of a specific variational inequality satisfying  $v \geq \varphi$  in  $D$ . For example in the case of minimizing energy

$$\int_D |\nabla v|^2 dx = \inf \left\{ \int_D |\nabla u|^2 dx : u|_{\partial D} = 0, u \geq \varphi \text{ in } D \right\}.$$

In [26] it was shown that the set of coincidence,

$$\Lambda = \{x \in D : v(x) = \varphi(x)\},$$

has the following structure in 2 dimensions:

Theorem. If  $\Delta\varphi \in C^{k, \alpha}$ ,  $0 < \alpha < 1$ ,  $k \geq 1$ , then each component of the interior of  $\Lambda$  is composed of a finite number of Jordan arcs each having a nondegenerate  $C^{k+1, \alpha}$  parametrization. Moreover if  $\Delta\varphi$  is real analytic the Jordan arcs are real analytic.

It was later shown by Caffarelli (even for the general  $n$ -dimensional case) that if  $x_0 \in \Lambda$  is a point of positive density of  $\Lambda$  then there exists a ball,  $B(x_0)$ , about  $x_0$  such that  $\partial\Lambda \cap B(x_0)$  is a  $C^1$  curve. In [31] Rivièrè and Caffarelli studied the case when  $x_0$  is a point of zero density and showed the existence of a neighborhood,  $B(x_0)$ , of  $x_0$  in which  $\Lambda \cap B(x_0)$  is contained between two tangent  $C^1$  curves. In fact, with a proper choice of

coordinates  $\Lambda \cap B(x_0)$  is contained between the curves

$$y = \pm C_1 |x| \exp \{-C_2 (\log |x|)^{1/2}\} .$$

The final work submitted for publication by Rivière and his co-authors was [34]. Here A.P. Calderón's recent results concerning the Cauchy integral over a  $C^1$ -curve were used to solve the Dirichlet and Neumann problems for Laplace's equation in a  $C^1$ -domain,  $D$ , contained in  $R^n$ . The data were assumed to belong to  $L^p(\partial D)$ ,  $1 < p < \infty$ , and the solutions were written respectively in the form of the classical double and single layer potentials. In the Dirichlet case the nontangential maximal function associated with the solution was shown to belong to  $L^p(\partial D)$  and, as a consequence, the solution converged nontangentially to the data at almost every point of the boundary. Similarly in the Neumann problem the nontangential maximal function associated to the gradient of the solution was shown to belong to  $L^p(\partial D)$  and again the data was assumed in a pointwise nontangential sense at almost every point of the boundary.

On November 23, 1977, during an informal gathering of harmonic analysts from the Midwest at the University of Chicago, Nestor spoke of some open problems which he considered exceptionally interesting. These problems are listed in this proceedings.

The last three years of Nestor's life were years of great personal growth. For each new crisis in his illness he found in himself new resources of courage. His sensitivity to other people increased, and his mathematical work continued unabated to the end. The grace he had shown under the most relentless pressure one has to face was his last, and greatest achievement.

Alberto Calderón  
University of Chicago  
Chicago, Illinois 60637

Eugene Fabes  
University of Minnesota  
Minneapolis, Minnesota 55455

Yoram Sagher  
University of Illinois at Chicago Circle  
Chicago, Illinois 60680

## Bibliography of N.M. Rivière

1. Singular integrals with mixed homogeneity (with E.B. Fabes), *Studia Math.* 27(1966), 19-38.
2. Commutators of singular integrals (with E.B. Fabes), *Studia Math.* 26 (1966), 225-232.
3. Symbolic calculus of kernels with mixed homogeneity (with E.B. Fabes), *Proc. Sump. Pure Math.* 10, Singular Integrals, (1967), 106-127.
4. Systems of parabolic equations with uniformly continuous coefficients (with E.B. Fabes), *J. D'Anal. Math.* 17 (1966), 305-334.
5. The converse of Wiener-Levy-Marcinkiewicz theorem (with Y. Sagher), *Studia Math.* 27 (1966), 133-138.
6.  $L^p$ -multiplier theorems (with W. Littman and C. McCarthy), *Studia Math.* 30 (1968), 193-217.
7. The nonexistence of  $L^p$ -estimates for certain translation-invariant operators (with W. Littman and C. McCarthy), *Studia Math.* 30 (1968), 219-229.
8. Vector valued multipliers and applications, *Bull. Amer. Math. Soc.* 74 (1968), 946-948.
9. Some recursive formulas on free distributive lattices, *J. Comb. Theory* 5 (1968), 229-234.
10. On singular integrals, *Bull. Amer. Math. Soc.* 75 (1969), 843-847.
11. Multipliers of trigonometric series and pointwise convergence (with Y. Sagher), *Trans. Amer. Math. Soc.* 140 (1969), 301-308.
12.  $L^p$ -estimates ( $1 < p \leq \infty$ ) near the boundary for solutions of the Dirichlet problem (with E.B. Fabes), *SIAM J. Adv. Diff. Eqns.* (special issue)(1969), Proofs appeared in *Ann. de Pisa* 24 (1970), 491-553.
13. Singular integrals and multiplier operators, *Arkiv Math.* 9 (1971), 243-278.
14. Interpolacion a la Marcinkiewicz, *Revista de la Union Matematica Argentina*, Volume dedicated to Professor G. Dominquez, 25 (1971), 363-377.
15. The initial value problem for the Navier-Stokes equations with data in  $L^p$  (with E.B. Fabes and B.F. Jones), *Arch. Rat. Mech. Anal.* 45 (1972), 222-240
16. Estimates for translation invariant operators on spaces with mixed norms (with C.S. Herz), *Studia Math.* 44 (1972), 511-515.
17. Interpolation between  $H^1$  and  $L^\infty$ , the real method (with Y. Sagher), *J. Funct. Anal.* 14 (1973), 401-409.
18. On two theorems of Paley (with Y. Sagher), *Proc. Amer. Math. Soc.* 42 (1974), 238-242.
19. Interpolation between  $H^p$  spaces, the real method (with C. Fefferman and Y. Sagher), *Trans. Amer. Math. Soc.* 19 (1974), 75-82.

20. On Hilbert transformations along curves (with A. Nagel and S. Wainger), Bull. Amer. Math. Soc. 80 (1974), 106-108.
21. Maximal smoothing operators (with C.P. Calderon and E.B. Fabes), Indiana Math. J. 23 (1974), 889-898.
22. Commutators of singular integral operators with  $C^1$ -kernels (with E.B. Fabes and W. Littman), Proc. Amer. Math. Soc. 48 (1975), 397-402.
23. Multipliers of the Hölder classes (with W. Madych), J. Funct. Anal. 21 (1976), 369-379.
24. On Hilbert transforms along curves II (with A. Nagel and W. Wainger), Amer. J. Math. 98 (1976), 395-403.
25. On the rectifiability of domains with finite perimeter (with L.A. Caffarelli), Ann. Scuola Norm. Sup. Pisa, Series IV, 3 (1976), 177-186.
26. Smoothness and analyticity of free boundaries (with L.A. Caffarelli), Ann. Scuola Norm. Sup. Pisa, Series IV, 3 (1976), 289-310.
27. A maximal function associated to the curve  $(t, t^2)$  (with A. Nagel and S. Wainger), Proc. Nat. Acad. Sci. 73 (1976), 1416-1417.
28. Singular integrals and hydrodynamical potentials (with E.B. Fabes and J. Lewis), Amer. J. Math. 99 (1977), 601-625.
29. Boundary value problems for the Navier-Stokes equation (with E.B. Fabes and J. Lewis), Amer. J. Math. 99 (1977), 626-668.
30. The smoothness of the elastic-plastic free boundary of a twisted bar (with L.A. Caffarelli), Proc. Amer. Math. Soc., Vol 63, No. 1 (1977), 56-58.
31. Asymptotic behavior of free boundaries at their singular points (with L.A. Caffarelli), Ann. of Math., 106 (1977), 309-317.
32. The Cauchy integral in Lipschitz domains and applications (with A.P. Calderon, C.P. Calderon, E.B. Fabes, and M. Jodeit) Bull. Amer. Math. Soc., Vol. 84, No. 2, (1978), 287-290.
33. On the Lipschitz character of the stress tensor, when twisting an elastic-plastic bar (with L.A. Caffarelli), Arch. Rat. Mech. Anal., to appear.
34. Potential techniques for boundary value problems on  $C^1$ -domains (with E.B. Fabes, M. Jodeit), Acta Mathematica, to appear.

Some Open Questions

1. Let  $\{\mathcal{U}_t : t > 0\}$  be a family of open bounded convex sets containing 0 such that  $\mathcal{U}_s \subset \mathcal{U}_t$  for  $s < t$ ,  $\bigcap \bar{\mathcal{U}}_t = \{0\}$ . If  $\mu$  and  $\nu$  are finite regular Borel measures is it true that

$$\lim_{t \rightarrow 0^+} \frac{\mu(x + \mathcal{U}_t)}{\nu(x + \mathcal{U}_t)} \text{ exists almost everywhere with respect to } \nu \text{ ?}$$

2. Consider the fundamental solution,  $m(x,y) = \frac{1}{x^2 - y + i}$  of the Schrödinger operator  $-\frac{\partial^2}{\partial x^2} + i\frac{\partial}{\partial y} + i$ . As a Fourier multiplier on  $L^p(\mathbb{R}^2)$   $m(x,y)$  is unbounded on  $L^p$  for  $p > 4$ . What can be said for the range  $4/3 \leq p \leq 4$  ?

3. Suppose  $P(x)$  and  $Q(x)$  are polynomials on  $\mathbb{R}^n$  such that  $\frac{P(x)}{Q(x)}$  is bounded. Does it follow that  $\frac{P(x)}{Q(x)}$  is Fourier multiplier on  $L^p(\mathbb{R}^n)$  for some intervals of  $p$ 's around 2 .

4. Assume  $k_1(x)$  and  $k_2(x)$ ,  $x \in \mathbb{R}^n$ , are smooth functions on  $\mathbb{R}^n \setminus \{0\}$  such that  $k_1(x)$  is an elliptic singular kernel and  $k_2(x)$  is a parabolic kernel, i.e.  $k_1(\lambda x) = \lambda^{-n} k_1(x)$ ,  $\lambda > 0$ ,  $x \neq 0$  and its mean value over the unit sphere is zero;  $k_2(\lambda x_1, \dots, \lambda x_{n-1}, \lambda^2 x_n) = \lambda^{n-1} k_2(x)$  and its appropriate mean value on the unit sphere is zero. Set  $K_1 f = k_1 * f$ . Does the composition  $K_1 K_2$  map  $L^1 \rightarrow L^{1,\infty}$  ? (see [23]).

5. Suppose  $T$  is a translation invariant operator mapping  $L^p \rightarrow L^{p,\infty}$  for a given  $p$ ,  $1 < p \leq 2$ . Does this imply  $T : L^p \rightarrow L^{p,p'}$ ,  $\frac{1}{p} + \frac{1}{p'} = 1$  ? (The unknown cases are  $1 < p < 2$  .)

6. Let  $Kf = k * f$  where  $k(x)$ ,  $x \in \mathbb{R}^n$ , is homogeneous of degree  $-n$ , mean value zero over the unit sphere,  $\Sigma$ , in  $\mathbb{R}^n$ , and in  $L \log^+ L(\Sigma)$ . Does  $K : L^\infty \rightarrow BMO$  ?

7. For a given bounded  $C^1$ -domain  $D \subset \mathbb{R}^n$ , consider the boundary value problems

$$\Delta^2 u(x) = 0, x \in D, \text{ with } u|_{\partial D}, \frac{\partial u}{\partial n}|_{\partial D} \text{ given}$$

$$\Delta^2 u(x) = 0, x \in D, \text{ with } \frac{\partial u}{\partial n}|_{\partial D}, \frac{\partial^2 u}{\partial n^2}|_{\partial D} \text{ given}$$

$$\Delta^2 u(x) = 0, x \in D, \text{ with } \frac{\partial^2 u}{\partial n^2}|_{\partial D}, \frac{\partial^3 u}{\partial n^3}|_{\partial D} \text{ given. Here } \frac{\partial^j u}{\partial n^j}|_{\partial D} \text{ denotes}$$

the  $j^{\text{th}}$  normal derivative of  $u$  on  $\partial D$ . Prescribe classes of boundary data which give existence and uniqueness.

Since the meeting in Williamstown Carlos Kenig and Peter Tomas have answered problem 2 and, as a consequence, also problem 3. They have proved that  $m(x,y)$  is only a multiplier on  $L^2(\mathbb{R}^2)$ .


## CONTENTS OF PART 1

Volume contents.....	v
Photograph of Rivière.....	vi
Nestor M. Rivière (1940–1978).....	vii
Bibliography of N. M. Rivière.....	xv
Some open questions.....	xvii
Photographs of participants.....	xxiii
Preface.....	xxv
 <b>Chapter 1. Real harmonic analysis</b>	
Some problems in harmonic analysis.....	3
E. M. STEIN	
On operators of harmonic analysis which are not convolutions.....	21
R. R. COIFMAN	
Maximal functions, covering lemmas and Fourier multipliers.....	29
ANTONIO CÓRDOBA	
Covering lemmas, maximal functions and multiplier operators in Fourier analysis .....	51
R. FEFFERMAN	
Besicovitch theory of linearly measurable sets and Fourier analysis.....	61
MIGUEL DE GUZMÁN	
Weighted norm inequalities for classical operators.....	69
BENJAMIN MUCKENHOUPT	
Applications of Fourier transforms to averages over lower dimensional sets.....	85
STEPHEN WAINGER	
Hilbert transforms and maximal functions related to variable curves.....	95
ALEXANDER NAGEL, ELIAS M. STEIN and STEPHEN WAINGER	
Regularity of spherical means.....	99
JACQUES PEYRIÈRE	
Restriction theorems for the Fourier transforms to some manifolds in $\mathbf{R}^n$ .....	101
ELENA PRESTINI	
Restriction theorems for the Fourier transform.....	111
PETER A. TOMAS	
Riesz potentials and Fourier multipliers.....	115
RICHARD J. BAGBY	

Fourier multipliers vanishing at infinity.....	121
LEONEDE DE MICHELE and IAN R. INGLIS	
Weighted norm inequalities for the Littlewood-Paley function $g_\lambda^*$ .....	125
ALBERTO TORCHINSKY	
Weighted norm inequalities for multipliers.....	133
WO-SANG YOUNG	
Non-equivalence between two kinds of conditions on weight functions.....	141
JAN-OLOV STRÖMBERG	
Some remarks about Beckner's inequality.....	149
DANIEL W. STROOCK	
Hypercontractive estimates for semigroups.....	159
FRED B. WEISSLER	
Singular integrals near $L^1$ .....	163
WILLIAM C. CONNETT	
On Carleson's convergence theorem for $L^2$ functions.....	167
BOGDAN M. BAISHANSKI	
Multiple Fourier series of functions of generalized bounded variation.....	171
DANIEL WATERMAN	
Some inequalities for Riesz potentials of trigonometric polynomials of several variables.....	175
G. WILMES	
A note on Sobolev spaces.....	183
BJÖRN E. J. DAHLBERG	
<b>Chapter 2. Hardy spaces and BMO</b>	
Some problems in the theory of Hardy spaces.....	189
GUIDO WEISS	
Weak-type inequalities for $H^p$ and BMO.....	201
COLIN BENNET and ROBERT SHARPLEY	
Singular integral characterizations of nonisotropic $H^p$ spaces and the F. and M. Riesz' theorem.....	231
R. R. COIFMAN and BJÖRN DAHLBERG	
A maximal theory for generalized Hardy spaces.....	235
ROBERTO A. MACÍAS and CARLOS SEGOVIA	
Local Hardy spaces.....	245
DAVID GOLDBERG	
Distributions with strong maximal functions $L^p(\mathbb{R}^n)$ .....	249
W. R. MADYCH	

Weighted Hardy spaces.....	253
JOSÉ GARCÍA-CUERVA	
Weighted Hardy spaces on Lipschitz domains.....	263
CARLOS E. KENIG	
The atomic decomposition of Hardy spaces.....	275
ROBERT H. LATTER	
The molecular characterization of Hardy spaces.....	281
MITCHELL H. TAIBLESON and GUIDO WEISS	
A characterization of $H^1(\Sigma_{n-1})$ .....	289
FULVIO RICCI and GUIDO WEISS	
Two constructions in BMO.....	295
JOHN B. GARNETT	
Invariant subspaces and subnormal operators.....	303
JAMES E. BRENNAN	
<b>Chapter 3. Harmonic functions, potential theory and theory of functions of one complex variable</b>	
Harmonic functions in Lipschitz domains.....	313
BJÖRN E. J. DAHLBERG	
A survey of harmonic functions on symmetric spaces.....	323
ADAM KORANYI	
Positive harmonic functions vanishing on the boundary of certain domains in $\mathbb{R}^{n+1}$ .....	345
MICHAEL BENEDICKS	
Positive harmonic functions with zero boundary values.....	349
HARRY KESTEN	
Harmonic functions with BMO boundary values.....	353
UMBERTO NERI	
$L^p$ -capacitary integrals with some applications.....	359
DAVID R. ADAMS	
Sobolev spaces, the Navier-Stokes equations and capacity.....	369
VICTOR L. SHAPIRO and GRANT V. WELLAND	
Approximation in $L^p$ by analytic and harmonic functions.....	377
LARS INGE HEDBERG	
On the Helson-Szegő theorem and a related class of modified Toeplitz kernels..	383
MISCHA COTLAR and CORA SADOSKY	
Some sharp inequalities for conjugate functions.....	409
ALBERT BAERNSTEIN II	

Constructions for $BMO(\mathbb{R})$ and $A_p(\mathbb{R}^n)$ .....	417
PETER W. JONES	
Structure of some subalgebra of $L^\infty$ of the torus.....	421
SUN-YUNG A. CHANG	
A geometric conditions which implies $BMOA$ .....	427
DAVID A. STEGENGA	
Proof of the Beurling-Malliavin theorem by duality and harmonic estimation....	431
PAUL KOOSIS	
Zero sets of absolutely convergent Taylor series.....	439
R. KAUFMAN	
Capacity and uniform algebras.....	445
J. WERMER	
Following functions of class $H^2$ .....	451
DOUGLAS N. CLARK	
A Hankel type operator arising in deformation theory.....	457
RICHARD ROCHBERG	
Representation theorems for holomorphic and harmonic functions.....	459
R. R. COIFMAN and R. ROCHBERG	



**Eli Stein and some of his students attending the Conference.**

*First row (left to right): David Goldberg, Eli Stein, Steve Wainger, Juan Peral;*

*Second row: Steve Krantz, Mitchell Taibleson, Daryl Geller, David Jerison;*

*Top row: Bill Beckner, Norman Weiss, Bob Fefferman, Charlie Fefferman.*



**A. Zygmund and some of his students attending the Conference.**

*First row (left to right): Antoni Zygmund, Mischa Cotlar, Eli Stein, Marshall Ash;*

*Second row: Guido Weiss, Eugene Fabes, Marvin Kohn;*

*Third row: Ben Muckenhoupt and Yoram Sagher;*

*Top row: Bill Connett, Dick Wheeden, Dan Waterman.*



**Conference participants who have been students or faculty  
at Washington University in St. Louis**

*First row:* Yoram Sagher, Alberto de la Torre, Jon Cohen;

*Second row:* Ronald Coifman and John Chao;

*Third row:* Michael Cwickel and Dick Hunt;

*Fourth row:* Yves Meyer, Antoni Zygmund (standing), Roberto Macias;

*Fifth row:* Guido Weiss, Pepe Garcia-Cuerva, Richard Rochberg, Ray Kunze;

*Sixth row:* Michel de Guzman and Ken Gross;

*Seventh row:* Rich Rubin, Al Baernstein, Mike Hemler, Fulvio Ricci, Mitch Taibleson;

*Top center:* Richard Bagby.

## PREFACE

A considerable development of harmonic analysis in  $\mathbb{R}^n$ , and related fields, occurred in the decade of the 1970's. New ideas emerged, old techniques were applied in novel ways, and the types of interactions of harmonic analysis with other parts of mathematics increased considerably. A comprehensive description of this activity would be quite long. It would have to include the following items:

A new function space, the functions of bounded mean oscillations (BMO), became as important as the Lebesgue and Hardy spaces. The latter, which once represented much of what was labelled as the "complex methods" in harmonic analysis, were given several real-variable characterizations and were studied by "real methods". This permitted one to introduce "Hardy spaces" in very general settings (spaces of homogeneous type). On the other hand, complex variable methods played a role in the study of real variable problems concerning Lipschitz domains. The Fourier transform became a powerful tool for obtaining estimates for operators which are neither of convolution type nor linear, such as Littlewood-Paley-Stein functions, various maximal operators, pseudo-differential operators, commutators and Fourier integral operators. Weighted norm inequalities became increasingly more significant. The increased interaction between harmonic analysis and other fields of mathematics involved the theory of martingales, nilpotent groups, questions of hypoellipticity for linear differential operators and for the study of function theory in strictly pseudo-convex domains in  $\mathbb{C}^n$ .

The 1978 American Mathematical Society Summer Institute in Williamstown was devoted to this development and these topics. These two volumes make up the proceedings of this conference; they contain three types of presentations: (i) papers suggesting problems for future research; (ii) expository papers that explain some of the new ideas and techniques introduced in harmonic analysis during the last decade; (iii) technical papers giving the latest reports on progress in various different topics.

We have organized this material in six chapters distributed over two volumes. The first volume contains, for the most part, papers dealing with analysis in  $\mathbb{R}^n$  and is divided into Chapter I, Real Harmonic Analysis, Chapter II, Hardy Spaces and BMO and Chapter III, Complex Harmonic Functions, Potential Theory and Functions of One Complex Variable. The second volume contains mostly papers on analysis in other settings. These papers give us a good picture of the strong interaction that has occurred between the various fields we mentioned above. This volume, again, is divided into three chapters: Chapter IV, Several Complex Variables, Chapter V, Pseudo Differential Operators and Partial Differential Equations and Chapter VI, Harmonic Analysis in Other Settings: Probability, Martingales, Local Fields, Lie Groups and Functional Analysis. The organization could well have been different since, in many cases, an individual article is connected with several topics discussed in other chapters so that it could easily have been placed elsewhere. We hope, however, that our selection makes these volumes reasonably accessible. We have also placed articles of a more expository nature at the beginning of the various sections or subsections.

We wish to thank the various people that have made this Summer Institute a success: Donald Sarason, Elias M. Stein, Mitchell H. Taibleson and Richard Wheeden, who have organized the institute's seminars; Dottie Smith, who did everything; Anna Pauline Bailey, who was particularly helpful during the organization period preceding the three weeks of the institute; our colleague, Ronald R. Coifman, who helped us throughout the organization and during the running of the institute; the very helpful staff supplied by Williams College: Peter Andrews, Marie Seitz, Eileen Sprague, and Jim Peck; the American Mathematical Society, who sponsored this meeting and the National Science Foundation who provided funds for the institute.

Stephen Wainger  
Department of Mathematics  
University of Wisconsin  
Madison, Wisconsin

Guido Weiss  
Department of Mathematics  
Washington University in St. Louis  
St. Louis, Missouri

