

Quantum curves for simple Hurwitz numbers of an arbitrary base curve

Xiaojun Liu, Motohico Mulase, and Adam Sorkin

ABSTRACT. Various generating functions of simple Hurwitz numbers of the projective line are known to satisfy many properties. They include a heat equation, a cut-and-join recursion, an infinite-order differential equation called a quantum curve equation, and a Schrödinger like partial differential equation. In this paper we generalize these properties to simple Hurwitz numbers with an arbitrary base curve. For projective line case, the equivalency between the cut-and-join recursion and the Chekhov-Eynard-Orantin topological recursion has been proved. However, the relation between these two recursions are not discussed for the simple Hurwitz number we considered here.

1. Introduction and the main results

The purpose of this paper is to determine functional properties of various generating functions of simple Hurwitz numbers with an arbitrary fixed base curve B . We derive a partial differential equation for the Laplace transform of these Hurwitz numbers. The equation is completely analogous to the result of [28] for the usual simple Hurwitz numbers based on \mathbb{P}^1 . We also obtain an infinite-order differential equation, or a *quantum curve*, for the case of an arbitrary base curve, that is parallel to the various Hurwitz problems studied in [3, 25, 26] with the base curve \mathbb{P}^1 and its orbifold version $\mathbb{P}^1[a]$.

Our main motivation is to examine whether the topological recursion of Chekhov-Eynard-Orantin [14, 15] and the existence of a quantum curve of [1, 7–9, 19] hold for the enumeration problem of simple Hurwitz numbers over an arbitrary curve B . Consider a holomorphic map $f : C \rightarrow B$ of a non-singular algebraic curve C onto a fixed base curve B . We choose a general point $0 \in B$ and fix it once for all. The quantity we are interested in this paper is the base B Hurwitz number $H_{g,n}^B(\mu_1, \dots, \mu_n)$, which counts the automorphism weighted number of the topological types of holomorphic maps f of a genus g domain curve with n labeled

2010 *Mathematics Subject Classification*. Primary 14H15, 14N35, 05C30, 11P21; Secondary 81T30.

The first author received the China Scholarship Council grant CSC-2010811063, which allowed him to conduct research at the Department of Mathematics, University of California, Davis. He is also supported by the National Science Foundation of China grants No.11201477, 11171175, 11471182 and the Chinese Universities Scientific Fund No.2011JS041.

The research of the second author is supported by NSF grant DMS-1104734.

The third author received support from the U.S. Government.

preimages of a fixed point of B with multiplicity $(\mu_1, \dots, \mu_n) \in \mathbb{Z}_+^n$, such that f is simply ramified apart from at these n points. Define its discrete Laplace transform

$$\begin{aligned}
 (1.1) \quad F_{g,n}^B(x_1, \dots, x_n) &= \sum_{\vec{\mu} \in \mathbb{Z}_+^n} H_{g,n}^B(\mu_1, \dots, \mu_n) \prod_{i=1}^n e^{-w_i \mu_i} \\
 &= \sum_{\vec{\mu} \in \mathbb{Z}_+^n} H_{g,n}^B(\mu_1, \dots, \mu_n) \prod_{i=1}^n x_i^{\mu_i}, \quad x_i = e^{-w_i},
 \end{aligned}$$

which we call the *free energy* of type (g, n) . Since \mathbb{P}^1 does not cover the base curve B of genus $g(B) > 0$,

$$F_{0,n}^B(\mu_1, \dots, \mu_n) = 0$$

for any $n \geq 1$ and any value of $(\mu_1, \dots, \mu_n) \in \mathbb{Z}_+^n$ in this case. Our main result is the following.

THEOREM 1.1. *For $2g - 2 + n > 0$, the free energies $F_{g,n}^B(x_1, \dots, x_n)$ satisfy the following partial differential equation:*

$$\begin{aligned}
 (1.2) \quad & \left(2g - 2 + n - (1 - \chi(B)) \sum_{i=1}^n x_i \frac{\partial}{\partial x_i} \right) F_{g,n}^B(x_{[n]}) \\
 &= \frac{1}{2} \sum_{i \neq j} \frac{x_i x_j}{x_i - x_j} \left(\frac{\partial}{\partial x_i} F_{g,n-1}^B(x_{[\hat{j}]}) - \frac{\partial}{\partial x_j} F_{g,n-1}^B(x_{[\hat{i}]}) \right) \\
 & \quad + \frac{1}{2} \sum_{i=1}^n u_1 \frac{\partial}{\partial u_1} u_2 \frac{\partial}{\partial u_2} \Big|_{u_1=u_2=x_i} \\
 & \quad \left[F_{g-1,n+1}^B(u_1, u_2, x_{[\hat{i}]}) + \sum_{\substack{g_1+g_2=g \\ I \sqcup J = [\hat{i}]}} F_{g_1,|I|+1}^B(u_1, x_I) F_{g_2,|J|+1}^B(u_2, x_J) \right].
 \end{aligned}$$

Here $[n] = \{1, \dots, n\}$ is an index set, $[\hat{i}] = [n] \setminus \{i\}$, and for any subset $I \subset [n]$, we denote $x_I = (x_i)_{i \in I}$. We denote by $\chi(B) = 2 - 2g(B)$ the Euler characteristic of the base curve B . Note that the complexity $2g - 2 + n$ is reduced by 1 on the right-hand side of the recursion when $g(B) \geq 1$, similar to the Chekhov-Eynard-Orantin integral recursion of [14, 15].

Let us define the **wave function** of the base B Hurwitz numbers by

$$(1.3) \quad Z^B(x, \hbar) = \exp \left(\sum_{g=1}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^B(x, x, \dots, x) \right).$$

Then it satisfies a **quantum curve** of infinite-order differential equation

$$(1.4) \quad \hbar x \frac{d}{dx} \left[1 - \hbar^{1-\chi(B)} x e^{\hbar x \frac{d}{dx}} \left(\frac{d}{dx} x \right)^{1-\chi(B)} \right] Z^B(x, \hbar) = 0.$$

If we introduce

$$y = \hbar x \frac{d}{dx}$$

and regard it as a commuting variable, then the total symbol of the above operator produces

$$(1.5) \quad x = y^{\chi(B)-1} e^{-y}.$$

We call (1.5) a **Lambert curve**, due to its similarity to the usual Lambert curve $x = ye^{-y}$. The wave function also satisfies a **Schrödinger** like partial differential equation

$$(1.6) \quad \left[\frac{\partial}{\partial \hbar} - \frac{1}{2} \left(x \frac{\partial}{\partial x} \right)^2 + \left(\frac{1}{2} - \frac{1 - \chi(B)}{\hbar} \right) x \frac{\partial}{\partial x} \right] Z^B(x, \hbar) = 0.$$

If we denote the above operators by P and Q , respectively, then they satisfy a commutation relation:

$$(1.7) \quad [P, Q] = -\frac{1}{\hbar} P.$$

This shows that the system of equations (1.4) and (1.6) are compatible. Moreover, the partition function has the following simple expression

$$(1.8) \quad Z^B(x, \hbar) = \sum_{m=0}^{\infty} (m!)^{1-\chi(B)} e^{\frac{1}{2}m(m-1)\hbar} \left(\hbar^{1-\chi(B)} x \right)^m.$$

REMARK 1.2. It is interesting to note here that the exact same formulas (1.2), (1.5), (1.6), and (1.8) hold for the case of an orbifold base $B = \mathbb{P}^1[a]$, $a > 0$, if we evaluate $\chi(B) = 1 + \frac{1}{a}$, interpret the sum in (1.8) as running over non-negative multiples am of a , and replace $(am)!$ by $(am)! \mapsto m!a^m$. Since the orbifold case requires a different set of preparations (see [3, 10, 25]), we will report the generalization to the case of higher-genus orbifold curve as a base elsewhere.

When the base curve B is an elliptic curve E , a special case of simple Hurwitz numbers exhibits a *quasi-modular* properties. Dijkgraaf [6] considered a generating function

$$(1.9) \quad F_g(q) = \sum_{n=1}^{\infty} \frac{1}{n!} H_{g,n}^E(1, 1, \dots, 1) q^n$$

for $g > 1$, and

$$(1.10) \quad F_1(q) = -\frac{1}{24} \log q + \sum_{n=1}^{\infty} \frac{1}{n!} H_{1,n}^E(1, 1, \dots, 1) q^n.$$

The significant fact here is that $F_g(q)$ for $g > 1$ is a quasi-modular form [6, 21]. The relation between the free energies of type (g, n) for all n and $F_g(q)$ is given by

$$(1.11) \quad F_g(q) = \lim_{\lambda \rightarrow 0} \sum_{n=1}^{\infty} \frac{1}{n!} \frac{1}{\lambda^n} F_{g,n}^E(\lambda q, \lambda q, \dots, \lambda q).$$

To motivate the present work, let us recall the corresponding counting problem of simple Hurwitz number $H_{g,n}^{\mathbb{P}^1}(\mu_1, \dots, \mu_n)$ for a pointed projective line (\mathbb{P}^1, ∞) , which has a long history since Hurwitz [20]. Its modern interest is due, among other things, to its rich functional properties [16, 17, 22, 32, 33], its relation with linear Hodge integrals on $\overline{\mathcal{M}}_{g,n}$ [12, 18, 30], and the Bouchard-Mariño conjecture [5] and

its solutions [2, 13, 28]. The theorem established in [13, 28] shows that the discrete Laplace transform

$$(1.12) \quad F_{g,n}^{\mathbb{P}^1}(x_1, \dots, x_n) = \sum_{\vec{\mu} \in \mathbb{Z}^n_+} H_{g,n}^{\mathbb{P}^1}(\mu_1, \dots, \mu_n) \prod_{i=1}^n x_i^{\mu_i}$$

satisfies an integral recursion formula that was originally proposed by Chekhov, Eynard and Orantin [14, 15], as conjectured in [5]. The input curve for the recursion, the *spectral curve*, is shown to be the original Lambert curve

$$(1.13) \quad x = ye^{-y}.$$

Then in [26, 34] it is discovered that there is a quantum curve of [1, 7–9, 19], which is a generator of the holonomic system that characterizes the wave function of the simple Hurwitz numbers of \mathbb{P}^1 . The wave function for $B = \mathbb{P}^1$ is

$$Z^{\mathbb{P}^1}(x, \hbar) = \exp \left(\sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^{\mathbb{P}^1}(x, x, \dots, x) \right).$$

It is established in [26] that the \mathbb{P}^1 wave function has an expression

$$(1.14) \quad Z^{\mathbb{P}^1}(x, \hbar) = \sum_{m=0}^{\infty} \frac{1}{m!} e^{\frac{1}{2}m(m-1)\hbar} \left(\frac{x}{\hbar}\right)^m,$$

and that it satisfies two equations:

$$(1.15) \quad \left[\hbar x \frac{\partial}{\partial x} - x e^{\hbar x \frac{\partial}{\partial x}} \right] Z^{\mathbb{P}^1}(x, \hbar) = 0,$$

$$(1.16) \quad \left[\frac{\partial}{\partial \hbar} - \frac{1}{2} \left(x \frac{\partial}{\partial x} \right)^2 + \left(\frac{1}{2} + \frac{1}{\hbar} \right) x \frac{\partial}{\partial x} \right] Z^{\mathbb{P}^1}(x, \hbar) = 0.$$

If we denote the above operators as P_1 and Q_1 , respectively, then they satisfy a commutation relation

$$(1.17) \quad [P_1, Q_1] = -\frac{1}{\hbar} P_1.$$

The semi-classical limit of each of the equations (1.15) and (1.16) recovers the Lambert curve (1.13), as shown in [26]. Then from this curve as the spectral curve, and using the functions x and y on it, the Chekhov-Eynard-Orantin integral recursion [14] determines the differential forms

$$(1.18) \quad d_1 d_2 \cdots d_n F_{g,n}^{\mathbb{P}^1}$$

defined on Σ^n for all (g, n) , where Σ is the Lambert curve given by (1.13). It is a simple consequence of the results in [13, 26, 28] that the differential form (1.18) uniquely recovers the primitive $F_{g,n}^{\mathbb{P}^1}$ as a function in $(y_1, \dots, y_n) \in \Sigma^n$. From this point of view, the generating function $F_{g,n}^{\mathbb{P}^1}(x(y_1), \dots, x(y_n))$ is completely determined by the equation (1.15) or (1.16), where x as a function in y is also given by (1.13).

Our motivating question is, *do the similar properties hold when we consider the simple Hurwitz numbers with an arbitrary curve as the base?* Since the genus 0 base B Hurwitz numbers $H_{0,n}^E(\mu_1, \dots, \mu_n)$ do not exist for a base curve B with $g(B) > 0$, the philosophy of [11] does not produce any spectral curve of the Chekhov-Eynard-Orantin integral recursion for this counting problem. Yet we have a topological

recursion (1.2) in the form of a partial differential equation. We note the straightforward generalizations (1.4), (1.6), (1.8), and (1.5), that reduce to the \mathbb{P}^1 case (1.15), (1.16), (1.14), and (1.13), respectively, when $\chi(B) = 2$.

The paper is organized as follows. In Section 2, we derive the cut-and-join equation of simple Hurwitz numbers with an arbitrary base curve. Then in Section 3, we consider the Laplace transform of the cut-and-join equation, which is exactly (1.2). Using this result we derive the Schrödinger equation (1.6) in Section 4. Section 5 is devoted to explaining the heat equation expression of the cut-and-join equation [16, 22, 33] and deriving its consequences. In Section 6 we first prove the expansion (1.8), derive the quantum curve (1.4), and then establish the commutator relation (1.7). In the final section we deduce the Lambert curve (1.5) as the semi-classical limit.

2. A cut-and-join equation for simple Hurwitz numbers

The enumeration problem we consider in this paper is the number of topological types of holomorphic maps $f : C \rightarrow B$ from a nonsingular curve C of genus g to a fixed base curve B , with an arbitrary ramification over one point on B and simple ramification at all other critical points. Let us denote by $0 \in B$ a general point arbitrarily chosen and fixed. Such a holomorphic map f is referred to as a *simple Hurwitz cover* of B . We label each inverse image of $0 \in B$ via f , and denote by $(\mu_1, \dots, \mu_n) \in \mathbb{Z}_+^n$ the orders of f at each inverse image of 0 . The base B Hurwitz number $H_{g,n}^B(\mu_1, \dots, \mu_n)$ counts the automorphism weighted number of the topological types of simple Hurwitz covers.

The degree of the map f is given by

$$(2.1) \quad d = |\mu| = \sum_{i=1}^n \mu_i.$$

Here use the notation $|\mu|$, borrowing from the convention in the theory of symmetric functions. The Riemann-Hurwitz formula tells us that there are

$$(2.2) \quad r = r(g, \mu) = 2g - 2 + n - d(2g(B) - 1)$$

simple ramification points. Instead of counting the *topological types* of f , we can fix r simple branch points on B other than $0 \in B$ and count the automorphism weighted holomorphic maps f . We note that (2.1) and (2.2) imply a condition for the degree and the genus for a base curve with $g(B) \geq 1$:

$$(2.3) \quad n(2g(B) - 1) \leq d(2g(B) - 1) \leq 2g - 2 + n,$$

$$(2.4) \quad n(g(B) - 1) \leq g - 1.$$

In particular, $H_{g,n}(\vec{\mu}) = 0$ for any $n \geq 1$ if $g < g(B)$. This poses a sharp contrast to the Gromov-Witten theory of curves [31].

The map $f : C \rightarrow B$ of degree d is determined by a monodromy representation

$$\rho \in \text{Hom}(\pi_1(B \setminus \{0, 1, \dots, r\}), S_d)$$

of the fundamental group of an $(r + 1)$ -punctured curve into the symmetric group of d letters. Here $\{1, \dots, r\} \subset B$ is the label of r simple branched points on B . Let $\gamma_j, j = 0, \dots, r$, be based simple loops around each point $j \in B$. Then a simple Hurwitz cover is constructed by assigning each γ_j for $j \geq 1$ to a transposition and

γ_0 to a product of n disjoint cycles of length μ_1, \dots, μ_n , subject to the commutator relation

$$\rho(\gamma_1) \cdots \rho(\gamma_r) \rho(\gamma_0) = [\alpha_1, \beta_1] \cdots [\alpha_{g(B)}, \beta_{g(B)}],$$

where $\alpha_i, \beta_i \in S_d$ are some elements in the permutation group.

The *cut-and-join equation* [16, 17, 20, 32] is the result of an analysis of what happens when we multiply a transposition $\rho(\gamma_r)$ to a product of disjoint cycles $\rho(\gamma_0)$. Since the fundamental group of the base curve does not make any effect on this multiplication, the exact same formula for $H_{g,n}^B(\vec{\mu})$ holds, as for the simple Hurwitz numbers for \mathbb{P}^1 . The only difference is the number r of simple ramification points.

PROPOSITION 2.1 (The cut-and-join equation). *For all $g \geq 0, n \geq 1$, and $(\mu_1, \dots, \mu_n) \in \mathbb{Z}_+^n$ subject to*

$$(2.5) \quad 2g - 2 + n - (2g(B) - 1) \sum_{i=1}^n \mu_i > 0,$$

the simple Hurwitz numbers $H_{g,n}^B(\mu_1, \dots, \mu_n)$ of degree d satisfy the following equation:

$$(2.6) \quad (2g - 2 + n - (2g(B) - 1)d) H_{g,n}^B(\mu_{[n]}) = \frac{1}{2} \sum_{i \neq j} (\mu_i + \mu_j) H_{g,n-1}^B(\mu_i + \mu_j, \mu_{[\hat{i}, \hat{j}]}) + \frac{1}{2} \sum_{i=1}^n \sum_{\alpha + \beta = \mu_i} \alpha \beta \left(H_{g-1, n+1}^B(\alpha, \beta, \mu_{[\hat{i}]}) + \sum_{\substack{g_1 + g_2 = g \\ I \sqcup J = [\hat{i}]}} H_{g_1, |I|+1}^B(\alpha, \mu_I) H_{g_2, |J|+1}^B(\beta, \mu_J) \right).$$

Here and throughout the paper we use the following notational convention. $[n] = \{1, \dots, n\}$ is an index set, and $[\hat{i}] = [n] \setminus \{i\}$, etc. For a subset $I \subset [n]$, $\mu_I = (\mu_i)_{i \in I}$.

REMARK 2.2. Unlike the case of \mathbb{P}^1 , the cut-and-join equation does not determine all values of simple Hurwitz numbers. When the degree d takes its maximum value $\frac{2g-2+n}{2g(B)-1}$, the equation gives a trivial equality $0 = 0$.

The genus 1 simple Hurwitz numbers based on an elliptic curve $B = E$ are easy to calculate. From (2.3) we have $n = d$ and $r = 0$ when $g = g(B) = 1$. Therefore, the covering is unramified. If we allow disconnected domain, then the total number is equal to the number of partitions

$$p(d) = |\text{Hom}(\pi_1(E), S_d) // S_d|$$

of degree d (see for example, [6]). Let

$$\phi(q) = \prod_{m=1}^{\infty} (1 - q^m)$$

be the Euler function. Then we have

$$\sum_{n=1}^{\infty} \frac{1}{n!} H_{1,n}^E(1, \dots, 1) q^n = -\log \phi(q) = \sum_{n=1}^{\infty} \left(\sum_{m|n} \frac{1}{m} \right) q^n,$$

hence

$$(2.7) \quad H_{1,n}^E(1, \dots, 1) = n! \sum_{m|n} \frac{1}{m} = (n-1)! \sigma(n),$$

where σ is the sum of divisors function. This is an integer sequence, and its first ten terms are 1, 3, 8, 42, 144, 1440, 5760, 75600, 524160, 6531840. This sequence has many interesting properties (see OEIS, A038048).

More generally, let us consider the case when the equality holds in (2.4). We need this analysis in Section 4. Because of (2.3), $g-1 = n(g(B)-1)$ implies $d = n$ and $r = 0$, hence the covering $f : C \rightarrow B$ that is counted is totally unramified. Here again the number of disconnected unramified coverings is given by the classical dimension formula (see for example [27] for elementary derivations):

$$|\text{Hom}(\pi_1(B), S_d) // S_d| = \sum_{\lambda \vdash d} \left(\frac{\dim \lambda}{d!} \right)^{\chi(B)}.$$

Here $\lambda \vdash d$ parametrizes irreducible representations of S_d , and $\dim \lambda$ is its dimension.

3. The discrete Laplace transform

The discrete Laplace transform $F_{g,n}^B(x_1, \dots, x_n)$ of (1.1) is a polynomial of degree $\frac{2g-2+n}{2g(B)-1}$ with the lowest degree term $H_{g,n}^E(1, 1, \dots, 1)x_1 \cdots x_n$. The following proposition is an analogue of the case of simple Hurwitz numbers of \mathbb{P}^1 , and the proof is exactly the same as that of [3, Lemma 4.1].

PROPOSITION 3.1. *The discrete Laplace transform of the cut-and-join equation (2.6) is precisely the partial differential equation (1.2).*

Reflecting Remark 2.2, the differential equation (1.2) alone does not determine free energies $F_{g,n}^B(x_1, \dots, x_n)$. This is because every homogeneous function of degree $\frac{2g-2+n}{2g(B)-1}$ is in the kernel of the Euler differential operator

$$2g - 2 + n - (2g(B) - 1) \sum_{i=1}^n x_i \frac{\partial}{\partial x_i}$$

on the left-hand side of (1.2). Thus the highest degree terms of free energies are not determined by this differential equation. We will determine the homogeneous highest degree terms of the free energies in a different method in Section 5.

The genus 1 elliptic Hurwitz numbers (2.7) yields

$$(3.1) \quad F_{1,n}^E(x_1, \dots, x_n) = \left(n! \sum_{m|n} \frac{1}{m} \right) x_1 \cdots x_n.$$

4. A Schrödinger equation

In this section we prove (1.6), provided that $g(B) > 0$. We remark here that for $g(B) = 0$, the proof is rather different [26], yet the same formula holds.

THEOREM 4.1. *The wave function (1.3) satisfies the Schrödinger-type equation (1.6):*

$$\left[\frac{\partial}{\partial \hbar} - \frac{1}{2} \left(x \frac{\partial}{\partial x} \right)^2 + \left(\frac{1}{2} - \frac{2g(B) - 1}{\hbar} \right) x \frac{\partial}{\partial x} \right] Z^B(x, \hbar) = 0.$$

PROOF. We use the same method of the proof of [26, Theorem 5.1, Theorem 5.3, Appendix A]. When (2.3) holds, the diagonal evaluation of (1.2) yields

$$\begin{aligned} (4.1) \quad & \left(2g - 2 + n - (2g(B) - 1)x \frac{d}{dx} \right) F_{g,n}^B(x, \dots, x) \\ &= \frac{n(n-1)}{2} x^2 \frac{\partial^2}{\partial u^2} \Big|_{u=x} F_{g,n-1}^B(u, x, \dots, x) \\ &+ \frac{n}{2} x^2 \frac{\partial^2}{\partial u_1 \partial u_2} \Big|_{u_1=u_2=x} F_{g-1,n+1}^B(u_1, u_2, x, \dots, x) \\ &+ \frac{n!}{2} \sum_{\substack{g_1+g_2=g \\ n_1+n_2=n-1}} \frac{x^2}{(n_1+1)!(n_2+1)!} \frac{d}{dx} F_{g_1,n_1+1}^B(x, \dots, x) \cdot \frac{d}{dx} F_{g_2,n_2+1}^B(x, \dots, x). \end{aligned}$$

For $m \geq 2g(B)$, let us define

$$S_m(x) := \sum_{2g-2+n=m-1} \frac{1}{n!} F_{g,n}^B(x, \dots, x).$$

Note that S_m contains a contribution from domain curves with $g-1 = n(g(B)-1)$, or equivalently, $d = n$ and $r = 0$, for which the cut-and-join equation is not valid. Therefore, when we deduce an equation for S_m 's, we need to remove these boundary terms from the equation. Now from (4.1) we obtain

$$\begin{aligned} (4.2) \quad & \left(m - (2g(B) - 1)x \frac{d}{dx} \right) S_{m+1}(x) \\ & - \frac{1}{2} x^2 \frac{d^2}{dx^2} S_m(x) - \frac{1}{2} \sum_{m_1+m_2=m+1} x \frac{d}{dx} S_{m_1}(x) \cdot x \frac{d}{dx} S_{m_2}(x) \\ &= \sum_{\substack{2g-2+n=m \\ g-1=n(g(B)-1)}} \frac{1}{n!} \left(m - (2g(B) - 1)x \frac{d}{dx} \right) F_{g,n}^B(x, \dots, x) \\ & - \frac{1}{2} \sum_{\substack{2g-2+n=m-1 \\ g-1=(n+1)(g(B)-1)}} \frac{1}{(n-1)!} x^2 \frac{\partial^2}{\partial u^2} \Big|_{u=x} F_{g,n}^B(u, x, \dots, x) \\ & - \frac{1}{2} \sum_{\substack{2g-2+n=m-1 \\ g=(n-1)(g(B)-1)}} \frac{1}{(n-2)!} x^2 \frac{\partial^2}{\partial u_1 \partial u_2} \Big|_{u_1=u_2=x} F_{g,n}^B(u_1, u_2, x, \dots, x) \\ & - \frac{1}{2} \sum_{\substack{2g-2+n=m \\ g-1=n(g(B)-1)}} \sum_{\substack{g_1+g_2=g \\ n_1+n_2=n+1}} \frac{x^2}{n_1! n_2!} \left(\frac{d}{dx} F_{g_1,n_1}^B(x, \dots, x) \right) \left(\frac{d}{dx} F_{g_2,n_2}^B(x, \dots, x) \right). \end{aligned}$$

We note here that in the boundary contribution $g-1 = n(g(B)-1)$, since $n = d$, the free energies are single monomials proportional to $x_1 x_2 \cdots x_n$. Let us look

at the right-hand side of (4.2). The first line of the right-hand side is 0 because $F_{g,n}^B(x, \dots, x) = H_{g,n}^B(1, \dots, 1)x^n$, and $m = 2g - 2 + n = n(2g(B) - 1)$, since $r = 0$. The second line is also 0, because $F_{g,n}^B(u, x, \dots, x)$ is linear in u .

The third summation on the right-hand side is empty, because we need $n(g(B) - 1) \leq g - 1 = (n - 1)(g(B) - 1) - 1$, which does not happen. Similarly, the fourth line summation is also empty, because we are requiring $g = g_1 + g_2, n + 1 = n_1 + n_2$, and

$$\begin{aligned} n_1(g(B) - 1) &\leq g_1 - 1 \\ n_2(g(B) - 1) &\leq g_2 - 1 \\ n(g(B) - 1) &= g - 1. \end{aligned}$$

We have thus obtained a recursion equation

$$(4.3) \quad \left(m - (2g(B) - 1)x \frac{d}{dx} \right) S_{m+1}(x) = \frac{1}{2}x^2 \frac{d^2}{dx^2} S_m(x) + \frac{1}{2} \sum_{m_1+m_2=m+1} x \frac{d}{dx} S_{m_1}(x) \cdot x \frac{d}{dx} S_{m_2}(x).$$

In terms of the generating function

$$F(x, \hbar) = \sum_{m=1}^{\infty} \hbar^{m-1} S_m(x),$$

(4.3) becomes

$$(4.4) \quad \left(\frac{\partial}{\partial \hbar} - \frac{2g(B) - 1}{\hbar} x \frac{d}{dx} \right) F(x, \hbar) = \frac{1}{2}x^2 \frac{d^2}{dx^2} F(x, \hbar) + \frac{1}{2} \left(x \frac{d}{dx} F(x, \hbar) \right)^2.$$

Since $Z^B(x, \hbar) = \exp F(x, \hbar)$, (1.6) follows directly from (4.4). This completes the proof. □

Using the Schrödinger equation (1.6), we can determine the form of the solution $Z^B(x, \hbar)$.

LEMMA 4.2. *The wave function has the following expansion:*

$$(4.5) \quad Z^B(x, \hbar) = \sum_{m=0}^{\infty} c_m e^{\frac{1}{2}m(m-1)\hbar} \left(\hbar^{1-\chi(B)} x \right)^m,$$

where c_m is a constant.

PROOF. The wave function $Z^B(x, \hbar)$ of (1.3) is a formal power series in x and \hbar . Thus it has an expansion of the form

$$Z^B(x, \hbar) = \sum_{m=0}^{\infty} c_m f_m(\hbar) x^m$$

with $f_m(\hbar) \in \mathbb{C}[[\hbar]]$. Then the Schrödinger equation (1.6) yields an ordinary differential equation

$$f'_m = \left(\frac{1}{2}m(m - 1) + m \frac{1 - \chi(B)}{\hbar} \right) f_m,$$

whose solution is

$$f_m(\hbar) = c e^{\frac{1}{2}m(m-1)\hbar} \hbar^m (1 - \chi(B)).$$

This completes the proof. □

5. The heat equation and its consequences

As we have seen in Section 4, the Schrödinger equation (1.6) determines the solution $Z^B(x, \hbar)$ only up to the form (4.5). To determine the coefficients, we need another technique.

In this section we use a heat equation technique of [22]. First we remark that the cut-and-join equation gives rise to a heat equation for another generating function of base B Hurwitz numbers. To determine a solution of a heat equation, we need to identify the initial value. We will show that the initial condition exactly corresponds to determining the highest degree terms of $F_{g,n}^B(x_1, \dots, x_n)$. The exponential generating function of these highest degree terms can be determined by a character formula of [31]. Thus we obtain the unique solution of the heat equation, which in turn gives all base B Hurwitz numbers.

The generating function for base B Hurwitz numbers we consider is

$$(5.1) \quad \mathbf{H}(t, \mathbf{p}) = \sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \mathbf{H}_{g,n}(t, \mathbf{p}),$$

$$(5.2) \quad \mathbf{H}_{g,n}(t, \mathbf{p}) = \frac{1}{n!} \sum_{\vec{\mu} \in \mathbb{Z}_+^n} H_{g,n}^B(\vec{\mu}) \mathbf{p}_{\mu} t^{r(g,\mu)},$$

where $r(g, \mu) = 2g - 2 + n - |\mu|(1 - \chi(B))$ is the number of simple ramification points, and $\mathbf{p}_{\mu} = p_{\mu_1} \cdots p_{\mu_n}$. The same argument of [16, 22, 23, 29, 33] shows that $e^{\mathbf{H}(t, \mathbf{p})}$ satisfies a *heat equation*, that is obtained from the cut-and-join equation. For a partition $\mu = (\mu_1 \geq \mu_2 \geq \dots)$ of a finite length $\ell(\mu)$, we define the **shifted power-sum function** by

$$(5.3) \quad \mathbf{p}_r[\mu] := \sum_{i=1}^{\infty} \left[\left(\mu_i - i + \frac{1}{2} \right)^r - \left(-i + \frac{1}{2} \right)^r \right].$$

This is a finite sum of $\ell(\mu)$ terms. Then we have [16, 33]

$$(5.4) \quad \sum_{i,j \geq 1} \left((i+j)p_i p_j \frac{\partial}{\partial p_{i+j}} + ij p_{i+j} \frac{\partial^2}{\partial p_i \partial p_j} \right) s_{\mu}(\mathbf{p}) = \mathbf{p}_2[\mu] \cdot s_{\mu}(\mathbf{p}),$$

where $s_{\mu}(\mathbf{p})$ is the Schur function defined by

$$(5.5) \quad s_{\mu}(\mathbf{p}) = \sum_{|\lambda|=|\mu|} \frac{\chi_{\mu}(\lambda)}{z_{\lambda}} \mathbf{p}_{\lambda}, \quad z_{\mu} = \prod_{i=1}^{\ell(\mu)} m_i! i^{m_i},$$

m_i = the number of parts in μ of length i , and $\chi_{\mu}(\lambda)$ is the value of the irreducible character of the representation μ of the symmetric group evaluated on any element of the conjugacy class described by λ . Let us denote the *cut-and-join* operator by

$$\Delta = \frac{1}{2} \sum_{i,j \geq 1} \left((i+j)p_i p_j \frac{\partial}{\partial p_{i+j}} + ij p_{i+j} \frac{\partial^2}{\partial p_i \partial p_j} \right).$$

Then Schur functions are eigenfunctions of the operator:

$$\Delta s_{\mu}(\mathbf{p}) = \frac{1}{2} \mathbf{p}_2[\mu] \cdot s_{\mu}(\mathbf{p}),$$

and the cut-and-join equation of simple Hurwitz numbers (2.6) yields a heat equation

$$(5.6) \quad \frac{\partial}{\partial t} e^{\mathbf{H}(t, \mathbf{p})} = \Delta e^{\mathbf{H}(t, \mathbf{p})}.$$

We can solve a heat equation by the method of eigenfunction expansion. Thus we have an expression

$$(5.7) \quad e^{\mathbf{H}(t, \mathbf{p})} = \sum_{\mu} a_{\mu} s_{\mu}(\mathbf{p}) e^{\frac{1}{2} \mathbf{p}_2[\mu] t}$$

for a constant a_{μ} associated with every partition μ . These constants are determined by the initial value $t = 0$. From (5.2) we see that the initial value comes from the cases when $r(g, \mu) = 0$.

First we note that (2.2) implies that simple Hurwitz numbers with $r = 0$ correspond to the case with only 1 branched point on the base curve B . If we allow disconnected domain curves to cover B , then the number of coverings of degree d with a prescribed ramification data given by a partition $\mu \vdash d$ over $0 \in B$ with no other ramification points is easy to calculate. Let us denote by $H_d^{B\bullet}(\mu)$ such a number, where we do not label the inverse images of $0 \in B$ this time. Then from [31, Eq.(0.10)] we obtain

$$(5.8) \quad H_d^{B\bullet}(\mu) = \sum_{\lambda \vdash d} \left(\frac{\dim \lambda}{d!} \right)^{\chi(B)} |C_{\mu}| \frac{\chi_{\lambda}(\mu)}{\dim \lambda}.$$

Here C_{μ} is the conjugacy class of the permutation group S_d determined by the cycle type μ , a partition $\lambda \vdash d$ is a label of an irreducible representation of S_d , and $\dim \lambda$ is its dimension. The cardinality of the conjugacy class is given by

$$(5.9) \quad |C_{\mu}| = \frac{d!}{\prod_i m_i! i^{m_i}} = \frac{d!}{z_{\mu}}.$$

The generating function of these disconnected simple Hurwitz numbers can be calculated, appealing to (5.5) and (5.9), as follows:

$$(5.10) \quad \begin{aligned} \sum_{d=1}^{\infty} \sum_{\mu \vdash d} H_d^{B\bullet}(\mu) \mathbf{p}_{\mu} &= \sum_{d=1}^{\infty} \sum_{\mu \vdash d} \sum_{\lambda \vdash d} \left(\frac{\dim \lambda}{d!} \right)^{\chi(B)} |C_{\mu}| \frac{\chi_{\lambda}(\mu)}{\dim \lambda} \mathbf{p}_{\mu} \\ &= \sum_{d=1}^{\infty} \sum_{\lambda \vdash d} \frac{d!}{\dim \lambda} \left(\frac{\dim \lambda}{d!} \right)^{\chi(B)} \sum_{\mu \vdash d} \frac{\chi_{\lambda}(\mu)}{z_{\mu}} \mathbf{p}_{\mu} \\ &= \sum_{\lambda} \left(\frac{|\lambda|!}{\dim \lambda} \right)^{1-\chi(B)} s_{\lambda}(\mathbf{p}), \end{aligned}$$

where the last sum runs over all partitions λ .

Note that $e^{\mathbf{H}(t, \mathbf{p})}$ is the generating function of simple Hurwitz numbers allowing disconnected domain curves. Therefore, the initial value $e^{\mathbf{H}(0, \mathbf{p})}$ counts the disconnected base B Hurwitz numbers with only one branched point at $0 \in B$. In other words, we have determined the initial condition by (5.10). The result is

$$(5.11) \quad e^{\mathbf{H}(0, \mathbf{p})} = \sum_{\mu} \left(\frac{|\mu|!}{\dim \mu} \right)^{1-\chi(B)} s_{\mu}(\mathbf{p}) = \sum_{\mu} a_{\mu} s_{\mu}(\mathbf{p}).$$

Since Schur functions are linear basis for symmetric functions, we establish the following.

THEOREM 5.1. *The exponential generating function of the base B simple Hurwitz numbers is given by*

$$(5.12) \quad e^{\mathbf{H}(t, \mathbf{p})} = \sum_{\mu} \left(\frac{|\mu|!}{\dim \mu} \right)^{1-\chi(B)} s_{\mu}(\mathbf{p}) e^{\frac{1}{2} \mathbf{p}_2[\mu]t}.$$

6. The quantum curve

As noted in [26], the diagonal specialization $F_{g,n}^E(x, x, \dots, x)$ corresponds to substituting the power-sum symmetric function

$$(6.1) \quad p_j = x_1^j + x_2^j + x_3^j + \dots$$

by its principal specialization $p_j = x^j$. More precisely, we have the following.

LEMMA 6.1. *Let us define the **principal specialization** of the power-sum symmetric functions by*

$$p_j(s) = \left(s^{1-\chi(B)} x \right)^j$$

$$\mathbf{p}_{\mu}(s) = p_{\mu_1}(s) \cdots p_{\mu_n}(s) = \left(s^{1-\chi(B)} x \right)^{|\mu|}.$$

Then we have

$$(6.2) \quad Z^B(x, \hbar) = e^{\mathbf{H}(\hbar, \mathbf{p}(\hbar))}.$$

PROOF. We have

$$\begin{aligned} \mathbf{H}(\hbar, \mathbf{p}(\hbar)) &= \sum_{g,n \geq 1} \sum_{\mu \in \mathbb{Z}_+^n} \frac{1}{n!} \hbar^{2g-2+n-|\mu|} \binom{1-\chi(B)}{1-\chi(B)} H_{g,n}^B(\mu_1, \dots, \mu_n) \mathbf{p}_{\mu}(\hbar) \\ &= \sum_{g,n \geq 1} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^E(x, x, \dots, x), \end{aligned}$$

which yields (6.2). □

From the expansion formulas (5.12) and (4.5), together with the equality (6.2), we obtain

$$(6.3) \quad \sum_{\mu} \left(\frac{|\mu|!}{\dim \mu} \right)^{1-\chi(B)} e^{\frac{1}{2} \mathbf{p}_2[\mu] \hbar} s_{\mu}(\mathbf{p}(\hbar)) = \sum_{m=0}^n c_m e^{\frac{1}{2} m(m-1) \hbar} \left(\hbar^{1-\chi(B)} x \right)^m.$$

As explained in [26, Section 6] and also in [25], this equality is exactly the effect of the principal specialization of Lemma 6.1, which turns the summation over all partitions into the sums over just one-row partitions. We have thus determined the coefficients c_m in the expansion (4.5). It is given by

$$(6.4) \quad c_m = (m!)^{1-\chi(B)}.$$

This completes the proof of (1.8).

The expansion (1.8) allows us to derive the quantum curve-type equation (1.4). Since

$$\begin{aligned} \left[\hbar^{1-\chi(B)} x e^{\hbar x \frac{d}{dx}} \left(\frac{d}{dx} x \right)^{1-\chi(B)} \right] (m!)^{1-\chi(B)} e^{\frac{1}{2} m(m-1)\hbar} \left(\hbar^{1-\chi(B)} x \right)^m \\ = ((m+1)!)^{1-\chi(B)} e^{\frac{1}{2} m(m+1)\hbar} \left(\hbar^{1-\chi(B)} x \right)^{m+1} \end{aligned}$$

for every $m \geq 0$, we have

$$(6.5) \quad \left(1 - \hbar^{1-\chi(B)} x e^{\hbar x \frac{d}{dx}} \left(\frac{d}{dx} x \right)^{1-\chi(B)} \right) Z^B(x, \hbar) = 1.$$

By differentiating (6.5) we obtain (1.4).

LEMMA 6.2. *Let*

$$(6.6) \quad P = \hbar x \frac{d}{dx} \left(1 - \hbar^{1-\chi(B)} x e^{\hbar x \frac{d}{dx}} \left(\frac{d}{dx} x \right)^{1-\chi(B)} \right),$$

$$(6.7) \quad Q = \frac{\partial}{\partial \hbar} - \frac{1}{2} \left(x \frac{\partial}{\partial x} \right)^2 + \left(\frac{1}{2} - \frac{1-\chi(B)}{\hbar} \right) x \frac{\partial}{\partial x}.$$

Then

$$[P, Q] = -\frac{1}{\hbar} P.$$

PROOF. We first note that Q commutes with $x \frac{d}{dx}$ and $\frac{d}{dx} x$. Since

$$\begin{aligned} & \left[\hbar^{1-\chi(B)} x e^{\hbar x \frac{d}{dx}}, Q \right] \\ &= \left[\hbar^{1-\chi(B)} x e^{\hbar x \frac{d}{dx}}, \frac{\partial}{\partial \hbar} \right] + \left[\hbar^{1-\chi(B)} x, -\frac{1}{2} \left(x \frac{\partial}{\partial x} \right)^2 + \left(\frac{1}{2} - \frac{1-\chi(B)}{\hbar} \right) x \frac{\partial}{\partial x} \right] e^{\hbar x \frac{d}{dx}} \\ &= -(1-\chi(B)) \hbar^{-\chi(B)} x e^{\hbar x \frac{d}{dx}} - \hbar^{1-\chi(B)} x^2 \frac{d}{dx} e^{\hbar x \frac{d}{dx}} \\ & \quad + \hbar^{1-\chi(B)} x^2 \frac{d}{dx} e^{\hbar x \frac{d}{dx}} + \frac{1}{2} \hbar^{1-\chi(B)} x e^{\hbar x \frac{d}{dx}} - \hbar^{1-\chi(B)} \left(\frac{1}{2} - \frac{1-\chi(B)}{\hbar} \right) x e^{\hbar x \frac{d}{dx}} \\ &= 0, \end{aligned}$$

(1.7) follows from

$$0 = \left[\frac{1}{\hbar} P, Q \right] = \frac{1}{\hbar} [P, Q] + \left[\frac{1}{\hbar}, Q \right] P = \frac{1}{\hbar} [P, Q] + \frac{1}{\hbar^2} P.$$

□

7. Semi-classical limit

The semi-classical analysis of the operators P and Q are performed in the following way. Suppose our counting problem had genus 0 contributions $F_{0,n}^B(x_1, \dots, x_n)$. Then we define

$$(7.1) \quad S_m(x) := \sum_{2g-2+n=m-1} \frac{1}{n!} F_{g,n}^B(x, \dots, x)$$

as before, and consider a formal expression

$$(7.2) \quad \overline{Z}^B(x, \hbar) = e^{\sum_{m=0}^{\infty} \hbar^{m-1} S_m(x)} = e^{\frac{1}{\hbar} S_0(x) + S_1(x)} Z^B(x, \hbar).$$

Let us introduce a variable u such that $x = e^u$, and regard the coefficients S_m as functions in u . Since

$$x \frac{d}{dx} = \frac{d}{du},$$

we have

$$P = \hbar \frac{d}{du} \left(1 - \hbar^{1-\chi(B)} e^u e^{\hbar \frac{d}{du}} \left(1 + \frac{d}{du} \right)^{1-\chi(B)} \right).$$

Then

$$(7.3) \quad \begin{aligned} e^{-S_1} e^{-\frac{1}{\hbar} S_0} P e^{\frac{1}{\hbar} S_0} e^{S_1} &= S'_0 - e^u (S'_0)^{2-\chi(B)} e^{\frac{1}{\hbar} (S_0(u+\hbar) - S_0(u))} + O(\hbar) \\ &= S'_0 - e^u (S'_0)^{2-\chi(B)} e^{S'_0} + O(\hbar), \end{aligned}$$

where $' = \frac{d}{du} = x \frac{d}{dx}$, and by $O(\hbar)$ we mean an operator whose application to the wave function $Z^B(x, \hbar)$ produces a function of order 1 or higher in \hbar . Now define a new variable

$$(7.4) \quad y = S'_0.$$

Then the semi-classical limit $\hbar \rightarrow 0$ of (7.3) yields an equation

$$(7.5) \quad y - xy^{2-\chi(B)} e^y = y \left(1 - xy^{1-\chi(B)} e^y \right) = 0,$$

since $Z^B(x, 0) = 1$. Note that this is exactly the *total symbol* of the operator P , where $\hbar x \frac{d}{dx}$ is represented by a commutative variable y . The second factor of (7.5) also recovers the Lambert curve (1.5).

Although the formal manipulation seems to work, however, we have to remember that we have derived the operator P assuming the shape of the solution $Z^B(x, \hbar)$ as in (1.8). Since we are imposing

$$P \overline{Z}^B(x, \hbar) = 0,$$

it forces that $y = 0$, which makes (7.5) trivially correct.

On the other hand, since the kernel of Q assumes only the expansion (4.5), where the summation index m can be negative, the semi-classical analysis of Q does go through.

$$(7.6) \quad \begin{aligned} &e^{-S_1} e^{-\frac{1}{\hbar} S_0} Q e^{\frac{1}{\hbar} S_0} e^{S_1} \\ &= e^{-S_1} e^{-\frac{1}{\hbar} S_0} \left[\frac{\partial}{\partial \hbar} - \frac{1}{2} \frac{\partial^2}{\partial u^2} + \left(\frac{1}{2} - \frac{1-\chi(B)}{\hbar} \right) \frac{\partial}{\partial u} \right] e^{\frac{1}{\hbar} S_0} e^{S_1} \\ &= -\frac{1}{\hbar^2} \left(S_0 + \frac{1}{2} (S'_0)^2 + (1-\chi(B)) S'_0 \right) \\ &\quad - \frac{1}{\hbar} \left(\frac{1}{2} S''_0 - \frac{1}{2} S'_0 + S'_0 S'_1 + (1-\chi(B)) S'_1 \right) + O(1). \end{aligned}$$

For the $\hbar \rightarrow 0$ limit to exist, we need

$$(7.7) \quad S_0 + \frac{1}{2}(S'_0)^2 + (1 - \chi(B))S'_0 = 0,$$

$$(7.8) \quad \frac{1}{2}S''_0 - \frac{1}{2}S'_0 + S'_0S'_1 + (1 - \chi(B))S'_1 = 0.$$

From (7.7) we obtain

$$(7.9) \quad S_0 = -\frac{1}{2}y^2 - (1 - \chi(B))y,$$

or equivalently,

$$(7.10) \quad y = -(y + 1 - \chi(B))\frac{dy}{du}.$$

Its solution is

$$y + \log y^{1-\chi(B)} = -u + \text{const.}$$

If we take the constant of integration to be 0, then we obtain

$$e^u = x = y^{\chi(B)-1}e^{-y},$$

recovering (1.5). From (7.8) we have

$$\frac{1}{2}y' - \frac{1}{2}y + (y + 1 - \chi(B))S'_1 = 0,$$

or equivalently

$$\frac{dS_1}{du} = -\frac{1}{2}\frac{1}{y + 1 - \chi(B)}\frac{dy}{du} + \frac{1}{2}\frac{y}{y + 1 - \chi(B)}\frac{dy}{dy}.$$

Since we know du/dy from (7.10), we can integrate the above equation to obtain

$$(7.11) \quad S_1 = -\frac{1}{2}y + \frac{1}{2}\log(y + 1 - \chi(B)) + \text{const.}$$

The above derivation of the semi-classical limit of the operator Q is valid if $Z^B(x, \hbar)$ contains negative powers of \hbar . If we assume the expansion (1.8), then such a situation occurs when the base curve B satisfies $\chi(B) > 1$. Indeed, our formulas (7.9) and (7.11) agree with those of [26, Section 5] for $B = \mathbb{P}^1$ with $y = z$, and [3, Section 6] for $B = \mathbb{P}^1[a]$ with $y = z^a$.

References

- [1] Mina Aganagic, Robbert Dijkgraaf, Albrecht Klemm, Marcos Mariño, and Cumrun Vafa, *Topological strings and integrable hierarchies*, *Comm. Math. Phys.* **261** (2006), no. 2, 451–516, DOI 10.1007/s00220-005-1448-9. MR2191887
- [2] Gaëtan Borot, Bertrand Eynard, Motohico Mulase, and Brad Safnuk, *A matrix model for simple Hurwitz numbers, and topological recursion*, *J. Geom. Phys.* **61** (2011), no. 2, 522–540, DOI 10.1016/j.geomphys.2010.10.017. MR2746135
- [3] Vincent Bouchard, Daniel Hernández Serrano, Xiaojun Liu, and Motohico Mulase, *Mirror symmetry for orbifold Hurwitz numbers*, *J. Differential Geom.* **98** (2014), no. 3, 375–423. MR3263522
- [4] Vincent Bouchard, Albrecht Klemm, Marcos Mariño, and Sara Pasquetti, *Remodeling the B-model*, *Comm. Math. Phys.* **287** (2009), no. 1, 117–178, DOI 10.1007/s00220-008-0620-4. MR2480744
- [5] Vincent Bouchard and Marcos Mariño, *Hurwitz numbers, matrix models and enumerative geometry*, From Hodge theory to integrability and TQFT tt*-geometry, *Proc. Sympos. Pure Math.*, vol. 78, Amer. Math. Soc., Providence, RI, 2008, pp. 263–283, DOI 10.1090/pspum/078/2483754. MR2483754

- [6] Robbert Dijkgraaf, *Mirror symmetry and elliptic curves*, The moduli space of curves (Texel Island, 1994), Progr. Math., vol. 129, Birkhäuser Boston, Boston, MA, 1995, pp. 149–163, DOI 10.1007/978-1-4612-4264-2.5. MR1363055
- [7] Robbert Dijkgraaf, Lotte Hollands, and Piotr Sulkowski, *Quantum curves and \mathcal{D} -modules*, J. High Energy Phys. **11** (2009), 047, 59, DOI 10.1088/1126-6708/2009/11/047. MR2628886
- [8] Robbert Dijkgraaf, Lotte Hollands, Piotr Sulkowski, and Cumrun Vafa, *Supersymmetric gauge theories, intersecting branes and free fermions*, J. High Energy Phys. **2** (2008), 106, 57, DOI 10.1088/1126-6708/2008/02/106. MR2385939
- [9] R. Dijkgraaf and C. Vafa, *Two Dimensional Kodaira-Spencer Theory and Three Dimensional Chern-Simons Gravity*, arXiv:0711.1932 (2007).
- [10] Norman Do, Oliver Leigh, and Paul Norbury, *Orbifold Hurwitz numbers and Eynard-Orantin invariants*, Math. Res. Lett. **23** (2016), no. 5, 1281–1327, DOI 10.4310/MRL.2016.v23.n5.a3. MR3601067
- [11] Olivia Dumitrescu, Motohico Mulase, Brad Safnuk, and Adam Sorkin, *The spectral curve of the Eynard-Orantin recursion via the Laplace transform*, Algebraic and geometric aspects of integrable systems and random matrices, Contemp. Math., vol. 593, Amer. Math. Soc., Providence, RI, 2013, pp. 263–315, DOI 10.1090/conm/593/11867. MR3087960
- [12] Torsten Ekedahl, Sergei Lando, Michael Shapiro, and Alek Vainshtein, *Hurwitz numbers and intersections on moduli spaces of curves*, Invent. Math. **146** (2001), no. 2, 297–327, DOI 10.1007/s002220100164. MR1864018
- [13] Bertrand Eynard, Motohico Mulase, and Bradley Safnuk, *The Laplace transform of the cut-and-join equation and the Bouchard-Mariño conjecture on Hurwitz numbers*, Publ. Res. Inst. Math. Sci. **47** (2011), no. 2, 629–670, DOI 10.2977/PRIMS/47. MR2849645
- [14] B. Eynard and N. Orantin, *Invariants of algebraic curves and topological expansion*, Commun. Number Theory Phys. **1** (2007), no. 2, 347–452, DOI 10.4310/CNTP.2007.v1.n2.a4. MR2346575
- [15] Leonid Chekhov, Bertrand Eynard, and Nicolas Orantin, *Free energy topological expansion for the 2-matrix model*, J. High Energy Phys. **12** (2006), 053, 31, DOI 10.1088/1126-6708/2006/12/053. MR2276699
- [16] I. P. Goulden, *A differential operator for symmetric functions and the combinatorics of multiplying transpositions*, Trans. Amer. Math. Soc. **344** (1994), no. 1, 421–440, DOI 10.2307/2154724. MR1249468
- [17] I. P. Goulden and D. M. Jackson, *Transitive factorisations into transpositions and holomorphic mappings on the sphere*, Proc. Amer. Math. Soc. **125** (1997), no. 1, 51–60, DOI 10.1090/S0002-9939-97-03880-X. MR1396978
- [18] Tom Graber and Ravi Vakil, *Hodge integrals and Hurwitz numbers via virtual localization*, Compositio Math. **135** (2003), no. 1, 25–36, DOI 10.1023/A:1021791611677. MR1955162
- [19] Sergei Gukov and Piotr Sulkowski, *A-polynomial, B-model, and quantization*, Homological mirror symmetry and tropical geometry, Lect. Notes Unione Mat. Ital., vol. 15, Springer, Cham, 2014, pp. 87–151, DOI 10.1007/978-3-319-06514-4_4. MR3330786
- [20] A. Hurwitz, *Ueber Riemann'sche Flächen mit gegebenen Verzweigungspunkten* (German), Math. Ann. **39** (1891), no. 1, 1–60, DOI 10.1007/BF01199469. MR1510692
- [21] M. Kaneko and D. Zagier, *A generalized Jacobi theta function and quasi-modular forms*, in: R.H. Dijkgraaf et al. (eds) The Moduli Space of Curves. Progress in Mathematics **129**, Birkhäuser Boston, 165–172 (1995).
- [22] M. Kazarian, *KP hierarchy for Hodge integrals*, Adv. Math. **221** (2009), no. 1, 1–21, DOI 10.1016/j.aim.2008.10.017. MR2509319
- [23] M. E. Kazarian and S. K. Lando, *An algebro-geometric proof of Witten's conjecture*, J. Amer. Math. Soc. **20** (2007), no. 4, 1079–1089, DOI 10.1090/S0894-0347-07-00566-8. MR2328716
- [24] Marcos Mariño, *Open string amplitudes and large order behavior in topological string theory*, J. High Energy Phys. **3** (2008), 060, 34, DOI 10.1088/1126-6708/2008/03/060. MR2391060
- [25] M. Mulase, S. Shadrin, and L. Spitz, *The spectral curve and the Schrödinger equation of double Hurwitz numbers and higher spin structures*, Commun. Number Theory Phys. **7** (2013), no. 1, 125–143, DOI 10.4310/CNTP.2013.v7.n1.a4. MR3108774
- [26] Motohico Mulase and Piotr Sulkowski, *Spectral curves and the Schrödinger equations for the Eynard-Orantin recursion*, Adv. Theor. Math. Phys. **19** (2015), no. 5, 955–1015, DOI 10.4310/ATMP.2015.v19.n5.a2. MR3487649

- [27] M. Mulase and J. Yu, *A generating function of the number of homomorphisms from a surface group into a finite group*, [arXiv:math.QA/0209008](https://arxiv.org/abs/math/0209008), (2002).
- [28] Motohico Mulase and Naizhen Zhang, *Polynomial recursion formula for linear Hodge integrals*, *Commun. Number Theory Phys.* **4** (2010), no. 2, 267–293, DOI 10.4310/CNTP.2010.v4.n2.a1. MR2725053
- [29] Andrei Okounkov, *Toda equations for Hurwitz numbers*, *Math. Res. Lett.* **7** (2000), no. 4, 447–453, DOI 10.4310/MRL.2000.v7.n4.a10. MR1783622
- [30] A. Okounkov and R. Pandharipande, *Gromov-Witten theory, Hurwitz numbers, and matrix models*, *Algebraic geometry—Seattle 2005. Part 1*, *Proc. Sympos. Pure Math.*, vol. 80, Amer. Math. Soc., Providence, RI, 2009, pp. 325–414, DOI 10.1090/pspum/080.1/2483941. MR2483941
- [31] A. Okounkov and R. Pandharipande, *Gromov-Witten theory, Hurwitz theory, and completed cycles*, *Ann. of Math. (2)* **163** (2006), no. 2, 517–560, DOI 10.4007/annals.2006.163.517. MR2199225
- [32] R. Vakil, *Enumerative Geometry of Curves via Degeneration Methods*, Ph.D. thesis, Harvard University (1997).
- [33] J. Zhou, *Hodge integrals, Hurwitz numbers, and symmetric groups*, [arXiv:math/0308024](https://arxiv.org/abs/math/0308024), (2003).
- [34] J. Zhou, *Quantum Mirror Curves for \mathbb{C}^3 and the Resolved Conifold*, [arXiv:1207.0598v1](https://arxiv.org/abs/1207.0598v1), (2012).

DEPARTMENT OF APPLIED MATHEMATICS, CHINA AGRICULTURAL UNIVERSITY, BEIJING, 100083, CHINA; AND DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, CALIFORNIA 95616–8633

Email address: xjliu@cau.edu.cn

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, CALIFORNIA 95616–8633

Email address: mulase@math.ucdavis.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, CALIFORNIA 95616–8633

Email address: azsorkin@math.ucdavis.edu