

**Mathematical
Surveys
and
Monographs**
Volume 227

Bounded Cohomology of Discrete Groups

Roberto Frigerio




American Mathematical Society

Bounded Cohomology of Discrete Groups

**Mathematical
Surveys
and
Monographs**
Volume 227

Bounded Cohomology of Discrete Groups

Roberto Frigerio



American Mathematical Society
Providence, Rhode Island

EDITORIAL COMMITTEE

Robert Guralnick
Michael A. Singer, Chair
Benjamin Sudakov
Constantin Teleman
Michael I. Weinstein

2010 *Mathematics Subject Classification*. Primary 18G60, 20J06, 55N10, 57N65;
Secondary 37E10, 37C85, 53C23, 57M07, 57R20.

For additional information and updates on this book, visit
www.ams.org/bookpages/surv-227

Library of Congress Cataloging-in-Publication Data

Names: Frigerio, Roberto, author.

Title: Bounded cohomology of discrete groups / Roberto Frigerio.

Description: Providence, Rhode Island : American Mathematical Society, [2017] | Series: Mathematical surveys and monographs ; volume 227 | Includes bibliographical references and index.

Identifiers: LCCN 2017023394 | ISBN 9781470441463 (alk. paper)

Subjects: LCSH: Intersection homology theory. | Homology theory. | Algebra, Homological. | AMS: Category theory; homological algebra – Homological algebra – Other (co)homology theories. msc | Group theory and generalizations – Connections with homological algebra and category theory – Cohomology of groups. msc | Algebraic topology – Homology and cohomology theories – Singular theory. msc | Manifolds and cell complexes – Topological manifolds – Algebraic topology of manifolds. msc | Dynamical systems and ergodic theory – Low-dimensional dynamical systems – Maps of the circle. msc | Dynamical systems and ergodic theory – Smooth dynamical systems: general theory – Dynamics of group actions other than \mathbb{Z} . msc | Differential geometry – Global differential geometry – Global geometric and topological methods (Gromov); differential geometric analysis on metric spaces. msc | Manifolds and cell complexes – Low-dimensional topology – Topological methods in group theory. msc | Manifolds and cell complexes – Differential topology – Characteristic classes and numbers. msc

Classification: LCC QA612.3 .F75 2017 | DDC 512/.55–dc23

LC record available at <https://lcn.loc.gov/2017023394>

Copying and reprinting. Individual readers of this publication, and nonprofit libraries acting for them, are permitted to make fair use of the material, such as to copy select pages for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgment of the source is given.

Republication, systematic copying, or multiple reproduction of any material in this publication is permitted only under license from the American Mathematical Society. Permissions to reuse portions of AMS publication content are handled by Copyright Clearance Center's RightsLink® service. For more information, please visit: <http://www.ams.org/rightslink>.

Send requests for translation rights and licensed reprints to reprint-permission@ams.org.

Excluded from these provisions is material for which the author holds copyright. In such cases, requests for permission to reuse or reprint material should be addressed directly to the author(s). Copyright ownership is indicated on the copyright page, or on the lower right-hand corner of the first page of each article within proceedings volumes.

© 2017 by the American Mathematical Society. All rights reserved.

The American Mathematical Society retains all rights
except those granted to the United States Government.
Printed in the United States of America.

∞ The paper used in this book is acid-free and falls within the guidelines
established to ensure permanence and durability.

Visit the AMS home page at <http://www.ams.org/>

10 9 8 7 6 5 4 3 2 1 22 21 20 19 18 17

Contents

Introduction	ix
Chapter 1. (Bounded) cohomology of groups	1
1.1. Cohomology of groups	1
1.2. The topological interpretation of group cohomology	3
1.3. Bounded cohomology of groups	3
1.4. The comparison map and exact bounded cohomology	5
1.5. The bar resolution	5
1.6. Topology and bounded cohomology	6
1.7. Further readings	6
Chapter 2. (Bounded) cohomology of groups in low degree	9
2.1. (Bounded) group cohomology in degree zero and one	9
2.2. Group cohomology in degree two	9
2.3. Bounded group cohomology in degree two: quasimorphisms	12
2.4. Homogeneous quasimorphisms	13
2.5. Quasimorphisms on abelian groups	14
2.6. The bounded cohomology of free groups in degree 2	15
2.7. Homogeneous 2-cocycles	16
2.8. The image of the comparison map	18
2.9. Further readings	20
Chapter 3. Amenability	23
3.1. Abelian groups are amenable	25
3.2. Other amenable groups	26
3.3. Amenability and bounded cohomology	27
3.4. Johnson's characterization of amenability	28
3.5. A characterization of finite groups via bounded cohomology	29
3.6. Further readings	30
Chapter 4. (Bounded) group cohomology via resolutions	33
4.1. Relative injectivity	33
4.2. Resolutions of Γ -modules	35
4.3. The classical approach to group cohomology via resolutions	38
4.4. The topological interpretation of group cohomology revisited	39
4.5. Bounded cohomology via resolutions	40
4.6. Relatively injective normed Γ -modules	41
4.7. Resolutions of normed Γ -modules	41
4.8. More on amenability	44
4.9. Amenable spaces	45

4.10.	Alternating cochains	48
4.11.	Further readings	49
Chapter 5.	Bounded cohomology of topological spaces	53
5.1.	Basic properties of bounded cohomology of spaces	53
5.2.	Bounded singular cochains as relatively injective modules	54
5.3.	The aspherical case	56
5.4.	Ivanov's contracting homotopy	56
5.5.	Gromov's Theorem	58
5.6.	Alternating cochains	59
5.7.	Relative bounded cohomology	60
5.8.	Further readings	62
Chapter 6.	ℓ^1 -homology and duality	65
6.1.	Normed chain complexes and their topological duals	65
6.2.	ℓ^1 -homology of groups and spaces	66
6.3.	Duality: first results	67
6.4.	Some results by Matsumoto and Morita	68
6.5.	Injectivity of the comparison map	70
6.6.	The translation principle	71
6.7.	Gromov equivalence theorem	73
6.8.	Further readings	75
Chapter 7.	Simplicial volume	77
7.1.	The case with non-empty boundary	77
7.2.	Elementary properties of the simplicial volume	78
7.3.	The simplicial volume of Riemannian manifolds	79
7.4.	Simplicial volume of gluings	80
7.5.	Simplicial volume and duality	82
7.6.	The simplicial volume of products	83
7.7.	Fiber bundles with amenable fibers	83
7.8.	Further readings	84
Chapter 8.	The proportionality principle	87
8.1.	Continuous cohomology of topological spaces	87
8.2.	Continuous cochains as relatively injective modules	88
8.3.	Continuous cochains as strong resolutions of \mathbb{R}	90
8.4.	Straightening in non-positive curvature	92
8.5.	Continuous cohomology versus singular cohomology	92
8.6.	The transfer map	93
8.7.	Straightening and the volume form	95
8.8.	Proof of the proportionality principle	97
8.9.	The simplicial volume of hyperbolic manifolds	97
8.10.	Hyperbolic straight simplices	98
8.11.	The seminorm of the volume form	99
8.12.	The case of surfaces	100
8.13.	The simplicial volume of negatively curved manifolds	100
8.14.	The simplicial volume of flat manifolds	101
8.15.	Further readings	101

Chapter 9. Additivity of the simplicial volume	105
9.1. A cohomological proof of subadditivity	105
9.2. A cohomological proof of Gromov additivity theorem	107
9.3. Further readings	110
Chapter 10. Group actions on the circle	113
10.1. Homeomorphisms of the circle and the Euler class	113
10.2. The bounded Euler class	114
10.3. The (bounded) Euler class of a representation	115
10.4. The rotation number of a homeomorphism	116
10.5. Increasing degree one map of the circle	119
10.6. Semi-conjugacy	120
10.7. Ghys' Theorem	122
10.8. The canonical real bounded Euler cocycle	126
10.9. Further readings	129
Chapter 11. The Euler class of sphere bundles	131
11.1. Topological, smooth and linear sphere bundles	131
11.2. The Euler class of a sphere bundle	133
11.3. Classical properties of the Euler class	136
11.4. The Euler class of oriented vector bundles	138
11.5. The Euler class of circle bundles	140
11.6. Circle bundles over surfaces	142
11.7. Further readings	143
Chapter 12. Milnor-Wood inequalities and maximal representations	145
12.1. Flat sphere bundles	145
12.2. The bounded Euler class of a flat circle bundle	149
12.3. Milnor-Wood inequalities	151
12.4. Flat circle bundles on surfaces with boundary	154
12.5. Maximal representations	162
12.6. Further readings	166
Chapter 13. The bounded Euler class in higher dimensions and the Chern conjecture	169
13.1. Ivanov-Turaev cocycle	169
13.2. Representing cycles via simplicial cycles	173
13.3. The bounded Euler class of a flat linear sphere bundle	174
13.4. The Chern conjecture	178
13.5. Further readings	179
Index	181
List of Symbols	185
Bibliography	187

Introduction

Bounded cohomology of groups was first defined by Johnson [Joh72] and Trauber during the seventies in the context of Banach algebras. As an independent and very active research field, however, bounded cohomology started to develop in 1982, thanks to the pioneering paper “Volume and Bounded Cohomology” by M. Gromov [Gro82], where the definition of bounded cohomology was extended to deal also with topological spaces.

Let $C^\bullet(M, \mathbb{R})$ denote the complex of real singular cochains with values in the topological space M . A cochain $\varphi \in C^n(M, \mathbb{R})$ is *bounded* if it takes uniformly bounded values on the set of singular n -simplices. Bounded cochains provide a sub-complex $C_b^\bullet(M, \mathbb{R})$ of singular cochains, and the bounded cohomology $H_b^\bullet(M, \mathbb{R})$ of M (with trivial real coefficients) is just the (co)homology of the complex $C_b^\bullet(M, \mathbb{R})$. An analogous definition of boundedness applies to group cochains with (trivial) real coefficients, and the bounded cohomology $H_b^\bullet(\Gamma, \mathbb{R})$ of a group Γ (with trivial real coefficients) is the (co)homology of the complex $C_b^\bullet(\Gamma, \mathbb{R})$ of the bounded cochains on Γ . A classical result which dates back to the forties ensures that the singular cohomology of an aspherical CW-complex is canonically isomorphic to the cohomology of its fundamental group. In the context of bounded cohomology a stronger result holds: the bounded cohomology of a topological space is canonically isomorphic to the bounded cohomology of its fundamental group, even without any assumption on the asphericity of the space [Gro82, Bro81, Iva87, FM11, Iva]. For example, the bounded cohomology of spheres is trivial in positive degree. On the other hand, the bounded cohomology of the wedge of two circles is infinite-dimensional in degrees 2 and 3, and still unknown in any degree bigger than 3. As we will see in this monograph, this phenomenon eventually depends on the fact that higher homotopy groups are abelian, and abelian groups are invisible to bounded cohomology, while “negatively curved” groups, such as non-abelian free groups, tend to have very big bounded cohomology modules.

The bounded cohomology of a group Γ can be defined with coefficients in any normed (e.g. Banach) Γ -module, where the norm is needed to make sense of the notion of boundedness of cochains. Moreover, if Γ is a topological group, then one may restrict to considering only *continuous* bounded cochains, thus obtaining the *continuous* bounded cohomology of Γ . In this monograph, we will often consider arbitrary normed Γ -modules, but we will restrict our attention to bounded cohomology of discrete groups. The reason for this choice is twofold. First, the (very powerful) theory of continuous bounded cohomology is based on a quite sophisticated machinery, which is not needed in the case of discrete groups. Secondly, Monod’s book [Mon01] and Burger-Monod’s paper [BM02] already provide an excellent introduction to continuous bounded cohomology, while to the author’s knowledge no reference is available where the fundamental properties of bounded

cohomology of discrete groups are collected and proved in detail. However, we should emphasize that the theory of continuous bounded cohomology is essential in proving important results also in the context of discrete groups: many vanishing theorems for the bounded cohomology of lattices in Lie groups can be obtained by comparing the bounded cohomology of the lattice with the continuous bounded cohomology of the ambient group.

This monograph is devoted to provide a self-contained introduction to bounded cohomology of discrete groups and topological spaces. Several (by now classical) applications of the theory will be described in detail, while many others will be completely omitted. Of course, the choice of the topics discussed here is largely arbitrary, and based on the taste (and on the knowledge) of the author.

Before describing the content of each chapter, let us provide a brief overview on the relationship between bounded cohomology and other research fields.

Geometric group theory and quasification. The bounded cohomology of a closed manifold is strictly related to the curvature of the metrics that the manifold can support. For example, if the closed manifold M is flat or positively curved, then the fundamental group of M is amenable, and $H_b^n(M, \mathbb{R}) = H_b^n(\pi_1(M), \mathbb{R}) = 0$ for every $n \geq 1$. On the other hand, if M is negatively curved, then it is well-known that the comparison map $H_b^\bullet(M, \mathbb{R}) \rightarrow H^\bullet(M, \mathbb{R})$ induced by the inclusion $C_b^\bullet(M, \mathbb{R}) \rightarrow C^\bullet(M, \mathbb{R})$ is surjective in every degree bigger than one.

In fact, it turns out that the surjectivity of the comparison map is related in a very clean way to the notion of Gromov hyperbolicity, which represents the coarse geometric version of negative curvature. Namely, a group Γ is Gromov hyperbolic if and only if the comparison map $H_b^n(\Gamma, V) \rightarrow H^n(\Gamma, V)$ is surjective for every $n \geq 2$ and for every Banach Γ -module V [Min01, Min02].

Coarse geometry comes into play also when studying the (non-)injectivity of the comparison map. In fact, let $EH_b^n(\Gamma, V)$ denote the kernel of the comparison map in degree n . It follows by the very definitions that an element of $H_b^n(\Gamma, V)$ lies in $EH_b^n(\Gamma, V)$ if and only if any of its representatives is the coboundary of a (possibly unbounded) cocycle. More precisely, a cochain is usually called a *quasi-cocycle* if its differential is bounded, and a quasi-cocycle is *trivial* if it is the sum of a cocycle and a bounded cochain. Then $EH_b^n(\Gamma, V)$ is canonically isomorphic to the space of $(n-1)$ -quasi-cocycles modulo trivial $(n-1)$ -quasi-cocycles. When $V = \mathbb{R}$, quasi-cocycles of degree one are usually called *quasimorphisms*. There exists a large literature which concerns the construction of non-trivial quasimorphisms in presence of (weaker and weaker versions of) negative curvature. Brooks [Bro81] first constructed infinitely many quasimorphisms on the free group with two generators F_2 , that were shown to define linearly independent elements in $EH^2(F_2, \mathbb{R})$ by Mitsumatsu [Mit84]. In particular, this proved that $EH_b^2(F_2, \mathbb{R})$ (which coincides with $H_b^2(F_2, \mathbb{R})$) is infinite-dimensional.

Quasi-cocycles are cochains which satisfy the cocycle equation only up to a finite error, and geometric group theory provides tools that are particularly well-suited to study notions which involve finite errors in their definition. Therefore, it is not surprising that Brooks' and Mitsumatsu's result has been generalized to larger and larger classes of groups, which now include the class of non-elementary relatively hyperbolic groups, and most mapping class groups. We refer the reader to Section 2.9 for a more detailed account on this issue. It is maybe worth mentioning that, even if in the cases cited above $EH_b^2(\Gamma, \mathbb{R})$ is always infinite-dimensional, there

exist lattices Γ in non-linear Lie groups for which $EH_b^2(\Gamma, \mathbb{R})$ is of finite non-zero dimension [MR06].

We have mentioned the fact that $H_b^2(\Gamma, \mathbb{R})$ is infinite-dimensional for negatively curved groups according to a suitable notion of negative curvature for groups. On the other hand, bounded cohomology vanishes for “positively curved” (i.e. finite) or “flat” (i.e. virtually abelian) groups. In fact, $H_b^n(\Gamma, \mathbb{R}) = 0$ for any $n \geq 1$ and any Γ belonging to the distinguished class of *amenable* groups. The class of amenable groups contains all virtually solvable groups, it is closed under quasi-isometries, and it admits a nice characterization in terms of bounded cohomology (see Section 3.4). These facts provide further evidence that bounded cohomology detects coarse non-positive curvature as well as coarse non-negative curvature.

Simplicial volume. The ℓ^∞ -norm of an n -cochain is the supremum of the values it takes on single singular n -simplices (or on single $(n+1)$ -tuples of elements of the group, when dealing with group cochains rather than with singular cochains). So a cochain φ is bounded if and only if it has a finite ℓ^∞ -norm, and the ℓ^∞ -norm induces a natural quotient ℓ^∞ -seminorm on bounded cohomology. The ℓ^∞ -norm on singular cochains arises as the dual of a natural ℓ^1 -norm on singular *chains*. This ℓ^1 -norm induces an ℓ^1 -seminorm on homology. If M is a closed oriented manifold, then the *simplicial volume* $\|M\|$ of M is the ℓ^1 -seminorm of the real fundamental class of M [Gro82]. Even if it depends only on the homotopy type of the manifold, the simplicial volume is deeply related to the geometric structures that a manifold can carry. As one of the main motivations for its definition, Gromov himself showed that the simplicial volume provides a lower bound for the *minimal volume* of a manifold, which is the infimum of the volumes of the Riemannian metrics that are supported by the manifold and that satisfy suitable curvature bounds.

An elementary duality result relates the simplicial volume of an n -dimensional manifold M to the bounded cohomology module $H_b^n(M, \mathbb{R})$. For example, $\|M\| = 0$ provided that $H_b^n(M, \mathbb{R}) = 0$. In particular, the simplicial volume of simply connected manifolds (or, more in general, of manifolds with an amenable fundamental group) is vanishing. It is worth stressing that no homological proof of this statement is available: in many cases, the fact that $\|M\| = 0$ cannot be proved by exhibiting fundamental cycles with arbitrarily small norm. Moreover, the exact value of non-vanishing simplicial volumes (of compact manifolds) is known only in the following very few cases: hyperbolic manifolds [Gro82, Thu79], some classes of 3-manifolds with boundary [BFP15], and the product of two surfaces [BK08b]. In the last case, it is not known to the author any description of a sequence of fundamental cycles whose ℓ^1 -norms approximate the simplicial volume. In fact, Bucher’s computation of the simplicial volume of the product of surfaces heavily relies on deep properties of bounded cohomology that have no counterpart in the context of singular homology.

Characteristic classes. A fundamental theorem by Gromov [Gro82] states that, if G is an algebraic subgroup of $GL_n(\mathbb{R})$, then every characteristic class of flat G -bundles lies in the image of the comparison map (i.e. it can be represented by a bounded cocycle). (See [Buc04] for an alternative proof of a stronger result.) Several natural questions arise from this result. First of all, one may ask whether such characteristic classes admit a canonical representative in bounded cohomology: since the comparison map is often non-injective, this would produce more refined

invariants. Even when one is not able to find natural bounded representatives for a characteristic class, the seminorm on bounded cohomology (which induces a seminorm on the image of the comparison map by taking the infimum over the bounded representatives) can be used to produce numerical invariants, or to provide useful estimates.

In this context, the most famous example is certainly represented by the Euler class. To every oriented circle bundle there is associated its Euler class, which arises as an obstruction to the existence of a section, and completely classifies the topological type of the bundle. When restricting to *flat* circle bundles, a *bounded* Euler class can be defined, which now depends on the flat structure of the bundle (and, in fact, classifies the isomorphism type of flat bundles with minimal holonomy [Ghy87, Ghy01]). Moreover, the seminorm of the bounded Euler class is equal to $1/2$. Now the Euler number of a circle bundle over a surface is obtained by evaluating the Euler class on the fundamental class of the surface. As a consequence, the Euler number of any flat circle bundle over a surface is bounded above by the product of $1/2$ times the simplicial volume of the surface. This yields the celebrated *Milnor-Wood inequalities* [Mil58a, Woo71], which provided the first explicit and easily computable obstructions for a circle bundle to admit a flat structure. Of course, Milnor's and Wood's original proofs did not explicitly use bounded cohomology, which was not defined yet. However, their arguments introduced ideas and techniques which are now fundamental in the theory of quasimorphisms, and they were implicitly based on the construction of a bounded representative for the Euler class.

These results also extend to higher dimensions. It was proved by Sullivan [Sul76] that the ℓ^∞ -seminorm of the (simplicial) Euler class of an oriented flat vector bundle is bounded above by 1. A clever trick due to Smillie allowed to sharpen this bound from 1 to 2^{-n} , where n is the rank of the bundle. Then, Ivanov and Turaev [IT82] gave a different proof of Smillie's result, also constructing a natural bounded Euler class in every dimension. A very clean characterization of this bounded cohomology class, as well as the proof that its seminorm is equal to 2^{-n} , have recently been provided by Bucher and Monod in [BM12].

Actions on the circle. If X is a topological space with fundamental group Γ , then any orientation-preserving topological action of Γ on the circle gives rise to a flat circle bundle over X . Therefore, we can associate to every such action a bounded Euler class. It is a fundamental result of Ghys [Ghy87, Ghy01] that the bounded Euler class encodes the most essential features of the dynamics of an action. For example, an action admits a global fixed point if and only if its bounded Euler class vanishes. Furthermore, in the almost opposite case of *minimal* actions (i.e. of actions every orbit of which is dense), the bounded Euler class provides a complete conjugacy invariant: two minimal circle actions share the same bounded Euler class if and only if they are topologically conjugate [Ghy87, Ghy01]. These results establish a deep connection between bounded cohomology and a fundamental problem in one-dimensional dynamics.

Representations and Rigidity. Bounded cohomology has been very useful in proving rigidity results for representations. It is known that an epimorphism between discrete groups induces an injective map on 2-dimensional bounded cohomology with real coefficients (see Theorem 2.17). As a consequence, if $H_b^2(\Gamma, \mathbb{R})$ is

finite-dimensional and $\rho: \Gamma \rightarrow G$ is any representation, then the second bounded cohomology of the image of ρ must also be finite-dimensional. In some cases, this information suffices to ensure that ρ is almost trivial. For example, if Γ is a uniform irreducible lattice in a higher rank semisimple Lie group, then by work of Burger and Monod we have that $H_b^2(\Gamma, \mathbb{R})$ is finite-dimensional [BM99]. On the contrary, non-virtually-abelian subgroups of mapping class groups of hyperbolic surfaces admit many non-trivial quasimorphisms [BF02], whence an infinite-dimensional second bounded cohomology group. As a consequence, the image of any representation of a higher rank lattice into a mapping class group is virtually abelian, whence finite. This provides an independent proof of a result by Farb, Kaimanovich and Masur [FM98, KM96].

Rigidity results of a different nature arise when exploiting bounded cohomology to get more refined invariants with respect to the ones provided by classical cohomology. For example, we have seen that the norm of the Euler class can be used to bound the Euler number of flat circle bundles. When considering representations into $PSL(2, \mathbb{R})$ of the fundamental group of closed surfaces of negative Euler characteristic, a celebrated result by Goldman [Gol80] implies that a representation has maximal Euler number if and only if it is faithful and discrete, i.e. if and only if it is the holonomy of a hyperbolic structure (in this case, one usually says that the representation is *geometric*). A new proof of this fact (in the more general case of surfaces with punctures) has been recently provided in [BIW10] via a clever use of the bounded Euler class of a representation (see also [Ioz02] for another proof of Goldman’s Theorem based on bounded cohomology). In fact, one may define maximal representations of surface groups into a much more general class of Lie groups (see e.g. [BIW10]). This strategy has been followed e.g. in [BI07, BI09] to establish deformation rigidity for representations into $SU(m, 1)$ of lattices in $SU(n, 1)$ also in the non-uniform case (the uniform case having being settled by Goldman and Millson in [GM87]). Finally, bounded cohomology has proved very useful in studying rigidity phenomena also in the context of orbit equivalence and measure equivalence for countable groups (see e.g. [MS06] and [BFS13]).

Content of the book. Let us now briefly outline the content of each chapter. In the **first chapter** we introduce the basic definitions about the cohomology and the bounded cohomology of groups. We introduce the comparison map between the bounded cohomology and the classical cohomology of a group, and we briefly discuss the relationship between the classical cohomology of a group and the singular cohomology of its classifying space, postponing to Chapter 5 the investigation of the same topic in the context of bounded cohomology.

In **Chapter 2** we study the bounded cohomology of groups in low degrees. When working with trivial coefficients, bounded cohomology in degree 0 and 1 is completely understood, so our attention is mainly devoted to degree 2. We recall that degree-2 classical cohomology is related to group extensions, while degree-2 bounded cohomology is related to the existence of quasimorphisms. We exhibit many non-trivial quasimorphisms on the free group, thus showing that $H_b^2(F, \mathbb{R})$ is infinite-dimensional for every non-abelian free group F . We also describe a somewhat neglected result by Bouarich, which states that every class in $H_b^2(\Gamma, \mathbb{R})$ admits a canonical “homogeneous” representative, and, following [Bou04], we use this fact to show that any group epimorphism induces an injective map on bounded cohomology (with trivial real coefficients) in degree 2. A stronger result (with more

general coefficients allowed, and where the induced map is shown to be an isometric embedding) may be found in [Hub12].

Chapter 3 is devoted to amenability, which represents a fundamental notion in the theory of bounded cohomology. We briefly review some results on amenable groups, also providing a complete proof of von Neumann’s Theorem, which ensures that abelian groups are amenable. Then we show that bounded cohomology of amenable groups vanishes (at least for a very wide class of coefficients), and we describe Johnson’s characterization of amenability in terms of bounded cohomology.

In **Chapter 4** we introduce the tools from homological algebra which are best suited to deal with bounded cohomology. Following [Iva87] and [BM99, BM02], we define the notion of relatively injective Γ -module, and we establish the basic results that allow to compute bounded cohomology via relatively injective strong resolutions. We also briefly discuss how this part of the theory interacts with amenability, also defining the notion of amenable action (in the very restricted context of discrete groups acting on discrete spaces).

We come back to the topological interpretation of bounded cohomology of groups in **Chapter 5**. By exploiting the machinery developed in the previous chapter, we describe Ivanov’s proof of a celebrated result due to Gromov, which asserts that the bounded cohomology of a space is isometrically isomorphic to the bounded cohomology of its fundamental group. We also introduce the relative bounded cohomology of topological pairs, and prove that $H_b^n(X, Y)$ is isometrically isomorphic to $H_b^n(X)$ whenever every component of Y has an amenable fundamental group.

In **Chapter 6** we introduce ℓ^1 -homology of groups and spaces. Bounded cohomology naturally provides a dual theory to ℓ^1 -homology. Following some works by Löh, we prove several statements on duality. As an application, we describe Löh’s proof of the fact that the ℓ^1 -homology of a space is canonically isomorphic to the ℓ^1 -homology of its fundamental group. We also review some results by Matsumoto and Morita, showing for example that in degree 2 the seminorm on bounded cohomology is always a norm, and providing a characterization of the injectivity of the comparison map in terms of the so-called *uniform boundary condition*. Following [BBF⁺14], we also make use of duality (and of the results on relative bounded cohomology proved in the previous chapter) to obtain a proof of Gromov equivalence Theorem.

Chapter 7 is devoted to an important application of bounded cohomology: the computation of simplicial volume. The simplicial volume of a closed oriented manifold is equal to the ℓ^1 -seminorm of its real fundamental class. Thanks to the duality between ℓ^1 -homology and bounded cohomology, the study of the simplicial volume of a manifold often benefits from the study of its bounded cohomology. Here we introduce the basic definitions and the most elementary properties of the simplicial volume, and we state several results concerning it, postponing the proofs to the subsequent chapters.

Gromov’s proportionality principle states that, for closed Riemannian manifolds admitting the same Riemannian universal covering, the ratio between the simplicial volume and the Riemannian volume is a constant only depending on the universal covering. In **Chapter 8** we prove a simplified version of the proportionality principle, which applies only to non-positively curved manifolds. The choice of restricting to the non-positively curved context allows us to avoid a lot of technicalities, while introducing the most important ideas on which also the proof of

the general statement is based. In particular, we introduce a bit of continuous cohomology (of topological spaces) and the transfer map, which relates the (bounded) cohomology of a lattice in a Lie group to the continuous (bounded) cohomology of the ambient group. Even if our use of the transfer map is very limited, we feel worth introducing it, since this map plays a very important role in the theory of continuous bounded cohomology of topological groups, as developed by Burger and Monod [BM99, BM02]. As an application, we carry out the computation of the simplicial volume of closed hyperbolic manifolds following the strategy described in [BK08a].

In **Chapter 9** we prove that the simplicial volume is additive with respect to gluings along π_1 -injective boundary components with an amenable fundamental group. Our proof of this fundamental theorem (which is originally due to Gromov) is based on a slight variation of the arguments described in [BBF⁺14]. In fact, we deduce additivity of the simplicial volume from a result on bounded cohomology, together with a suitable application of duality.

As mentioned above, bounded cohomology has found interesting applications in the study of the dynamics of homomorphisms of the circle. In **Chapter 10** we introduce the Euler class and the bounded Euler class of a circle action, and we review a fundamental result due to Ghys [Ghy87, Ghy99, Ghy01], who proved that semi-conjugacy classes of circle actions are completely classified by their bounded Euler class. The bounded Euler class is thus a much finer invariant than the classical Euler class, and this provides a noticeable instance of a phenomenon mentioned above: passing from classical to bounded cohomology often allows to refine classical invariants. We also relate the bounded Euler class of a cyclic group of homeomorphisms of the circle to the classical *rotation number* of the generator of the group, and prove some properties of the rotation number that will be used in the next chapters. Finally, following Matsumoto [Mat86] we describe the canonical representative of the *real* bounded Euler class, also proving a characterization of semi-conjugacy in terms of the real bounded Euler class.

Chapter 11 is devoted to a brief digression from the theory of bounded cohomology. The main aim of the chapter is a detailed description of the Euler class of a sphere bundle. However, our treatment of the subject is a bit different from the usual one, in that we define the Euler cocycle as a *singular* (rather than cellular) cocycle. Of course, cellular and singular cohomology are canonically isomorphic for every CW-complex. However, cellular cochains are not useful to compute the *bounded* cohomology of a cellular complex: for example, it is not easy to detect whether the singular coclass corresponding to a cellular one admits a bounded representative or not, and this motivates us to work directly in the singular context, even if this choice could seem a bit costly at first glance.

In **Chapter 12** we specialize our study of sphere bundles to the case of *flat* bundles. The theory of flat bundles builds a bridge between the theory of fiber bundles and the theory of representations. For example, the Euler class of a flat circle bundle corresponds (under a canonical morphism) to the Euler class of a representation canonically associated to the bundle. This leads to the definition of the *bounded* Euler class of a flat circle bundle. By putting together an estimate on the norm of the bounded Euler class and the computation of the simplicial volume of surfaces carried out in the previous chapters, we are then able to prove *Milnor-Wood inequalities*, which provide sharp estimates on the possible

Euler numbers of flat circle bundles over surfaces. We then concentrate our attention on *maximal* representations, i.e. on representations of surface groups which attain the extremal value allowed by Milnor-Wood inequalities. A celebrated result by Goldman [Gol80] states that maximal representations of surface groups into the group of orientation-preserving isometries of the hyperbolic plane are exactly the holonomies of hyperbolic structures. Following [BIW10], we will give a proof of Goldman’s theorem based on the use of bounded cohomology. In doing this, we will describe the Euler number of flat bundles over surfaces with boundary, as defined in [BIW10].

Chapter 13 is devoted to higher-dimensional generalizations of Milnor-Wood inequalities. We introduce Ivanov-Turaev’s bounded Euler cocycle in dimension $n \geq 2$, and from an estimate (proved to be optimal in [BM12]) on its norm we deduce several inequalities on the possible Euler numbers of flat vector bundles over closed manifolds. We also discuss the relationship between this topic and the Chern conjecture, which predicts that the Euler characteristic of a closed affine manifold should vanish.

Acknowledgements. This book originates from the notes of a Ph. D. course I taught at the University of Rome “La Sapienza” in Spring 2013. I would like to thank Gabriele Mondello, who asked me to teach the course: without his invitation this book would have never been written. Many thanks go to the mathematicians with whom I discussed many topics touched in this monograph; writing down a list may be dangerous, because I am certainly forgetting someone, but I would like to mention at least Michelle Bucher, Marc Burger, Chris Connell, Stefano Francaviglia, Federico Franceschini, Tobias Hartnick, Alessandra Iozzi, Jean-François Lafont, Clara Löh, Bruno Martelli, Cristina Pagliantini, Maria Beatrice Pozzetti, Roman Sauer, Ben Schmidt, Alessandro Sisto. Finally, special thanks go to Elia Fioravanti, Federico Franceschini and Cristina Pagliantini, who pointed out to me some inaccuracies in previous versions of this book.

Index

- G -atlas, 132
 - flat -, 146
 - maximal -, 132
- $R[\Gamma]$ -
 - complex, 35
 - map, 1
 - map of normed modules, 4
 - module, 1
 - augmented normed - complex, 42
 - dual normed - module, 27
 - normed - complex, 41
 - normed - module, 3
- Γ -
 - complex, 35
 - homotopy, 35
 - homotopy for normed complexes, 41
 - invariants, 1
 - augmented - complex, 35
 - augmented normed - complex, 42
- k -tuple
 - positively oriented -, 119
 - weakly positively oriented -, 119
- 2-cocycle
 - homogeneous -, 16
- affine
 - manifold, 178
 - space-form, 178
 - complete - manifold, 178
 - special - manifold, 180
- alternating
 - group cochain, 48
 - singular cochain, 59
- amenable
 - action, 46
 - set, 46
 - group, 23
 - elementary - group, 26
- bar resolution, 6
- central extension, 9
- chain
 - ℓ^1 -, 66
 - singular -, 3
- circle bundle, 131
- class
 - fundamental -, 77
- cochain
 - alternating -, 48
 - alternating singular -, 59
 - bounded continuous -, 88
 - continuous -, 87
 - singular -, 3
 - singular bounded -, 53
 - special -, 61
- coclass
 - fundamental -, 82
- cohomology
 - module of a group, 2
 - bounded - module, 4
 - bounded singular - module, 53
 - exact bounded -, 5
 - relative bounded -, 60
 - relative singular -, 60
 - singular -, 3
- commensurable
 - Riemannian manifolds, 79
- comparison map, 5, 53, 70
- complex, 35
 - $R[\Gamma]$ -, 35
 - Banach chain -, 65
 - homogeneous -, 1
 - inhomogeneous -, 6
 - normed $R[\Gamma]$ -, 41
 - normed chain -, 65
 - normed dual -, 65
- contracting homotopy, 35, 42
- defect, 12
- Eilenberg-MacLane space, 3
- Euler class
 - of $\text{Homeo}^+(S^1)$, 113
 - of a representation, 115
 - of a sphere bundle, 136
 - of a vector bundle, 138
 - bounded -, 173

- bounded - of $\text{Homeo}^+(S^1)$, 115
- bounded - of a flat circle bundle, 151
- bounded real - of $\text{Homeo}^+(S^1)$, 126
- canonical representative of the bounded real -, 127
- real - of $\text{Homeo}^+(S^1)$, 126
- real - of a sphere bundle, 151
- real bounded - of a flat circle bundle, 151
- Euler cochain, 171
- Euler number
 - of a sphere bundle, 137
 - of representations of punctured surface groups, 155
- extension
 - central -, 9
 - equivalent -, 9
- foliation
 - transverse -, 147
- fundamental
 - class, 77
 - coclass, 82
- general position, 170
- good lift, 120
- group
 - amenable -, 23
 - elementary amenable -, 26
- Haar measure, 93
- holonomy
 - of a flat bundle, 147
 - representation, 163
- homogeneous
 - 2-cocycle, 16
 - quasimorphism, 13
- homogeneous complex, 1
- homology
 - ℓ^1 -, 66
 - relative singular -, 60
 - singular -, 3
- homotopy
 - Γ -, 35
 - contracting -, 35, 42
- inhomogeneous complex, 6
- injective
 - relatively -, 33, 41
 - strongly -, 33, 41
- Kronecker product, 66
- manifold
 - affine -, 178
 - complete affine -, 178
 - special affine -, 180
 - tangentially flat -, 178
- map
 - chain - between Γ -complexes, 35
 - change of coefficients -, 2
 - comparison -, 5, 53, 70
 - increasing of degree one -, 119
- mean, 23
 - bi-invariant -, 24
 - invariant -, 23
 - left-invariant -, 23
- Milnor-Wood inequalities, 153
 - for punctured surfaces, 162
- minimal representation, 121
- minimal volume, 84
- module
 - $R[\Gamma]$ -, 1
 - dual normed $R[\Gamma]$ -, 27
- norm
 - ℓ^1 -, 66, 67
 - ℓ^∞ -, 4
- projective bundle, 138
- quasimorphism, 12
 - bushy -, 21
 - homogeneous -, 13
 - split -, 15
- relatively injective, 33, 41
- representation
 - geometric -, 163
 - holonomy -, 163
 - maximal -, 162
 - minimal -, 121
 - semi-conjugate -, 121
- resolution, 35
 - of a normed $R[\Gamma]$ -module, 42
 - bar -, 6
 - relatively injective -, 35, 42
 - standard -, 36
 - strong -, 35, 42
- rotation number, 116
- semi-conjugate
 - representations, 121
- seminorm
 - ℓ^1 -, 67
 - ℓ^∞ -, 4
 - canonical -, 4
- simplex
 - smooth -, 95
 - straight -, 92, 96
- simplicial volume, 77
- stable integral -, 84
- integral -, 84
- integral foliated -, 85
- Lipschitz -, 86
- space
 - Eilenberg-MacLane -, 3
 - continuously uniquely geodesic -, 91
- sphere G -bundle, 132
 - flat -, 145

- isomorphic -, 132, 146
- sphere bundle, 131
 - associated with a vector bundle, 133
 - with structure group, 131
- flat - with structure group G , 145
- flat linear -, 146
- flat smooth -, 146
- flat topological -, 146
- isomorphic -, 131
- linear -, 132
- linearly isomorphic -, 132
- smooth -, 132
- smoothly isomorphic -, 132
- topological -, 132
- stable commutator length, 20
- straight simplex, 92, 96
- straightening, 92
- strongly injective, 33, 41
- theorem
 - Banach-Alaouglu -, 25
 - Markov-Kakutani -, 25
- uniform boundary condition, 68
- vector bundle
 - flat, 146

List of Symbols

- $\|\cdot\|_\infty$, 4, 5, 53, 66
 $\|\cdot\|_1$, 66, 67
 $\|\cdot\|_1(\theta)$, 73, 74
 $\|\cdot\|_\infty(\theta)$, 74
 $[x]$, 13
 $|\Gamma|$, 25
 alt_b^n , 49
 alt^n , 60
 $B^n(\Gamma, V)$, 2
 $B_b^n(\Gamma, V)$, 4
 $B\Gamma$, 3
 $B^n(E^\bullet)$, 35
 $\overline{B}^n(\Gamma, R)$, 9
 $\overline{B}_b^n(\Gamma, R)$, 9
 $C^n(\Gamma, V)$, 1
 $C_n(X, R)$, 3
 $C^n(X, R)$, 3
 $C_b^n(\Gamma, V)$, 4
 $\overline{C}^n(\Gamma, V)$, 5
 $C_{\text{alt}}^n(\Gamma, V)$, 48
 $C_{b,\text{alt}}^n(\Gamma, V)$, 48
 $C_b^n(X, R)$, 53
 $C_{\text{alt}}^n(X, \mathbb{R})$, 59
 $C_{b,\text{alt}}^n(X, \mathbb{R})$, 59
 $C^n(X, Y)$, 60
 $C_b^n(X, Y)$, 60
 $C_{bs}^n(\tilde{X}, \tilde{Y})$, 61
 $C_{bs}^n(X, Y)$, 61
 $C_n^{\ell_1}(X)$, 66
 $C_n(\Gamma)$, 66
 $C_i^{\ell_1}(\Gamma)$, 67
 $C_c^n(M)$, 87
 $C_{b,c}^n(M)$, 88
 c_{x_0} , 114
 $D(f)$, 12
 $EH_b^n(\Gamma, V)$, 5
 $(E^\bullet, \delta^\bullet)$, 35
 $(E, E^\bullet, \delta^\bullet)$, 35, 42
 $e(\mathcal{C})$, 10
 e , 113
 e_b , 115
 $e(\rho)$, 115
 $e_b(\rho)$, 115
 $e_b^{\mathbb{R}}$, 126
 $e_b^{\mathbb{R}}$, 126
 $e_b^{\mathbb{R}}(\rho)$, 126
 $e_b^{\mathbb{R}}(\rho)$, 126
 $\text{eu}(E)$, 136, 137, 138
 $\text{eu}_b(E)$, 151
 $\text{eu}^{\mathbb{R}}(E)$, 151
 $\text{eu}_b^{\mathbb{R}}(E)$, 151, 174
 E_ρ , 148
 $e_b^{\mathbb{R}}(\rho, \Sigma_{g,n}, \partial\Sigma_{g,n})$, 155
 $e(\rho, \Sigma_{g,n}, \partial\Sigma_{g,n})$, 155
 eul , 171
 $[\text{eul}]_b$, 173
 $H^n(\Gamma, V)$, 2
 $H_n(X, R)$, 3
 $H^n(X, R)$, 3
 $H_b^n(\Gamma, V)$, 4
 $H^n(E^\bullet)$, 35
 $H_b^n(E^\bullet)$, 42
 $H_b^n(X, R)$, 53
 $H^n(X, Y)$, 60
 $H_b^n(X, Y)$, 60
 $H_n(C)$, 65
 $H_b^n(C')$, 65
 $H_b^n(\Gamma)$, 66
 $H_b^n(X)$, 66
 $H_n^{\ell_1}(X)$, 66
 $H_n(\Gamma)$, 66
 $H_i^{\ell_1}(\Gamma)$, 67
 $H_c^n(M)$, 88
 $H_{b,c}^n(M)$, 88

\mathbb{H}^n , 98	$\overline{Z}_b^n(\Gamma, R)$, 9
$\partial\mathbb{H}^n$, 98	$Z^n(E^\bullet)$, 35
$\widetilde{\mathbb{H}^n}$, 98	
$\text{Homeo}_+(S^1)$, 113	
$\text{Homeo}^+(S^1)$, 113	
$\ell^\infty(\Gamma)$, 23	
$\ell^\infty(\Gamma)'$, 23	
$\ell^1(\Gamma)$, 29	
$\ell_0^1(\Gamma)$, 29	
$\ell^\infty(S^{n+1}, V)$, 46	
$K(\Gamma, 1)$, 3	
$[M]_{\mathbb{Z}}$, 77	
$[M]_{\mathbb{R}}$, 77	
$[M]$, 77	
$[M, \partial M]_{\mathbb{Z}}$, 77	
$[M, \partial M]_{\mathbb{R}}$, 77	
$[M, \partial M]_{\mathbb{Z}}$, 77	
$[M, \partial M]^*$, 82	
$\ M\ $, 77	
$\ M, \partial M\ $, 77	
$\ M\ _{\mathbb{Z}}$, 84	
$\ M\ _{\mathbb{Z}}^\infty$, 84	
μ_G , 93	
$\mathbb{P}(E)$, 138	
$Q(\Gamma, R)$, 12	
$Q^h(\Gamma, R)$, 13	
$q\text{-UBC}$, 68	
$R[\Gamma]$, 1	
res^n , 93	
rot , 116	
$\widetilde{\text{rot}}$, 117	
$S_n(Y)$, 39	
${}_sS_k(M)$, 95	
$S(E)$, 133	
Σ_g , 100	
$\Sigma_{g,n}$, 155	
Σ_C , 157	
trans^n , 93	
τ , 127	
V^Γ , 1	
v_n , 78, 79, 98	
Vol_M , 96	
$[\text{Vol}_M]$, 96	
$[\text{Vol}_M]_c$, 96	
$[\widetilde{\text{Vol}}_M]_c^G$, 97	
$Z^n(\Gamma, V)$, 2	
$Z_b^n(\Gamma, V)$, 4	
$\overline{Z}^n(\Gamma, R)$, 9	

Bibliography

- [AEG94] S. Adams, G. A. Elliott, and T. Giordano, *Amenable actions of groups*, Trans. Amer. Math. Soc. **344** (1994), 803–822.
- [AH61] M. F. Atiyah and F. Hirzebruch, *Bott periodicity and the parallelizability of the spheres*, Proc. Cambridge Philos. Soc. **57** (1961), 223–226.
- [Bav91] C. Bavard, *Longueur stable des commutateurs*, Enseign. Math. **37** (1991), 109–150.
- [BBF⁺14] M. Bucher, M. Burger, R. Frigerio, A. Iozzi, C. Pagliantini, and M. B. Pozzetti, *Isometric properties of relative bounded cohomology*, J. Topol. Anal. **6** (2014), no. 1, 1–25.
- [BBF16] M. Bestvina, K. Bromberg, and K. Fujiwara, *Bounded cohomology with coefficients in uniformly convex Banach spaces*, Comment. Math. Helv. **91** (2016), 203–218.
- [BBH⁺17] G. Ben Simon, M. Burger, T. Hartnick, A. Iozzi, and A. Wienhard, *On weakly maximal representations of surface groups*, J. Differential Geom. **105** (2017), 375–404.
- [BBI13] M. Bucher, M. Burger, and A. Iozzi, *A dual interpretation of the Gromov–Thurston proof of Mostow rigidity and volume rigidity for representations of hyperbolic lattices*, Trends in harmonic analysis, Springer INdAM Ser., Springer, Milan, 2013, pp. 47–76.
- [BCG96] G. Besson, G. Courtois, and S. Gallot, *Minimal entropy and mostow’s rigidity theorems*, Ergodic Theory Dynam. Systems **16** (1996), 623–649.
- [BCL] M. Bucher, C. Connell, and J. F. Lafont, *Vanishing simplicial volume for certain affine manifolds*, arXiv:1610.00832, to appear in Proc. Amer. Math. Soc.
- [BE78] R. Bieri and B. Eckmann, *Relative homology and Poincaré duality for group pairs*, J. Pure Appl. Algebra **13** (1978), 277–319.
- [Bea95] A. F. Beardon, *The geometry of discrete groups. corrected reprint of the 1983 original*, Graduate Texts in Mathematics, vol. 91, Springer-Verlag, New York, 1995.
- [BF02] M. Bestvina and K. Fujiwara, *Bounded cohomology of subgroups of mapping class groups*, Geom. Topol. **6** (2002), 69–89.
- [BFH] M. Bucher, R. Frigerio, and T. Hartnick, *A note on semi-conjugacy for circle actions*, Enseign. Math. **62** (2016), 317–360.
- [BFP15] M. Bucher, R. Frigerio, and C. Pagliantini, *The simplicial volume of 3-manifolds with boundary*, J. Topol. **8** (2015), 457–475.
- [BFP17] M. Bucher, R. Frigerio, and C. Pagliantini, *A quantitative version of a theorem by Jungreis*, Geom. Dedicata **187** (2017), 199–218.
- [BFS13] U. Bader, A. Furman, and R. Sauer, *Integrable measure equivalence and rigidity of hyperbolic lattices*, Invent. Math. **194** (2013), 313–379.
- [BG88] J. Barge and E. Ghys, *Surfaces et cohomologie bornée*, Invent. Math. **92** (1988), 509–526.
- [BG92] ———, *Cocycles d’Euler et de Maslov*, Math. Ann. **294** (1992), 235–265.
- [BG11] M. Bucher and T. Gelander, *The generalized Chern conjecture for manifolds that are locally a product of surfaces*, Adv. Math. **228** (2011), 1503–1542.
- [BG13] ———, *Milnor–Wood inequalities for products*, Algebr. Geom. Topol. **13** (2013), 1733–1742.
- [BI07] M. Burger and A. Iozzi, *Bounded differential forms, generalized Milnor–Wood inequality and an application to deformation rigidity*, Geom. Dedicata **125** (2007), 1–23.
- [BI09] ———, *A useful formula from bounded cohomology*, Géométries à courbure négative ou nulle, groupes discrets et rigidités, Sémin. Congr., vol. 18, Soc. Math. France, Paris, 2009, pp. 243–292.
- [BIMW08] M. Burger, A. Iozzi, N. Monod, and A. Wienhard, *Bounds for cohomology classes*, Enseign. Math. **54** (2008), 52–54.

- [BIW10] M. Burger, A. Iozzi, and A. Wienhard, *Surface group representations with maximal Toledo invariant*, Ann. of Math. (2) **172** (2010), 517–566.
- [BIW14] ———, *Higher Teichmüller spaces: from $SL(2, \mathbb{R})$ to other Lie groups*, Handbook of Teichmüller theory. Vol. IV, IRMA Lect. Math. Theor. Phys., vol. 19, Eur. Math. Soc., Zürich, 2014, pp. 539–618.
- [BK07] M. Bucher-Karlsson, *Simplicial volume of locally symmetric spaces covered by $SL_3\mathbb{R}/SO(3)$* , Geom. Dedicata **125** (2007), 203–224.
- [BK08a] M. Bucher-Karlsson, *The proportionality constant for the simplicial volume of locally symmetric spaces*, Colloq. Math. **111** (2008), 183–198.
- [BK08b] ———, *The simplicial volume of closed manifolds covered by $\mathbb{H}^2 \times \mathbb{H}^2$* , J. Topol. **1** (2008), 584–602.
- [BK14] M. Bucher and I. Kim, *Proportionality principle for the simplicial volume of families of \mathbb{Q} -rank 1 locally symmetric spaces*, Math. Z. **276** (2014), 153–172.
- [Bla16] M. Blank, *Relative bounded cohomology for groupoids*, Geom. Dedicata **184** (2016), 27–66.
- [BM99] M. Burger and N. Monod, *Bounded cohomology of lattices in higher rank Lie groups*, J. Eur. Math. Soc. **1** (1999), 199–235.
- [BM02] ———, *Continuous bounded cohomology and applications to rigidity theory*, Geom. Funct. Anal. **12** (2002), 219–280.
- [BM12] M. Bucher and N. Monod, *The norm of the Euler class*, Math. Ann. **353** (2012), 523–544.
- [Bot75] R. Bott, *Some remarks on continuous cohomology*, Manifolds - Tokyo 1973 (Proc. Internat. Conf., Tokyo, 1973) (Univ. Tokyo Press, ed.), vol. 8, 1975, pp. 161–170.
- [Bou63] N. Bourbaki, *Intégration VII et VIII. Éléments de mathématique*, no. 1306, Hermann, Paris, 1963.
- [Bou95] A. Bouarich, *Suites exactes en cohomologie bornée réelle des groupes discrets*, C. R. Acad. Sci. Paris Sér. I Math. **320** (1995), 1355–1359.
- [Bou04] ———, *Théorèmes de Zilber-Eilenberg et de Brown en homologie l_1* , Proyecciones **23** (2004), 151–186.
- [Bou12] ———, *Expression de la différentielle d_3 de la suite spectrale de Hochschild-Serre en cohomologie bornée réelle*, Extracta Math. **27** (2012), 245–300.
- [BP92] R. Benedetti and C. Petronio, *Lectures on hyperbolic geometry*, Universitext, Springer-Verlag, Berlin, 1992.
- [Bro81] R. Brooks, *Some remarks on bounded cohomology*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978), Ann. of Math. Stud., vol. 97, Princeton Univ. Press, Princeton, N.J., 1981, pp. 53–63.
- [Bro82] K. S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, vol. 87, Springer-Verlag, New York-Berlin, 1982.
- [BS84] R. Brooks and C. Series, *Bounded cohomology for surface groups*, Topology **23** (1984), 29–36.
- [BT82] R. Bott and L. W. Tu, *Differential forms in algebraic topology*, Graduate Texts in Mathematics, vol. 82, Springer-Verlag, 1982.
- [Buc04] M. Bucher, *Characteristic classes and bounded cohomology*, Ph.D. thesis, ETH Zürich, 2004.
- [Buc09] M. Bucher, *Simplicial volume of products and fiber bundles*, Discrete groups and geometric structures, Contemp. Math., vol. 501, Amer. Math. Soc., Providence, RI, 2009, pp. 79–86.
- [Büh11] ———, *On the algebraic foundations of bounded cohomology*, Mem. Amer. Math. Soc. **214** (2011), xxii+97.
- [Bur11] M. Burger, *An extension criterion for lattice actions on the circle*, Geometry, rigidity, and group actions, Chicago Lectures in Math., Univ. Chicago Press, Chicago, IL, 2011, pp. 3–31.
- [Cal07] D. Calegari, *Foliations and the geometry of 3-manifolds*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2007.
- [Cal09a] ———, *Faces of the scl norm ball*, Geom. Topol. **13** (2009), 1313–1336.
- [Cal09b] D. Calegari, *scl*, MSJ series, Mathematical Society of Japan, 2009.

- [CFI16] I. Chatterji, T. Fernós, and A. Iozzi, *The median class and superrigidity of actions on CAT(0) cube complexes*, J. Topol. **9** (2016), 349–400.
- [Cho80] C. Chou, *Elementary amenable groups*, Illinois J. Math. **24** (1980), 396–407.
- [CØ03] J.-L. Clerc and R. Ørsted, *The Gromov norm of the Kaehler class and the Maslov index*, Asian J. Math. **7** (2003), 269–295.
- [CW11] D. Calegari and A. Walker, *Zigurrats and rotation numbers*, J. Mod. Dyn. **5** (2011), 711–746.
- [Day57] M. M. Day, *Amenable semigroups*, Illinois J. Math. **1** (1957), 509–544.
- [DT58] A. Dold and R. Thom, *Quasifaserungen und unendliche symmetrische produkte*, Ann. of Math. **67** (1958), 239–281.
- [DT87] A. Domic and D. Toledo, *The Gromov norm of the Kaehler class of symmetric domains*, Math. Ann. **276** (1987), 425–432.
- [Dug66] J. Dugundji, *Topology*, Allyn and Bacon, Inc., Boston, Mass, 1966.
- [Dup79] J. L. Dupont, *Bounds for characteristic numbers of flat bundles*, Algebraic topology, Aarhus 1978, Lecture Notes in Math, vol. 763, Springer, Berlin, 1979, 1979, pp. 109–119.
- [EF97] D. Epstein and K. Fujiwara, *The second bounded cohomology of word-hyperbolic groups*, Topology **36** (1997), 1275–1289.
- [FFM12] S. Francaviglia, R. Frigerio, and B. Martelli, *Stable complexity and simplicial volume of manifolds*, J. Topology **5** (2012), 977–1010.
- [FFPS] F. Franceschini, R. Frigerio, M.B. Pozzetti, and A. Sisto, *The zero norm subspace of bounded cohomology of acylindrically hyperbolic groups*, arXiv:1703.03752.
- [FK16] K. Fujiwara and M. Kapovich, *On quasihomomorphisms with noncommutative targets*, Geom. Funct. Anal. **26** (2016), 478–519.
- [FLPS16] R. Frigerio, C. Löh, C. Pagliantini, and R. Sauer, *Integral foliated simplicial volume of aspherical manifolds*, Israel J. Math. **216** (2016), 707–751.
- [FM] R. Frigerio and M. Moraschini, *Gromov’s theory of multicomplexes with applications to bounded cohomology and simplicial volume*, in preparation.
- [FM98] B. Farb and H. Masur, *Superrigidity and mapping class groups*, Topology **37** (1998), 1169–1176.
- [FM11] K. Fujiwara and J. K. Manning, *Simplicial volume and fillings of hyperbolic manifolds*, Algebr. Geom. Topol. **11** (2011), 2237–2264.
- [FP10] R. Frigerio and C. Pagliantini, *The simplicial volume of hyperbolic manifolds with geodesic boundary*, Algebr. Geom. Topol. **10** (2010), 979–1001.
- [FP12] ———, *Relative measure homology and continuous bounded cohomology of topological pairs*, Pacific J. Math. **257** (2012), 95–130.
- [Fra] F. Franceschini, *A characterization of relatively hyperbolic groups via bounded cohomology*, arXiv:1505.04465, to appear in Groups Geom. Dyn.
- [Fra04] S. Francaviglia, *Hyperbolic volume of representations of fundamental groups of cusped 3-manifolds*, Int. Math. Res. Not. **9** (2004), 425–459.
- [Fra16] F. Franceschini, *Proportionality principle for the Lipschitz simplicial volume*, Geom. Dedicata **182** (2016), 287–306.
- [Fri11] R. Frigerio, *(Bounded) continuous cohomology and Gromov’s proportionality principle*, Manuscripta Math. **134** (2011), 435–474.
- [FS02] K. Fujiwara and T. Soma, *Bounded classes in the cohomology of manifolds*, Geom. Dedicata **92** (2002), 73–85.
- [Fuj98] K. Fujiwara, *The second bounded cohomology of a group acting on a Gromov-hyperbolic space*, Proc. London Math. Soc. **76** (1998), 70–94.
- [Fuj00] ———, *The second bounded cohomology of an amalgamated free product of groups*, Trans. Amer. Math. Soc. **352** (2000), 1113–1129.
- [Ger92] A. M. Gersten, *Bounded cocycles and combings of groups*, Internat. J. Algebra Comput. **2** (1992), 307–326.
- [Ghy87] E. Ghys, *Groupes d’homéomorphismes du cercle et cohomologie bornée*, The Lefschetz centennial conference, Part III (Mexico City, 1984), Contemp. Math., vol. 58, Amer. Math. Soc., Providence, RI, 1987, pp. 81–106.
- [Ghy99] ———, *Actions de réseaux sur le cercle*, Invent. Math. **137** (1999), 199–231.
- [Ghy01] ———, *Groups acting on the circle*, Enseign. Math. (2) **47** (2001), 329–407.

- [GM87] W. M. Goldman and J. J. Millson, *Local rigidity of discrete groups acting on complex hyperbolic space*, *Invent. Math.* **88** (1987), 495–520.
- [GM07] Y. Glasner and N. Monod, *Amenable actions, free products and a fixed point property*, *Bull. Lond. Math. Soc.* **39** (2007), 138–150.
- [Gol80] W. Goldman, *Discontinuous groups and the Euler class*, Ph.D. thesis, University of California, Berkeley, 1980.
- [Gol88] W. M. Goldman, *Topological components of spaces of representations*, *Invent. Math.* **93** (1988), 557–607.
- [Gol14] ———, *Two papers which changed my life: Milnor’s seminal work on flat manifolds and bundles*, *Frontiers in complex dynamics*, Princeton Math. Ser., vol. 51, Princeton Univ. Press, Princeton, NJ, 2014, pp. 679–703.
- [Gre69] F. P. Greenleaf, *Amenable actions of locally compact groups*, *J. Functional Analysis* **4** (1969), 295–315.
- [Gri95] R. I. Grigorchuk, *Some results on bounded cohomology*, *Combinatorial and geometric group theory* (Edinburgh, 1993), London Math. Soc. Lecture Note Ser., vol. 204, Cambridge Univ. Press, Cambridge, 1995, pp. 111–163.
- [Gro82] M. Gromov, *Volume and bounded cohomology*, *Inst. Hautes Études Sci. Publ. Math.* **56** (1982), 5–99.
- [Gro93] M. Gromov, *Asymptotic invariants of infinite groups*, *Geometric group theory, vol. 2*, London Math. Soc. Lectures Notes Ser., vol. 182, Cambridge Univ. Press, Cambridge, 1993.
- [Gro99] ———, *Metric structures for Riemannian and non-Riemannian spaces*, *Progress in Mathematics*, vol. 152, Birkhäuser, 1999, with appendices by M. Katz, P. Pansu, and S. Semmes, translated by S.M. Bates.
- [Gro09] M. Gromov, *Singularities, expanders and topology of maps. part i: Homology versus volume in the spaces of cycles*, *Geom. Funct. Anal.* **19** (2009), 743–841.
- [GSW02] L. Guijarro, T. Schick, and G. Walschap, *Bundles with spherical Euler class*, *Pacific J. Math.* **207** (2002), 377–391.
- [Ham08] U. Hamenstädt, *Bounded cohomology and isometry groups of hyperbolic spaces*, *J. Eur. Math. Soc.* **10** (2008), 315–349.
- [Han98] S.K. Hansen, *Measure homology*, *Math. Scand.* **83** (1998), 205–219.
- [Hat02] A. Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002.
- [HK01] M. Hoster and D. Kotschick, *On the simplicial volumes of fiber bundles*, *Proc. Amer. Math. Soc.* **129** (2001), 1229–1232.
- [HM81] U. Haagerup and H. J. Munkholm, *Simplices of maximal volume in hyperbolic n -space*, *Acta Math.* **147** (1981), 1–11.
- [HO12] T. Hartnick and A. Ott, *Surjectivity of the comparison map in bounded cohomology for Hermitian Lie groups*, *Int. Math. Res. Not.* (2012), no. 9, 2068–2093.
- [HO13] M. Hull and D. Osin, *Induced quasicocycles on groups with hyperbolically embedded subgroups*, *Algebr. Geom. Topol.* **13** (2013), 2635–2665.
- [HO15] T. Hartnick and A. Ott, *Bounded cohomology via partial differential equations, I*, *Geom. Topol.* **19** (2015), 3603–3643.
- [HS16] T. Hartnick and P. Schweitzer, *On quasiisomorphism groups of free groups and their transitivity properties*, *J. Algebra* **450** (2016), 242–281.
- [HT75] M. W. Hirsch and W. P. Thurston, *Foliated bundles, invariant measures and flat manifolds*, *Ann. Math.* **101** (1975), 369–390.
- [Hub12] T. Huber, *Rotation quasimorphisms for surfaces*, Ph.D. thesis, ETH Zürich, 2012, available at <http://e-collection.library.ethz.ch/eserv/eth:7059/eth-7059-02.pdf>.
- [Hus94] D. Husemoller, *Fibre bundles. Third edition.*, Graduate Texts in Mathematics, vol. 20, Springer-Verlag, New York, 1994.
- [Ioz02] A. Iozzi, *Bounded cohomology, boundary maps, and representations into $\text{Homeo}_+(S^1)$ and $SU(n, 1)$* , *Rigidity in Dynamics and Geometry* (Cambridge, 2000), Springer, Berlin, 2002.
- [IT82] N. V. Ivanov and V. G. Turaev, *The canonical cocycle for the Euler class of a flat vector bundle*, *Dokl. Akad. Nauk SSSR* **265** (1982), 521–524.
- [Iva] N. V. Ivanov, *Notes on the bounded cohomology theory*, arXiv:1708.05150.
- [Iva87] N. V. Ivanov, *Foundation of the theory of bounded cohomology*, *J. Soviet. Math.* **37** (1987), 1090–1114.

- [Iva90] ———, *The second bounded cohomology group*, J. Soviet Math. **52** (1990), 2822–2824.
- [IY82] H. Inoue and K. Yano, *The Gromov invariant of negatively curved manifolds*, Topology **21** (1982), 83–89.
- [Joh72] B. E. Johnson, *Cohomology in Banach algebras*, Mem. Amer. Math. Soc. **127** (1972).
- [Jun97] D. Jungreis, *Chains that realize the Gromov invariant of hyperbolic manifolds*, Ergod. Th. Dynam. Sys. **17** (1997), 643–648.
- [Kai03] V. A. Kaimanovich, *Double ergodicity of the Poisson boundary and applications to bounded cohomology*, Geom. Funct. Anal. **13** (2003), 852–861.
- [KK12] S. Kim and I. Kim, *Simplicial volume of \mathbb{Q} -rank one locally symmetric spaces covered by the product of \mathbb{R} -rank one symmetric spaces*, Algebr. Geom. Topol. **12** (2012), 1165–1181.
- [KK15] S. Kim and T. Kuessner, *Simplicial volume of compact manifolds with amenable boundary*, J. Topol. Anal. (2015), no. 7, 23–46.
- [KL01] B. Kleiner and B. Leeb, *Groups quasi-isometric to symmetric spaces*, Comm. Anal. Geom. **9** (2001), 239–260.
- [Kli] B. Klingler, *Chern’s conjecture for special affine manifolds*, Ann. of Math. (2) **186** (2017), 69–95.
- [KM96] V. A. Kaimanovich and H. Masur, *The Poisson boundary of the mapping class group*, Invent. Math. **125** (1996), 221–264.
- [KS69] R. Kirby and L. C. Siebenmann, *On the triangulation of manifolds and the Hauptvermutung*, Bull. Amer. Math. Soc. **75** (1969), 742–749.
- [KS75] B. Kostant and D. Sullivan, *The Euler characteristic of an affine space form is zero*, Bull. Amer. Math. Soc. **8** (1975), 937–938.
- [KT68] F. Kamber and P. Tondeur, *Flat manifolds*, Lecture notes in mathematics **67** (1968), iv + 53 pp.
- [Kue15] T. Kuessner, *Multicomplexes, bounded cohomology and additivity of simplicial volume*, Bull. Korean Math. Soc. (2015), no. 52, 1855–1899.
- [Lee03] J. M. Lee, *Introduction to smooth manifolds*, Graduate Texts in Mathematics, no. 218, Springer-Verlag, New York, 2003.
- [Löh] C. Löh, *Group cohomology and bounded cohomology*, available at http://www.math.uni-muenster.de/u/clara.loeh/topologie3_ws0910/prelim.pdf.
- [Löh06] C. Löh, *Measure homology and singular homology are isometrically isomorphic*, Math. Z. **253** (2006), 197–218.
- [Löh07] C. Löh, *Homology and simplicial volume*, Ph.D. thesis, WWU Münster, 2007, available online at <http://nbn-resolving.de/urn:nbn:de:hbz:6-37549578216>.
- [Löh08] C. Löh, *Isomorphisms in l^1 -homology*, Münster J. Math. **1** (2008), 237–266.
- [Löh16] ———, *Finite functorial semi-norms and representability*, Int. Math. Res. Notices **2016** (12) (2016), 3616–3638.
- [Löh17] C. Löh, *A note on bounded-cohomological dimension of discrete groups*, J. Math. Soc. Japan **69** (2017), 715–734.
- [LP16] C. Löh and C. Pagliantini, *Integral foliated simplicial volume of hyperbolic 3-manifolds*, Groups Geom. Dyn. **10** (2016), 825–865.
- [LS06] J. F. Lafont and B. Schmidt, *Simplicial volume of closed locally symmetric spaces of non-compact type*, Acta Math. **197** (2006), 129–143.
- [LS09a] C. Löh and R. Sauer, *Degree theorems and Lipschitz simplicial volume for non-positively curved manifolds of finite volume*, J. Topology **2** (2009), 193–225.
- [LS09b] ———, *Simplicial volume of Hilbert modular varieties*, Comment. Math. Helv. **84** (2009), 457–470.
- [Luo06] F. Luo, *Continuity of the volume of simplices in classical geometry*, Commun. Cont. Math. **8** (2006), 211–231.
- [LW] J.-F. Lafont and S. Wang, *Barycentric straightening and bounded cohomology*, arXiv:1503.06369, to appear in J. Eur. Math. Soc.
- [Man05] J. F. Manning, *Geometry of pseudocharacters*, Geom. Topol. **9** (2005), 1147–1185.
- [Man15] K. Mann, *Spaces of surface group representations*, Invent. Math. **201** (2015), 669–710.
- [Mat86] S. Matsumoto, *Numerical invariants for semiconjugacy of homeomorphisms of the circle*, Proc. Amer. Math. Soc. **98** (1986), 163–168.
- [Mat87] S. Matsumoto, *Some remarks on foliated S^1 -bundles*, Invent. Math. **90** (1987), 343–358.

- [Mat12] D. V. Mathews, *The hyperbolic meaning of the Milnor-Wood inequality*, Expo. Math. **30** (2012), 49–68.
- [Mat16] S. Matsumoto, *Basic partitions and combination of group actions on the circle: A new approach to a theorem of Kathryn Mann*, Enseign. Math. **62** (2016), 15–47.
- [Mdz09] L. Mdzinarishvili, *Continuous singular cohomology*, Georgian Math. J. **16** (2009), 321–342.
- [Mil58a] J. Milnor, *On the existence of a connection with curvature zero*, Comment. Math. Helv. **21** (1958), 215–223.
- [Mil58b] J. Milnor, *Some consequences of a theorem of Bott*, Ann. of Math. (2) **68** (1958), 444–449.
- [Min01] I. Mineyev, *Straightening and bounded cohomology of hyperbolic groups*, Geom. Funct. Anal. **11** (2001), 807–839.
- [Min02] ———, *Bounded cohomology characterizes hyperbolic groups*, Q. J. Math. **53** (2002), 59–73.
- [Mit84] Y. Mitsumatsu, *Bounded cohomology and l^1 -homology of surfaces*, Topology **23** (1984), 465–471.
- [MM85] S. Matsumoto and S. Morita, *Bounded cohomology of certain groups of homeomorphisms*, Proc. Amer. Math. Soc. **94** (1985), 539–544.
- [Mon01] N. Monod, *Continuous bounded cohomology of locally compact groups*, Lecture notes in Mathematics, no. 1758, Springer-Verlag, Berlin, 2001.
- [Mon06] N. Monod, *An invitation to bounded cohomology*, International Congress of Mathematicians. Vol. II, Eur. Math. Soc., Zürich, 2006, pp. 1183–1211.
- [Mos76] M. A. Mostow, *Continuous cohomology of spaces with two topologies*, Mem. Amer. Math. Soc. **7** (1976), 1–142.
- [MR06] N. Monod and B. Rémy, *Boundedly generated groups with pseudocharacters*, J. London Math. Soc. (2) **73** (2006), 84–108, Appendix to: J. F. Manning, Quasi-actions on trees and property (QFA).
- [MS74] J. W. Milnor and J. D. Stasheff, *Characteristic classes*, Annals of Mathematics Studies, no. 76, Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1974.
- [MS04] N. Monod and Y. Shalom, *Cocycle superrigidity and bounded cohomology for negatively curved spaces*, J. Differential Geom. **67** (2004), 395–455.
- [MS06] ———, *Orbit equivalence rigidity and bounded cohomology*, Ann. of Math. (2) **164** (2006), 825–878.
- [MY07] I. Mineyev and A. Yaman, *Relative hyperbolicity and bounded cohomology*, available online at <http://www.math.uiuc.edu/mineyev/math/art/rel-hyp.pdf>, 2007.
- [Nos91] G. A. Noskov, *Bounded cohomology of discrete groups with coefficients*, Leningrad Math. J. (1991), no. 5, 1067–1084.
- [Ol'80] A. Y. Ol'shanskii, *On the question of the existence of an invariant mean on a group*, Uspekhi Mat. Nauk **35** (1980), no. 4(214), 199–200.
- [OS02] A. Y. Ol'shanskii and M. V. Sapir, *Non-amenable finitely presented torsion by-cyclic groups*, Publ. Math. Inst. Hautes Études Sci. **96** (2002), 43–169.
- [Osi16] D. Osin, *Acylindrically hyperbolic groups*, Trans. Amer. Math. Soc. **368** (2016), 851–888.
- [Par03] H. S. Park, *Relative bounded cohomology*, Topology Appl. **131** (2003), 203–234.
- [Par04] H. Park, *Foundations of the theory of l^1 homology*, J. Korean Math. Soc. **41** (2004), 591–615.
- [Pat88] A. L. T. Paterson, *Amenability*, Mathematical Surveys and Monographs, no. 29, American Mathematical Society, Providence, RI, 1988.
- [Pey02] N. Peyerimhoff, *Simplices of maximal volume or minimal total edge length in hyperbolic space*, J. London Math. Soc. **66** (2002), 753–768.
- [Pie84] J.-P. Pier, *Amenable locally compact groups*, Pure and Applied Mathematics, Wiley, 1984.
- [Poi81] H. Poincaré, *Mémoire sur les courbes définies par une équation différentielle*, Journal de Mathématiques **7 et 8** (1881), 375–422 et 251–296, Oeuvres des Henri Poincaré. Tome I. Gauthier Villars.
- [Rat94] John G. Ratcliffe, *Foundations of hyperbolic manifolds*, Graduate Texts in Mathematics, no. 149, Springer-Verlag, New York, 1994.

- [Rola] P. Rolli, *Quasi-morphisms on free groups*, arXiv:0911.4234.
- [Rolb] ———, *Split quasicocycles*, arXiv:1305.0095.
- [Rud91] W. Rudin, *Functional analysis. second edition*, International Series in Pure and Applied Mathematics, McGraw-Hill, Inc., New York, 1991.
- [Sam99] A. Sambusetti, *Minimal entropy and simplicial volume*, Manuscripta Math. **99** (1999), 541–560.
- [Sau02] R. Sauer, *l^2 -invariants of groups and discrete measured groupoids*, Ph.D. thesis, 2002, http://www.math.uni-muenster.de/u/lueck/homepages/roman_sauer/publ/sauer.pdf.
- [Sch05] M. Schmidt, *l^2 -Betti numbers of \mathcal{R} -spaces and the integral foliated simplicial volume*, Ph.D. thesis, Westfälische Wilhelms-Universität Münster, 2005.
- [Sco78] P. Scott, *Subgroups of surface groups are almost geometric*, J. London Math. Soc. (2) **17** (1978), 555–565.
- [Smi] J. Smillie, *The Euler characteristic of flat bundles*, unpublished preprint.
- [Smi77] J. Smillie, *Flat manifolds with non-zero Euler characteristics*, Comment. Math. Helv. **52** (1977), 453–455.
- [Som81] T. Soma, *The Gromov invariant of links*, Invent. Math. **64** (1981), 445–454.
- [Som98] ———, *Existence of non-Banach bounded cohomology*, Topology **37** (1998), 179–193.
- [Ste51] N. Steenrod, *The topology of fiber bundles*, Princeton University Press, 1951.
- [Sul76] D. Sullivan, *A generalization of Milnor's inequality concerning affine foliations and affine manifolds*, Comment. Math. Helv. **51** (1976), 183–189.
- [Tak] M. Takamura, *Semi-conjugacy and a theorem of Ghys*, unpublished.
- [Thu79] W. P. Thurston, *The geometry and topology of 3-manifolds*, Princeton, 1979, mimeographed notes.
- [vN29] J. von Neumann, *Zur allgemeinen theorie der massen*, Fund. Math. **13** (1929), 73–116.
- [Woo71] J. W. Wood, *Bundles with totally disconnected structure group*, Comment. Math. Helv. **46** (1971), 257–273.
- [Zas98] A. Zastrow, *On the (non)-coincidence of Milnor-Thurston homology theory with singular homology theory*, Pacific J. Math. (1998), 369–396.
- [Zim78] R. J. Zimmer, *Amenable ergodic group actions and an application to Poisson boundaries of random walks*, J. Funct. Anal. **27** (1978), 350–372.
- [Zim84] ———, *Ergodic theory and semisimple groups*, Monographs in Mathematics, vol. 81, Birkhäuser Verlag, Basel, 1984.

Selected Published Titles in This Series

- 227 **Roberto Frigerio**, Bounded Cohomology of Discrete Groups, 2017
- 222 **Alexandru Buium**, Foundations of Arithmetic Differential Geometry, 2017
- 221 **Dennis Gaitsgory and Nick Rozenblyum**, A Study in Derived Algebraic Geometry, 2017
- 219 **Richard Evan Schwartz**, The Projective Heat Map, 2017
- 218 **Tushar Das, David Simmons, and Mariusz Urbański**, Geometry and Dynamics in Gromov Hyperbolic Metric Spaces, 2017
- 217 **Benoit Fresse**, Homotopy of Operads and Grothendieck–Teichmüller Groups, 2017
- 216 **Frederick W. Gehring, Gaven J. Martin, and Bruce P. Palka**, An Introduction to the Theory of Higher-Dimensional Quasiconformal Mappings, 2017
- 215 **Robert Bieri and Ralph Strebel**, On Groups of PL-homeomorphisms of the Real Line, 2016
- 214 **Jared Speck**, Shock Formation in Small-Data Solutions to 3D Quasilinear Wave Equations, 2016
- 213 **Harold G. Diamond and Wen-Bin Zhang (Cheung Man Ping)**, Beurling Generalized Numbers, 2016
- 212 **Pandelis Dodos and Vassilis Kanellopoulos**, Ramsey Theory for Product Spaces, 2016
- 211 **Charlotte Hardouin, Jacques Sauloy, and Michael F. Singer**, Galois Theories of Linear Difference Equations: An Introduction, 2016
- 210 **Jason P. Bell, Dragos Ghioca, and Thomas J. Tucker**, The Dynamical Mordell–Lang Conjecture, 2016
- 209 **Steve Y. Oudot**, Persistence Theory: From Quiver Representations to Data Analysis, 2015
- 208 **Peter S. Ozsváth, András I. Stipsicz, and Zoltán Szabó**, Grid Homology for Knots and Links, 2015
- 207 **Vladimir I. Bogachev, Nicolai V. Krylov, Michael Röckner, and Stanislav V. Shaposhnikov**, Fokker–Planck–Kolmogorov Equations, 2015
- 206 **Bennett Chow, Sun-Chin Chu, David Glickenstein, Christine Guenther, James Isenberg, Tom Ivey, Dan Knopf, Peng Lu, Feng Luo, and Lei Ni**, The Ricci Flow: Techniques and Applications: Part IV: Long-Time Solutions and Related Topics, 2015
- 205 **Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik**, Tensor Categories, 2015
- 204 **Victor M. Buchstaber and Taras E. Panov**, Toric Topology, 2015
- 203 **Donald Yau and Mark W. Johnson**, A Foundation for PROPs, Algebras, and Modules, 2015
- 202 **Shiri Artstein-Avidan, Apostolos Giannopoulos, and Vitali D. Milman**, Asymptotic Geometric Analysis, Part I, 2015
- 201 **Christopher L. Douglas, John Francis, André G. Henriques, and Michael A. Hill, Editors**, Topological Modular Forms, 2014
- 200 **Nikolai Nadirashvili, Vladimir Tkachev, and Serge Vlăduț**, Nonlinear Elliptic Equations and Nonassociative Algebras, 2014
- 199 **Dmitry S. Kaliuzhnyi-Verbovetskyi and Victor Vinnikov**, Foundations of Free Noncommutative Function Theory, 2014
- 198 **Jörg Jahnel**, Brauer Groups, Tamagawa Measures, and Rational Points on Algebraic Varieties, 2014
- 197 **Richard Evan Schwartz**, The Octagonal PETs, 2014

For a complete list of titles in this series, visit the
AMS Bookstore at www.ams.org/bookstore/survseries/.

The theory of bounded cohomology, introduced by Gromov in the late 1980s, has had powerful applications in geometric group theory and the geometry and topology of manifolds, and has been the topic of active research continuing to this day. This monograph provides a unified, self-contained introduction to the theory and its applications, making it accessible to a student who has completed a first course in algebraic topology and manifold theory. The book can be used as a source for research projects for master's students, as a thorough introduction to the field for graduate students, and as a valuable landmark text for researchers, providing both the details of the theory of bounded cohomology and links of the theory to other closely related areas.



© www.rccandoguidi.com

The first part of the book is devoted to settling the fundamental definitions of the theory, and to proving some of the (by now classical) results on low-dimensional bounded cohomology and on bounded cohomology of topological spaces. The second part describes applications of the theory to the study of the simplicial volume of manifolds, to the classification of circle actions, to the analysis of maximal representations of surface groups, and to the study of flat vector bundles with a particular emphasis on the possible use of bounded cohomology in relation with the Chern conjecture. Each chapter ends with a discussion of further reading that puts the presented results in a broader context.

The author manages a near perfect equilibrium between necessary technicalities (always well motivated) and geometric intuition, leading the readers from the first simple definition to the most striking applications of the theory in 13 very pleasant chapters. This book can serve as an ideal textbook for a graduate topics course on the subject and become the much-needed standard reference on Gromov's beautiful theory.

—Michelle Bucher



For additional information
and updates on this book, visit

www.ams.org/bookpages/surv-227

AMS on the Web
www.ams.org

ISBN 978-1-4704-4146-3



9 781470 441463

SURV/227