

**Mathematical
Surveys
and
Monographs**

Volume 233



Sobolev and Viscosity Solutions for Fully Nonlinear Elliptic and Parabolic Equations

N. V. Krylov



**Mathematical
Surveys
and
Monographs**

Volume 233

Sobolev and Viscosity Solutions for Fully Nonlinear Elliptic and Parabolic Equations

N. V. Krylov



EDITORIAL COMMITTEE

2010 *Mathematics Subject Classification.* Primary 35-02; Secondary 35J60, 35K55.

For additional information and updates on this book, visit
www.ams.org/bookpages/surv-233

Library of Congress Cataloging-in-Publication Data

Names: Krylov, N. V. (Nikolai Vladimirovich), author.

Title: Sobolev and viscosity solutions for fully nonlinear elliptic and parabolic equations / N.V. Krylov.

Description: Providence, Rhode Island: American Mathematical Society, [2018] | Series: Mathematical surveys and monographs; volume 233 | Includes bibliographical references and index.

Identifiers: LCCN 2018018540 | ISBN 9781470447403 (alk. paper)

Subjects: LCSH: Viscosity solutions. | Elliptic functions. | Differential equations, Parabolic. | AMS: Partial differential equations – Research exposition (monographs, survey articles). msc | Partial differential equations – Elliptic equations and systems – Nonlinear elliptic equations. msc | Partial differential equations – Parabolic equations and systems – Nonlinear parabolic equations. msc

Classification: LCC QA316 .K6945 2018 | DDC 515/.353--dc23

LC record available at <https://lccn.loc.gov/2018018540>

Copying and reprinting. Individual readers of this publication, and nonprofit libraries acting for them, are permitted to make fair use of the material, such as to copy select pages for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgment of the source is given.

Republication, systematic copying, or multiple reproduction of any material in this publication is permitted only under license from the American Mathematical Society. Requests for permission to reuse portions of AMS publication content are handled by the Copyright Clearance Center. For more information, please visit www.ams.org/publications/pubpermissions.

Send requests for translation rights and licensed reprints to reprint-permission@ams.org.

© 2018 by the American Mathematical Society. All rights reserved.

The American Mathematical Society retains all rights

those granted to the United States Government.

Printed in the United States of America.

∞ The paper used in this book is acid-free and falls within the guidelines established to ensure permanence and durability.

Visit the AMS home page at <https://www.ams.org/>

10 9 8 7 6 5 4 3 2 1 23 22 21 20 19 18

Contents

Preface	vii
Chapter 1. Bellman's equations with constant “coefficients” in the whole space	1
1. Finite-difference equations	1
2. Solvability of Bellman's equations	5
Chapter 2. Estimates in L_p for solutions of the Monge-Ampère type equations	15
1. Some properties of λ -concave functions	15
2. Monge-Ampère type elliptic and parabolic equations in the whole space	22
Chapter 3. The Aleksandrov estimates	31
1. Parabolic uniformly nondegenerate case	33
2. General parabolic case	43
3. Elliptic operators	47
Chapter 4. First results for fully nonlinear equations	55
1. The maximum principle and uniqueness of solutions in Sobolev classes	55
1:1. Parabolic operators	60
1:2. Elliptic operators	64
2. Passage to the limit for nonlinear operators in the Sobolev classes	67
Chapter 5. Finite-difference equations of elliptic type	77
1. Some elementary technical tools	77
2. Solvability of elliptic finite-difference equations	78
3. The way we proceed further	83
4. Some estimates for finite-difference elliptic equations with constant coefficients	85
5. Estimates for finite-difference elliptic equations of cut-off type with variable coefficients	92
Chapter 6. Elliptic differential equations of cut-off type	101
1. A particular case of equations in pure derivatives	101
2. General elliptic equations with Lipschitz continuous H	108
3. Minimax representation of certain functions	111
4. One step closer to nonlinear partial-differential operators	116
5. Proof of Theorem 6.2.6	120
6. The case of H measurable in space variables	122

6:1. Some generalizations	124
6:2. Proof of Theorem 6.6.3	126
7. Proof of Theorem 6.6.20	130
 Chapter 7. Finite-difference equations of parabolic type	135
1. Solvability of parabolic finite-difference equations	135
2. Some estimates for second-order finite-differences of solutions to parabolic equations with constant coefficients	138
3. Maximum principle for semi-discrete parabolic equations	142
4. Estimates for finite-difference parabolic equations with variable coefficients	144
 Chapter 8. Parabolic differential equations of cut-off type	151
1. A particular case of equations in pure derivatives	151
2. General parabolic equations with Lipschitz continuous H	155
3. The case of H measurable in time and space variables	158
4. Proof of Theorems 8.3.10 and 6.6.5	162
 Chapter 9. A priori estimates in C^α for solutions of linear and nonlinear equations	165
1. Estimating the integrals of Green's functions over small sets	165
2. Auxiliary results	168
3. Proof of Theorems 9.1.1 and 9.1.2	173
4. Fang-Hua Lin's type estimates for elliptic and parabolic equations	177
5. Estimating functions that are \mathcal{L} -harmonic outside small sets	182
6. The Harnack inequality and the interior Hölder continuity for \mathcal{L} -harmonic functions	187
7. On the global modulus of continuity of solutions of nonlinear elliptic equations	196
8. On the global modulus of continuity of solutions of nonlinear parabolic equations	199
 Chapter 10. Solvability in $W_{p,\text{loc}}^2$ of fully nonlinear elliptic equations	207
1. Main results	207
2. A priori estimates in $C_{\text{loc}}^{2+\alpha}$ for equations $F(D^2u) = 0$	216
3. Solvability in $C_{\text{loc}}^{2+\alpha}$ for equations $F(D^2v) = 0$ and estimates of sharp functions of D^2u	222
4. A priori estimates in $W_{p,\text{loc}}^2$	225
5. Proof of Theorems 10.1.10, 10.1.14, and 10.1.17	230
 Chapter 11. Nonlinear elliptic equations in $C_{\text{loc}}^{2+\alpha}(\Omega) \cap C(\bar{\Omega})$	235
1. Main results	235
2. Characterizations of the spaces $C^{s+\alpha}$	236
3. Proof of Theorems 1.4 and 1.6	246
 Chapter 12. Solvability in $W_{p,\text{loc}}^{1,2}$ of fully nonlinear parabolic equations	251
1. Main results	251
2. A priori estimates in $C_{\text{loc}}^{2+\alpha}$ for equations $\partial_t u + F(D^2u) = 0$	256
3. Solvability in $C_{\text{loc}}^{2+\alpha}$ for equations $\partial_t v + F(D^2v) = 0$ and estimates of the integral oscillations of D^2u	261

4. A priori estimates in $W_{p,\text{loc}}^{1,2}$	263
5. Proof of Theorems 12.1.7 and 12.1.10	266
 Chapter 13. Elements of the $C^{2+\alpha}$ -theory of fully nonlinear elliptic and parabolic equations	 271
1. Introduction	271
2. Characterizations of the spaces $C^{s+\alpha}(B_R^+)$	274
3. Solvability of model linear equations in $\hat{C}^{2+\alpha}(\mathcal{B}_R)$	276
4. Solvability of the Laplace equation in $\hat{C}^{2+\alpha}(\mathcal{B}_R^+)$	281
5. A priori estimates in $\hat{C}^{2+\alpha}(B_R)$ and $\hat{C}^{2+\alpha}(\mathcal{B}_R^+)$ for linear equations with variable coefficients	286
6. Solvability in $\hat{C}^{2+\alpha}(\mathcal{B}_R^+)$ of linear equations with variable coefficients	289
7. About the oscillation of v/x^1 near the flat boundary of B_R^+	291
8. A priori estimates in $\hat{C}^{2+\alpha}(\mathcal{B}_1^+)$ for equations $F(D^2u, x) = 0$	296
9. The method of continuity	300
10. Proof of Theorem 13.1.4	303
11. Solvability in $\hat{C}^{2+\alpha}(\mathcal{B}_1^+)$ for $F(D^2u) = f(x)$	305
12. Proof of Theorem 13.1.6	307
 Chapter 14. Nonlinear elliptic equations in $W_p^2(\Omega)$	 311
1. Main results	311
2. Boundary a priori estimates in the simplest case	315
3. A priori estimates in W_p^2 near the boundary and the proof of Theorem 14.1.3	320
4. Investigating when $W_{p,\text{loc}}^2(\tilde{B}_R^+)$ -solutions belong to $W_p^2(\tilde{B}_\rho^+)$	325
5. Proof of Theorem 14.1.5	326
 Chapter 15. Nonlinear parabolic equations in $W_p^{1,2}$	 329
1. Main results	329
2. Boundary a priori estimates in the simplest case	331
3. A priori estimates in $W_p^{1,2}$ near the boundary and the proof of Theorem 15.1.3	335
4. Proof of Theorem 15.1.4	339
 Chapter 16. $C^{1+\alpha}$ -regularity of viscosity solutions of general parabolic equations	 341
1. Generalities	341
2. Main results	343
3. Auxiliary results about linear equations	348
4. Estimating the difference of solutions of two different equations	351
5. Main estimate for solutions of an auxiliary cut-off equation	358
6. Proof of Theorem 16.2.8	360
 Chapter 17. $C^{1+\alpha}$ -regularity of L_p -viscosity solutions of the Isaacs parabolic equations with almost VMO coefficients	 367
1. Main results	367
2. Auxiliary facts	370
3. Main estimate for solutions of (17.2.1)	372

4. Estimating the C^κ -norm of solutions of (8.3.1)	376
5. Proof of Theorem 17.1.5	379
Chapter 18. Uniqueness and existence of extremal viscosity solutions for parabolic equations	381
1. Uniqueness of C -viscosity solutions of parabolic equations	381
2. Proof of Lemma 18.1.7	385
3. Existence of maximal and minimal L_p -viscosity solutions	395
4. Uniqueness of L_p -viscosity solutions for parabolic Isaacs equations	400
Appendix A. Proof of Theorem 6.2.1	405
Appendix B. Proof of Lemma 9.2.6	409
Appendix C. Some tools from Real Analysis	413
1. Partitions, maximal and sharp functions	413
1:1. Partitions and stopping times	413
1:2. Maximal functions	416
1:3. Sharp functions	418
2. Sharp functions in a more classical settings	421
3. Restricted sharp functions in the presence of lateral boundaries	427
Bibliography	431
Index	439

Preface

The main sources of nonlinear second-order elliptic and parabolic equations are the theory of differential equations and its numerous applications. At present, the class of quasilinear equations, which are linear with respect to the second-order derivatives of the unknown function, is studied almost as thoroughly as the class of linear equations. The results of these studies are summarized in many articles and books; see, for instance, Ladyzhenskaya and Ural'tseva [86], Ladyženskaja, Solonnikov, and Ural'ceva [87], Ivanov [49], Gilbarg and Trudinger [45], and Lieberman [89].

For the fully nonlinear equations, the main subject of this book, only sparse results were available for a long time, not allowing one to build any general theory. However, the equations on the plane were an exception. There, using some specific features available only for two-dimensional space, one was able to develop the solvability theory in $W_p^{1,2}$ and W_p^2 for any p close to 2 (see, for example, [53]).

With no general theory at hand, some classes of fully nonlinear equations were investigated in the areas where they arose by using available methods in these areas. The first class of such equations is related to the famous Monge-Ampère equation which arose in geometry and was studied for quite some time by using the theory of convex surfaces. Aleksandrov [1] in 1958 introduced the notion of generalized solutions of Monge-Ampère equations. Its smoothness, before 1971, was proved only in dimension two (see Bakel'man [7]). Then, Pogorelov [98, 99] proved *interior* smoothness for any dimension (see also Cheng and Yau [19], who pointed out a vicious circle in some arguments of Pogorelov). In the multidimensional case, the smoothness up to the boundary was proved only after an approach to *general* nonlinear equations was developed in the early 1980s.

Another class of nonlinear elliptic and parabolic equations, so-called Bellman's equations, arose from the probabilistic theory of controlled stochastic diffusion processes. When the diffusion is not controlled, Bellman's equations become quasilinear (see Fleming and Rishel [43] and the reference therein). It turned out that by using probabilistic methods, it was possible to prove the solvability of general degenerate nonlinear elliptic Bellman's equations in the whole space in the class of functions with bounded second-order spacial derivatives. Some of these results are presented in [57]. Probably the latest results about general possibly degenerate Bellman's equations in smooth domains are presented in [58] and [63]. In [63], in particular, the solvability of possibly degenerate complex Monge-Ampère equations in strictly pseudo-convex domains is established (this result was earlier obtained by PDE methods in [17] for *nondegenerate* case). Part of [63] is translated into the pure PDE language in [64].

These probabilistic results enabled us to deal with a very large class of fully nonlinear equations. The technique was based to a large degree on the fact that the

Monge-Ampère equation is a particular case of Bellman's equations. However, this theory was very hard to comprehend for people from pure PDEs: it was all presented in probabilistic terms. Up until 1979 no other approaches were found. The probabilistic approach was developed by the author, Nisio, Pragarauskas, Safonov, Lions, Menaldi, and many others.

In 1979 Brézis and Evans [11] proved the existence of a $C^{2+\alpha}$ -solution for Bellman's equation with two elliptic operators. The probabilistic approach only provided solutions with two *bounded* derivatives. The results of Evans and Friedman [36], Evans and Lions [38], and Lions [91] published in about 1979 were obtained by the PDE methods, but the solutions there had only bounded derivatives, and the only novelty in comparison with earlier results was that purely analytical methods were used.

A real breakthrough came in 1982 when Evans [34] and the author [59] independently proved the solvability in $C^{2+\alpha}$ of a wide class of fully nonlinear equations. In [34] elliptic equations are considered like

$$H(u(x), Du(x), D^2u(x), x) = 0, \quad (1)$$

and in [59] both elliptic equations like (1) and parabolic ones like

$$\partial_t u(t, x) + H(u(t, x), Du(t, x), D^2u(t, x), t, x) = 0 \quad (2)$$

are investigated. The proofs were presented in pure PDE terms. The starting point for these proofs was the fact that the solutions of *linear* equations with measurable coefficients are Hölder continuous. This result was obtained by Safonov and the author in [83] and [84]. We present the results of [84] in this book in almost the same way as it is done in [61]; however, we go slightly deeper, which enables us to obtain Fang-Hua Lin's type estimates for elliptic and parabolic equations. These estimates play a crucial role in our approach to the Sobolev space theory.

The results in [84] opened the way to build a solvability theory in classes $C^{2+\alpha}$ of equations like (1) and (2) with H which is *convex or concave* with respect to D^2u and is in C^2 with respect to x or in $C^{1,2}$ with respect to (t, x) . We refer the reader to [45] and [61] and the references therein for information on what was achieved in this area.

The next breakthrough came when Safonov ([102], [103], and [104]) came up with a technique allowing one to treat the H 's that are only Hölder continuous with respect to the independent variables, which made the whole $C^{2+\alpha}$ -theory of fully nonlinear equations as satisfactory as in the case of linear equations. Safonov's approach is also remarkable in the respect that it works equally well for linear and fully nonlinear equations. In our exposition here we also use these techniques.

Then, naturally arose the interest in equations when H is neither convex nor concave with respect to D^2u . These equations often appear as the Isaacs equations in the theory of stochastic differential games. By now we know that generally one can only hope that their solutions are in $C^{1+\alpha}$ for an $\alpha \in (0, 1)$ (see [96]). In these circumstances the notion of the so-called viscosity solutions introduced by Lions [92] turned out to be quite effective. This notion does not require the solutions to have any derivatives. Viscosity solutions later turned out to exist and be unique in a wide variety of cases. At present the theory of viscosity solutions is quite a popular subject. The reader will find three chapters in this book devoted to this theory.

The next dramatic turn of the theory came from groundbreaking results by Caffarelli [12] in 1989 (see also [14]) about a priori estimates in Sobolev W_p^2 classes. His discovery is comparable with the discoveries in [34] and [59], which were the starting points of the $C^{2+\alpha}$ -theory. He works in the framework of the theory of viscosity solutions and proves that under some conditions any viscosity solution is in $W_{p,\text{loc}}^2$. The reader is referred to [23], [51], [52], and [25] for more detailed information about the results obtained by many authors in this theory. We also provide some additional information later in the text while comparing our results with the ones obtained by using the theory of viscosity solutions. The major difference is well seen when the equations are just linear. Then the main coefficients in the theory of viscosity solutions are assumed to be uniformly close to uniformly continuous functions (usual requirement in the classical L_p -theory for linear equations), and in our theory it quite suffices for them to only be in VMO or almost in VMO; that is, to just have sufficiently small integral oscillations over small balls (elliptic case) or cylinders (parabolic case).

The main object of studies in this book is the first boundary-value problem for general fully nonlinear second-order uniformly elliptic and parabolic equations like (1) or (2). Apart from the simplest degenerate Bellman's equations in the whole space with constant coefficients, like equations of Monge-Ampère type, which admit solutions with bounded second-order derivatives, we do not say much about any *specific* class of equation degenerate or not, although we provide quite a few examples of equations to illustrate our assumptions and results. These are given in 1.2.9, 4.1.20, 4.1.21, 6.6.11, 6.6.12, 10.1.1, 10.1.2, 10.1.5, 10.1.18 10.1.24, 10.1.25, 12.1.1, 12.1.14, 12.1.20, 16.2.10, 17.1.6, 18.1.4, C.1.3, C.1.4. Some examples are also given at the end of Section 1.2.

We are looking for solutions in Sobolev classes or for C -viscosity or L_p -viscosity solutions and present *basic solvability results* about such *general* equations. In particular, we do not discuss the remarkable results of Caffarelli [13] about Sobolev solutions of the Monge-Ampère equations continued by many researchers (see [28] and the references therein).

Most of the auxiliary results are taken from old sources ([61], [103], and [104]), and the core of the main new results was obtained in the last few years. Exposition of these results is based on a generalization of the Fefferman-Stein theorem, Fang-Hua Lin's type estimates, and the so-called “ersatz” existence theorems, saying that one can slightly modify “any” equation and get a “cut-off” equation that will have solutions with bounded derivatives. Here is an example of such “ersatz” existence theorem (cf. Theorem 6.6.3).

1. ASSUMPTION. We are given a domain $\Omega \subset \mathbb{R}^d$ which is bounded and which satisfies the exterior ball condition (see Definition 5.5.3). We are also given a function $g \in C^{1,1}(\mathbb{R}^d)$.

Let \mathbb{S} be the set of symmetric $d \times d$ matrices. Suppose that we are given a Borel measurable function $H(\mathbf{u}, x)$,

$$\mathbf{u} = (\mathbf{u}', \mathbf{u}''), \quad \mathbf{u}' = (\mathbf{u}'_0, \mathbf{u}'_1, \dots, \mathbf{u}'_d) \in \mathbb{R}^{d+1}, \quad \mathbf{u}'' \in \mathbb{S}, \quad x \in \mathbb{R}^d.$$

2. ASSUMPTION. (i) The function $H(\mathbf{u}, x)$ is Lipschitz continuous with respect to \mathbf{u}'' , and at all points of differentiability of H with respect to \mathbf{u}'' we have $D_{\mathbf{u}''} H \in \mathbb{S}_\delta$ (a bounded set of strictly positive symmetric $d \times d$ matrices),

(ii) For a constant $K_0 \in [0, \infty)$, the number

$$\bar{H} := \sup_{\mathbf{u}', x} (|H(\mathbf{u}', 0, x)| - K_0 |\mathbf{u}'|)$$

is finite,

(iii) The function $H(\mathbf{u}, x)$ is continuous with respect to \mathbf{u} for any x ,

(iv) The function $H(\mathbf{u}, x)$ is nonincreasing with respect to \mathbf{u}'_0 .

3. THEOREM. *Under the above assumptions, there exists a function $P(\mathbf{u}'')$ which is positive homogeneous of order one and convex with respect to \mathbf{u}'' such that for any $K > 0$ the equation*

$$\max \left[H(u(x), Du(x), D^2u(x), x), P(D^2u(x)) - K \right] = 0$$

(a.e.) in Ω with boundary condition $u = g$ on $\partial\Omega$ has a solution which is continuous in $\bar{\Omega}$, has bounded gradient, and whose second-order derivatives are locally bounded in Ω .

Theorems of that kind allow us to prove the solvability in Sobolev classes for equations that are quite far from the ones that are convex or concave with respect to the Hessian of the unknown functions. They play the role of starting points, which in the $C^{2+\alpha}$ -theory play linear equations allowing one by using the method of continuity to go from linear to fully nonlinear equations. We will see how this works in Chapter 13. The method of continuity consists of moving step-by-step along the parameter connecting a linear equation to the one of interest. It does not work in Sobolev space setting, and the results like Theorem 3 allow one not to use any steps but just send K to infinity and having a priori estimates arrive at solutions of (1).

In the framework of viscosity solutions these approximation theorems also allow us to deal with classical approximating solutions, thus avoiding sometimes very involved constructions from the usual theory of viscosity solutions.

Comments on the book's structure

One of fundamental results of the book is the above Theorem 3 and its parabolic counterpart, Theorem 8.3.3. Their particular cases, when H depends not on the full Hessian D^2u but on finitely many pure second-order derivatives in fixed directions, are given as Theorems 6.1.5 and 8.1.4, respectively. For understanding the proofs of these particular cases the reader need not know *anything* from the theory of linear or nonlinear partial differential equations. Everything is based on rather simple arithmetical manipulations with finite-differences presented in Chapters 5 and 7. Therefore, there was a strong temptation to start the book from scratch with Chapter 5 and in some 35 pages to arrive at a solvability result like Theorem 3 for particular operators H .

Actually, it could even be a good idea for the reader to start with Chapter 5 in order to see why the previous material is needed if one wants to go to more general H .

After we prove the solvability of “cut-off” equations, we send K to infinity, but first we need to obtain some estimates of solutions uniform with respect to K . First, in Chapter 9, come C^α estimates of Krylov-Safonov and Fang-Hua Lin’s type estimates of the integrals of small powers of the magnitude of the second-order derivatives of solutions. The Krylov-Safonov estimates and the fact that we know

the solvability of “cut-off” equations allow us in Chapter 10 to prove Evans-Krylov $C^{2+\alpha}$ -estimates for model equations like $F(D^2u) = 0$ and further, by combining them with Fang-Hua Lin’s type estimates from Section 9.4 and a generalization of the Fefferman-Stein theorem from Appendix C for not necessarily even locally summable functions, to establish in Theorem 10.1.14 the solvability in $W_{p,\text{loc}}^2$ for elliptic equations under rather weak conditions without assuming that H is convex or concave with respect to D^2u . This line of arguments:

Evans-Krylov $C^{2+\alpha}$ -estimates, Fang-Hua Lin’s
type estimates, Fefferman-Stein theorem

is repeated several times like in rondo in different circumstances, each time requiring some new initial information.

For instance, Chapter 11, “Nonlinear elliptic equations in $C_{\text{loc}}^{2+\alpha}(\Omega) \cap C(\bar{\Omega})$ ”, appeared, in particular, because of the need to obtain Evans-Krylov $C^{2+\alpha}$ -estimates for model parabolic equations like $\partial_t u + F(D^2u) = 0$. There were two possibilities: either to reprove Evans-Krylov $C^{2+\alpha}$ -estimates for parabolic equations (following what was done in [61]), or to rewrite the equation as $F(D^2u) = -\partial_t u$, first estimate the C^α -norm of $\partial_t u$, and then consider the equation for any fixed t as an elliptic one. We preferred the second choice because it is more elegant and richer in ideas, but then we needed a theory of solvability of equations $F(D^2u) = f$ in $C^{2+\alpha}$ -spaces, which is insightful in its own rights.

After that in Chapter 12 we are able to basically repeat the reasoning in Chapter 10 and develop the solvability theory for *parabolic* equations in $W_{p,\text{loc}}^{1,2}$.

Then comes the need for solvability theory in global Sobolev spaces W_p^2 for elliptic and $W_p^{1,2}$ for parabolic equations. Everything here depends on a priori estimates near the boundary, so model equations should be considered in Hölder classes of functions in half balls or half cylinders, and this gave rise to a comparatively long Chapter 13, “Elements of the $C^{2+\alpha}$ -theory of fully nonlinear elliptic and parabolic equations”.

Once the boundary a priori estimates are obtained for model equations, again mimicking the line of arguments in Chapter 10, we can then prove the solvability results in W_p^2 spaces in Chapter 14 and in $W_p^{1,2}$ spaces in Chapter 15.

The remaining chapters contain applications of the parabolic counterpart of Theorem 3 to various issues in the theory of viscosity solutions for parabolic equations. It would be interesting to know if all of them, especially Theorem 17.1.5 on the $C^{1+\alpha}$ -regularity of the L_p -viscosity solutions for Isaacs equations with almost VMO leading coefficients, can be proved by means of the theory of viscosity solution.

There are no historical notes or comments after each chapter. These are spread throughout the text and only a few times presented in a more condensed form in Remarks 3.0.3, 3.1.7, 3.3.8, 4.2.10, 6.6.22, 9.1.3, 9.1.7, 9.4.8, 9.6.9, 10.1.23, 10.1.26, 10.2.5, 11.2.5, 12.1.12, 12.1.21, 14.1.9, 15.1.6, 16.2.9, 17.1.7, 18.1.3, 18.3.6, and 18.4.3; and Examples 10.1.24, 16.2.10, and 18.1.4.

Comments on the exposition

The theorems, lemmas, remarks, and such which are part of the main body of the text are numbered serially in a single system that proceeds by section. Theorem 1.7.6 is the sixth numbered unit in the seventh section in the first chapter. In the course of Chapter 1, this theorem is referred to as Theorem 7.6, and in the course

of the seventh section of Chapter 1, it is referred to as Theorem 6. Similarly and independently of these units formulas are numbered and cross-referenced.

In the proofs of various results in this book we use the symbol N sometimes with indices to denote finite nonnegative constants which may change from one occurrence to another, and we do not always specify on which data these constants depend. In these cases the reader should remember that, if in the statement of a result there are constants called N which are claimed to depend only on certain parameters, then in the proof of the result the constants N also depend only on the same parameters unless specifically stated otherwise. Of course, if we write

$$N = N(\dots),$$

this means that N depends only on what is inside the parentheses.

A complete reference list of notation can be found in the index at the end of the book.

In the case of elliptic equations we concentrate only on equations in the Euclidean space

$$\mathbb{R}^d = \{x = (x^1, \dots, x^d) : x^i \in (-\infty, \infty)\}$$

or in domains $\Omega \subset \mathbb{R}^d$. By domains we mean general open sets. If Γ is a measurable subset of \mathbb{R}^d , by $|\Gamma|$ we denote its Lebesgue measure.

The scalar product in \mathbb{R}^d is denoted by

$$\langle \cdot, \cdot \rangle.$$

If a is a matrix of any dimension (for instance, column-vector), we set

$$|a| = [\text{tr } aa^*]^{1/2},$$

where a^* is the transpose of a .

On some occasions, we allow ourselves to use different symbols for the same objects, for example,

$$\begin{aligned} u_{x^i} &= \frac{\partial u}{\partial x^i} = D_i u, & u_{x^i x^j} &= \frac{\partial^2 u}{\partial x^i \partial x^j} = D_{ij} u, \\ u_x &= Du = (D_1 u, \dots, D_d u), & u_{xx} &= D^2 u = (D_{ij} u)_{i,j=1}^d. \end{aligned}$$

Somewhat ambiguously we set

$$D_i u v := (D_i u) v.$$

In the case of parabolic equations we work with

$$\mathbb{R}^{d+1} = \{(t, x) : t \in \mathbb{R}, x \in \mathbb{R}^d\}.$$

If Γ is a measurable subset of \mathbb{R}^{d+1} , by $|\Gamma|$ we denote its Lebesgue measure. If $z_1 = (t_1, x_1), z_2 = (t_2, x_2) \in \mathbb{R}^{d+1}$, we set

$$\rho(z_1, z_2) = |x_1 - x_2| + |t_1 - t_2|^{1/2}$$

and call $\rho(z_1, z_2)$ the parabolic distance between z_1 and z_2 .

For functions $u(t, x)$ given on subdomains of \mathbb{R}^{d+1} we use the above notation only for the derivatives in x ; sometimes we write $D_x u, D_x^2 u$ to emphasize this, and denote

$$\partial_t u = \frac{\partial u}{\partial t} = u_t, \quad \partial_t D_k u = u_{tx^k} = u_{x^k t},$$

and so on.

In various parts of the book we speak about measurable functions and sets: Borel measurable and Lebesgue measurable, and if we say that a particular function or a set is *measurable*, we always mean that it is Lebesgue measurable. On many occasions we consider functions like $F(u, x)$, where u belongs to a Euclidean space, say \mathfrak{U} , and $x \in \mathbb{R}^d$, and require them to be Borel measurable in (u, x) . This is done for simplicity of presentation. Actually, what is needed is for them to be measurable with respect to the product of the Borel σ -field in \mathfrak{U} and the Lebesgue σ -field in \mathbb{R}^d . Generally, it is useful to keep in mind that for any Lebesgue measurable function there exists a Borel measurable one coinciding with it almost everywhere.

If Ω is a domain in \mathbb{R}^d and $p \in [1, \infty]$, by $L_p(\Omega)$ we mean the set of all (Lebesgue) measurable real-valued functions f for which

$$\|f\|_{L_p(\Omega)} := \left(\int_{\Omega} |f(x)|^p dx \right)^{1/p} < \infty$$

with the standard extension of this formula if $p = \infty$. One knows that $L_p(\Omega)$ is a Banach space.

By $C_0^\infty(\Omega)$ we mean the set of all infinitely differentiable functions on Ω with compact support (contained) in Ω . By the *support* of a function we mean the closure of the set where the function is different from zero. We call a subset of \mathbb{R}^d *compact* if it is closed and bounded. Of course, saying “compact support” is the same as saying “bounded support”; we keep “compact” just to remind us that we are talking about *closed* sets. We set

$$C_0^\infty = C_0^\infty(\mathbb{R}^d).$$

Similar notation is introduced for domains in \mathbb{R}^{d+1} and for functions of (t, x) rather than x alone.

We always use the summation convention with respect to repeated indices no matter whether they are at the same level or at different ones, and if there are signs of inf or sup followed by the sums of products of terms with repeated indices, then the summation is performed first, inside the signs of inf or sup, for instance,

$$\inf_{\alpha} [a_i^{\alpha} u_i + b_j^{\alpha} v_j] := \inf_{\alpha} \left[\sum_i a_i^{\alpha} u_i + \sum_j b_j^{\alpha} v_j \right].$$

By \mathbb{S} we mean the set of $d \times d$ symmetric matrices. If $a \in \mathbb{S}$ we usually represent its entries by a^{ij} . However, on many occasions a itself has superscripts, and then we use a_{ij} for the entries of a . In the same vein for numbers u we use the notation

$$u^{\pm} = u_{\pm} = (1/2)(|u| \pm u).$$

We remind the reader that

$$a \wedge b = \min(a, b), \quad a \vee b = \max(a, b).$$

One more general stipulation is that

$$\begin{aligned} \text{positive} &= \text{nonnegative}, & \text{negative} &= \text{nonpositive}, \\ \text{decreasing} &= \text{nonincreasing}, & \text{increasing} &= \text{nondecreasing}. \end{aligned}$$

Acknowledgments

My deepest gratitude is due to Sungbin Lim and Hyunwoo Kwon, who on their own initiative typeset in L^AT_EX my book [61] and sent me the T_EX file, allowing me to use it if I ever go back to the subject of [61]. I used their file in a few places. I am also grateful to the NSF for supporting me in the early stages of writing the book through grant DMS-1160569. Finally, the referees made some comments, which certainly helped improve the presentation.

Nicolai Krylov
Minneapolis, MN
August 2017

Bibliography

- [1] A. D. Aleksandrov, *Dirichlet's problem for the equation $\text{Det}||z_{ij}|| = \varphi(z_1, \dots, z_n, z, x_1, \dots, x_n)$. I* (Russian, with English summary), Vestnik Leningrad. Univ. Ser. Mat. Meh. Astr. **13** (1958), no. 1, 5–24. MR0096903
- [2] A. D. Aleksandrov, *Certain estimates for the Dirichlet problem*, Soviet Math. Dokl. **1** (1961), 1151–1154. English translation in Soviet Math. Dokl., Vol. 1 (1961), 1151–1154. MR0147776
- [3] A. D. Aleksandrov, *Uniqueness conditions and bounds for the solution of the Dirichlet problem* (Russian, with English summary), Vestnik Leningrad. Univ. Ser. Mat. Meh. Astronom. **18** (1963), no. 3, 5–29. English translation in Amer. Mat. Soc. Transl., Vol. 68 (1968), no. 2, 89–119. MR0164135
- [4] A. D. Aleksandrov, *Majorants of solutions of linear equations of order two* (Russian, with English summary), Vestnik Leningrad. Univ. **21** (1966), no. 1, 5–25. English translation in Amer. Mat. Soc. Transl., Vol. 68 (1968), no. 2, 120–143. MR0199540
- [5] S. Agmon, A. Douglis, and L. Nirenberg, *Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I*, Comm. Pure Appl. Math. **12** (1959), 623–727, DOI 10.1002/cpa.3160120405. MR0125307
- [6] I. Ja. Bakel'man, *On the theory of quasilinear elliptic equations* (Russian), Sibirsk. Mat. Ž. **2** (1961), 179–186. MR0126604
- [7] I. Ja. Bakel'man, *Geometrical methods for solving elliptic equations*, Moscow, Nauka, 1965 in Russian.
- [8] J. M. Ball and L. C. Evans, *Weak convergence theorems for nonlinear partial differential equations of first and second order*, J. London Math. Soc. (2) **25** (1982), no. 2, 332–346, DOI 10.1112/jlms/s2-25.2.332. MR653391
- [9] J.-M. Bony, *Principe du maximum dans les espaces de Sobolev* (French), C. R. Acad. Sci. Paris Sér. A-B **265** (1967), A333–A336. MR0223711
- [10] M. Bramanti and M. C. Cerutti, *$W_p^{1,2}$ solvability for the Cauchy-Dirichlet problem for parabolic equations with VMO coefficients*, Comm. Partial Differential Equations **18** (1993), no. 9–10, 1735–1763, DOI 10.1080/03605309308820991. MR1239929
- [11] H. Brézis and L. C. Evans, *A variational inequality approach to the Bellman-Dirichlet equation for two elliptic operators*, Arch. Rational Mech. Anal. **71** (1979), no. 1, 1–13, DOI 10.1007/BF00250667. MR522704
- [12] L. A. Caffarelli, *Interior a priori estimates for solutions of fully nonlinear equations*, Ann. of Math. (2) **130** (1989), no. 1, 189–213, DOI 10.2307/1971480. MR1005611
- [13] L. A. Caffarelli, *Interior $W^{2,p}$ estimates for solutions of the Monge-Ampère equation*, Ann. of Math. (2) **131** (1990), no. 1, 135–150, DOI 10.2307/1971510. MR1038360
- [14] L. A. Caffarelli and X. Cabré, *Fully nonlinear elliptic equations*, American Mathematical Society Colloquium Publications, vol. 43, American Mathematical Society, Providence, RI, 1995. MR1351007
- [15] L. Caffarelli, M. G. Crandall, M. Kocan, and A. Święch, *On viscosity solutions of fully nonlinear equations with measurable ingredients*, Comm. Pure Appl. Math. **49** (1996), no. 4, 365–397, DOI 10.1002/(SICI)1097-0312(199604)49:4<365::AID-CPA3>3.3.CO;2-V. MR1376656
- [16] L. Caffarelli and L. Silvestre, *On the Evans-Krylov theorem*, Proc. Amer. Math. Soc. **138** (2010), no. 1, 263–265, DOI 10.1090/S0002-9939-09-10077-1. MR2550191
- [17] L. Caffarelli, J. J. Kohn, L. Nirenberg, and J. Spruck, *The Dirichlet problem for nonlinear second-order elliptic equations. II. Complex Monge-Ampère, and uniformly elliptic, equations*, Comm. Pure Appl. Math. **38** (1985), no. 2, 209–252, DOI 10.1002/cpa.3160380206. MR780073

- [18] L. Caffarelli and L. Silvestre, *The Evans-Krylov theorem for nonlocal fully nonlinear equations*, Ann. of Math. (2) **174** (2011), no. 2, 1163–1187, DOI 10.4007/annals.2011.174.2.9. MR2831115
- [19] S. Y. Cheng and S. T. Yau, *On the regularity of the Monge-Ampère equation $\det(\partial^2 u / \partial x_i \partial x_j) = F(x, u)$* , Comm. Pure Appl. Math. **30** (1977), no. 1, 41–68, DOI 10.1002/cpa.3160300104. MR0437805
- [20] F. Chiarenza, M. Frasca, and P. Longo, *Interior $W^{2,p}$ estimates for nondivergence elliptic equations with discontinuous coefficients*, Ricerche Mat. **40** (1991), no. 1, 149–168. MR1191890
- [21] F. Chiarenza, M. Frasca, and P. Longo, *$W^{2,p}$ -solvability of the Dirichlet problem for nondivergence elliptic equations with VMO coefficients*, Trans. Amer. Math. Soc. **336** (1993), no. 2, 841–853, DOI 10.2307/2154379. MR1088476
- [22] M. G. Crandall, K. Fok, M. Kocan, and A. Święch, *Remarks on nonlinear uniformly parabolic equations*, Indiana Univ. Math. J. **47** (1998), no. 4, 1293–1326, DOI 10.1512/iumj.1998.47.1561. MR1687138
- [23] M. G. Crandall, H. Ishii, and P.-L. Lions, *User's guide to viscosity solutions of second order partial differential equations*, Bull. Amer. Math. Soc. (N.S.) **27** (1992), no. 1, 1–67, DOI 10.1090/S0273-0979-1992-00266-5. MR1118699
- [24] M. G. Crandall, M. Kocan, P. L. Lions, and A. Święch, *Existence results for boundary problems for uniformly elliptic and parabolic fully nonlinear equations*, Electron. J. Differential Equations (1999), no. 24, 22 pp. MR1696765
- [25] M. G. Crandall, M. Kocan, and A. Święch, *L^p -theory for fully nonlinear uniformly parabolic equations*, Comm. Partial Differential Equations **25** (2000), no. 11–12, 1997–2053, DOI 10.1080/03605300008821576. MR1789919
- [26] J. V. da Silva and E. V. Teixeira, *Sharp regularity estimates for second order fully nonlinear parabolic equations*, Math. Ann. **369** (2017), no. 3–4, 1623–1648, DOI 10.1007/s00208-016-1506-y. MR3713553
- [27] E. De Giorgi, *Sulla differenziabilità e l'analiticità delle estremali degli integrali multipli regolari* (Italian), Mem. Accad. Sci. Torino. Cl. Sci. Fis. Mat. Nat. (3) **3** (1957), 25–43. MR0093649
- [28] G. De Philippis, A. Figalli, and O. Savin, *A note on interior $W^{2,1+\varepsilon}$ estimates for the Monge-Ampère equation*, Math. Ann. **357** (2013), no. 1, 11–22, DOI 10.1007/s00208-012-0895-9. MR3084340
- [29] H. Dong and N. V. Krylov, *On the existence of smooth solutions for fully nonlinear parabolic equations with measurable “coefficients” without convexity assumptions*, Comm. Partial Differential Equations **38** (2013), no. 6, 1038–1068, DOI 10.1080/03605302.2012.756013. MR3053993
- [30] H. Dong, N. V. Krylov, and X. Li, *On fully nonlinear elliptic and parabolic equations with VMO coefficients in domains*, Algebra i Analiz **24** (2012), no. 1, 53–94, DOI 10.1090/S1061-0022-2012-01231-8; English transl., St. Petersburg Math. J. **24** (2013), no. 1, 39–69. MR3013294
- [31] N. Dunford and J. T. Schwartz, *Linear Operators. I. General Theory*, With the assistance of W. G. Bade and R. G. Bartle. Pure and Applied Mathematics, Vol. 7, Interscience Publishers, Inc., New York; Interscience Publishers, Ltd., London, 1958. MR0117523
- [32] L. Escauriaza, *$W^{2,n}$ a priori estimates for solutions to fully nonlinear equations*, Indiana Univ. Math. J. **42** (1993), no. 2, 413–423, DOI 10.1512/iumj.1993.42.42019. MR1237053
- [33] L. C. Evans, *A convergence theorem for solutions of nonlinear second-order elliptic equations*, Indiana Univ. Math. J. **27** (1978), no. 5, 875–887, DOI 10.1512/iumj.1978.27.27059. MR503721
- [34] L. C. Evans, *Classical solutions of fully nonlinear, convex, second-order elliptic equations*, Comm. Pure Appl. Math. **35** (1982), no. 3, 333–363, DOI 10.1002/cpa.3160350303. MR649348
- [35] L. C. Evans, *Some estimates for nondivergence structure, second order elliptic equations*, Trans. Amer. Math. Soc. **287** (1985), no. 2, 701–712, DOI 10.2307/1999671. MR768735
- [36] L. C. Evans and A. Friedman, *Optimal stochastic switching and the Dirichlet problem for the Bellman equation*, Trans. Amer. Math. Soc. **253** (1979), 365–389, DOI 10.2307/1998203. MR536953

- [37] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*, Revised edition, Textbooks in Mathematics, CRC Press, Boca Raton, FL, 2015. MR3409135
- [38] L. C. Evans and P.-L. Lions, *Résolution des équations de Hamilton-Jacobi-Bellman pour des opérateurs uniformément elliptiques* (French, with English summary), C. R. Acad. Sci. Paris Sér. A-B **290** (1980), no. 22, A1049–A1052. MR582494
- [39] E. Fabes, N. Garofalo, and S. Salsa, *A control on the set where a Green's function vanishes*, Colloq. Math. **60/61** (1990), no. 2, 637–647, DOI 10.4064/cm-60-61-2-637-647. MR1096402
- [40] E. Fabes and C. Kenig, *Examples of singular parabolic measures and singular transition probability densities*, Duke Math. J. **48** (1981), no. 4, 845–856, DOI 10.1215/S0012-7094-81-04846-8. MR782580
- [41] E. Fabes and D. Stroock, *The L^p -integrability of Green's functions and fundamental solutions for elliptic and parabolic equations*, Duke Math. J. **51** (1984), no. 4, 997–1016, DOI 10.1215/S0012-7094-84-05145-7. MR771392
- [42] W. Feller, *An Introduction to Probability Theory and Its Applications. Vol. I*, John Wiley & Sons, Inc., New York, N.Y., 1950. MR0038583
- [43] W. H. Fleming and R. W. Rishel, *Deterministic and stochastic optimal control*, Springer-Verlag, Berlin-New York, 1975. Applications of Mathematics, no. 1. MR0454768
- [44] A. Figalli, Y.-H. Kim, and R. J. McCann, *Hölder continuity and injectivity of optimal maps*, Arch. Ration. Mech. Anal. **209** (2013), no. 3, 747–795, DOI 10.1007/s00205-013-0629-5. MR3067826
- [45] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition. MR1814364
- [46] K. K. Golovkin, *On equivalent normalizations of fractional spaces* (Russian), Trudy Mat. Inst. Steklov. **66** (1962), 364–383. MR0154028
- [47] K. K. Golovkin and V. A. Solonnikov, *Estimates for integral operators in translation invariant norms. II* (Russian), Trudy Mat. Inst. Steklov. **92** (1966), 5–30. English translation in Proc. Steklov Inst. Math., 92 (1968), 3–32. MR0205005
- [48] M. de Guzmán, *Differentiation of integrals in R^n* , Lecture Notes in Mathematics, Vol. 481, Springer-Verlag, Berlin-New York, 1975. With appendices by Antonio Córdoba, and Robert Fefferman, and two by Roberto Moriyón. MR0457661
- [49] A. V. Ivanov, *Second-order quasilinear degenerate and nonuniformly elliptic and parabolic equations* (Russian), Trudy Mat. Inst. Steklov. **160** (1982), 285. English translation by American Mathematical Soc., 1984. MR702172
- [50] R. Jensen and A. Święch, *Uniqueness and existence of maximal and minimal solutions of fully nonlinear elliptic PDE*, Commun. Pure Appl. Anal. **4** (2005), no. 1, 199–207. MR2126286
- [51] N. Katzourakis, *An introduction to viscosity solutions for fully nonlinear PDE with applications to calculus of variations in L^∞* , SpringerBriefs in Mathematics, Springer, Cham, 2015. MR3289084
- [52] S. Koike, A beginner guide to the theory of viscosity solutions, June 28, 2012, 2nd edition, available online at <http://www.math.tohoku.ac.jp/~koike/evis2012version.pdf>.
- [53] N. V. Krylov, *Boundedly inhomogeneous nonlinear elliptic and parabolic equations on the plane* (Russian), Mat. Sb. (N.S.) **82** (124) (1970), 99–110. English translation in Math. USSR Sbornik, Vol. 11 (1970), no. 1, 89–99. MR0262662
- [54] N. V. Krylov, *The uniqueness of the solution of Bellman's equation* (Russian), Izv. Akad. Nauk SSSR Ser. Mat. **35** (1971), 1377–1388. English translation in Math. USSR Izvestija, Vol. 5 (1971), no. 6, 1387–1398. MR0292571
- [55] N. V. Krylov, *Certain estimates in the theory of stochastic integrals*, Teoriya Veroyatnostei i ee Primeneniya, Vol. 18 (1973), no. 1, 56–65, Russian; English translation in Theor. Probability Appl., Vol. 18 (1973), no. 1, 54–63.
- [56] N. V. Krylov, *Some estimates for the density of the distribution of a stochastic integral* (Russian), Izv. Akad. Nauk SSSR Ser. Mat. **38** (1974), 228–248. English translation in Math. USSR Izvestija, Vol. 8 (1974), no. 1, 233–254. MR0345206
- [57] N. V. Krylov, *Controlled diffusion processes*, Applications of Mathematics, vol. 14, Springer-Verlag, New York-Berlin, 1980. Translated from the Russian by A. B. Aries. MR601776
- [58] N. V. Krylov, *Some new results in the theory of controlled diffusion processes* (Russian), Mat. Sb. (N.S.) **109(151)** (1979), no. 1, 146–164, 166. English translation in Math. USSR Sbornik, Vol. 37 (1980), no. 1, 133–149. MR538554

- [59] N. V. Krylov, *Boundedly inhomogeneous elliptic and parabolic equations* (Russian), Izv. Akad. Nauk SSSR Ser. Mat. **46** (1982), no. 3, 487–523, 670. English translation in Math. USSR Izvestija, Vol. 20 (1983), no. 3, 459–492. MR661144
- [60] N. V. Krylov, *G-convergence of elliptic operators in nondivergence form* (Russian), Mat. Zametki **37** (1985), no. 4, 522–527, 600. English translation in Math. Notes, Vol. 37 (1985), no. 3-4, 290–292. MR790976
- [61] N. V. Krylov, *Nonlinear elliptic and parabolic equations of the second order*, Mathematics and its Applications (Soviet Series), vol. 7, D. Reidel Publishing Co., Dordrecht, 1987. Translated from the Russian by P. L. Buzytsky [P. L. Buzytskij]. MR901759
- [62] N. V. Krylov, *Estimates of the maximum of the solution of a parabolic equation and estimates of the distribution of a semimartingale* (Russian), Mat. Sb. (N.S.) **130**(172) (1986), no. 2, 207–221, 284. English translation in Math. USSR Sbornik, Vol. 58 (1987), no. 1, 207–222. MR854972
- [63] N. V. Krylov, *Smoothness of the payoff function for a controllable diffusion process in a domain* (Russian), Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 1, 66–96, DOI 10.1070/IM1990v03n01ABEH000603; English transl., Math. USSR-Izv. **34** (1990), no. 1, 65–95. MR992979
- [64] N. V. Krylov, *A theorem on degenerate elliptic Bellman equations in bounded domains*, Differential Integral Equations **8** (1995), no. 5, 961–980. MR1325541
- [65] N. V. Krylov, *Lectures on elliptic and parabolic equations in Hölder spaces*, Graduate Studies in Mathematics, vol. 12, American Mathematical Society, Providence, RI, 1996. Russian translation, “Nauchnaya kniga”, Novosibirsk, 1998. MR1406091
- [66] N. V. Krylov, *Fully nonlinear second order elliptic equations: recent development*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **25** (1997), no. 3-4, 569–595 (1998). Dedicated to Ennio De Giorgi. MR1655532
- [67] N. V. Krylov, *On factorizations of smooth nonnegative matrix-values functions and on smooth functions with values in polyhedra*, Appl. Math. Optim. **58** (2008), no. 3, 373–392, DOI 10.1007/s00245-008-9040-2. MR2456852
- [68] N. V. Krylov, *Lectures on elliptic and parabolic equations in Sobolev spaces*, Graduate Studies in Mathematics, vol. 96, American Mathematical Society, Providence, RI, 2008. MR2435520
- [69] N. V. Krylov, *Second-order elliptic equations with variably partially VMO coefficients*, J. Funct. Anal. **257** (2009), no. 6, 1695–1712, DOI 10.1016/j.jfa.2009.06.014. MR2540989
- [70] N. V. Krylov, *On Bellman’s equations with VMO coefficients*, Methods Appl. Anal. **17** (2010), no. 1, 105–121, DOI 10.4310/MAA.2010.v17.n1.a4. MR2735102
- [71] N. V. Krylov, *On a representation of fully nonlinear elliptic operators in terms of pure second order derivatives and its applications*, Problemy Matemat. Analiza, Vol. 59, July 2011, p. 3–24 in Russian; English translation in Journal of Mathematical Sciences, New York, Vol. 177 (2011), no. 1, 1–26.
- [72] N. V. Krylov, *On the existence of smooth solutions for fully nonlinear elliptic equations with measurable “coefficients” without convexity assumptions*, Methods Appl. Anal. **19** (2012), no. 2, 119–146, DOI 10.4310/MAA.2012.v19.n2.a2. MR3011597
- [73] N. V. Krylov, *Some L_p -estimates for elliptic and parabolic operators with measurable coefficients*, Discrete Contin. Dyn. Syst. Ser. B **17** (2012), no. 6, 2073–2090, DOI 10.3934/dcdsb.2012.17.2073. MR2924453
- [74] N. V. Krylov, *An ersatz existence theorem for fully nonlinear parabolic equations without convexity assumptions*, SIAM J. Math. Anal. **45** (2013), no. 6, 3331–3359, DOI 10.1137/120903762. MR3124896
- [75] N. V. Krylov, *On the existence of W_p^2 solutions for fully nonlinear elliptic equations under relaxed convexity assumptions*, Comm. Partial Differential Equations **38** (2013), no. 4, 687–710, DOI 10.1080/03605302.2012.741177. MR3040680
- [76] N. V. Krylov, *On the rate of convergence of difference approximations for uniformly non-degenerate elliptic Bellman’s equations*, Appl. Math. Optim. **69** (2014), no. 3, 431–458, DOI 10.1007/s00245-013-9228-y. MR3197305
- [77] N. V. Krylov, *On $C^{1+\alpha}$ regularity of solutions of Isaacs parabolic equations with VMO coefficients*, NoDEA Nonlinear Differential Equations Appl. **21** (2014), no. 1, 63–85, DOI 10.1007/s00030-013-0238-x. MR3158801

- [78] N. V. Krylov, *To the theory of viscosity solutions for uniformly parabolic Isaacs equations*, Methods Appl. Anal. **22** (2015), no. 3, 259–280, DOI 10.4310/MAA.2015.v22.n3.a2. MR3404762
- [79] N. V. Krylov, *On the existence of W_p^2 solutions for fully nonlinear elliptic equations under either relaxed or no convexity assumptions*, Commun. Contemp. Math. **19** (2017), no. 6, 1750009, 39, DOI 10.1142/S0219199717500092. MR3691503
- [80] N. V. Krylov, *On the existence of $W_p^{1,2}$ solutions for fully nonlinear parabolic equations under either relaxed or no convexity assumptions*, to appear in Harvard University, Center of Mathematical Sciences and Applications, Nonlinear Equations Publication, <http://arxiv.org/abs/1705.02400>
- [81] N. V. Krylov, *Uniqueness for L_p -viscosity solutions for uniformly parabolic Isaacs equations with measurable lower order terms*, Comm. on Pure Appl. Analysis **17** (2018), no. 6, 2495–2516. DOI:3934/cpaa.2018119.
- [82] N. V. Krylov, *$C^{1+\alpha}$ -regularity of viscosity solutions of general nonlinear parabolic equations*, Problemy Matematicheskogo Analiza, no. 93, June 2018, 3–23; English translation in Journal of Mathematical Sciences, New York (Springer), Vol. 232 (2018), no. 4, 403–427.
- [83] N. V. Krylov and M. V. Safonov, *An estimate for the probability of a diffusion process hitting a set of positive measure* (Russian), Dokl. Akad. Nauk SSSR **245** (1979), no. 1, 18–20. English translation in Soviet Math. Dokl., Vol. 20 (1979), 253–255. MR525227
- [84] N. V. Krylov and M. V. Safonov, *A property of the solutions of parabolic equations with measurable coefficients* (Russian), Izv. Akad. Nauk SSSR Ser. Mat. **44** (1980), no. 1, 161–175, 239. English translation in Math. USSR Izvestija, Vol. 16 (1981), no. 1, 151–164. MR563790
- [85] H. J. Kuo and N. S. Trudinger, *Discrete methods for fully nonlinear elliptic equations*, SIAM J. Numer. Anal. **29** (1992), no. 1, 123–135, DOI 10.1137/0729008. MR1149088
- [86] O. A. Ladyženskaya and N. N. Ural'tseva, *Lineinyye i kvazilineinyye uravneniya èllipticheskogo tipa*. (Russian), Izdat. “Nauka”, Moscow, 1973. Second edition, revised. MR0509265
- [87] O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Ural'ceva, *Linear and quasilinear equations of parabolic type*, Translated from the Russian by S. Smith. Translations of Mathematical Monographs, Vol. 23, American Mathematical Society, Providence, R.I., 1968. MR0241822
- [88] E. M. Landis, *Second order equations of elliptic and parabolic type*, Translations of Mathematical Monographs, vol. 171, American Mathematical Society, Providence, RI, 1998. English translation by Amer. Math. Soc., Providence, RI, 1997. MR1487894
- [89] G. M. Lieberman, *Second order parabolic differential equations*, World Scientific Publishing Co., Inc., River Edge, NJ, 1996. MR1465184
- [90] F.-H. Lin, *Second derivative L^p -estimates for elliptic equations of nondivergent type*, Proc. Amer. Math. Soc. **96** (1986), no. 3, 447–451, DOI 10.2307/2046592. MR822437
- [91] P.-L. Lions, *Équations de Hamilton-Jacobi-Bellman dégénérées* (French, with English summary), C. R. Acad. Sci. Paris Sér. A-B **289** (1979), no. 5, A329–A332. MR552287
- [92] P.-L. Lions, *Optimal control of diffusion processes and Hamilton-Jacobi-Bellman equations. II. Viscosity solutions and uniqueness* (English, with French summary), Comm. Partial Differential Equations **8** (1983), no. 11, 1229–1276, DOI 10.1080/03605308308820301. MR709162
- [93] J. Moser, *A Harnack inequality for parabolic differential equations*, Comm. Pure Appl. Math. **17** (1964), 101–134, DOI 10.1002/cpa.3160170106. MR0159139
- [94] A. I. Nazarov, *Interpolation of linear spaces and estimates for the maximum of a solution for parabolic equations* (Russian), Partial differential equations (Russian), Akad. Nauk SSSR Sibirsk. Otdel., Inst. Mat., Novosibirsk, 1987, pp. 50–72, 132. English translation in <https://arxiv.org/abs/1507.05232>. MR994027
- [95] A. I. Nazarov and N. N. Ural'tseva, *Convex-monotone hulls and an estimate of the maximum of the solution of a parabolic equation* (Russian, with English summary), Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **147** (1985), 95–109, 204–205. English translation in Journal of Soviet Mathematics April 1987, Vol. 37, no. 1, 851–859. MR821477
- [96] N. Nadirashvili and S. Vlăduț, *Singular viscosity solutions to fully nonlinear elliptic equations* (English, with English and French summaries), J. Math. Pures Appl. (9) **89** (2008), no. 2, 107–113, DOI 10.1016/j.matpur.2007.10.004. MR2391642

- [97] E. A. Pimentel and E. V. Teixeira, *Sharp Hessian integrability estimates for nonlinear elliptic equations: an asymptotic approach*, J. Math. Pures Appl. (9) **106** (2016), no. 4, 744–767, DOI 10.1016/j.matpur.2016.03.010. MR3539473
- [98] A. V. Pogorelov, *The Dirichlet problem for the multidimensional analogue of the Monge-Ampère equation* (Russian), Dokl. Akad. Nauk SSSR **201** (1971), 790–793. MR0293228
- [99] A. V. Pogorelov, *The Minkowski multidimensional problem*, V. H. Winston & Sons, Washington, D.C.; Halsted Press [John Wiley & Sons], New York-Toronto-London, 1978. Translated from the Russian by Vladimir Oliker; Introduction by Louis Nirenberg; Scripta Series in Mathematics. MR0478079
- [100] M. V. Safonov, *Harnack's inequality for elliptic equations and Hölder property of their solutions* (Russian), Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **96** (1980), 272–287, 312. English translation in Journal of Soviet Mathematics, Vol. 21 (March 1983), no. 5, 851–863. MR579490
- [101] M. V. Safonov, *An example of the diffusion processes with singular distribution at a fixed time instant*, Abstr. Comm. Third Intern. Vilnius Conf. Probability Theory and Math. Statist., 1981, 133–134.
- [102] M. V. Safonov, *The classical solution of the elliptic Bellman equation* (Russian), Dokl. Akad. Nauk SSSR **278** (1984), no. 4, 810–813. English translation in Soviet Math. Dokl., Vol. 30 (1984), no. 2, 482–485. MR765302
- [103] M. V. Safonov, *Classical solution of second-order nonlinear elliptic equations* (Russian), Izv. Akad. Nauk SSSR Ser. Mat. **52** (1988), no. 6, 1272–1287, 1328, DOI 10.1070/IM1989v03n03ABEH000858; English transl., Math. USSR-Izv. **33** (1989), no. 3, 597–612. MR984219
- [104] M. V. Safonov, *On the boundary value problems for fully nonlinear elliptic equations of second order*, The Australian National University, Mathematics Research Report no. MRR 049-94, Canberra, 1994. <http://www-users.math.umn.edu/~safon002/NOTES/notes.html>
- [105] M. V. Safonov, *Non-divergence elliptic equations of second order with unbounded drift*, Nonlinear partial differential equations and related topics, Amer. Math. Soc. Transl. Ser. 2, vol. 229, Amer. Math. Soc., Providence, RI, 2010, pp. 211–232, DOI 10.1090/trans2/229/13. MR2667641
- [106] E. Schmutz, *Rational points on the unit sphere*, Cent. Eur. J. Math. **6** (2008), no. 3, 482–487, DOI 10.2478/s11533-008-0038-4. MR2425007
- [107] L. Silvestre and B. Sirakov, *Boundary regularity for viscosity solutions of fully nonlinear elliptic equations*, Comm. Partial Differential Equations **39** (2014), no. 9, 1694–1717, DOI 10.1080/03605302.2013.842249. MR3246039
- [108] L. Silvestre and E. V. Teixeira, *Regularity estimates for fully non linear elliptic equations which are asymptotically convex*, Contributions to nonlinear elliptic equations and systems, Progr. Nonlinear Differential Equations Appl., vol. 86, Birkhäuser/Springer, Cham, 2015, pp. 425–438, DOI 10.1007/978-3-319-19902-3-25. MR3494911
- [109] J. Streets and M. Warren, *Evans-Krylov estimates for a nonconvex Monge-Ampère equation*, Math. Ann. **365** (2016), no. 1–2, 805–834, DOI 10.1007/s00208-015-1293-x. MR3498927
- [110] A. Świech, *$W^{1,p}$ -interior estimates for solutions of fully nonlinear, uniformly elliptic equations*, Adv. Differential Equations **2** (1997), no. 6, 1005–1027. MR1606359
- [111] N. S. Trudinger, *Comparison principles and pointwise estimates for viscosity solutions of nonlinear elliptic equations*, Rev. Mat. Iberoamericana **4** (1988), no. 3–4, 453–468, DOI 10.4171/RMI/80. MR1048584
- [112] N. S. Trudinger, *On regularity and existence of viscosity solutions of nonlinear second order, elliptic equations*, Partial differential equations and the calculus of variations, Vol. II, Progr. Nonlinear Differential Equations Appl., vol. 2, Birkhäuser Boston, Boston, MA, 1989, pp. 939–957. MR1034037
- [113] L. Wang, *On the regularity theory of fully nonlinear parabolic equations*, Bull. Amer. Math. Soc. (N.S.) **22** (1990), no. 1, 107–114, DOI 10.1090/S0273-0979-1990-15854-9. MR999619
- [114] L. Wang, *On the regularity theory of fully nonlinear parabolic equations. I*, Comm. Pure Appl. Math. **45** (1992), no. 1, 27–76, DOI 10.1002/cpa.3160450103. MR1135923
- [115] L. Wang, *On the regularity theory of fully nonlinear parabolic equations. II*, Comm. Pure Appl. Math. **45** (1992), no. 2, 141–178, DOI 10.1002/cpa.3160450202. MR1139064

- [116] N. Winter, *$W^{2,p}$ and $W^{1,p}$ -estimates at the boundary for solutions of fully nonlinear, uniformly elliptic equations*, Z. Anal. Anwend. **28** (2009), no. 2, 129–164, DOI 10.4171/ZAA/1377. MR2486925
- [117] V. V. Zhikov and M. M. Sirazhudinov, *G-compactness of a class of second-order nondivergence elliptic operators* (Russian), Izv. Akad. Nauk SSSR Ser. Mat. **45** (1981), no. 4, 718–733, 927. English translation in Mathematics of the USSR-Izvestiya, Vol. 19 (1982), no. 1, 27–40. MR631435
- [118] V. V. Zhikov, S. M. Kozlov, and O. A. Olešnik, *G-convergence of parabolic operators* (Russian), Uspekhi Mat. Nauk **36** (1981), no. 1(217), 11–58, 248. English translation in Russian Mathematical Surveys, Vol. 36 (1981), no. 1, 9–60. MR608940

Index

Sets of functions	
$B(\Omega)$, 80	C_r^+ , 193
$B(\mathbb{R}^d)$, 2	$C_{\tau,r}^+$, 273
$C^{0,1}(G)$, 92, 101	$C^{1,1}$ -domain, 313
$C^{1+\alpha}(Q)$, 236	$C_R^o(t,x)$, 172
$C^{1,1}(G)$, 92, 101	$C_{T,R}^o$, 172
$C^{1,2}(Q)$, 42	$C_{T,R}^o(t,x)$, 172
$C^{2+\alpha}$, 216	C_R , 165
$C^{2+\alpha}(\bar{B}_\rho^+)$, 271	$C_R(t,x)$, 165
$C^{2+\alpha}(C_{\tau,r}^+)$, 273	$C_{T,R}$, 36
$C_{loc}^{2+\alpha}(\mathbb{C}_r^+)$, 274	$C_{T,R}(t,x)$, 165
$C_{loc}^{2+\alpha}(\mathbb{B}_R^+)$, 271	C_r^+ , 181
$C^2(Q)$, 24	D'_H , 111
$C^\infty(Q)$, 31	$E(u)$, 111
$C^\infty(\bar{Q})$, 31	$E_{H,\beta}(v)$, 116
C_0^∞ , xiii	$G(t_0, x_0)$, 33
$C_0^\infty(\Omega)$, xiii	$Q_{ x }^o$, 142
$C_{loc}^\infty(\mathbb{B}_R^+)$, 281	$Q_{h,\Lambda}^o$, 4
$C^\kappa(\Pi)$, 343	Λ , 78, 85, 101
$C_{loc}^\kappa(\Pi)$, 343	Λ_∞^h , 85
$C^n(Q)$, 50	Λ_∞ , 85
$C^n(\bar{Q})$, 50	Λ_n , 85
$L_{1,loc}$, 413	Ω^ρ , 78
L_p , xiii	Π , 31
$W_{p,loc}^{1,2}(Q)$, 31	Π^ρ , 253
$W_p^{1,2}(Q)$, 31	Π_h , 135
$W_{p,loc}^2(\Omega)$, 32	\mathbb{C}_r^+ , 273
$W_p^2(\Omega)$, 32	$\mathbb{C}_{\tau,r}^+$, 273
$W_\infty^{1,2}(\mathbb{R}^{d+1})$, 144	\mathbb{R}^{d+1} , xii, 60
${}^0 W_{d+1}^{1,2}$, 69	\mathbb{R}_+^{d+1} , 331
\mathbb{L} , 413	\mathbb{R}^d , xii
\mathbb{U}_q^p , 185	\mathbb{R}_+^d , 180, 271
\mathbb{U}_q^m , 168	\mathbb{R}_+ , 264
$\mathbb{U}_q^{t,M}$, 166	\mathbb{S} , xiii
$\mathbb{U}_q^{t,m}$, 166	\mathbb{S}_+^1 , 22
$\mathbb{U}_q^{t,p}$, 182	$\mathbb{S}_+^{\leq 1}$, 22
\mathcal{L} -harmonic functions, 276	\mathbb{S}_+ , 22
\mathcal{P}_2^0 , 281	\mathbb{S}_δ , 108
\mathcal{P}_s , 240	\mathbb{Z} , 413
$\mathfrak{H}_{E,\beta}$, 116	\mathcal{E} , 116
\mathfrak{H}_E , 111	\mathcal{F}^0 , 413
$\hat{C}^\alpha(B_R)$, 272	\mathcal{F}_n , 413
$\hat{C}^\alpha(\mathbb{B}_R^+)$, 272	\check{B}_r^+ , 314
$\hat{C}^{2+\alpha}(B_R)$, 272	$\check{B}_r^+(z)$, 314
λ -concave function, 15	\check{C}_R^+ , 333
Sets	$\check{C}_r^+(z)$, 333
B_R , 19	\mathfrak{U}_r , 64
$B_R(x_0)$, 8	\mathfrak{U}^t , 60
B_R^+ , 180, 271	\hat{Q}^o , 138
$B_R^+(x)$, 271	\mathbb{B}_R^+ , 271
	$\mathbb{B}_R^+(x)$, 271
	$\partial'_h \Pi$, 135

- $\partial_\rho \Omega$, 78
 $\partial_{h,\Lambda} Q$, 4
 $\partial_{x^1} B_r^+$, 193
 $\partial_{x^1} B_R^+(x)$, 271
 $\partial_x C_r^+$, 193
 $\partial_x C_{\tau,r}^+$, 273
 Operators
 $D^2 u$, xii
 D_t^s , 237
 $D_{(l)}$, 7
 $D_{(l)}$, 237
 $D_{(l_1)(l_2)}$, 7
 $D_{u_{ij}'}$, 61
 D_l , 79
 D_u , xii
 $H[u]$, 110, 123, 159
 $P[u]$, 110, 123, 159
 $T_{h,l}$, 1
 $\Delta_{h,l}^{(s)}$, 237
 $\Delta_{h,k}$, 85
 $\Delta_{h,l}$, 1
 Δ_k , 85
 Mh , 227, 265
 $\mathcal{H}[u]$, 102, 152
 $\mathcal{H}_{K,h}$, 93
 \mathcal{H}_K , 93
 $\mathcal{H}_h[v]$, 79
 \mathcal{L} -type operator, 68
 \mathcal{L}_h^α , 1
 $\mathcal{M}f$, 416
 $\mathcal{P}[u]$, 85, 102, 152
 $\mathcal{P}_h[u]$, 85
 f , 211, 413
 δ -nondegenerate operator, 68
 $\delta_h^2 v$, 79
 $\delta_{h,k}$, 85
 $\delta_{h,l}$, 1
 $\delta_h v$, 79
 δ_k , 85
 $\partial_t u$, xii
 $c^m[u]$, 16
 $c_{ij}^m[u]$, 16
 u_t , xii
 $u_{x^i x^j}$, xii
 u_{x^i} , xii
 u_{xx} , xii
 u_x , xii
 Sets of operators
 $\mathbb{L}_{\delta,K}^\infty$, 168
 $\mathbb{L}_{\delta,K}^{t,\infty}$, 166
 $\mathbb{L}_{\delta,K}^t$, 165
 $\mathbb{L}_{\delta,K}$, 168
 Functions
 I_Γ , 33
 $M(t,x,q)$, 166
 $P(u'')$, 109
 $P_{m,\delta/2,\Lambda}$, 109
 $\mathcal{D}(u'')$, 16
 $\mathcal{P}(z'')$, 92
 $\mathcal{P}_{m,\delta/2}(z'')$, 92
 $\omega_{F,u,\Omega}(\rho)$, 209
 $\omega_{F,u,\Pi}(\rho)$, 252
 $\omega_n(x)$, 413
 $\omega_u(\Omega, \rho)$, 209
 $\omega_u(\Pi, \rho)$, 252
 $\rho_T(t, x)$, 146
 ρ_Ω , 78
 $c^m(u'')$, 16
 $c_{ij}^m(u'')$, 16
 g_γ^\sharp , 421
 $h_{\gamma,\rho}^\#$, 227
 $h_{G,\gamma,\rho}^\#$, 422
 $h_{Q,\gamma,\rho}^\#$, 265
 $h_{Q,\gamma}^\#$, 421
 $h_{\gamma,\rho}^\#$, 423
 $h_\gamma^\#$, 421
 $h_{\Omega,\gamma,\rho}^\#$, 333
 $m(t, x, q)$, 166
 $m(x, q)$, 168
 $p(t, x, q)$, 182
 $p(x, q)$, 185
 u_γ^\sharp , 420
 Norms
 $[f]_{C^\alpha(C_{\tau,r}^+)}^{}$, 274
 $[u]_{C^\alpha}$, 216
 $|a|$, xii, 31
 $|u''|$, 207, 368
 $||\cdot||_{W_\infty^{1,2}(\mathbb{R}^{d+1})}$, 144
 $||\phi||_{C^\kappa(\Pi)}$, 343
 Miscellanea
 (x^1, x') , 180
 $M_{\alpha,R}(a)$, 286
 $[u']$, 62, 367
 α_0 , 189, 216, 291
 $\bar{\alpha}$, 223
 $\bar{\delta}$, 109
 \bar{F}_B , 368
 $\int_\Gamma f(t, x) dx dt$, 367
 γ_0 , 177
 κ_d , 19
 $\langle \cdot, \cdot \rangle$, xii
 $\mu^{(0)}$, 7
 μ_R , 372
 $\text{osc}(g, \Gamma)$, 189
 $\text{osc}_\Gamma g$, 189
 ∂' , 33
 $\rho(z_1, z_2)$, xii
 $\rho_0(\Omega)$, 313
 $\rho_{\text{ext}}(\Omega)$, 92
 $\rho_{\text{int}}(\Omega)$, 78
 u , ix
 u' , ix

- u'' , ix
- u'_k , ix
- $|\Gamma|$, xii, 166, 252
- $|\omega|$, 413
- $a \vee b$, xiii
- $a \wedge b$, xiii
- a^* , xii
- f_A , 413
- $f_{|n}$, 414
- $f_{|\tau}$, 414
- $p(l)$, 116
- $r(\alpha)$, 22
- u^\pm , xiii

- \mathcal{R} -viscosity solution, 341

- Aleksandrov-Busemann-Feller theorem, 7
- almost in VMO, ix, 212
- anti-positive operator, 67

- boundedly inhomogeneous functions, 111

- characterization of $C^{1+\alpha}$ -functions, 360
- compact set, xiii

- domain, xii
- dyadic cubes, 414
- dyadic sharp function, 421

- exterior ball condition, 92

- Fefferman-Stein theorem, 420, 422
- filtration of partitions, 413

- Isaacs equation, 213

- maximal function, 416
- maximal inequality, 416
- maximum principle, 47
- measurable function/set, xiii
- multi-index, 276

- parabolic boundary, 33
- parabolic distance, xii
- parabolic dyadic cubes, 414

- Rademacher's theorem, 55

- scalar product, xii
- set dominating (t_0, x_0) , 33
- stopping time, 414
- support, xiii

SELECTED PUBLISHED TITLES IN THIS SERIES

- 233 **N. V. Krylov**, Sobolev and Viscosity Solutions for Fully Nonlinear Elliptic and Parabolic Equations, 2018
- 230 **Stephen D. Smith**, Applying the Classification of Finite Simple Groups, 2018
- 229 **Alexander Molev**, Sugawara Operators for Classical Lie Algebras, 2018
- 228 **Zhenbo Qin**, Hilbert Schemes of Points and Infinite Dimensional Lie Algebras, 2018
- 227 **Roberto Frigerio**, Bounded Cohomology of Discrete Groups, 2017
- 226 **Marcelo Aguiar and Swapneel Mahajan**, Topics in Hyperplane Arrangements, 2017
- 225 **Mario Bonk and Daniel Meyer**, Expanding Thurston Maps, 2017
- 224 **Ruy Exel**, Partial Dynamical Systems, Fell Bundles and Applications, 2017
- 223 **Guillaume Aubrun and Stanisław J. Szarek**, Alice and Bob Meet Banach, 2017
- 222 **Alexandru Buium**, Foundations of Arithmetic Differential Geometry, 2017
- 221 **Dennis Gaitsgory and Nick Rozenblyum**, A Study in Derived Algebraic Geometry, 2017
- 220 **A. Shen, V. A. Uspensky, and N. Vereshchagin**, Kolmogorov Complexity and Algorithmic Randomness, 2017
- 219 **Richard Evan Schwartz**, The Projective Heat Map, 2017
- 218 **Tushar Das, David Simmons, and Mariusz Urbański**, Geometry and Dynamics in Gromov Hyperbolic Metric Spaces, 2017
- 217 **Benoit Fresse**, Homotopy of Operads and Grothendieck–Teichmüller Groups, 2017
- 216 **Frederick W. Gehring, Gaven J. Martin, and Bruce P. Palka**, An Introduction to the Theory of Higher-Dimensional Quasiconformal Mappings, 2017
- 215 **Robert Bieri and Ralph Strelbel**, On Groups of PL-homeomorphisms of the Real Line, 2016
- 214 **Jared Speck**, Shock Formation in Small-Data Solutions to 3D Quasilinear Wave Equations, 2016
- 213 **Harold G. Diamond and Wen-Bin Zhang (Cheung Man Ping)**, Beurling Generalized Numbers, 2016
- 212 **Pandelis Dodos and Vassilis Kanellopoulos**, Ramsey Theory for Product Spaces, 2016
- 211 **Charlotte Hardouin, Jacques Sauloy, and Michael F. Singer**, Galois Theories of Linear Difference Equations: An Introduction, 2016
- 210 **Jason P. Bell, Dragos Ghioca, and Thomas J. Tucker**, The Dynamical Mordell–Lang Conjecture, 2016
- 209 **Steve Y. Oudot**, Persistence Theory: From Quiver Representations to Data Analysis, 2015
- 208 **Peter S. Ozsváth, András I. Stipsicz, and Zoltán Szabó**, Grid Homology for Knots and Links, 2015
- 207 **Vladimir I. Bogachev, Nicolai V. Krylov, Michael Röckner, and Stanislav V. Shaposhnikov**, Fokker–Planck–Kolmogorov Equations, 2015
- 206 **Bennett Chow, Sun-Chin Chu, David Glickenstein, Christine Guenther, James Isenberg, Tom Ivey, Dan Knopf, Peng Lu, Feng Luo, and Lei Ni**, The Ricci Flow: Techniques and Applications: Part IV: Long-Time Solutions and Related Topics, 2015
- 205 **Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik**, Tensor Categories, 2015
- 204 **Victor M. Buchstaber and Taras E. Panov**, Toric Topology, 2015
- 203 **Donald Yau and Mark W. Johnson**, A Foundation for PROPs, Algebras, and Modules, 2015

For a complete list of titles in this series, visit the
AMS Bookstore at www.ams.org/bookstore/survseries/.

This book concentrates on first boundary-value problems for fully nonlinear second-order uniformly elliptic and parabolic equations with discontinuous coefficients. We look for solutions in Sobolev classes, local or global, or for viscosity solutions. Most of the auxiliary results, such as Aleksandrov's elliptic and parabolic estimates, the Krylov–Safonov and the Evans–Krylov theorems, are taken from old sources, and the main results were obtained in the last few years.

Presentation of these results is based on a generalization of the Fefferman–Stein theorem, on Fang–Hua Lin's like estimates, and on the so-called “ersatz” existence theorems, saying that one can slightly modify “any” equation and get a “cut-off” equation that has solutions with bounded derivatives. These theorems allow us to prove the solvability in Sobolev classes for equations that are quite far from the ones which are convex or concave with respect to the Hessians of the unknown functions. In studying viscosity solutions, these theorems also allow us to deal with classical approximating solutions, thus avoiding sometimes heavy constructions from the usual theory of viscosity solutions.

ISBN 978-1-4704-4740-3



9 781470 447403

SURV/233



For additional information
and updates on this book, visit
www.ams.org/bookpages/surv-233



AMS
www.ams.org