

## LECTURE 1

# Introduction

When a group  $G$  acts on a manifold  $M$ , one would like to understand the relation between:

- algebraic properties of the group  $G$ ,
- the topology of  $M$ ,
- the  $G$ -invariant geometric structures on  $M$ , and
- dynamical properties of the action (such as dense orbits, invariant measures, etc.).

If we assume that  $G$  is a connected Lie group, then the structure theory (A1.3) tells us there are two main cases to consider:

- *solvable*  
Solvable groups are usually studied by starting with  $\mathbb{R}^n$  and proceeding by induction.
- *semisimple*  
There is a classification that provides a list of the semisimple groups ( $\mathrm{SL}(n, \mathbb{R})$ ,  $\mathrm{SO}(p, q)$ , etc.), so a case-by-case analysis is possible. Alternatively, the groups can sometimes be treated via other categorizations, e.g., by real rank.

The emphasis in these lectures is on the semisimple case.

(1.1) **Assumption.** *In this lecture,  $G$  always denotes a connected, noncompact, semisimple Lie group.*

### 1A. Discrete versions of $G$

A connected Lie group may have discrete subgroups that approximate it. There are two notions of this that play very important roles in these lectures:

- *lattice*  
This is a discrete subgroup  $\Gamma$  of  $G$  such that  $G/\Gamma$  is compact (or, more generally, such that  $G/\Gamma$  has finite volume).
- *arithmetic subgroup*  
Suppose  $G$  is a (closed) subgroup of  $\mathrm{GL}(n, \mathbb{R})$ , and let  $\Gamma = G \cap \mathrm{GL}(n, \mathbb{Z})$  be the set of “integer points” of  $G$ . If  $\Gamma$  is Zariski dense in  $G$  (or, equivalently, if  $G \cap \mathrm{GL}(n, \mathbb{Q})$  is dense in  $G$ ), then we say  $\Gamma$  is an *arithmetic* subgroup of  $G$ .

Discrete versions inherit important algebraic properties of  $G$ :

(1.2) **Example.** Suppose  $\Gamma$  is a discrete version of  $G$  (that is,  $\Gamma$  is either a lattice or an arithmetic subgroup).

- 1) Because Assumption 1.1 tells us that  $G$  is semisimple, we know  $G$  has no solvable, normal subgroups (or, more precisely, none that are connected and nontrivial). One can show that  $\Gamma$  also has no solvable, normal subgroups (or, more precisely, none that are infinite, if  $G$  has finite center). This follows from the Borel Density Theorem (A6.1), which tells us that  $\Gamma$  is “Zariski dense” in (a large subgroup of)  $G$ .
- 2) If  $G$  is simple, then, by definition,  $G$  has no normal subgroups (or, more precisely, none that are connected, nontrivial, and proper). If we furthermore assume  $\mathbb{R}\text{-rank}(G) \geq 2$  (i.e.,  $G$  has a subgroup of dimension  $\geq 2$  that is diagonalizable over  $\mathbb{R}$ ) and the center of  $G$  is finite, then theorems of G. A. Margulis link  $\Gamma$  and  $G$  more tightly:
  - (a)  $\Gamma$  has no normal subgroups (or, more precisely, none that are infinite and of infinite index), and
  - (b) roughly speaking, every lattice in  $G$  is arithmetic.
 (See Theorems A7.3 and A7.9 for precise statements.)

(1.3) *Remark.* Both of the conclusions of Example 1.2(2) can fail if we eliminate the assumption that  $\mathbb{R}\text{-rank}(G) \geq 2$ . For example, in  $\mathrm{SL}(2, \mathbb{R})$ , the free group  $F_2$  is a lattice, and some other lattices have homomorphisms onto free groups.

(1.4) **Example** (Linear actions). Let  $M = V$ , where  $V$  is a finite-dimensional vector space (over  $\mathbb{R}$  or  $\mathbb{C}$ ). A linear action of  $G$  on  $M$  (that is, a homomorphism  $G \rightarrow \mathrm{GL}(V)$ ) is known as a (finite-dimensional) *representation* of  $G$ . These representations are classified by the well-known theory of *highest weights*, so we can think of the linear actions of  $G$  as being known.

If  $\mathbb{R}\text{-rank}(G) \geq 2$  (and  $G$  is simple), then the linear actions of  $\Gamma$  are closely related to the linear actions of  $G$ . Namely, the spirit, but not exactly the statement, of the Margulis Superrigidity Theorem (A7.4) is that if  $\varphi: \Gamma \rightarrow \mathrm{GL}(V)$ , then either

- $\varphi$  extends to a representation of  $G$ , or
- the image of  $\varphi$  is contained in a compact subgroup of  $\mathrm{GL}(V)$ .

In other words, any representation of  $\Gamma$  extends to a representation of  $G$ , modulo compact groups. This is another illustration of the close connection between  $G$  and its discrete versions.

(1.5) *Remark.* An alternative formulation (A7.7) of the Margulis Superrigidity Theorem says that (in spirit) each representation of  $\Gamma$  either extends to  $G$  or extends to  $G$  after composing with some field automorphism of  $\mathbb{C}$ .

## 1B. Nonlinear actions

These lectures explore some of what is known about the nonlinear actions of  $G$  and its discrete versions. (As was mentioned in Example 1.4, the theory of highest weights provides a largely satisfactory theory of the linear actions.) The connections between  $G$  and  $\Gamma$  are of particular interest.

We begin the discussion with two basic examples of  $G$ -actions.

(1.6) **Example.**  $G$  acts on  $G/\Gamma$ , where  $\Gamma$  is any lattice. More generally, if  $G \hookrightarrow H$ , and  $\Lambda$  is a lattice in  $H$ , then  $G$  acts by translations on  $H/\Lambda$ .

(1.7) **Example.** The linear action of  $\mathrm{SL}(n, \mathbb{R})$  on  $\mathbb{R}^n$  factors through to a (nonlinear) action of  $\mathrm{SL}(n, \mathbb{R})$  on the projective space  $\mathbb{R}P^{n-1}$ . More generally,  $\mathrm{SL}(n, \mathbb{R})$  acts on Grassmannians, and other flag varieties.

Generalizing further, any semisimple Lie group  $G$  has transitive actions on some projective varieties. They are of the form  $G/Q$ , where  $Q$  is a “parabolic” subgroup of  $G$ . (We remark that there are only finitely many of these actions, up to isomorphism, because there are only finitely many parabolic subgroups of  $G$ , up to conjugacy.)

(1.8) *Remark.* In the case of  $\mathrm{SL}(n, \mathbb{R})$ , any parabolic subgroup  $Q$  is block upper-triangular:

$$Q = \begin{bmatrix} \boxed{*} & & * \\ & \boxed{*} & \\ 0 & & \boxed{*} \end{bmatrix}$$

A natural question that guides research in the area is:

*To what extent is every action on a compact manifold (or perhaps a more general space) built out of the above two basic examples?* (1.9)

Unfortunately, there are methods to construct actions that seem to be much less amenable to classification. The two known methods are *Induction* and *Blowing up*; we will briefly describe each of them.

**1B(a). Induction.** Suppose some subgroup  $H$  of  $G$  acts on a space  $Y$ . Then

- 1)  $H$  acts on  $G \times Y$  by

$$h \cdot (g, y) = (gh^{-1}, hy),$$

and

- 2)  $G$  acts on the quotient  $X = (G \times Y)/H$  by

$$a \cdot [(g, y)] = [(ag, y)].$$

The action of  $G$  on  $X$  is said to be *induced* from the action of  $H$  on  $Y$ .

(1.10) *Remark.*

- 1) Ignoring the second coordinate yields a  $G$ -equivariant map  $X \rightarrow G/H$ , so we see that  $X$  is a fiber bundle over  $G/H$  with fiber  $Y$ . Note that the fiber over  $[e] = eH$  is  $H$ -invariant, and the action of  $H$  on this fiber is isomorphic to the action of  $H$  on  $Y$ .
- 2) Conversely, if  $G$  acts on  $X$ , and there is a  $G$ -equivariant map from  $X$  to  $G/H$ , then  $X$  is  $G$ -equivariantly isomorphic to an action induced from  $H$ . Namely, if we let  $Y$  be the fiber of  $X$  over  $[e]$ , then  $Y$  is  $H$ -invariant, and the map  $(g, y) \mapsto g \cdot y$  factors through to a  $G$ -equivariant bijection  $(G \times Y)/H \rightarrow X$ .

(1.11) **Example.** Let  $H$  be the stabilizer in  $\mathrm{SL}(n, \mathbb{R})$  of a point in  $\mathbb{R}P^{n-1}$ , so

$$H = \left\{ h = \begin{bmatrix} \lambda & * & \cdots & * \\ 0 & \boxed{A} & & \\ \vdots & & & \\ 0 & & & \end{bmatrix} \mid \det A = \lambda^{-1} \right\}$$

Then  $h \mapsto \log |\lambda|$  is a homomorphism from  $H$  onto  $\mathbb{R}$ , so every  $\mathbb{R}$ -action yields an action of  $H$ ; hence, by induction, an action of  $\mathrm{SL}(n, \mathbb{R})$ .

The upshot is that every vector field on any compact manifold  $M$  yields an  $\mathrm{SL}(n, \mathbb{R})$ -action on a compact manifold  $M'$ . Thus, all the complications that arise for  $\mathbb{R}$ -actions also arise for  $\mathrm{SL}(n, \mathbb{R})$ -actions.

It is hopeless to classify all  $\mathbb{R}$ -actions on compact manifolds, so Example 1.11 puts a damper on the hope for a complete classification of actions of simple Lie groups. The following example discourages the belief in a classification theorem even more.

(1.12) **Example.** Let  $\Gamma$  be a lattice in  $\mathrm{PSL}(2, \mathbb{R})$ , such that  $\Gamma$  has a homomorphism onto a nonabelian free group  $F$ . By the same argument as in Example 1.11, we see that every action of  $F$  on a compact manifold yields an action of  $\mathrm{PSL}(2, \mathbb{R})$  on a manifold (and the manifold is compact if  $\mathrm{PSL}(2, \mathbb{R})/\Gamma$  is compact).

In  $\mathrm{SL}(n, \mathbb{R})$ , let

$$Q = \begin{bmatrix} \boxed{\begin{matrix} * & * \\ * & * \end{matrix}} & * & \cdots & * \\ 0 & 0 & \boxed{*} & \\ \vdots & \vdots & & \\ 0 & 0 & & \end{bmatrix},$$

where the box in the top left corner is  $2 \times 2$ , so  $Q$  has a homomorphism onto  $\mathrm{PSL}(2, \mathbb{R})$ . Thus,

$$F\text{-action} \rightarrow \mathrm{PSL}(2, \mathbb{R})\text{-action} \rightarrow Q\text{-action} \rightarrow \mathrm{SL}(n, \mathbb{R})\text{-action}.$$

So the free-group problem arises for every  $\mathrm{SL}(n, \mathbb{R})$ , not just  $\mathrm{SL}(2, \mathbb{R})$ .

**1B(b). Blowing up** (Katok-Lewis, Benveniste). Let  $\Gamma$  be a finite-index subgroup of  $\mathrm{SL}(n, \mathbb{Z})$ , so  $\Gamma$  acts on  $\mathbb{T}^n$  by automorphisms, and assume the action has two distinct fixed points  $x$  and  $y$ .

- 1) First, *blow up* at these two fixed points. That is, letting  $T_p(\mathbb{T}^n)$  be the tangent space to  $\mathbb{T}^n$  at  $p$ ,
  - replace  $x$  with  $\{\text{rays in } T_x(\mathbb{T}^n)\}$ , and
  - replace  $y$  with  $\{\text{rays in } T_y(\mathbb{T}^n)\}$ .

Thus,  $x$  and  $y$  have each been replaced with a sphere.

- 2) Now, glue the two spheres together.

The upshot is that the union of two fixed points can be replaced by a projective space with the usual action of  $\mathrm{SL}(n, \mathbb{Z})$  on  $\mathbb{R}P^{n-1}$ .

(1.13) *Remark.*

- 1) Blowing up results in a different manifold; the fundamental group is different. (This will be discussed in Example 4.18 below.)
- 2) The construction can embed the action on a projective space (which is not volume preserving) into a volume-preserving action.
- 3) The change in the manifold is on a submanifold of positive codimension.

The above construction is due to A. Katok and J. Lewis. It was generalized by J. Benveniste along the following lines.

**(1.14) Example.**

- 1) Embed  $G$  in a larger group  $H$ , let  $M = H/\Gamma$ , for some lattice  $\Lambda$  in  $H$ .
- 2) Assume  $G$  is contained in a subgroup  $L$  of  $H$ , such that  $L$  has a closed orbit on  $H/\Gamma$ .

For some pairs of closed  $L$ -orbits, we can blow up transversally (that is, take the set of rays in a subspace transverse to the  $L$ -orbit), and then glue to make a new action. Varying the gluing results in actions that have nontrivial perturbations.

**1C. Open questions**

The examples in §1B(a) and §1B(b) suggest there is a limit to what can be done, so we present a few problems that are likely to be approachable. Some will be discussed in later lectures.

- 1) Are there actions of  $\Gamma$  on low-dimensional manifolds?  
 Low-dimensional can either mean low in an absolute sense, as in dimension  $\leq 3$ , or it can mean low relative to  $\Gamma$ , which is often taken to be less than  
 (lowest dimension of a representation of  $G$ )  $- 1$   
 For example, if  $\Gamma = \mathrm{SL}(n, \mathbb{Z})$ , then “low-dimensional” means either  $\leq 3$  or  $< n - 1$ .
- 2) When are actions locally rigid?  
 Not all actions are locally rigid. For example, if the action is induced from a vector field, then it may be possible to perturb the vector field. (Note that perturbations may be non-linear actions, even if the original action is linear.)
- 3) Suppose  $G$  preserves a geometric structure on  $M$ , defined by a structure group  $H$ . Then what is the relation between  $G$  and  $H$ ?  
 Sometimes, assuming that  $G$  preserves a suitable geometric structure eliminates the examples constructed above.
- 4) Does every action have either:
  - an invariant geometric structure of rigid type (at least, on an open set), or
  - an equivariant quotient  $G$ -manifold with such a structure?
- 5) If  $G$  acts on  $M$ , what can be said about the fundamental group  $\pi_1(M)$ ?  
 What about other aspects of the topology of  $M$ ?
- 6) What are the consequences of assuming there is a  $G$ -invariant volume form on  $M$ ?

Approaches to these questions must bear in mind the constructions of §1B(a) and §1B(b) that provide counterexamples to many naive conjectures.

**Comments**

Proofs of the fundamental theorems of G. A. Margulis mentioned in Examples 1.2 and 1.4 can be found in [9, 10, 11].

The blowing-up construction of §1B(b) is due to A. Katok and J. Lewis [7]. Example 1.14 appeared in the unpublished Ph.D. thesis of E. J. Benveniste [1].

See [4] for details of a stronger result. A few additional examples created by a somewhat different method of gluing appear in [6, §4].

Some literature on the open questions in Section 1C:

- 1) Actions on manifolds of low dimension  $\leq 2$  are discussed in Lecture 2 (and the comments at the end).
- 2) See [3] for a recent survey of the many results on local rigidity of group actions.
- 3) Actions with an invariant geometric structure are discussed in Lectures 3, 5 and 6.
- 4) The examples of Katok-Lewis and Benveniste do not have a rigid geometric structure that is invariant [2]. On the other hand, the notion of “almost-rigid structure” is introduced in [2], and all known volume-preserving, smooth actions of higher-rank simple Lie groups have an invariant structure of that type.
- 5) Results on the fundamental group of  $M$  are discussed in Lectures 4, 5 and 7.
- 6) Many of the results discussed in these lectures apply only to actions that are volume preserving (or, at least, have an invariant probability measure). Only Lecture 9 is specifically devoted to actions that are *not* volume preserving.

See the survey of D. Fisher [5] (and other papers in the same volume) and the ICM talk of F. Labourie [8] for a different view of several of the topics that will be discussed in these lectures.

## References

- [1] E. J. Benveniste: Rigidity and deformations of lattice actions preserving geometric structures. Ph. D. thesis, Mathematics, University of Chicago, 1996.
- [2] E. J. Benveniste and D. Fisher: Nonexistence of invariant rigid structures and invariant almost rigid structures, *Comm. Anal. Geom.* 13 (2005), no. 1, 89–111. MR 2154667 (2006f:53056)
- [3] D. Fisher: Local Rigidity: Past, Present, Future, in: B. Hasselblatt, ed., *Dynamics, Ergodic Theory and Geometry*, Cambridge U. Press, New York, 2007, pp. 45–98. ISBN 9780521875417, MR 2369442
- [4] D. Fisher: Deformations of group actions, *Trans. Amer. Math. Soc.* 360 (2008), no. 1, 491–505. MR 2342012
- [5] D. Fisher: Groups acting on manifolds: around the Zimmer program, in preparation for B. Farb and D. Fisher, eds., *Geometry, Rigidity and Group Actions*, U. of Chicago Press, in preparation.
- [6] D. Fisher and K. Whyte: Continuous quotients for lattice actions on compact spaces, *Geom. Dedicata* 87 (2001), no. 1-3, 181–189. MR 1866848 (2002j:57070)
- [7] A. Katok and J. Lewis: Global rigidity results for lattice actions on tori and new examples of volume-preserving actions, *Israel J. Math.* 93 (1996), 253–280. MR 1380646 (96k:22021)

- [8] F. Labourie: Large groups actions on manifolds, Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998). *Doc. Math.* 1998, Extra Vol. II, 371–380. MR 1648087 (99k:53069)
- [9] G. A. Margulis: *Discrete Subgroups of Semisimple Lie Groups*. Springer, Berlin Heidelberg New York, 1991. ISBN 3-540-12179-X, MR 1090825 (92h:22021)
- [10] D. W. Morris: *Introduction to Arithmetic Groups* (preprint).  
<http://arxiv.org/abs/math/0106063>
- [11] R. J. Zimmer: *Ergodic Theory and Semisimple Groups*. Birkhäuser, Basel, 1984. ISBN 3-7643-3184-4, MR 0776417 (86j:22014)