

## Generalized Argument Principle and Rouché's Theorem

In this chapter we review the results of Gohberg and Sigal in [114] concerning the generalization to operator-valued functions of two classical results in complex analysis, the *argument principle* and *Rouché's theorem*.

To state the argument principle, we first observe that if  $f$  is holomorphic and has a zero of order  $n$  at  $z_0$ , we can write  $f(z) = (z - z_0)^n g(z)$ , where  $g$  is holomorphic and nowhere vanishing in a neighborhood of  $z_0$ , and therefore

$$\frac{f'(z)}{f(z)} = \frac{n}{z - z_0} + \frac{g'(z)}{g(z)}.$$

Then the function  $f'/f$  has a simple pole with residue  $n$  at  $z_0$ . A similar fact also holds if  $f$  has a pole of order  $n$  at  $z_0$ , that is, if  $f(z) = (z - z_0)^{-n} h(z)$ , where  $h$  is holomorphic and nowhere vanishing in a neighborhood of  $z_0$ . Then

$$\frac{f'(z)}{f(z)} = -\frac{n}{z - z_0} + \frac{h'(z)}{h(z)}.$$

Therefore, if  $f$  is holomorphic, the function  $f'/f$  will have simple poles at the zeros and poles of  $f$ , and the residue is simply the order of the zero of  $f$  or the negative of the order of the pole of  $f$ .

The argument principle results from an application of the residue formula. It asserts the following.

**THEOREM 1.1** (Argument principle). *Let  $V \subset \mathbb{C}$  be a bounded domain with smooth boundary  $\partial V$  positively oriented and let  $f(z)$  be a meromorphic function in a neighborhood of  $\bar{V}$ . Let  $P$  and  $N$  be the number of poles and zeros of  $f$  in  $V$ , counted with their multiplicities. If  $f$  has no poles and never vanishes on  $\partial V$ , then*

$$(1.1) \quad \frac{1}{2\pi\sqrt{-1}} \int_{\partial V} \frac{f'(z)}{f(z)} dz = N - P.$$

Rouché's theorem is a consequence of the argument principle [237]. It is in some sense a continuity statement. It says that a holomorphic function can be perturbed slightly without changing the number of its zeros. It reads as follows.

**THEOREM 1.2** (Rouché's theorem). *With  $V$  as above, suppose that  $f(z)$  and  $g(z)$  are holomorphic in a neighborhood of  $\bar{V}$ . If  $|f(z)| > |g(z)|$  for all  $z \in \partial V$ , then  $f$  and  $f + g$  have the same number of zeros in  $V$ .*

In order to explain the main results of Gohberg and Sigal in [114], we begin with the finite-dimensional case which was first considered by Keldyš in [152]; see also [183]. We proceed to generalize formula (1.1) in this case as follows. If a

matrix-valued function  $A(z)$  is holomorphic in a neighborhood of  $\overline{V}$  and is invertible in  $\overline{V}$  except possibly at a point  $z_0 \in V$ , then by Gaussian eliminations we can write

$$(1.2) \quad A(z) = E(z)D(z)F(z) \quad \text{in } V,$$

where  $E(z), F(z)$  are holomorphic and invertible in  $V$  and  $D(z)$  is given by

$$D(z) = \begin{pmatrix} (z - z_0)^{k_1} & & 0 \\ & \ddots & \\ 0 & & (z - z_0)^{k_n} \end{pmatrix}.$$

Moreover, the powers  $k_1, k_2, \dots, k_n$  are uniquely determined up to a permutation.

Let  $\text{tr}$  denote the trace. By virtue of the factorization (1.2), it is easy to produce the following identity:

$$\begin{aligned} & \frac{1}{2\pi\sqrt{-1}} \text{tr} \int_{\partial V} A(z)^{-1} \frac{d}{dz} A(z) dz \\ &= \frac{1}{2\pi\sqrt{-1}} \text{tr} \int_{\partial V} \left( E(z)^{-1} \frac{d}{dz} E(z) + D(z)^{-1} \frac{d}{dz} D(z) + F(z)^{-1} \frac{d}{dz} F(z) \right) dz \\ &= \frac{1}{2\pi\sqrt{-1}} \text{tr} \int_{\partial V} D(z)^{-1} \frac{d}{dz} D(z) dz \\ &= \sum_{j=1}^n k_j, \end{aligned}$$

which generalizes (1.1).

In the next sections, we will extend the above identity as well as the factorization (1.2) to infinite-dimensional spaces under some natural conditions.

### 1.1. Definitions and Preliminaries

In this section we introduce the notation we will use in the text, gather a few definitions, and present some basic results, which are useful for the statement of the generalized Rouché theorem.

**1.1.1. Compact Operators.** If  $\mathcal{B}$  and  $\mathcal{B}'$  are two Banach spaces, we denote by  $\mathcal{L}(\mathcal{B}, \mathcal{B}')$  the space of bounded linear operators from  $\mathcal{B}$  into  $\mathcal{B}'$ . An operator  $K \in \mathcal{L}(\mathcal{B}, \mathcal{B}')$  is said to be compact provided  $K$  takes any bounded subset of  $\mathcal{B}$  to a relatively compact subset of  $\mathcal{B}'$ , that is, a set with compact closure.

The operator  $K$  is said to be of finite rank if  $\text{Im}(K)$ , the range of  $K$ , is finite-dimensional. Clearly every operator of finite rank is compact.

The next result is called the Fredholm alternative. See, for example, [164].

**PROPOSITION 1.3 (Fredholm alternative).** *Let  $K$  be a compact operator on the Banach space  $\mathcal{B}$ . For  $\lambda \in \mathbb{C}, \lambda \neq 0$ ,  $(\lambda I - K)$  is surjective if and only if it is injective.*

**1.1.2. Fredholm Operators.** An operator  $A \in \mathcal{L}(\mathcal{B}, \mathcal{B}')$  is said to be *Fredholm* provided the subspace  $\text{Ker } A$  is finite-dimensional and the subspace  $\text{Im } A$  is closed in  $\mathcal{B}'$  and of finite codimension. Let  $\text{Fred}(\mathcal{B}, \mathcal{B}')$  denote the collection of all Fredholm operators from  $\mathcal{B}$  into  $\mathcal{B}'$ . We can show that  $\text{Fred}(\mathcal{B}, \mathcal{B}')$  is open in  $\mathcal{L}(\mathcal{B}, \mathcal{B}')$ .

Next, we define the index of  $A \in \text{Fred}(\mathcal{B}, \mathcal{B}')$  to be

$$\text{ind } A = \dim \text{Ker } A - \text{codim } \text{Im } A.$$

In finite dimensions, the index depends only on the spaces and not on the operator.

The following proposition shows that the index is stable under compact perturbations [164].

PROPOSITION 1.4. *If  $A : \mathcal{B} \rightarrow \mathcal{B}'$  is Fredholm and  $K : \mathcal{B} \rightarrow \mathcal{B}'$  is compact, then their sum  $A + K$  is Fredholm, and*

$$\text{ind}(A + K) = \text{ind } A.$$

Proposition 1.4 is a consequence of the following fundamental result about the index of Fredholm operators.

PROPOSITION 1.5. *The mapping  $A \mapsto \text{ind } A$  is continuous in  $\text{Fred}(\mathcal{B}, \mathcal{B}')$ ; i.e.,  $\text{ind}$  is constant on each connected component of  $\text{Fred}(\mathcal{B}, \mathcal{B}')$ .*

**1.1.3. Characteristic Value and Multiplicity.** We now introduce the notions of characteristic values and root functions of analytic operator-valued functions, with which the readers might not be familiar. We refer, for instance, to the book by Markus [175] for the details.

Let  $\mathfrak{U}(z_0)$  be the set of all operator-valued functions with values in  $\mathcal{L}(\mathcal{B}, \mathcal{B}')$  which are holomorphic in some neighborhood of  $z_0$ , except possibly at  $z_0$ .

The point  $z_0$  is called a *characteristic value* of  $A(z) \in \mathfrak{U}(z_0)$  if there exists a vector-valued function  $\phi(z)$  with values in  $\mathcal{B}$  such that

- (i)  $\phi(z)$  is holomorphic at  $z_0$  and  $\phi(z_0) \neq 0$ ,
- (ii)  $A(z)\phi(z)$  is holomorphic at  $z_0$  and vanishes at this point.

Here,  $\phi(z)$  is called a *root function* of  $A(z)$  associated with the characteristic value  $z_0$ . The vector  $\phi_0 = \phi(z_0)$  is called an *eigenvector*. The closure of the linear set of eigenvectors corresponding to  $z_0$  is denoted by  $\text{Ker}A(z_0)$ .

Suppose that  $z_0$  is a characteristic value of the function  $A(z)$  and  $\phi(z)$  is an associated root function. Then there exists a number  $m(\phi) \geq 1$  and a vector-valued function  $\psi(z)$  with values in  $\mathcal{B}'$ , holomorphic at  $z_0$ , such that

$$A(z)\phi(z) = (z - z_0)^{m(\phi)}\psi(z), \quad \psi(z_0) \neq 0.$$

The number  $m(\phi)$  is called the *multiplicity* of the root function  $\phi(z)$ .

For  $\phi_0 \in \text{Ker}A(z_0)$ , we define the rank of  $\phi_0$ , denoted by  $\text{rank}(\phi_0)$ , to be the maximum of the multiplicities of all root functions  $\phi(z)$  with  $\phi(z_0) = \phi_0$ .

Suppose that  $n = \dim \text{Ker}A(z_0) < +\infty$  and that the ranks of all vectors in  $\text{Ker}A(z_0)$  are finite. A system of eigenvectors  $\phi_0^j$ ,  $j = 1, \dots, n$ , is called a *canonical system of eigenvectors* of  $A(z)$  associated to  $z_0$  if their ranks possess the following property: for  $j = 1, \dots, n$ ,  $\text{rank}(\phi_0^j)$  is the maximum of the ranks of all eigenvectors in the direct complement in  $\text{Ker}A(z_0)$  of the linear span of the vectors  $\phi_0^1, \dots, \phi_0^{j-1}$ . We call

$$N(A(z_0)) := \sum_{j=1}^n \text{rank}(\phi_0^j)$$

the *null multiplicity* of the characteristic value  $z_0$  of  $A(z)$ . If  $z_0$  is not a characteristic value of  $A(z)$ , we put  $N(A(z_0)) = 0$ .

Suppose that  $A^{-1}(z)$  exists and is holomorphic in some neighborhood of  $z_0$ , except possibly at  $z_0$ . Then the number

$$M(A(z_0)) = N(A(z_0)) - N(A^{-1}(z_0))$$

is called the *multiplicity* of  $z_0$ . If  $z_0$  is a characteristic value and not a pole of  $A(z)$ , then  $M(A(z_0)) = N(A(z_0))$  while  $M(A(z_0)) = -N(A^{-1}(z_0))$  if  $z_0$  is a pole and not a characteristic value of  $A(z)$ .

**1.1.4. Normal Points.** Suppose that  $z_0$  is a pole of the operator-valued function  $A(z)$  and the Laurent series expansion of  $A(z)$  at  $z_0$  is given by

$$(1.3) \quad A(z) = \sum_{j \geq -s} (z - z_0)^j A_j.$$

If in (1.3) the operators  $A_{-j}$ ,  $j = 1, \dots, s$ , have finite-dimensional ranges, then  $A(z)$  is called *finitely meromorphic* at  $z_0$ .

The operator-valued function  $A(z)$  is said to be of *Fredholm type* (of index zero) at the point  $z_0$  if the operator  $A_0$  in (1.3) is Fredholm (of index zero).

If  $A(z)$  is holomorphic and invertible at  $z_0$ , then  $z_0$  is called a *regular point* of  $A(z)$ . The point  $z_0$  is called a *normal point* of  $A(z)$  if  $A(z)$  is finitely meromorphic, of Fredholm type at  $z_0$ , and regular in a neighborhood of  $z_0$  except at  $z_0$  itself.

**1.1.5. Trace.** Let  $A$  be a finite-dimensional operator acting from  $\mathcal{B}$  into itself. There exists a finite-dimensional invariant subspace  $\mathcal{C}$  of  $A$  such that  $A$  annihilates some direct complement of  $\mathcal{C}$  in  $\mathcal{B}$ . We define the trace of  $A$  to be that of  $A|_{\mathcal{C}}$ , which is given in the usual way. It is desirable to recall some results about the trace operator.

PROPOSITION 1.6. *The following results hold:*

- (i)  $\text{tr } A$  is independent of the choice of  $\mathcal{C}$ , so that it is well-defined.
- (ii)  $\text{tr}$  is linear.
- (iii) If  $B$  is a finite-dimensional operator from  $\mathcal{B}$  to itself, then

$$\text{tr } AB = \text{tr } BA.$$

- (iv) If  $M$  is a finite-dimensional operator from  $\mathcal{B} \times \mathcal{B}'$  to itself, given by

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix},$$

then  $\text{tr } M = \text{tr } A + \text{tr } D$ .

Recall that if an operator-valued function  $C(z)$  is finitely meromorphic in the neighborhood  $V$  of  $z_0$ , which contains no poles of  $C(z)$  except possibly  $z_0$ , then  $\int_{\partial V} C(z) dz$  is a finite-dimensional operator. The following identity will also be used frequently.

PROPOSITION 1.7. *Let  $A(z)$  and  $B(z)$  be two operator-valued functions which are finitely meromorphic in the neighborhood  $\bar{V}$  of  $z_0$ , which contains no poles of  $A(z)$  and  $B(z)$  other than  $z_0$ . Then we have*

$$(1.4) \quad \text{tr} \int_{\partial V} A(z)B(z) dz = \text{tr} \int_{\partial V} B(z)A(z) dz.$$

### 1.2. Factorization of Operators

We say that  $A(z) \in \mathfrak{U}(z_0)$  admits a factorization at  $z_0$  if  $A(z)$  can be written as

$$(1.5) \quad A(z) = E(z)D(z)F(z),$$

where  $E(z), F(z)$  are regular at  $z_0$  and

$$(1.6) \quad D(z) = P_0 + \sum_{j=1}^n (z - z_0)^{k_j} P_j.$$

Here,  $P_j$ 's are mutually disjoint projections,  $P_1, \dots, P_n$  are one-dimensional operators, and  $I - \sum_{j=0}^n P_j$  is a finite-dimensional operator.

**THEOREM 1.8.**  *$A(z) \in \mathfrak{U}(z_0)$  admits a factorization at  $z_0$  if and only if  $A(z)$  is finitely meromorphic and of Fredholm type of index zero at  $z_0$ .*

**PROOF.** Suppose that  $A(z)$  is finitely meromorphic and of Fredholm type of index zero at  $z_0$ . We shall construct  $E, F$ , and  $D$  such that (1.5) holds. Write the Laurent series expansion of  $A(z)$  as follows:

$$A(z) = \sum_{j=-\nu}^{+\infty} (z - z_0)^j A_j$$

in some neighborhood  $U$  of  $z_0$ . Since  $\text{ind} A_0 = 0$ , then by the Fredholm alternative  $B_0 := A_0 + K_0$  is invertible for some finite-dimensional operator  $K_0$ . Consequently,

$$B(z) := K_0 + \sum_{j=0}^{+\infty} (z - z_0)^j A_j$$

is invertible in some neighborhood  $U_1$  of  $z_0$  and

$$(1.7) \quad A(z) = C(z) + B(z) = B(z)[I + B^{-1}(z)C(z)],$$

where

$$C(z) = \sum_{j=-\nu}^{-1} (z - z_0)^j A_j - K_0.$$

Since  $K(z) := B^{-1}(z)C(z)$  is finitely meromorphic, we can write  $K(z)$  in the form

$$K(z) = \sum_{j=1}^{\nu} (z - z_0)^{-j} K_j + T_1(z),$$

where  $K_j, j = 1, \dots, \nu$ , are finite-dimensional and  $T_1$  is holomorphic.

Since the operators  $A_j$  and  $K_j$  are finite-dimensional, there exists a subspace  $\mathfrak{N}$  of  $\mathcal{B}$  of finite codimension such that

$$\begin{cases} \mathfrak{N} \subset \text{Ker } A_j, & j = -\nu, \dots, -1, \\ \mathfrak{N} \subset \text{Ker } K_j, & j = 0, \dots, \nu, \\ \mathfrak{N} \cap \text{Im } K_j = \{0\}, & j = 1, \dots, \nu. \end{cases}$$

Let  $\mathfrak{C}$  be a direct finite-dimensional complement of  $\mathfrak{N}$  in  $\mathcal{B}$  and let  $P$  be the projection onto  $\mathfrak{C}$  satisfying  $P(I - P) = 0$ . Set  $P_0 := I - P$ . We have

$$\begin{aligned} I + K(z) &= I + PK(z)P + P_0K(z)P \\ &= I + PK(z)P + P_0T_1(z)P, \end{aligned}$$

and therefore,

$$(1.8) \quad I + K(z) = (I + PK(z)P)(I + P_0T_1(z)P).$$

Since  $P(I + K(z))P$  can be viewed as an operator from  $\mathfrak{C}$  into itself and  $\mathfrak{C}$  is finite-dimensional, it follows from Gaussian elimination that

$$P(I + K(z))P = E_1(z)D_1(z)F_1(z),$$

where  $D_1(z)$  is diagonal and  $E_1(z)$  and  $F_1(z)$  are holomorphic and invertible. In view of (1.8), this implies that

$$\begin{aligned} A(z) &= B(z)(P_0 + P(I + K(z))P)(I + P_0T_1(z)P) \\ &= B(z)(P_0 + E_1(z)D_1(z)F_1(z))(I + P_0T_1(z)P) \\ &= B(z)(P_0 + E_1(z))(P_0 + D_1(z))(P_0 + F_1(z))(I + P_0T_1(z)P). \end{aligned}$$

Here  $I + P_0T_1(z)P$  is holomorphic and invertible with inverse  $I - P_0T_1(z)P$ . Thus, taking

$$E(z) := B(z)(P_0 + E_1(z)), \quad F(z) := (P_0 + F_1(z))(I + P_0T_1(z)P)$$

yields the desired factorization for  $A$  since  $E(z)$  and  $F(z)$ , given by the above formulas, are holomorphic and invertible at  $z_0$ .

The converse result, that  $A(z) = E(z)D(z)F(z)$  with  $E(z), F(z)$  regular at  $z_0$  and  $D(z)$  satisfying (1.6) is finitely meromorphic and of Fredholm type of index zero at  $z_0$ , is easy.  $\square$

**COROLLARY 1.9.**  *$A(z)$  is normal at  $z_0$  if and only if  $A(z)$  admits a factorization such that  $I = \sum_{j=0}^n P_j$  in (1.6). Moreover, we have*

$$M(A(z_0)) = k_1 + \cdots + k_n$$

for  $k_1, \dots, k_n$ , given by (1.6).

**COROLLARY 1.10.** *Every normal point of  $A(z)$  is a normal point of  $A^{-1}(z)$ .*

### 1.3. Main Results of the Gohberg and Sigal Theory

We now tackle our main goal of this chapter, which is to generalize the argument principle and Rouché's theorem to operator-valued functions.

**1.3.1. Argument Principle.** Let  $V$  be a simply connected bounded domain with rectifiable boundary  $\partial V$ . An operator-valued function  $A(z)$  which is finitely meromorphic and of Fredholm type in  $V$  and continuous on  $\partial V$  is called *normal* with respect to  $\partial V$  if the operator  $A(z)$  is invertible in  $\overline{V}$ , except for a finite number of points of  $V$  which are normal points of  $A(z)$ .

**LEMMA 1.11.** *An operator-valued function  $A(z)$  is normal with respect to  $\partial V$  if it is finitely meromorphic and of Fredholm type in  $V$ , continuous on  $\partial V$ , and invertible for all  $z \in \partial V$ .*

PROOF. To prove that  $A$  is normal with respect to  $\partial V$ , it suffices to prove that  $A(z)$  is invertible except at a finite number of points in  $V$ . To this end choose a connected open set  $U$  with  $\overline{U} \subset V$  so that  $A(z)$  is invertible in  $V \setminus U$ . Then, for each  $\xi \in U$ , there exists a neighborhood  $U_\xi$  of  $\xi$  in which the factorization (1.5) holds. In  $U_\xi$ , the kernel of  $A(z)$  has a constant dimension except at  $\xi$ . Since  $\overline{U}$  is compact, we can find a finite covering of  $\overline{U}$ , *i.e.*,

$$\overline{U} \subset U_{\xi_1} \cup \cdots \cup U_{\xi_k},$$

for some points  $\xi_1, \dots, \xi_k \in U$ . Therefore,  $\dim \text{Ker } A(z)$  is constant in  $V \setminus \{\xi_1, \dots, \xi_k\}$ , and so  $A(z)$  is invertible in  $\overline{V} \setminus \{\xi_1, \dots, \xi_k\}$ .  $\square$

Now, if  $A(z)$  is normal with respect to the contour  $\partial V$  and  $z_i, i = 1, \dots, \sigma$ , are all its characteristic values and poles lying in  $V$ , we put

$$(1.9) \quad \mathcal{M}(A(z); \partial V) = \sum_{i=1}^{\sigma} M(A(z_i)).$$

The full multiplicity  $\mathcal{M}(A(z); \partial V)$  of  $A(z)$  in  $V$  is the number of characteristic values of  $A(z)$  in  $V$ , counted with their multiplicities, minus the number of poles of  $A(z)$  in  $V$ , counted with their multiplicities.

THEOREM 1.12 (Generalized argument principle). *Suppose that the operator-valued function  $A(z)$  is normal with respect to  $\partial V$ . Then we have*

$$(1.10) \quad \mathcal{M}(A(z); \partial V) = \frac{1}{2\sqrt{-1}\pi} \text{tr} \int_{\partial V} A^{-1}(z) \frac{d}{dz} A(z) dz.$$

PROOF. Let  $z_j, j = 1, \dots, \sigma$ , denote all the characteristic values and all the poles of  $A$  lying in  $V$ . The key of the proof lies in using the factorization (1.5) in each of the neighborhoods of the points  $z_j$ . We have

$$(1.11) \quad \frac{1}{2\sqrt{-1}\pi} \text{tr} \int_{\partial V} A^{-1}(z) \frac{d}{dz} A(z) dz = \sum_{j=1}^{\sigma} \frac{1}{2\sqrt{-1}\pi} \text{tr} \int_{\partial V_j} A^{-1}(z) \frac{d}{dz} A(z) dz,$$

where, for each  $j$ ,  $V_j$  is a neighborhood of  $z_j$ . Moreover, in each  $V_j$ , the following factorization of  $A$  holds:

$$A(z) = E^{(j)}(z) D^{(j)}(z) F^{(j)}(z), \quad D^{(j)}(z) = P_0^{(j)} + \sum_{i=1}^{n_j} (z - z_j)^{k_{ij}} P_i^{(j)}.$$

As for the matrix-valued case at the beginning of this chapter, it is readily verified that

$$\begin{aligned} \frac{1}{2\sqrt{-1}\pi} \text{tr} \int_{\partial V_j} A^{-1}(z) \frac{d}{dz} A(z) dz &= \frac{1}{2\sqrt{-1}\pi} \text{tr} \int_{\partial V_j} (D^{(j)}(z))^{-1} \frac{d}{dz} D^{(j)}(z) dz \\ &= \sum_{i=1}^{n_j} k_{ij} = M(A(z_j)). \end{aligned}$$

Now, (1.10) follows by using (1.11).  $\square$

The following is an immediate consequence of Lemma 1.11, identity (1.10), and (1.4).

COROLLARY 1.13. *If the operator-valued functions  $A(z)$  and  $B(z)$  are normal with respect to  $\partial V$ , then  $C(z) := A(z)B(z)$  is also normal with respect to  $\partial V$ , and*

$$\mathcal{M}(C(z); \partial V) = \mathcal{M}(A(z); \partial V) + \mathcal{M}(B(z); \partial V).$$

The following general form of the argument principle will be useful. It can be proven by the same argument as the one in Theorem 1.12.

THEOREM 1.14. *Suppose that  $A(z)$  is an operator-valued function which is normal with respect to  $\partial V$ . Let  $f(z)$  be a scalar function which is analytic in  $V$  and continuous in  $\bar{V}$ . Then*

$$\frac{1}{2\sqrt{-1}\pi} \operatorname{tr} \int_{\partial V} f(z) A^{-1}(z) \frac{d}{dz} A(z) dz = \sum_{j=1}^{\sigma} M(A(z_j)) f(z_j),$$

where  $z_j, j = 1, \dots, \sigma$ , are all the points in  $V$  which are either poles or characteristic values of  $A(z)$ .

**1.3.2. Generalization of Rouché's Theorem.** A generalization of Rouché's theorem to operator-valued functions is stated below.

THEOREM 1.15 (Generalized Rouché's theorem). *Let  $A(z)$  be an operator-valued function which is normal with respect to  $\partial V$ . If an operator-valued function  $S(z)$  which is finitely meromorphic in  $V$  and continuous on  $\partial V$  satisfies the condition*

$$\|A^{-1}(z)S(z)\|_{\mathcal{L}(\mathcal{B}, \mathcal{B})} < 1, \quad z \in \partial V,$$

then  $A(z) + S(z)$  is also normal with respect to  $\partial V$  and

$$\mathcal{M}(A(z); \partial V) = \mathcal{M}(A(z) + S(z); \partial V).$$

PROOF. Let  $C(z) := A^{-1}(z)S(z)$ . By Corollary 1.10,  $C(z)$  is finitely meromorphic in  $V$ . Suppose that  $z_1, z_2, \dots, z_n$ , are all of the poles of  $C(z)$  in  $V$  and that  $C(z)$  has the following Laurent series expansion in some neighborhood of each  $z_j$ :

$$C(z) = \sum_{k=-\nu_j}^{+\infty} (z - z_j)^k C_k^{(j)}.$$

Let  $\mathfrak{N}$  be the intersection of the kernels  $\operatorname{Ker} C_k^{(j)}$  for  $j = 1, \dots, n$  and  $k = 1, \dots, \nu_j$ . Then,  $\dim \mathcal{B}/\mathfrak{N} < +\infty$  and the restriction  $C(z)|_{\mathfrak{N}}$  of  $C(z)$  to  $\mathfrak{N}$  is holomorphic in  $V$ .

Let  $q := \max_{z \in \partial V} \|C(z)\|$ , which by assumption is less than 1. Since

$$\Delta_z \|C(z)|_{\mathfrak{N}}\|^2 = 4 \left\| \frac{\partial}{\partial z} C(z)|_{\mathfrak{N}} \right\|^2,$$

then  $\|C(z)|_{\mathfrak{N}}\|$  is subharmonic in  $V$ , and hence we have from the maximum principle

$$\max_{z \in V} \|C(z)|_{\mathfrak{N}}\| \leq q.$$

It then follows that

$$\|(I + C(z))x\| \geq (1 - q)\|x\|, \quad x \in \mathfrak{N}, \quad z \in V.$$

This implies that  $(I + C(z))|_{\mathfrak{H}}$  has a closed range and  $\text{Ker}(I + C(z))|_{\mathfrak{H}} = 0$ . Therefore,  $I + C(z)$  has a closed range and a kernel of finite dimension for  $z \in V \setminus \{z_1, \dots, z_n\}$ . By a slight extension of Proposition 1.5 [241],  $\mathcal{I}(z)$  defined by

$$\mathcal{I}(z) = \dim \text{Ker}(I + C(z)) - \text{codim Im}(I + C(z))$$

is continuous for  $z \in \overline{V} \setminus \{z_1, \dots, z_n\}$ . Thus,

$$\text{ind}(I + C(z)) = 0 \quad \text{for } z \in \overline{V} \setminus \{z_1, \dots, z_n\}.$$

Moreover, since the Laurent series expansion of  $(I + C(z))|_{\mathfrak{H}}$  in a neighborhood of  $z_j$  is given by

$$(1.12) \quad (I + C(z))|_{\mathfrak{H}} = I|_{\mathfrak{H}} + \sum_{k=0}^{+\infty} (z - z_j)^k C_k^{(j)}|_{\mathfrak{H}},$$

it follows that  $(I + C_0^{(j)})|_{\mathfrak{H}}$  has a closed range and a trivial kernel. Using Propositions 1.4 and 1.5, we have

$$\text{ind}(I + C_0^{(j)}) = \text{ind}\left(I + \sum_{k=0}^{+\infty} (z - z_j)^k C_k^{(j)}\right) = \text{ind}(I + C(z)) = 0.$$

Thus,  $(I + C_0^{(j)})$  is Fredholm. By Lemma 1.11, we deduce that  $I + C(z)$  is normal with respect to  $\partial V$ .

Now we claim that  $\mathcal{M}(I + C(z); \partial V) = 0$ . To see this, we note that  $I + tC(z)$  is normal with respect to  $\partial V$  for  $0 \leq t \leq 1$ . Let

$$f(t) := \mathcal{M}(I + tC(z); \partial V).$$

Then  $f(t)$  attains integers as its values. On the other hand, since

$$(1.13) \quad f(t) = \frac{1}{2\sqrt{-1}\pi} \text{tr} \int_{\partial V} t(I + tC(z))^{-1} \frac{d}{dz} C(z) dz$$

and  $(I + tC(z))^{-1}$  is continuous in  $[0, 1]$  in operator norm uniformly in  $z \in \partial V$ ,  $f(t)$  is continuous in  $[0, 1]$ . Thus,  $f(1) = f(0) = 0$ .

Finally, with the help of Corollary 1.13, we can conclude that the theorem holds.  $\square$

**1.3.3. Generalization of Steinberg's theorem.** Steinberg's theorem asserts that if  $K(z)$  is a compact operator on a Banach space, which is analytic in  $z$  for  $z$  in a region  $V$  in the complex plane, then  $I + K(z)$  is meromorphic in  $V$ . See [238]. A generalization of this theorem to finitely meromorphic operators was first given by Gohberg and Sigal in [114]. The following important result holds.

**THEOREM 1.16 (Generalized Steinberg's theorem).** *Suppose that  $A(z)$  is an operator-valued function which is finitely meromorphic and of Fredholm type in the domain  $V$ . If the operator  $A(z)$  is invertible at one point of  $V$ , then  $A(z)$  has a bounded inverse for all  $z \in V$ , except possibly for certain isolated points.*

#### 1.4. Concluding Remarks

In this chapter, we have reviewed the main results in the theory of Gohberg and Sigal on meromorphic operator-valued functions. These results concern the generalization of the argument principle and the Rouché theorem to meromorphic operator-valued functions. Some of these results have been extended to very general operator-valued functions in [46, 170] and with other types of spectrum than isolated eigenvalues in [174].

Throughout this book, the theory of Gohberg and Sigal will be applied to perturbation theory of eigenvalues. Other interesting applications include the investigation of scattering resonances and scattering poles [118, 57] and the study of the regularity of the solutions of elliptic boundary value problems near conical points [154].