

Index

- admissible operator, 131
- admissible set of operators, 142
- admissible set of symbols, 26

- Bernoulli numbers, 79
- Bernoulli polynomials, 80
- Beta function, 3
- bidegree of conformal covariance, 174

- canonical discrete sum, 94
- canonical integral, 28
- canonical trace, 123
- classical symbol on an open set, 106
- closed linear forms, 22
- conformal anomaly, 180
- conformal Laplacian, 175
- conformality, 174
- conformally covariant operators, 174
- conformally invariant functional, 177
- continuity of a weighted trace, 148
- continuity of the residue, 122
- convex (closed) polytope, 83
- covariant linear forms on symbols, 32
- covariant set of symbols, 32
- cut-off Gamma function, 11
- cut-off integral, 36, 86
- cut-off sum, 82, 89

- defect formula, 148
- dimensional regularisation, 72
- discrepancies, 12, 13, 39, 74, 139

- elliptic operator, 111
- energy-momentum tensor, 181
- Epstein's ζ -function, 102
- Euler's constant, 2
- Euler's lemma, 21
- Euler-Maclaurin formula, 80, 81
- excision function around the origin, 17
- expanded polytope, 84

- formal classical symbols, 19
- Fréchet topology on classical operators, 117
- Fréchet topology on classical symbols, 19

- Gamma function, 2

- Hadamard integral and method, 7, 36
- Hadamard extension of Gamma, 11
- Hadamard finite part method, 1, 7
- heat-kernel regularised trace, 158
- holomorphic regularisation on operators, 135
- holomorphic family in a Banach space, 56
- holomorphic family of operators, 56, 133
- holomorphic family of symbols, 57
- holomorphic regularisation on symbols, 70
- homogeneous distribution, 2, 16
- Hurwitz zeta function, 98

- index of D_+ , 159
- integral polytopes, 83

- kernel of an operator, 108
- Khovanskii-Pukhlikov formula, 84

- Laplace-Beltrami operator, 162, 175
- leading symbol, 18, 107, 111, 116
- Lichnerowicz formula, 162
- local conformal anomaly, 181
- local discrepancies, 139
- localisation subordinated to a chart, 114
- locality of the noncommutative residue, 119
- log-polyhomogeneous operator, 111
- log-polyhomogeneous order, 107
- log-polyhomogeneous symbol, 17, 107
- log-type, 17, 107

- minimal subtraction scheme, 54
- modified dimensional regularisation, 70
- multiplicative anomaly of a weighted determinant, 170
- multiplicative anomaly of the zeta determinant, 173

- noncommutative residue of a symbol, 21
- noncommutative residue of an operator, 121
- nonlocality of canonical traces, 123
- nonlocality of regularised traces, 136

- odd-class symbols, 18
- order of a symbol, 17
- order of an operator, 116

- Paneitz operators, 175
- period, 38
- pointwise conformally covariant of weight, 177
- Polyakov action, 181
- polyhomogeneous operator, 111
- polyhomogeneous symbol, 17, 85
- primitive lattice vectors, 83
- principal angle, 131
- properly supported symbol, 110
- properly supported operator, 110
- pseudodifferential operator, 108, 109, 115
- pseudolocality, 109

- Q -curvature, 175
- Q -weighted determinant, 167
- quotient Fréchet topology on formal
 - classical symbols, 19

- \mathbb{R}^d -translation invariant set/linear form, 30
- \mathcal{R} -regularised discrete sum, 96
- \mathcal{R} -regularised integral, 71
- \mathcal{R} -regularised trace, 135
- regular integral polytopes, 83
- regular linear form, 122
- regularised evaluator, 54
- regularised Euler-Maclaurin formula, 96
- rescaling invariant linear form, 41
- rescaling invariant set, 41
- residue determinant, 155
- Riesz extension of Gamma function, 7
- Riesz regularisation, 5, 70
- Rota-Baxter operator, 54
- Rota-Baxter relation, 54

- Schwartz functions on \mathbb{R}^d , 16
- Schwartz functions on \mathbb{R}^+ , 1
- simple integral polytopes, 83
- singular linear form on operators, 122
- singular linear form on symbols, 20
- singular support of a distribution, 109
- smoothing operator, 108, 115
- smoothing symbol, 16
- star-product, 112
- Stokes' property on symbols, 23
- Stokes' theorem, 23
- symbol valued differential forms, 22
- symbolically defined set of operators, 121
- symbols with constant coefficients, 15

- Todd function, 80
- trace of a smoothing operator, 115
- translation invariant linear form on
 - symbols, 30
- translation invariant set of symbols, 30

- twisted Dirac operator, 162
- twisted \mathbb{Z}_2 -graded spinor bundle, 162
- type k -log-polyhomogeneous, 17, 106

- weight for a trace, 136
- weight, conformal, 174
- weighted determinant, 157
- weighted supertraces, 158
- weighted trace, 136

- Yamabe operator, 175

- \mathbb{Z}^d -translation invariant, 91
- zeta function, 98
- zeta regularised traces, 135
- zeta-determinant, 168