

## ASYMPTOTICS OF SMALL EIGENVALUES OF RIEMANN SURFACES

MARC BURGER

Recently there has been a great deal of interest in geometric bounds on small eigenvalues of the Laplace operator on a Riemann surface [S.W.Y., D.P.R.S.]. Here we determine the precise asymptotic behaviour of these small eigenvalues. Let  $S_\delta$  be a compact Riemann surface of genus  $g \geq 2$  whose first  $k$  nonzero eigenvalues  $0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k$  are small, i.e.,  $\lambda_k \leq \delta$  and  $\lambda_{k+1} \geq c_1$ . Then by [S.W.Y.] there exists a constant  $a = a(g) > 0$  such that the closed geodesics  $\gamma_1 \cdots \gamma_r$  of length less than  $a \cdot \delta$  separate  $S_\delta$  into  $k + 1$  pieces  $S_1, \dots, S_{k+1}$  and all other closed geodesics of  $S_\delta$  have length greater than  $a(g)$ . Let  $\Lambda$  be the graph whose vertices are the pieces  $S_i$ . Suppose vertex  $S_i$  has mass  $v_i = \text{vol}(S_i)$  and the length  $L_{ij}$  of an edge joining  $S_i$  to  $S_j$  is the total length of the geodesics contained in  $S_i \cap S_j$ . Furthermore, let  $0 < \lambda_1(\Lambda) \leq \dots \leq \lambda_k(\Lambda)$  be the spectrum of the quadratic form  $\sum (F(S_i) - F(S_j))^2 L_{ij}$  with respect to the norm  $\sum F(S_i)^2 v_i$ . Then one has

THEOREM 1.

$$\lim_{\delta \rightarrow 0} \frac{\lambda_j(S_\delta)}{\lambda_j(\Lambda)} = \frac{1}{\pi} \quad \text{for all } 1 \leq j \leq k.$$

This convergence is uniform for all surfaces  $S_\delta$  with  $\lambda_{k+1}(S_\delta) \geq c_1$  and fixed genus.

REMARK. The fact that  $\limsup_{\delta \rightarrow 0} \lambda_j(S_\delta)/\lambda_j(\Lambda) \leq 1/\pi$  follows easily from [C.CdV]. This paper also shows the convergence of this ratio in the case that the lengths  $l(\gamma_i)$  all have the same behaviour near zero, i.e.  $l(\gamma_i) = d_i \varepsilon$  for  $\varepsilon \rightarrow 0$ , and fixed  $d_i$ .

SKETCH OF PROOF. Complete  $\gamma_1 \cdots \gamma_r$  to a set of geodesics  $\gamma_1 \cdots \gamma_{3g-3}$ , giving a decomposition of  $S$  into  $Y$ -pieces with length  $l(\gamma_i) \leq L_g$ , a constant depending only on  $g$  (see [Bu2, §13]). Then using a modified version of an argument of [B1] we show that  $\lambda_j \cdot (1 + o(\sqrt{\delta})) \geq \pi^{-1} \lambda_j(\Gamma)$ , where  $\Gamma$  is the graph of the  $Y$ -pieces, and the length of an edge corresponding to a small geodesic is  $l(\gamma)$ . The proof of this also uses the asymptotic of the first nonzero eigenvalue of  $Y_1 \cup Y_2$  for the Neumann problem, where  $Y_1, Y_2$  are  $Y$ -pieces,  $Y_1 \cap Y_2 = \gamma$  and  $l(\gamma)$  is small. This can be deduced from [C.CdV], because there is only one small geodesic separating  $Y_1 \cup Y_2$ . To finish the proof we have then to compare  $\lambda_j(\Gamma)$  with  $\lambda_j(\Lambda)$ . To do this we consider  $\Lambda$  to be the graph of the connected components of  $\Gamma$  after removing the small edges of  $\Gamma$ .

Received by the editors February 10, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 53C22, 58G25; Secondary 30F99.

