

PIERCING CONVEX SETS

NOGA ALON AND DANIEL J. KLEITMAN

ABSTRACT. A family of sets has the (p, q) property if among any p members of the family some q have a nonempty intersection. It is shown that for every $p \geq q \geq d + 1$ there is a $c = c(p, q, d) < \infty$ such that for every family \mathcal{F} of compact, convex sets in R^d that has the (p, q) property there is a set of at most c points in R^d that intersects each member of \mathcal{F} . This extends Helly's Theorem and settles an old problem of Hadwiger and Debrunner.

1. INTRODUCTION

For two integers $p \geq q$, a family of sets \mathcal{H} has the (p, q) property if among any p members of the family some q have a nonempty intersection. \mathcal{H} is k -pierceable if it can be split into k or fewer subfamilies, each having a nonempty intersection. The piercing number of \mathcal{H} , denoted by $P(\mathcal{H})$, is the minimum value of k such that \mathcal{H} is k -pierceable. (If no such finite k exists then $P(\mathcal{H}) = \infty$.)

The classical theorem of Helly [14] states that any family of compact convex sets in R^d that satisfies the $(d + 1, d + 1)$ -property is 1-pierceable. Hadwiger and Debrunner considered the more general problem of studying the piercing numbers of families \mathcal{F} of compact, convex sets in R^d that satisfy the (p, q) property. By considering the intersections of hyperplanes in general position in R^d with an appropriate box one easily checks that for $q \leq d$ the piercing number can be infinite, even if $p = q$. Thus we may assume that $p \geq q \geq d + 1$.

Let $M(p, q; d)$ denote the maximum possible piercing number (which is possibly infinity) of a family of compact convex sets in R^d with the (p, q) -property. By Helly's Theorem,

$$M(d + 1, d + 1; d) = 1$$

for all d and, trivially, $M(p, q; d) \geq p - q + 1$. Hadwiger and Debrunner [12] proved that for $p \geq q \geq d + 1$ satisfying

$$(1) \quad p(d - 1) < (q - 1)d,$$

this bound is tight, i.e., $M(p, q; d) = p - q + 1$. In all other cases, it is not even known if $M(p, q; d)$ is finite, and the question of deciding if this function is finite, raised by Hadwiger and Debrunner in 1957 in [12], remained open. This question,

1991 *Mathematics Subject Classification*. Primary 52A35.

Research supported in part by a United States Israel BSF Grant and by a Bergmann Memorial Grant

Supported in part by NSF grant DMS-9108403

Received by the editors August 23, 1991

which is usually referred to as the (p, q) -problem, is considered in various survey articles and books, including [13, 5, 8]. The smallest case in which finiteness is unknown, which is pointed out in all the above-mentioned articles, is the special case $p = 4, q = 3, d = 2$. We note, in all the cases where finiteness is known, that in fact $M(p, q; d) = p - q + 1$ and there are examples of Danzer and Grünbaum (cf. [13]) that show $M(4, 3; 2) \geq 3 > 4 - 3 + 1$.

The (p, q) -problem received a considerable amount of attention, and finiteness was proved for various restricted classes of convex sets, including the family of parallelotopes with edges parallel to the coordinate axes in R^d [13, 19, 6], families of homothetes of a convex set [19], and, using a similar approach, families of convex sets with a certain “squareness” property ([9], see also [21]).

Despite these efforts, the problem of deciding if $M(p, q; d)$ is finite remained open for all values of $p \geq q \geq d + 1$ that do not satisfy (1).

The purpose of the present note is to announce the solution of this problem, established in the following

Theorem 1.1. *For every $p \geq q \geq d + 1$ there is a $c = c(p, q, d) < \infty$ such that $M(p, q; d) \leq c$; i.e., for every family \mathcal{F} of compact, convex sets in R^d that has the (p, q) property there is a set of at most c points in R^d that intersects each member of \mathcal{F} .*

The detailed proof will appear in [3]. Here we briefly sketch the main ideas. Three tools are applied: a fractional version of Helly’s Theorem that was first proved in [15], Farkas’s Lemma (or Linear Programming Duality), and a recent result proved in [1].

It may seem that there are almost no interesting families of compact convex sets in R^d that satisfy the (p, q) -property for some $p \geq q \geq d + 1$. A large class of examples can be constructed as follows. Let μ be an arbitrary probability distribution on R^d , and let \mathcal{F} be the family of all compact convex sets F in R^d satisfying $\mu(F) \geq \varepsilon$. Since the sum of the measures of any set of more than d/ε such sets is greater than d , it follows that if p is the smallest integer strictly larger than d/ε then \mathcal{F} has the $(p, d + 1)$ property. It then follows that $P(\mathcal{F}) \leq M(p, d + 1; d + 1)$, i.e., for every probability measure in R^d there is a set X of at most $M(p, d + 1; d + 1)$ points such that any compact convex set in R^d whose measure exceeds ε intersects X .

The following theorem is an immediate consequence of Theorem 1.1.

Theorem 1.2. *Let \mathcal{F} be a family of compact convex sets in R^d , and suppose that for every subfamily \mathcal{F}' of cardinality x of \mathcal{F} the inequality $P(\mathcal{F}') < \lceil x/d \rceil$ holds; i.e., \mathcal{F}' can be pierced by less than x/d points. Then $P(\mathcal{F}) \leq M(x, d + 1; d + 1)$.*

Observe that in order to deduce a finite upper bound for the piercing number of \mathcal{F} , the assumption that $P(\mathcal{F}') < \lceil x/d \rceil$ cannot be replaced by $P(\mathcal{F}') \leq \lceil x/d \rceil$ as shown by an infinite family of hyperplanes in general position (intersected with an appropriate box), whose piercing number is infinite.

2. A SKETCH OF THE PROOFS

Since we do not try to optimize the constants here, and since obviously $M(p, q; d) \leq M(p, d + 1; d)$ for all $p \geq q \geq d + 1$, it suffices to prove an upper bound for $M(p, d + 1; d)$. Another simple observation is that by compactness we can restrict our attention to finite families of convex sets.

Let \mathcal{F} be a family of n convex sets in R^d , and suppose that \mathcal{F} has the $(p, d+1)$ property. Our objective is to find an upper bound for the piercing number $P(\mathcal{F})$ of \mathcal{F} , where the bound depends only on p and d . It is convenient to describe the ideas in three subsections.

2.1. A fractional version of Helly's Theorem. Katchalski and Liu [15] proved the following result, which can be viewed as a fractional version of Helly's Theorem.

Theorem 2.1 [15]. *For every $0 < \alpha \leq 1$ and every d there is a $\delta = \delta(\alpha, d) > 0$ such that for every $n \geq d+1$, every family of n convex sets in R^d that contains at least $\alpha \binom{n}{d+1}$ intersecting subfamilies of cardinality $d+1$ contains an intersecting subfamily of at least δn of the sets.*

Notice that Helly's Theorem is equivalent to the statement that in the above theorem $\delta(1, d) = 1$.

A sharp quantitative version of this theorem was proved by Kalai [16] and, independently, by Eckhoff [7]. See also [2] for a very short proof. All these proofs rely on Wegner's Theorem [20] that asserts that the nerve of a family of convex sets in R^d is d -collapsible.

Theorem 2.1, together with a simple probabilistic argument, can be applied to prove

Lemma 2.2. *For every $p \geq d+1$ there is a positive constant $\beta = \beta(p, d)$ with the following property. Let $\mathcal{F} = \{A_1, \dots, A_n\}$ be a family of n convex sets in R^d that has the $(p, d+1)$ property. Let a_i be nonnegative integers, define $m = \sum_{i=1}^n a_i$, and let \mathcal{G} be the family of cardinality m consisting of a_i copies of A_i for $1 \leq i \leq n$. Then there is a point x in R^d that belongs to at least βm members of \mathcal{G} .*

2.2. Farkas's Lemma and a lemma on hypergraphs. The following is a known variant of the well-known lemma of Farkas (cf. [17, p. 90]).

Lemma 2.3. *Let A be a real matrix and b a real (column) vector. Then the system $Ax \leq b$ has a solution $x \geq 0$ if and only if for every (row) vector $y \geq 0$ that satisfies $yA \geq 0$ the inequality $yb \geq 0$ holds.*

This lemma can be used to prove the following.

Corollary 2.4. *Let $H = (V, E)$ be a hypergraph and let $0 \leq \gamma \leq 1$ be a real. Then the following two conditions are equivalent:*

- (i) *There exists a weight function $f: V \mapsto R^+$ satisfying $\sum_{v \in V} f(v) = 1$ and $\sum_{v \in e} f(v) \geq \gamma$ for all $e \in E$.*
- (ii) *For every function $g: E \mapsto R^+$ there is a vertex $v \in V$ such that $\sum_{e: v \in e} g(e) \geq \gamma \sum_{e \in E} g(e)$.*

By the last corollary and Lemma 2.2 one can prove the following result.

Corollary 2.5. *Suppose $p \geq d+1$ and let $\beta = \beta(p, d)$ be the constant from Lemma 2.2. Then for every family $\mathcal{F} = \{A_1, \dots, A_n\}$ of n convex sets in R^d with the $(p, d+1)$ property there is a finite (multi-)set $Y \subset R^d$ such that $|Y \cap A_i| \geq \beta|Y|$ for all $1 \leq i \leq n$.*

2.3. Weak ε -nets for convex sets. The following result is proved in [1].

Theorem 2.6 [1]. *For every real $0 < \varepsilon < 1$ and every integer d there exists a constant $b = b(\varepsilon, d)$ such that the following holds:*

For every m and every multiset Y of m points in R^d , there is a subset X of at most b points in R^d such that the convex hull of any subset of εm members of Y contains at least one point of X .

Several arguments that supply various upper bounds for $b(\varepsilon, d)$ are given in [1]. The simplest one is based on a result of Bárány [4] whose proof is based on a deep result of Tverberg [8].

Theorem 1.1 follows from the above results quite easily. Let $\mathcal{F} = \{A_1, \dots, A_n\}$ be a family of n convex sets in R^d with the $(p, d+1)$ property, where $p \geq d+1$. By Corollary 2.5 there is a finite (multi-)set $Y \subset R^d$ such that $|Y \cap A_i| \geq \beta|Y|$ for all $1 \leq i \leq n$, where $\beta = \beta(p, d)$ is as in Lemma 2.2. By Theorem 2.6 there is a set X of at most $b(\beta, d)$ points in R^d such that the convex hull of any set of $\beta|Y|$ members of Y contains at least one point of X . Since each member of \mathcal{F} contains at least $\beta|Y|$ points in Y , it must contain at least one point of X . Therefore, $P(\mathcal{F}) \leq |X| \leq b(\beta(p, d), d)$, completing the proof.

The detailed proofs of the above lemmas and corollaries, as well as some methods to improve the estimates for the numbers $M(p, q; d)$, will appear in [3].

REFERENCES

1. N. Alon, I. Bárány, Z. Füredi, and D. J. Kleitman, *Point selections and weak ε -nets for convex hulls*, *Combin. Prob. and Computing* **1**, submitted.
2. N. Alon and G. Kalai, *A simple proof of the upper bound theorem*, *European J. Combinatorics* **6** (1985), 211–214.
3. N. Alon and D. J. Kleitman, *Piercing convex sets and the Hadwiger Debrunner (p, q) problem*, *Adv. Math.* (to appear).
4. I. Bárány, *A generalization of Carathéodory's theorem*, *Discrete Math.* **40** (1982), 141–152.
5. L. Danzer, B. Grünbaum, and V. Klee, *Helly's theorem and its relatives*, *Proc. Sympos. Pure Math.*, vol. 7, Amer. Math. Soc., Providence, RI, 1963, pp. 101–180.
6. V. L. Dol'nikov, *A coloring problem*, *Sibirsk. Mat. Zh.* **13** (1972), 1272–1283; English transl. in *Siberian Math J.* **13** (1972), 886–894.
7. J. Eckhoff, *An upper bound theorem for families of convex sets*, *Geom. Dedicata* **19** (1985), 217–227.
8. ———, *Helly, Radon, and Carathéodory type theorems*, *Handbook of Convex Geometry* (to appear).
9. B. E. Fullbright, *Intersectional properties of certain families of compact convex sets*, *Pacific J. Math.* **50** (1974), 57–62.
10. B. Grünbaum, *On intersections of similar sets*, *Portugal Math.* **18** (1959), 155–164.
11. ———, *Lectures on combinatorial geometry*, Mimeographed Notes, Univ. of Washington, Seattle, 1974.
12. H. Hadwiger and H. Debrunner, *Über eine Variante zum Helly'schen Satz*, *Arch. Math.* **8** (1957), 309–313.
13. H. Hadwiger, H. Debrunner, and V. Klee, *Combinatorial geometry in the plane*, Holt, Rinehart, and Winston, New York, 1964.
14. E. Helly, *Über Mengen konvexer Körper mit gemeinschaftlichen Punkten*, *Jahresber. Deutsch. Math. Verein.* **32** (1923), 175–176.
15. M. Katchalski and A. Liu, *A problem of geometry in R^d* , *Proc. Amer. Math. Soc.* **75** (1979), 284–288.
16. G. Kalai, *Intersection patterns of convex sets*, *Israel J. Math.* **48** (1984), 161–174.
17. A. Schrijver, *Theory of linear and integer programming*, Wiley, New York, 1986.
18. H. Tverberg, *A generalization of Radon's Theorem*, *J. London Math. Soc. (2)* **41** (1966), 123–128.
19. G. Wegner, *Über eine kombinatorisch-geometrische Frage von Hadwiger und Debrunner*, *Israel J. Math.* **3** (1965), 187–198.

20. _____, *d-collapsing and nerves of families of convex sets*, Arch. Math. **26** (1975), 317–321.
21. _____, *Über Helly-Gallaische Stichzahlprobleme*, 3, Koll. Discrete Geom. Salzburg (1985), 277–282.

DEPARTMENT OF MATHEMATICS, RAYMOND AND BEVERLY SACKLER FACULTY OF EXACT SCIENCES, TEL AVIV UNIVERSITY, TEL AVIV, ISRAEL AND BELLCORE, MORRISTOWN, NEW JERSEY 07960

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MASSACHUSETTS 02139