


CONTEMPORARY MATHEMATICS

342

Towards a Theory of Geometric Graphs

János Pach
Editor



American Mathematical Society

Towards a Theory of Geometric Graphs



Paul Turán (1910–1976)



Paul Erdős (1913–1996)

CONTEMPORARY MATHEMATICS

342

Towards a Theory of Geometric Graphs

János Pach
Editor



American Mathematical Society
Providence, Rhode Island

Editorial Board

Dennis DeTurck, managing editor

Andreas Blass Andy R. Magid Michael Vogelius

2000 *Mathematics Subject Classification*. Primary 05C62; Secondary 57M50, 90C27, 05C35, 68W35.

Library of Congress Cataloging-in-Publication Data

Towards a theory of geometric graphs / János Pach, editor.

p. cm. — (Contemporary mathematics, ISSN 0271-4132 ; 342)

Includes bibliographical references.

ISBN 0-8218-3484-3 (alk. paper)

1. Graph theory. I. Pach, János. II. Contemporary mathematics (American Mathematical Society) ; v. 342.

QA166.T68 2004

511'.5—dc22

2003063929

Copying and reprinting. Material in this book may be reproduced by any means for educational and scientific purposes without fee or permission with the exception of reproduction by services that collect fees for delivery of documents and provided that the customary acknowledgment of the source is given. This consent does not extend to other kinds of copying for general distribution, for advertising or promotional purposes, or for resale. Requests for permission for commercial use of material should be addressed to the Acquisitions Department, American Mathematical Society, 201 Charles Street, Providence, Rhode Island 02904-2294, USA. Requests can also be made by e-mail to reprint-permission@ams.org.

Excluded from these provisions is material in articles for which the author holds copyright. In such cases, requests for permission to use or reprint should be addressed directly to the author(s). (Copyright ownership is indicated in the notice in the lower right-hand corner of the first page of each article.)

© 2004 by the American Mathematical Society. All rights reserved.

The American Mathematical Society retains all rights
except those granted to the United States Government.

Printed in the United States of America.

∞ The paper used in this book is acid-free and falls within the guidelines
established to ensure permanence and durability.

Visit the AMS home page at <http://www.ams.org/>

10 9 8 7 6 5 4 3 2 1 09 08 07 06 05 04

To the memories of Paul Erdős and Paul Turán

Contents

| | |
|--|-----|
| Preface | ix |
| On the complexity of the linkage reconfiguration problem HELMUT ALT, CHRISTIAN KNAUER, GÜNTER ROTE, AND SUE WHITESIDES | 1 |
| Falconer conjecture, spherical averages and discrete analogs GEORGYI ARUTYUNYANTS AND ALEXANDER IOSEVICH | 15 |
| Turán-type extremal problems for convex geometric hypergraphs PETER BRASS | 25 |
| The thrackle conjecture for K_5 and $K_{3,3}$ GRANT CAIRNS, MARGARET MCINTYRE, AND YURY NIKOLAYEVSKY | 35 |
| Three-dimensional grid drawings with sub-quadratic volume VIDA DUJMOVIĆ AND D. R. WOOD | 55 |
| On a coloring problem for the integer grid ADRIAN DUMITRESCU AND RADOŠ RADOIČIĆ | 67 |
| Separating thickness from geometric thickness DAVID EPPSTEIN | 75 |
| Direction trees in centered polygons ROBERT E. JAMISON | 87 |
| Path coverings of two sets of points in the plane ATSUSHI KANEKO, MIKIO KANO, AND KAZUHIRO SUZUKI | 99 |
| Length of sums in a Minkowski space GYULA O. H. KATONA, RICHARD MAYER, AND WOJBOR A. WOYCZYNSKI | 113 |
| A new entropy inequality for the Erdős distance problem NETS HAWK KATZ AND GÁBOR TARDOS | 119 |
| Coloring intersection graphs of geometric figures with a given clique number ALEXANDER KOSTOCHKA | 127 |
| Convex quadrilaterals and k -sets LÁSZLÓ LOVÁSZ, KATALIN VESZTERGOMBI, ULI WAGNER, AND EMO WELZL | 139 |

| | |
|---|-----|
| Distance graphs and rigidity HIROSHI MAEHARA | 149 |
| A Ramsey property of planar graphs JAROSLAV NEŠETRIL, JÓZSEF SOLYMOSI, AND PAVEL VALTR | 169 |
| A generalization of quasi-planarity JÁNOS PACH, RADOŠ RADOIČIĆ, AND GÉZA TÓTH | 177 |
| Geometric incidences JÁNOS PACH AND MICHA SHARIR | 185 |
| Large sets must have either a k -edge or a $(k + 2)$ -edge MICHA PERLES AND ROM PINCHASI | 225 |
| Topological graphs with no self-intersecting cycle of length 4 ROM PINCHASI AND RADOŠ RADOIČIĆ | 233 |
| A problem on restricted sumsets IMRE Z. RUZSA | 245 |
| The gap between the crossing numbers and the convex crossing numbers FARHAD SHAHROKHI, ONDREJ SÝKORA, LÁSZLÓ SZÉKELY, AND IMRICH VRŤO | 249 |
| Distinct distances in high dimensional homogeneous sets JÓZSEF SOLYMOSI AND VAN H. VU | 259 |
| The biplanar crossing number of the random graph JOEL H. SPENCER | 269 |
| The unit distance problem on spheres KONRAD SWANEPOEL AND PAVEL VALTR | 273 |
| Short proof for a theorem of Pach, Spencer, and Tóth LÁSZLÓ SZÉKELY | 281 |

Preface

The early development of graph theory was heavily motivated and influenced by topological and geometric themes, such as the Königsberg Bridge Problem, Euler's Polyhedral Formula, or Kuratowski's characterization of planar graphs. When in 1936 Dénes König published his classical *Theorie der endlichen und unendlichen Graphen*, the first book ever written on the subject, he stressed this connection by adding the subtitle *Kombinatorische Topologie der Streckenkomplexe* (Combinatorial Topology of Systems of Segments). He wanted to emphasize that the subject of his investigations was very *concrete*: planar figures consisting of points connected by straight-line segments. However, in the second half of the twentieth century, graph theoretical research took an interesting turn. In the most popular, most rapidly growing areas (the theory of random graphs, Ramsey theory, extremal graph theory, algebraic graph theory, etc.), graphs were considered *abstract* binary relations rather than geometric objects. Many of the powerful techniques developed in these fields have been successfully applied in other areas of mathematics, including geometry.

In the past two decades we have witnessed a renaissance of geometry that has permeated virtually every part of mathematics. Graph theory has been no exception. Topological and geometric methods have proved to be instrumental in the solution of many important problems in graph and hypergraph theory. Meanwhile, a number of newly emerging fields of computer science have served as rich sources of exciting open geometric questions. These include computational geometry, robotics, pattern recognition, computerized tomography, VLSI design, graph drawing, computer graphics, computer-aided design (CAD), and geometric information systems (GIS), to mention only a few. Many of the questions arising were related to systems of segments in the plane or to simplices in higher dimensions. Some of them looked very familiar: they had been asked in different contexts much earlier by Paul Erdős, Paul Turán, Micha Perles, John Conway, and others. For instance, Erdős's notoriously difficult questions on the distribution of distances determined by finite point sets in Euclidean spaces boil down to problems about incidences between points and lines, circles, spheres, and other geometric objects (see the survey of Pach and Sharir in this volume). Similar questions turned out to play an important role in robotics. To solve Turán's Brick Factory Problem [T77], one has to find a drawing of the complete bipartite graph that minimizes the number of crossings between the edges. The "crossing number" of a graph is a central notion in VLSI design: it is closely related to the minimum area of a chip realizing the same network. According to Conway's Thrackle Conjecture [W69], if a graph G can be drawn in the plane so that any pair of its edges cross precisely once (where a common endpoint is counted as a crossing), then G has at most as many

edges as vertices. Perles found a very elegant proof for straight-line drawings, and generalized the question in other directions.

For lack of better understanding of the “Combinatorial Topology of Systems of Segments,” the traditional methods of graph theory have rarely provided satisfactory answers to questions of the above type. One can often associate a graph or hypergraph to the structure under investigation and observe that it satisfies certain (sometimes fairly simple) geometric conditions. Then one can completely forget about geometry, and finish the proof by establishing a theorem on *abstract* graphs with the given properties. A simple prototype of this approach is the following argument of Erdős [E46], which proves that any set P of n points in the plane contains at most $O(n^{3/2})$ point pairs whose distance is one. Connect two elements of P by an edge if they are at unit distance. Since two (unit) circles have at most two points in common, the resulting graph has no $K_{2,3}$ as a subgraph. ($K_{2,3}$ is the complete bipartite graph with 2 and 3 vertices in its classes.) The result now follows from the fact [KST54] that the number of edges of any (abstract) $K_{2,3}$ -free graph with n vertices is at most $O(n^{3/2})$. This last statement is asymptotically sharp, but the geometric result is not. Exploring further geometric properties, for instance, the structure of crossings, it can be shown [SSzT84, Sz97] that the number of unit distance pairs determined by n points is $O(n^{4/3})$. Even this bound is probably very far from being best possible.

A *geometric graph* is a graph drawn by straight-line edges in the plane. It seems that to tackle many problems arising in combinatorial and computational geometry and in the application areas mentioned above, one has to study *geometric* graphs satisfying certain relevant *geometric* conditions. We illustrate this phenomenon by Lovász’ proof [L71] showing that the number of “halving lines” determined by an n -element point set P in general position in the plane is $O(n^{3/2})$. Suppose n is even. A line ℓ connecting two elements $p, q \in P$ is called a halving line if either of the open half-planes bounded by ℓ contains precisely $\frac{n}{2} - 1$ other points of P . Define a geometric graph G on the vertex set P by connecting $p, q \in P$ with a straight-line segment (edge) if and only if they induce a halving line. It is not hard to argue that any line can cross at most n edges of G . On the other hand, every geometric graph on n vertices that satisfies this condition has $O(n^{3/2})$ edges. In fact, much more is true: Ajtai, Chvátal, Newborn, Szemerédi [ACN82] and, independently, Leighton [L83] proved that the number of crossings in any geometric graph with n vertices and $e > 4n$ edges is at least Ce^3/n^2 , for a suitable constant $C > 0$. Thus, if Lovász’ geometric graph G had more than $\gamma n^{3/2}$ edges, then it would have an edge crossing at least $2Ce^2/n^2 > 2C\gamma^2 n > n$ others, provided that $\gamma \geq 1/\sqrt{2C}$. This would contradict the property that every line crosses at most n edges of G . Using a more delicate argument based on the same idea, Dey [D98] showed that the number of halving lines is $O(n^{4/3})$. It is quite likely that this bound can be further improved.

The above examples suggest how to reformulate one of the fundamental questions of *extremal graph theory* for geometric graphs: What is the maximum number of edges that a geometric graph of n vertices can have if it satisfies certain geometric properties? The papers of Pach-Radoičić-Tóth, Székely, and Pinchasi-Radoičić focus on problems of this type. The last paper represents a real breakthrough: it is shown that the number of edges of a geometric graph with n vertices that contains no self-intersecting cycle of length 4 is $O(n^{8/5})$. This result has already found

several applications. Brass generalizes some extremal problems to geometric hypergraphs (e.g., to systems of triangles induced by a point set in convex position). The contributions of Cairns-McIntyre-Nikolayevsky and Perles-Pinchasi are related to Conway’s Thrackle conjecture and to the problem of halving lines, respectively. Lovász-Vesztergombi-Wagner-Welzl discover a beautiful connection between k -sets and crossing numbers. As a consequence, they get surprisingly close to determining the smallest number of crossings that a complete geometric graph with n vertices can have. The papers of Eppstein and Shahrokhi-Sýkora-Székely-Vrto analyze the relationship between other natural graph parameters related to edge crossings.

After the pioneering work of Székely [Sz97] and Solymosi-Tóth [ST01], there have been many important developments related to Erdős’s question about the minimum number of distinct distances determined by n points in the plane [E46]. The paper of Katz-Tardos included in this volume represents the state of the art in this area: the best lower bound presently known for this function. On the other hand, Ruzsa’s contribution indicates the theoretical limits of their approach. Solymosi-Vu make a big step towards an asymptotic solution of the analogous question in higher dimensions. Here they detail their elegant arguments only for “homogeneous” sets, but it is not hard to extend their proof to the general case. A construction of Swanepoel-Valtr improves the best known lower bound on the number of times the same distance can occur among n points on the sphere. The papers of Katona-Mayer-Woyczynski and Jamison address related questions for distance and slope distributions, respectively.

The basic problems of the *theory of random graphs* and *Ramsey theory* can also be rephrased for geometric graphs. Spencer’s paper describes the expected behavior of the biplanar crossing number of random graphs, while the works of Dumitrescu-Radoičić, Kaneko-Kano-Suzuki, Kostochka, and Nešetřil-Solymosi-Valtr address Ramsey-type questions. The last paper proves the following remarkable result that can be regarded as a far-reaching generalization of Fáry’s theorem that states that every planar graph permits a straight-line drawing: For any coloring with a finite number of colors of all segments induced by point pairs in the plane, there is a color such that every planar graph has a straight-line drawing in which all edges have that color. The papers of Alt-Knauer-Rote-Whitesides and Maehara are concerned with the mobility and rigidity of geometric graphs (linkages), respectively. In the work of Arutyunants-Iosevich, some deep and surprising connections are uncovered between distance geometry and measure theory. Roughly speaking, they prove that Falconer’s celebrated conjecture [F85], claiming that every d -dimensional set of Hausdorff dimension at least $d/2$ has positive Lebesgue measure, is almost surely true for random metrics. The paper of Dujmović-Wood applies Lovász’ Local Lemma to construct almost optimally compact crossing-free embeddings of geometric graphs in 3-space with the property that all vertices are mapped to integer points.

Our collection is dedicated to two outstanding mathematicians and lifelong friends: Paul Turán and Paul Erdős. Turán’s Brick Factory Problem was the first genuine optimization problem in graph drawing. The systematic study of extremal problems for geometric graphs was initiated many years ago by P. Erdős, M. Perles, and his student, Y. Kupitz [K79], and by S. Avital and H. Hanani [AH66].

The present volume is *neither* a conference proceedings (i.e., quasi-random collection of articles), *nor* a systematic review of the field. It is a careful selection of 25

invited and thoroughly refereed papers, reporting about exciting recent discoveries on the way “Towards a Theory of Geometric Graphs.”

János Pach
Berkeley, October 2003

References

- [ACN82] M. Ajtai, V. Chvátal, M. Newborn, and E. Szemerédi: Crossing-free subgraphs, *Annals of Discrete Mathematics* **12** (1982) 9–12.
- [AH66] S. Avital and H. Hanani: Graphs (in Hebrew), *Gilyonot Lematematika* **3** (1966) 2–8.
- [D98] T. K. Dey: Improved bounds on planar k -sets and related problems, *Discrete and Computational Geometry* **19** (1998), 373–382.
- [E46] P. Erdős: On sets of distances of n points, *American Mathematical Monthly* **53** (1946), 248–250.
- [F85] K. J. Falconer: On the Hausdorff dimension of distance sets, *Mathematika* **32** (1985), 206–212.
- [GO04] J. E. Goodman and J. O’Rourke: *Handbook of Discrete and Computational Geometry, 2nd ed.*, CRC Press Ser. Discrete Math. Appl., CRC, Boca Raton, FL, 2004, to appear.
- [KST54] T. Kövári, V. T. Sós, and P. Turán: On a problem of K. Zarankiewicz, *Colloquium Mathematicum* **3**, 50–57.
- [K79] Y. Kupitz: *Extremal Problems in Combinatorial Geometry, Aarhus University Lecture Notes Series* **53**, Aarhus University, Denmark, 1979.
- [L71] L. Lovász: On the number of halving lines, *Annales Universitatis Scientiarum Budapest, Eötvös, Sectio Mathematica* **14** (1971), 107–108.
- [L83] T. Leighton: *Complexity Issues in VLSI, Foundations of Computing Series*, MIT Press, Cambridge, MA, 1983.
- [ST01] J. Solymosi and Cs. Tóth: Distinct distances in the plane, *Discrete and Computational Geometry* **25** (2001), 629–634.
- [SSzT84] J. Spencer, E. Szemerédi, and W. T. Trotter: Unit distances in the Euclidean plane, in: *Graph Theory and Combinatorics* (B. Bollobás, ed.), Academic Press, New York, 1984, 293–303.
- [Sz97] L. Székely, Crossing numbers and hard Erdős problems in discrete geometry, *Combinatorics, Probability and Computing* **6** (1997), 353–358.
- [T77] P. Turán: A note of welcome, *J. Graph Theory* **1** (1977), 7–9.
- [W69] D. R. Woodall: Thrackles and deadlock, in: *Combinatorial Mathematics and Its Applications* (D.J.A. Welsh, ed.), Academic Press, London, 335–348.

Titles in This Series

- 343 **N. Sthanumoorthy and Kailash C. Misra, Editors**, Kac-Moody Lie algebras and related topics, 2004
- 342 **János Pach, Editor**, Towards a theory of geometric graphs, 2004
- 341 **Hugo Arizmendi, Carlos Bosch, and Lourdes Palacios, Editors**, Topological algebras and their applications, 2004
- 340 **Rafael del Río and Carlos Villegas-Blas, Editors**, Spectral theory of Schrödinger operators, 2004
- 339 **Peter Kuchment, Editor**, Waves in periodic and random media, 2003
- 338 **Pascal Auscher, Thierry Coulhon, and Alexander Grigor'yan, Editors**, Heat kernels and analysis on manifolds, graphs, and metric spaces, 2003
- 337 **Krishan L. Duggal and Ramesh Sharma, Editors**, Recent advances in Riemannian and Lorentzian geometries, 2003
- 336 **José González-Barrios, Jorge A. León, and Ana Meda, Editors**, Stochastic models, 2003
- 335 **Geoffrey L. Price, B. Mitchell Baker, Palle E.T. Jorgensen, and Paul S. Muhly, Editors**, Advances in quantum dynamics, 2003
- 334 **Ron Goldman and Rimvydas Krasauskas, Editors**, Topics in algebraic geometry and geometric modeling, 2003
- 333 **Giovanni Alessandrini and Gunther Uhlmann, Editors**, Inverse problems: Theory and applications, 2003
- 332 **John Bland, Kang-Tae Kim, and Steven G. Krantz, Editors**, Explorations in complex and Riemannian geometry, 2003
- 331 **Luchezar L. Avramov, Marc Chardin, Marcel Morales, and Claudia Polini, Editors**, Commutative algebra: Interactions with algebraic geometry, 2003
- 330 **S. Y. Cheng, C.-W. Shu, and T. Tang, Editors**, Recent advances in scientific computing and partial differential equations, 2003
- 329 **Zhangxin Chen, Roland Glowinski, and Kaitai Li, Editors**, Current trends in scientific computing, 2003
- 328 **Krzysztof Jarosz, Editor**, Function spaces, 2003
- 327 **Yulia Karpeshina, Günter Stolz, Rudi Weikard, and Yanni Zeng, Editors**, Advances in differential equations and mathematical physics, 2003
- 326 **Kenneth D. T-R McLaughlin and Xin Zhou, Editors**, Recent developments in integrable systems and Riemann-Hilbert problems, 2003
- 325 **Seok-Jin Kang and Kyu-Hwan Lee, Editors**, Combinatorial and geometric representation theory, 2003
- 324 **Caroline Grant Melles, Jean-Paul Brasselet, Gary Kennedy, Kristin Lauter, and Lee McEwan, Editors**, Topics in algebraic and noncommutative geometry, 2003
- 323 **Vadim Olshevsky, Editor**, Fast algorithms for structured matrices: theory and applications, 2003
- 322 **S. Dale Cutkosky, Dan Edidin, Zhenbo Qin, and Qi Zhang, Editors**, Vector bundles and representation theory, 2003
- 321 **Anna Kamińska, Editor**, Trends in Banach spaces and operator theory, 2003
- 320 **William Beckner, Alexander Nagel, Andreas Seeger, and Hart F. Smith, Editors**, Harmonic analysis at Mount Holyoke, 2003
- 319 **W. H. Schikhof, C. Perez-Garcia, and A. Escassut, Editors**, Ultrametric functional analysis, 2003
- 318 **David E. Radford, Fernando J. O. Souza, and David N. Yetter, Editors**, Diagrammatic morphisms and applications, 2003
- 317 **Hui-Hsiung Kuo and Ambar N. Sengupta, Editors**, Finite and infinite dimensional analysis in honor of Leonard Gross, 2003

TITLES IN THIS SERIES

- 316 **O. Cornea, G. Lupton, J. Oprea, and D. Tanré, Editors**, Lusternik-Schnirelmann category and related topics, 2002
- 315 **Theodore Voronov, Editor**, Quantization, Poisson brackets and beyond, 2002
- 314 **A. J. Berrick, Man Chun Leung, and Xingwang Xu, Editors**, Topology and Geometry: Commemorating SISTAG, 2002
- 313 **M. Zuhair Nashed and Otmar Scherzer, Editors**, Inverse problems, image analysis, and medical imaging, 2002
- 312 **Aaron Bertram, James A. Carlson, and Holger Kley, Editors**, Symposium in honor of C. H. Clemens, 2002
- 311 **Clifford J. Earle, William J. Harvey, and Sevín Recillas-Pishmish, Editors**, Complex manifolds and hyperbolic geometry, 2002
- 310 **Alejandro Adem, Jack Morava, and Yongbin Ruan, Editors**, Orbifolds in mathematics and physics, 2002
- 309 **Martin Guest, Reiko Miyaoka, and Yoshihiro Ohnita, Editors**, Integrable systems, topology, and physics, 2002
- 308 **Martin Guest, Reiko Miyaoka, and Yoshihiro Ohnita, Editors**, Differentiable geometry and integrable systems, 2002
- 307 **Ricardo Weder, Pavel Exner, and Benoit Grébert, Editors**, Mathematical results in quantum mechanics, 2002
- 306 **Xiaobing Feng and Tim P. Schulze, Editors**, Recent advances in numerical methods for partial differential equations and applications, 2002
- 305 **Samuel J. Lomonaco, Jr. and Howard E. Brandt, Editors**, Quantum computation and information, 2002
- 304 **Jorge Alberto Calvo, Kenneth C. Millett, and Eric J. Rawdon, Editors**, Physical knots: Knotting, linking, and folding geometric objects in \mathbb{R}^3 , 2002
- 303 **William Cherry and Chung-Chun Yang, Editors**, Value distribution theory and complex dynamics, 2002
- 302 **Yi Zhang, Editor**, Logic and algebra, 2002
- 301 **Jerry Bona, Roy Choudhury, and David Kaup, Editors**, The legacy of the inverse scattering transform in applied mathematics, 2002
- 300 **Sergei Vostokov and Yuri Zarhin, Editors**, Algebraic number theory and algebraic geometry: Papers dedicated to A. N. Parshin on the occasion of his sixtieth birthday, 2002
- 299 **George Kamberov, Peter Norman, Franz Pedit, and Ulrich Pinkall, Editors**, Quaternions, spinors, and surfaces, 2002
- 298 **Robert Gilman, Alexei G. Myasnikov, and Vladimir Shpilrain, Editors**, Computational and statistical group theory, 2002
- 297 **Stephen Berman, Paul Fendley, Yi-Zhi Huang, Kailash Misra, and Brian Parshall, Editors**, Recent developments in infinite-dimensional Lie algebras and conformal field theory, 2002
- 296 **Sean Cleary, Robert Gilman, Alexei G. Myasnikov, and Vladimir Shpilrain, Editors**, Combinatorial and geometric group theory, 2002
- 295 **Zhangxin Chen and Richard E. Ewing, Editors**, Fluid flow and transport in porous media: Mathematical and numerical treatment, 2002
- 294 **Robert Coquereaux, Ariel García, and Roberto Trincherro, Editors**, Quantum symmetries in theoretical physics and mathematics, 2002

For a complete list of titles in this series, visit the
AMS Bookstore at www.ams.org/bookstore/.

The early development of graph theory was heavily motivated and influenced by topological and geometric themes, such as the Königsberg Bridge Problem, Euler's Polyhedral Formula, or Kuratowski's characterization of planar graphs. In 1936, when Dénes König published his classical *Theory of Finite and Infinite Graphs*, the first book ever written on the subject, he stressed this connection by adding the subtitle *Combinatorial Topology of Systems of Segments*. He wanted to emphasize that the subject of his investigations was very *concrete*: planar figures consisting of points connected by straight-line segments. However, in the second half of the twentieth century, graph theoretical research took an interesting turn. In the most popular and most rapidly growing areas (the theory of random graphs, Ramsey theory, extremal graph theory, algebraic graph theory, etc.), graphs were considered as *abstract* binary relations rather than geometric objects. Many of the powerful techniques developed in these fields have been successfully applied in other areas of mathematics. However, the same methods were often incapable of providing satisfactory answers to questions arising in geometric applications.

In the spirit of König, *geometric graph theory* focuses on combinatorial and geometric properties of graphs drawn in the plane by straight-line edges (or more generally, by edges represented by simple Jordan arcs). It is an emerging discipline that abounds in open problems, but it has already yielded some striking results which have proved instrumental in the solution of several basic problems in combinatorial and computational geometry. The present volume is a careful selection of 25 invited and thoroughly refereed papers, reporting about important recent discoveries on the way *Towards a Theory of Geometric Graphs*.

ISBN 0-8218-3484-3



9 780821 834848

CONM/342

AMS on the Web
www.ams.org