

ON A REFINEMENT OF THE NON-ORIENTABLE 4-GENUS OF TORUS KNOTS

JOSHUA M. SABLOFF

(Communicated by Shelly Harvey)

ABSTRACT. In formulating a non-orientable analogue of the Milnor Conjecture on the 4-genus of torus knots, Batson [Math. Res. Lett. 21 (2014), pp. 423–436] developed an elegant construction that produces a smooth non-orientable spanning surface in B^4 for a given torus knot in S^3 . While Lobb [Math. Res. Lett. 26 (2019), pp. 1789] showed that Batson’s surfaces do not always minimize the non-orientable 4-genus, we prove that they do minimize among surfaces that share their normal Euler number. We also determine the possible pairs of normal Euler number and first Betti number for non-orientable surfaces whose boundary lies in a class of torus knots for which Batson’s surfaces are non-orientable 4-genus minimizers.

1. INTRODUCTION

The orientable 4-genus of a knot in S^3 and its significant role in low-dimensional topology have been extensively studied, but comparatively little is known about its non-orientable counterpart. For example, an important milestone in the study of the smooth orientable 4-genus was Kronheimer and Mrowka’s solution of the Milnor Conjecture, which showed that the 3- and 4-genera of torus knots coincide [11, 12]; see also [18]. On the other hand, the non-orientable 4-genus of torus knots is still not known in general, though the goal of this paper is to increase our understanding.

To specify the problem more precisely, say that, for a knot $K \subset S^3$, a properly embedded smooth (non-orientable) surface $F \subset B^4$ with $\partial F = K$ is a **(non-orientable) filling** of K . The **non-orientable genus** of K is

$$\gamma_4(K) = \min\{b_1(F) : F \text{ is a non-orientable filling of } K\}.$$

Lower bounds on the non-orientable 4-genus have proven to be more subtle to construct than those on the orientable 4-genus. In this paper, we will rely on lower bounds derived from the signature [8] and on the ν invariant from knot Floer homology [17]; see Section 3 for details. See [6, 16, 19, 20] for other efforts based on classical tools and [1–7] for more recent developments based on the Heegaard Floer package, gauge theory, or Khovanov homology.

To approach the question of the non-orientable 4-genus of torus knots, Batson [3] proposed a construction of minimal-genus fillings using **pinch moves**, which, briefly, consist of attaching a non-orientable band between two adjacent strands of a torus knot lying on a standard torus. The result of a pinch move is another torus knot, which, as we shall detail in Section 2.1, is strictly simpler. Thus,

Received by the editors July 4, 2022, and, in revised form, March 7, 2023.

2020 *Mathematics Subject Classification.* Primary 57K10.

In particular, this material is based upon work supported by the Institute for Advanced Study.

©2023 by the author(s) under Creative Commons
Attribution-NonCommercial-NoDerivatives 4.0 License (CC BY NC ND 4.0)

repeatedly pinching a torus knot $T(p, q)$ yields an unknot, and attaching the non-orientable bands to a disk bounded by the unknot yields a **pinch surface** $F(p, q)$. Batson conjectured that pinch surfaces realize $\gamma_4(T(p, q))$. Jabuka and Van Cott [10, Corollary 1.12] showed that Batson’s conjecture holds for the infinite family of torus knots for which the ν bound is sharp (cf. Corollary 3.5). We shall term such knots **JVC knots**; see Definition 3.7 for the precise characterization. Even so, Lobb [13] showed that, for $T(4, 9)$, the pinch surface does not realize the non-orientable 4-genus; see [4, 14] for further counterexamples.

To explore Batson’s conjecture and pinch surfaces more deeply, recall that, in contrast to the orientable case, a non-orientable filling F has an interesting additional invariant: the Euler class of its normal bundle, termed the **normal Euler number** $e(F)$. The normal Euler number may be used to refine the non-orientable 4-genus by defining, for any $e \in 2\mathbb{Z}$,

$$\gamma_4^e(K) = \min\{b_1(F) : F \text{ is a non-orientable filling of } K \text{ with } e(F) = e\}.$$

The main result of this paper shows that pinch surfaces do, indeed, minimize genus among fillings that share their normal Euler number.

Theorem 1.1 (Refined Batson Conjecture). *The pinch surface $F(p, q)$ realizes the minimal first Betti number among all non-orientable fillings F of $T(p, q)$ with $e(F) = e(F(p, q))$, i.e. $\gamma_4^e(T(p, q)) = b_1(F(p, q))$.*

To give the theorem above more context, we turn to Allen’s work on the “geography” of non-orientable fillings [1]. For a given knot $K \subset S^3$, Allen asked for the set $R(K)$ of pairs $(e, b) \in \mathbb{Z}^2$ realized by the normal Euler number and non-orientable genus, respectively, of a filling F of K . To state the second main theorem of the paper, we introduce the **wedge** $W_{(E, B)} \subset \mathbb{Z}^2$, which is defined by

$$W_{(E, B)} = \left\{ (e, b) \in 2\mathbb{Z} \times \mathbb{Z} : \begin{aligned} &\frac{1}{2}|e - E| \leq b - B, \\ &e \equiv 2b \pmod{4} \end{aligned} \right\}.$$

For a non-orientable filling F , we use the shorthand $W_F = W_{(e(F), b_1(F))}$.

Theorem 1.2. *For any JVC knot $T(p, q)$, the realizable set $R(T(p, q))$ coincides with $W_{F(p, q)}$.*

The theorem is illustrated in Figure 1. As a corollary, JVC knots satisfy Allen’s conjecture that torus knots realize their non-orientable 4-genus at a *unique* normal Euler number [1, Conjecture 1.6].

The remainder of the paper is organized as follows. In Section 2, we recall the construction of pinch surfaces and compute their normal Euler numbers. We review several lower bounds on the non-orientable genus and define the class of JVC torus knots in Section 3. Finally, in Sections 4 and 5, we prove Theorems 1.1 and 1.2, respectively.

2. PINCH SURFACES

2.1. Construction of pinch surfaces. As described in the introduction, a pinch surface $F(p, q)$ for a torus knot $T(p, q)$ is, roughly speaking, the result of attaching a specific sequence of non-orientable bands to $T(p, q)$ to produce an unknot, and then capping the unknot off with a disk. We will make this construction more




FIGURE 1. The set of realizable pairs $(e(F), b_1(F))$ for fillings F of a JVC knot $T(p, q)$ lie in the shaded wedge $W_{F(p,q)}$. The $T(3, 4)$ knot is illustrated here.

precise using ideas from [3] and [10], adapting notation to better dovetail with later arguments.

We start by defining a non-orientable band cobordism. An embedding h of the square $S = [0, 1]^2$ into S^3 is a **non-orientable band** for a knot K if $K \cap h(S) = h([0, 1] \times \{0, 1\})$, the orientation of K agrees with that of $h([0, 1] \times \{0\})$, and the orientation of K disagrees with that of $h([0, 1] \times \{1\})$ (or vice versa). A **non-orientable band move** on K along h is the result of the surgery that replaces the arcs $K \cap h(S)$ with the image of $\{0, 1\} \times [0, 1]$; we denote the result by $K\#h$. A **non-orientable band cobordism** for K is a properly embedded surface in $S^3 \times [0, 1]$ with boundary given by $K \subset S^3 \times \{1\}$ and $K\#h \subset S^3 \times \{0\}$ constructed by attaching a 1-handle to $K \times \{1\}$ along $K \cap h$ with framing specified by h . The direction of the cobordism has been chosen so that successive non-orientable band moves produce a surface in B^4 .

Next, we specialize to torus knots $T(p, q)$, which we understand to lie on a standard torus \mathbb{T} in S^3 . For the rest of the paper, we make the following assumption about the indices p and q :

Assumption. Assume that p and q are positive and relatively prime, and that q is odd with $p > q$ if p is also odd, reversing the roles of p and q if this were not the case.

We visualize $T(p, q)$ as a collection of parallel lines with slope p/q on a unit square with opposite sides identified. Up to isotopy, there is a unique choice of non-orientable band $h \subset \mathbb{T}$ for $T(p, q)$. We call a non-orientable band move on $T(p, q)$ along h a **pinch move**; see Figure 2. Note that $T(p, q)\#h$ also lies on \mathbb{T} , and hence is, itself, a torus knot $T(r, s)$. We use the term **pinch cobordism** for the cobordism between $T(r, s) \times \{0\}$ and $T(p, q) \times \{1\}$ resulting from a concatenation of the trace of a “straightening” isotopy from $T(r, s)$ to $T(p, q)\#h$ and the non-orientable band cobordism coming from the pinch move.

Lemma 2.1 (Lemma 2.1 of [10]). *The result of applying a pinch move to $T(p, q)$ is the torus knot $T(p - 2t, q - 2u)$, where t and u are the integers uniquely determined*




FIGURE 2. A non-orientable band move applied to $T(p, q)$ yields $T(r, s)$.

by the requirements that

$$\begin{aligned}
 t &\equiv -q^{-1} \pmod p \quad \text{and} \quad 0 \leq t \leq p - 1, \\
 u &\equiv p^{-1} \pmod q \quad \text{and} \quad 0 \leq u \leq q - 1.
 \end{aligned}$$

The **sign** of a pinch move is $\epsilon = \text{Sign}(q - 2u)$; by [10, Lemma 2.3], the signs of $p - 2t$ and $q - 2u$ agree when both quantities are nonzero. If the sign of a pinch move is negative, then replace $T(r, s)$ by $T(-r, -s)$ to maintain the assumption on the positivity of the indices. Denote a pinch move with sign ϵ by $T(p, q) \xrightarrow{\epsilon} T(r, s)$. Lemma 2.1 implies that $0 \leq |r| < p$ and $0 \leq |s| < q$. Hence, $T(p, q)$ must reduce to the unknot U after some n pinch moves, which we keep track of using the following notation:

$$(2.1) \quad T(p, q) = T(p_n, q_n) \xrightarrow{\epsilon_n} T(p_{n-1}, q_{n-1}) \xrightarrow{\epsilon_{n-1}} \dots \xrightarrow{\epsilon_1} T(p_0, q_0) = U.$$

By [10, Theorem 2.7], we have $q_0 = 1$.

Finally, the **pinch surface** $F(p, q) \subset B^4$ of $T(p, q)$ is the non-orientable filling of $T(p, q)$ constructed from concatenating the minimal number of pinch cobordisms necessary to transform $T(p, q)$ into an unknot $T(p_0, 1)$, and then gluing a disk to the unknot $T(p_0, 1)$.

2.2. Normal Euler numbers of pinch surfaces. A key ingredient in understanding the refined Batson Conjecture is a computation of the normal Euler number of a pinch surface. We begin by recalling a definition of the normal Euler number of a closed surface in a 4-manifold, then proceed to adjust it to a properly embedded surface. See [8] as well as [3, 17] for further discussion. The actual computation of the normal Euler number of a pinch surface, which will be accomplished in Lemma 2.3, will use Lemma 2.2 rather than the details of the definition.

As a first step, let F be a smooth closed surface in an oriented 4-manifold M ; the manifold M will be either B^4 or $S^3 \times [0, 1]$ in this paper. To concretely compute the Euler number of the normal bundle to F , let F' be a small transverse pushoff of F , i.e. a section of the normal bundle. At each point $x \in F \cap F'$, find compatible orientations of $T_x F$ and $T_x F'$, and use those orientations and an ambient orientation of M to assign a sign to each such x . The **normal Euler number** $e(F)$ is defined to be the sum of the signs of the intersection points $x \in F \cap F'$.

We next extend this definition to a filling $F \subset B^4$ of a knot $K \subset S^3$. In fact, it will prove useful to go further and define the normal Euler number for a non-orientable cobordism F in $S^3 \times [0, 1]$ between knots $K_0 \subset S^3 \times \{0\}$ and $K_1 \subset S^3 \times \{1\}$; a filling of K may be regarded as a cobordism from $K_0 = \emptyset$ to $K_1 = K$. To define the normal Euler number in this setting, we cap off F by

Seifert surfaces of the K_i in $S^3 \times \{i\}$ to form the closed surface \hat{F} , and we take the **normal Euler number** $e(F)$ to be the normal Euler number of the closed surface \hat{F} . This definition is equivalent to constructing a transverse pushoff F' of F so that the push-offs K'_i at the ends realize the Seifert framings of K_i , and then summing signed intersections of F and F' .

Some facts about the normal Euler number $e(F)$ to keep in mind are that it vanishes for orientable surfaces, that it is always even, that it satisfies

$$(2.2) \quad e(F) \equiv 2b_1(F) \pmod{4}$$

(see [15]), and that it is additive under concatenation of cobordisms. This last fact, combined with the computation of the normal Euler number of a band cobordism in the lemma below, may be used to compute the normal Euler number of a surface constructed via a sequence of band moves.

Lemma 2.2 (Lemma 4.2 of [17]). *Suppose h is a non-orientable band for a knot K . Choose a section s of the normal bundle of h and framings λ of K and λ_h of $K\#h$ that agree with s along $K \cap h(S)$. The normal Euler number of the non-orientable band cobordism F is then computed by*

$$e(F) = \text{lk}(K\#h, \lambda_h) - \text{lk}(K, \lambda).$$

Lemma 2.3, which computes the normal Euler number of a pinch surface, is the main technical contribution in this paper.

Lemma 2.3. *If $T(p, q)$ is transformed to $T(p_0, 1)$ in the construction of the pinch surface $F(p, q)$, then*

$$e(F(p, q)) = p_0 - pq.$$

Proof. Suppose that the pinch surface from $T(p, q)$ to $T(p_0, 1)$ is constructed from n pinches as in Equation (2.1). We claim that the normal Euler number of the non-orientable band cobordism F_k for $T(p_k, q_k) \rightarrow T(p_{k-1}, q_{k-1})$ is $p_{k-1}q_{k-1} - p_kq_k$. The lemma then follows from the claim, the additivity of the normal Euler number, and the fact that the normal Euler number of the final capping disk vanishes.

To prove the claim, we use Lemma 2.2 with $K = T(p_k, q_k)$ and $K\#h = T(p_{k-1}, q_{k-1})$. Let the section s and the framings λ and λ_h all come from a normal vector field to \mathbb{T} ; this choice clearly satisfies the hypothesis of the lemma. Isotope the knot λ to another copy of $T(p_k, q_k)$ on \mathbb{T} by rotating by $\frac{\pi}{2}$ in the direction normal to K . Thus, we see that $\text{lk}(K, \lambda)$ is the same as the linking number between the two components of the torus link $T(2p_k, 2q_k)$. It is straightforward to check that this linking number is p_kq_k . A similar computation for $K\#h$ and λ_h holds, and hence Lemma 2.2 implies that

$$e(F_k) = \text{lk}(K\#h, \lambda_h) - \text{lk}(K, \lambda) = p_{k-1}q_{k-1} - p_kq_k,$$

as required. □

3. LOWER BOUNDS ON THE NON-ORIENTABLE GENUS

Of particular importance for this paper are two lower bounds that involve both the non-orientable 4-genus and the normal Euler number. The first bound relies on the signature.

Theorem 3.1 ([8]). *For any filling F of a knot K , we have $|\sigma(K) - e(F)/2| \leq b_1(F)$. In particular, we have*

$$\left| \sigma(K) - \frac{e}{2} \right| \leq \gamma_4^e(K).$$

See [1, §2.3] or [20, Theorem 1.1] for further discussion of this theorem.

The second bound uses the v invariant, which is derived from an unoriented version of knot Floer homology.

Theorem 3.2 (Theorem 1.1 of [17]). *For any filling F of a knot K , we have $|2v(K) - e(F)/2| \leq b_1(F)$. In particular, we have*

$$\left| 2v(K) - \frac{e}{2} \right| \leq \gamma_4^e(K).$$

To apply Theorem 3.2 to the proof of Theorem 1.1, we will use Jabuka and Van Cott’s computation of the v invariant for a torus knot in terms of its pinching sequence (2.1).

Theorem 3.3 (Theorem 1.14 of [10]). *If $T(p, q)$ is transformed to $T(p_0, 1)$ in the construction of the pinch surface, then*

$$2v(T(p, q)) = b_1(F(p, q)) + \frac{1}{2}(p_0 - pq).$$

Combining Theorem 3.3 with Lemma 2.3 yields a relationship between v of a torus knot and the normal Euler number of its pinch surface.

Corollary 3.4. *Under the conditions of Theorem 3.3, we obtain:*

- (1) $2v(T(p, q)) = b_1(F(p, q)) + \frac{1}{2}e(F(p, q))$ and hence
- (2) $2v(T(p, q)) \geq \frac{1}{2}e(F(p, q))$.

Returning to the original Theorems 3.1 and 3.2, we may combine them obtain a bound on the non-orientable 4-genus independent of the normal Euler number.

Corollary 3.5 (Theorem 1.2 of [17]). *For any knot $K \subset S^3$, we have*

$$\left| v(K) - \frac{\sigma(K)}{2} \right| \leq \gamma_4(K).$$

Jabuka and Van Cott were able to determine the sign of the quantity on the left of the inequality above for torus knots.

Proposition 3.6 (Corollary 1.10 of [10]). *For any torus knot $T(p, q)$, we have $2v(T(p, q)) \geq \sigma(T(p, q))$.*

Jabuka and Van Cott also completely characterized which torus knots have a pinch surface that realizes the bound in Corollary 3.5. As the precise identification of the conditions on the pairs (p, q) that determine these knots is not necessary for this paper, we leave the technicalities for Theorem 1.11 in [10]. Nevertheless, we make Definition 3.7.

Definition 3.7. A torus knot $T(p, q)$ is a **JVC knot** if its pinch surface realizes the bound in Corollary 3.5, i.e. if

$$2v(T(p, q)) - \sigma(T(p, q)) = 2b_1(F(p, q)).$$

Example 3.8. As noted in Example 1.13 of [10], the knots $T(2k, 2k - 1)$ are JVC knots.




FIGURE 3. The wedges W_σ , W_v , and W_F for (left) an arbitrary knot K and filling F and (right) for a JVC knot $T(p, q)$ and $F = F(p, q)$

Example 3.9. The computations in the proof of Theorem 1.3 in [1, §5] show that $T(3, k)$ is a JVC knot if and only if k is equal to 4 or 5 modulo 6.

Example 3.10. Example 5.3 of [9], together with Theorem 1.11(b) of [10], shows that $T(km + 1, m)$ with $k \geq 1$, $m > 1$, and both m and k odd are JVC knots.

4. PROOF OF THE REFINED BATSON CONJECTURE

We now have all of the ingredients needed for the proof of Theorem 1.1. Given a torus knot $T(p, q)$ and its pinch surface $F(p, q)$, the theorem follows from the following short computation, in which we use the notation $e = e(F(p, q))$ for ease of reading:

$$\begin{aligned}
 b_1(F(p, q)) &\geq \gamma_4^e(T(p, q)) \\
 &\geq 2v(T(p, q)) - \frac{1}{2}e && \text{by Theorem 3.2} \\
 &= b_1(F(p, q)) && \text{by Corollary 3.4(1).}
 \end{aligned}$$

Hence, we obtain $\gamma_4^e(T(p, q)) = b_1(F(p, q))$, as required.

5. THE GEOGRAPHY OF THE JVC KNOTS

In this final section, we contextualize Theorem 1.1 using Allen’s geography question. In particular, we prove Theorem 1.2, which completely determines the geography of the JVC knots.

We begin by defining, for a given knot K , two special wedges, which we illustrate in Figure 3 (see also Figures 11 and 12 in [1]):

- (1) Let W_σ be the wedge $W_{(2\sigma(K), 0)}$, which contains all pairs (e, b) allowed by Theorem 3.1.
- (2) Let W_v be the wedge $W_{(4v(K), 0)}$, which contains all pairs (e, b) allowed by Theorem 3.2.

Thus, Theorems 3.1 and 3.2 yield the following:

Lemma 5.1. *For any knot K , $R(K) \subset W_\sigma \cap W_v$.*

Next, we formalize a fact hinted at in the introduction, namely that every point in the wedge W_F associated to a filling F is realizable.

Lemma 5.2. *If F is a non-orientable filling of a knot K , then $W_F \subset R(K)$.*




FIGURE 4. Concatenating with the non-orientable band cobordism pictured results in a new filling of a given knot with b_1 increased by 1 and e changed by 2. To change e by -2 , switch the crossing.

Proof. We may modify the filling F to a new filling F' with $b_1(F') = b_1(F) + 1$ and $e(F') = e(F) \pm 2$, either by taking a connect sum with a projective plane with normal Euler number ± 2 as in [1, §3.1] or by applying Lemma 2.2 as in Figure 4. Repeating this process appropriately yields fillings for all $(e, b) \in W_F$. \square

We now proceed to prove Theorem 1.2. For readability, we denote the torus knot $T(p, q)$ by K and the pinch surface for $T(p, q)$ by F . The proof is visualized in Figure 3(b), where we see that the pinch surface F lies at the intersection of the wedges W_σ and W_v .

Lemmas 5.1 and 5.2 imply that $W_F \subset R(K) \subset W_\sigma \cap W_v$. Thus, it suffices to show that $W_\sigma \cap W_v \subset W_F$. Suppose that $(e, b) \in W_\sigma \cap W_v$. To show that $(e, b) \in W_F$, we need to prove that $\frac{1}{2}|e - e(F)| \leq b - b_1(F)$ (the parity properties required by the wedge are automatically satisfied).

First, rearrange Corollary 3.4(1) to yield

$$(5.1) \quad e(F) = 4v(K) - 2b_1(F).$$

Second, combine the sharpness of Corollary 3.5 with Proposition 3.6 to obtain the relation

$$(5.2) \quad 4v(K) = 2\sigma(K) + 4b_1(F).$$

Plugging Equation (5.2) into Equation (5.1) produces

$$(5.3) \quad e(F) = 2\sigma(K) + 2b_1(F).$$

Returning to the goal of proving that $\frac{1}{2}|e - e(F)| \leq b - b_1(F)$, we may compute that


$$\begin{aligned} e(F) - e &= 4v(K) - e - 2b_1(F) && \text{by (5.1)} \\ &\leq 2b - 2b_1(F) && \text{since } (e, b) \in W_v \end{aligned}$$

and that

$$\begin{aligned} e(F) - e &= 2\sigma(K) - e + 2b_1(F) && \text{by (5.3)} \\ &\geq -2b + 2b_1(F) && \text{since } (e, b) \in W_\sigma. \end{aligned}$$

This completes the proof of Theorem 1.2.

Remark 5.3. The proof of Theorem 1.2 potentially applies more generally than to JVC knots, as what we really needed was the sharpness of the bounds in Theorem 3.2 and Corollary 3.5 at a given filling F of K , along with a guarantee that $\frac{\sigma(K)}{2}$

FIGURE 5. Two realizable (e, b) pairs for $T(4, 9)$

and $\frac{e(F)}{4}$ lie on the same side of $v(K)$, as in Proposition 3.6 and Corollary 3.4(2), respectively.

Remark 5.4. The torus knot $T(4, 9)$ investigated in [13] presents a different picture than the JVC knots analyzed above. Instead of intersecting in the wedge $W_{F(4,9)}$, the wedges W_σ and W_v coincide, and hence give less information than in the JVC knot case. In fact, it is straightforward, if delicate, to calculate that the pinch surface has $e(F(4, 9)) = -36$ and $b_1(F(4, 9)) = 2$, while Lobb's minimizing surface F_{min} has $e(F_{min}) = -34$ and $b_1(F_{min}) = 1$. While we already knew from Theorem 1.1 and Corollary 3.4(2) that the pinch surface $F(4, 9)$ lies on the lower left boundary of the wedge W_v , we see that F_{min} does as well; see Figure 5.

ACKNOWLEDGMENTS

The investigations leading to the main result grew out of discussions with Braeden Reinoso about when non-orientable Lagrangian fillings of Legendrian knots realize γ_4^e . The author thanks Slaven Jabuka and Cornelia Van Cott for encouragement and generative conversations that helped to contextualize the content of the paper, and the referee for comments that improved the exposition. The author further thanks the Institute for Advanced Study for hosting him during the preparation of the paper.

REFERENCES

- [1] Samantha Allen, *Nonorientable surfaces bounded by knots: a geography problem*, Preprint, [arXiv:2007.14332](https://arxiv.org/abs/2007.14332), 2020.
- [2] William Ballinger, *Concordance invariants from the $e(-1)$ spectral sequence on Khovanov homology*, Preprint, [arXiv:2004.10807](https://arxiv.org/abs/2004.10807), 2020.
- [3] Joshua Batson, *Nonorientable slice genus can be arbitrarily large*, *Math. Res. Lett.* **21** (2014), no. 3, 423–436, DOI 10.4310/MRL.2014.v21.n3.a1. MR3272020
- [4] Fraser Binns, Sungkyung Kang, Jonathan Simone, and Paula Truöl, *On the nonorientable four-ball genus of torus knots*, Preprint, [arXiv:2109.09187](https://arxiv.org/abs/2109.09187), 2021.
- [5] Aliakbar Daemi and Christopher Scaduto, *Chern-Simons functional, singular instantons, and the four-dimensional clasp number*, Preprint, [arXiv:2007.13160](https://arxiv.org/abs/2007.13160), 2020.
- [6] Patrick M. Gilmer and Charles Livingston, *The nonorientable 4-genus of knots*, *J. Lond. Math. Soc. (2)* **84** (2011), no. 3, 559–577, DOI 10.1112/jlms/jdr024. MR2855790
- [7] Marco Golla and Marco Marengon, *Correction terms and the nonorientable slice genus*, *Michigan Math. J.* **67** (2018), no. 1, 59–82, DOI 10.1307/mmj/1511924604. MR3770853
- [8] C. McA. Gordon and R. A. Litherland, *On the signature of a link*, *Invent. Math.* **47** (1978), no. 1, 53–69, DOI 10.1007/BF01609479. MR500905

- [9] Stanislav Jabuka and Cornelia A. Van Cott, *Comparing nonorientable three genus and nonorientable four genus of torus knots*, J. Knot Theory Ramifications **29** (2020), no. 3, 2050013, 15, DOI 10.1142/S0218216520500133. MR4101607
- [10] Stanislav Jabuka and Cornelia A. Van Cott, *On a nonorientable analogue of the Milnor conjecture*, Algebr. Geom. Topol. **21** (2021), no. 5, 2571–2625, DOI 10.2140/agt.2021.21.2571. MR4334520
- [11] P. B. Kronheimer and T. S. Mrowka, *Gauge theory for embedded surfaces. I*, Topology **32** (1993), no. 4, 773–826, DOI 10.1016/0040-9383(93)90051-V. MR1241873
- [12] P. B. Kronheimer and T. S. Mrowka, *Gauge theory for embedded surfaces. II*, Topology **34** (1995), no. 1, 37–97, DOI 10.1016/0040-9383(94)E0003-3. MR1308489
- [13] Andrew Lobb, *A counterexample to Batson’s conjecture*, Math. Res. Lett. **26** (2019), no. 6, 1789, DOI 10.4310/MRL.2019.v26.n6.a8. MR4078695
- [14] Vincent Longo, *An infinite family of counterexamples to Batson’s conjecture*, Preprint, [arXiv:2011.00122](https://arxiv.org/abs/2011.00122), 2020.
- [15] W. S. Massey, *Proof of a conjecture of Whitney*, Pacific J. Math. **31** (1969), 143–156. MR250331
- [16] Hitoshi Murakami and Akira Yasuhara, *Four-genus and four-dimensional clasp number of a knot*, Proc. Amer. Math. Soc. **128** (2000), no. 12, 3693–3699, DOI 10.1090/S0002-9939-00-05461-7. MR1690998
- [17] Peter S. Ozsváth, András I. Stipsicz, and Zoltán Szabó, *Unoriented knot Floer homology and the unoriented four-ball genus*, Int. Math. Res. Not. IMRN **17** (2017), 5137–5181, DOI 10.1093/imrn/rnw143. MR3694597
- [18] Jacob Rasmussen, *Khovanov homology and the slice genus*, Invent. Math. **182** (2010), no. 2, 419–447, DOI 10.1007/s00222-010-0275-6. MR2729272
- [19] O. Ja. Viro, *Positioning in codimension 2, and the boundary* (Russian), Uspehi Mat. Nauk **30** (1975), no. 1(181), 231–232. MR0420641
- [20] Akira Yasuhara, *Connecting lemmas and representing homology classes of simply connected 4-manifolds*, Tokyo J. Math. **19** (1996), no. 1, 245–261, DOI 10.3836/tjm/1270043232. MR1391941

DEPARTMENT OF MATHEMATICS AND STATISTICS, HAVERFORD COLLEGE, HAVERFORD, PENNSYLVANIA 19041

Email address: jsabloff@haverford.edu

URL: <https://jsabloff.sites.haverford.edu/>