

A CORRECTION TO "ON THE AVERAGE NUMBER OF REAL ROOTS OF A RANDOM ALGEBRAIC EQUATION"¹

M. KAC

Dr. R. P. Boas has kindly called my attention to the fact that the proofs of the results of Littlewood and Offord² have been published in the Proc. Cambridge Philos. Soc. vol. 35 (1939) pp. 133–148. It should also be mentioned that a similar problem was studied by entirely different methods by S. O. Rice in his paper *The distribution of the maxima of a random curve*, Amer. J. Math. vol. 61 (1939) pp. 409–416. Finally, I take this opportunity to state that under the conditions considered in my note the probability that equation (1) has a multiple root is zero. In fact, this probability is the probability that the discriminant vanish; this probability is clearly equal to zero since the discriminant is an analytic function of the coefficients and the measure considered is nonsingular.

CORNELL UNIVERSITY

Received by the editors June 9, 1943.

¹ Bull. Amer. Math. Soc. vol. 49 (1943) pp. 314–320.

² See the footnote on page 314 of my note.

A CORRECTION TO "A LINEAR TRANSFORMATION WHOSE VARIABLES AND COEFFICIENTS ARE SETS OF POINTS"

S. T. SANDERS, JR.

In my paper *A linear transformation whose variables and coefficients are sets of points*, published in Bull. Amer. Math. Soc. vol. 48 (1942), the fourth line, page 442, should read: "there is a subscript $k = k'$ such that $P_{k'}$ involves only factors of P_1 ."

SOUTHWESTERN LOUISIANA INSTITUTE

Received by the editors April 14, 1943.