

ON ISOMETRIES OF PRODUCT SETS

PAUL J. KELLY

1. **Introduction.** If A is a metric set, and A^2 its cartesian product with itself, it is possible, in terms of the metric A , to metrize A^2 in many different ways. Thus, the problem of determining when A^2 isometric to B^2 implies A isometric to B can be attacked by placing limitations on the sets A and B or on the product metrics. The present paper proves that the isometry is implied when A and B are bounded and linear and the metric in A^2 and B^2 is determined by the use of a modified Minkowski gauge.

Also stated, without proof, are certain conditions under which the isometry of two product sets implies the isometry of the factors. The proofs for these results, given in the author's doctoral thesis,¹ are long and differ only in detail from the one demonstrated.

2. Definitions.

(A) *Metric set.* A is a metric set if to each pair of its element a_i, a_j there corresponds a real, non-negative number $\rho(a_i, a_j)$ which is independent of the order of the elements, zero if and only if $a_i = a_j$, and which satisfies $\rho(a_i, a_j) + \rho(a_j, a_k) \geq \rho(a_i, a_k)$.

(B) *Isometry.* Metric sets A and B are isometric (indicated by $A \equiv B$) if there exists between them a 1-1 correspondence of elements, T , such that $\rho(a_i, a_j) = \rho(T(a_i), T(a_j))$. The metric in A and B need not be the same, but in writing $\rho(a_i, a_j)$ we shall understand that ρ is the metric for the set containing a_i and a_j .

(C) *Linear set.* Linear, here, will mean a set isometric to a subset of the euclidean line.

(D) *Bounded set.* A metric set A is bounded if $\rho(a_i, a_j)$ is equal to or less than some real number R for all couples in A . If there exists a couple a_i, a_j , such that $\rho(a_i, a_j) = R$, the set assumes its bound.

(E) *Modified Minkowski gauge.* In the first quadrant of the cartesian plane let C' be a curve having nonzero intercepts. A curve C consisting of C' together with the segments of the x and y axes which C' intercepts will be called a modified Minkowski gauge if the following properties hold: (1) C is a simply connected, closed, convex curve; (2) if $P_1(x_1, y_1), P_2(x_2, y_2)$ are any two points in the first quadrant and the lines joining the origin to P_1 and P_2 cut C in P'_1 and P'_2 , respectively, then the relations $x_1 \leq x_2$, and $y_1 \leq y_2$ imply OP_1/OP'_1

Received by the editors August 17, 1947.

¹ *On isometric transformations*, University of Wisconsin, 1942.

$\leq OP_2/OP'_2$. It is easily shown, and we shall need to use, the fact that (1) and (2) imply that: (3) when $x_1 < x_2$ and $y_1 < y_2$ then $OP_1/OP'_1 < OP_2/OP'_2$. In addition to (1) and (2) the gauge will be called symmetric if: (4) C is symmetric about the line $y=x$.

(F) *Minkowski metrizations*. Let A , with elements a_i , be any metric set. We use C to metrize A^2 as follows. To a pair of elements (a_i, a_j) , (a_k, a_l) in A^2 there corresponds a metrizing point P in the plane with coordinates $[\rho(a_i, a_k), \rho(a_j, a_l)]$. Let the line determined by O and P cut C in the point P' . Then $\rho[(a_i, a_j), (a_k, a_l)]$ is defined as the ratio OP/OP' . We shall also use the notation $\|P\|$ for OP/OP' . Thus condition (2) under (E) could be stated that when $x_1 \leq x_2$ and $y_1 \leq y_2$ then $\|(x_1, y_1)\| \leq \|(x_2, y_2)\|$.

It can be shown that if C has properties (1) and (2) under (E) then A^2 will satisfy definition (A) of a metric set.² Some condition similar to (2) is essential, for the metric defined this way with an ordinary Minkowski gauge permits examples in which A^2 is not a metric set. Symmetry about $y=x$ is not a necessary but a natural condition to assume in dealing with square sets. Since $\|P\|=1$ if and only if P lies on C , C is often referred to as a unit gauge. Various familiar metrics derive in a natural way from different choices of the gauge. If C' is the arc of a unit circle then $\|(x, y)\| = (x^2 + y^2)^{1/2}$. If C is a unit square, then $\|(x, y)\| = \text{Max}(x, y)$. If C' is the line joining (1, 0) and (0, 1) then $\|(x, y)\| = x + y$.

THEOREM 1. *If A and B are bounded linear sets which assume their bounds and if A^2, B^2 are metrized by the same symmetric gauge C , then $A^2 \equiv B^2$ implies $A \equiv B$.*

PROOF. (1) Let T be the correspondence establishing $A^2 \equiv B^2$. Let M_1 and M_2 be the bounds of A and B respectively. From linearity these bounds are assumed at one and only one pair of elements in each set. Let $\rho(\bar{a}, a^*) = M_1$, $\rho(\bar{b}, b^*) = M_2$.

(2) Let P_1, P_2, P_3, P_4 in A^2 be the points $(\bar{a}, \bar{a}), (a^*, \bar{a}), (a^*, a^*), (\bar{a}, a^*)$ and Q_1, Q_2, Q_3, Q_4 in B^2 have similar coordinates with "a" replaced by "b." Let $T(P_i) = S_i = (u_i, v_i)$, $i=1, 2, 3, 4$.

(3) The set $\{S_i\}$ is in some order the set $\{Q_i\}$, $i=1, 2, 3, 4$. We prove this by showing each u_i and each v_i is \bar{b} or b^* as follows.

(3.1) From the coordinates of P_i and the definition of the metric in A^2 we have $\rho(P_1, P_3) = \rho(P_2, P_4) = \|(M_1, M_1)\|$ and $\rho(Q_1, Q_3) = \rho(Q_2, Q_4) = \|(M_2, M_2)\|$. From property (2) under (E) these must be the largest distances which occur in A^2 and B^2 respectively. Then

² Ibid.

from $A^2 \equiv B^2$, we have $\|(M_1, M_1)\| = \|(M_2, M_2)\|$. Since (M_1, M_1) and (M_2, M_2) both lie on $y=x$ and this line cuts C' in only one point it follows that $M_1 = M_2$. Let $M_1 = M_2 = M$. From symmetry of C the x and y intercepts of C' are equal and for simplicity we may take them as $(M, 0)$ and $(0, M)$. (This could always be done by a radial expansion or contraction of C that would change all distances in A^2 and B^2 by the same multiple.) With this condition it follows from (1) and (2) that $\rho(P_1, P_2) = \rho(P_2, P_3) = \rho(P_3, P_4) = \rho(P_4, P_1) = 1$.

From (3.1) and $A^2 \equiv B^2$ we have

$$(3.2) \quad \rho(S_1, S_2) = \rho(S_2, S_3) = \rho(S_3, S_4) = \rho(S_4, S_1) = 1,$$

and

$$(3.3) \quad \rho(S_1, S_3) = \rho(S_2, S_4) = \|(M, M)\|.$$

Now $\|(M, M)\| \geq 1$ and we treat these two possibilities separately.

(3.4) *Case 1.* $\|(M, M)\| > 1$. Then if $\bar{P}(\bar{x}, \bar{x})$ is the gauge point on $y=x$ (that is, intersection of C' and $y=x$), we have $0 < \bar{x} < M$. From (1) we have

$$(3.5) \quad \rho(u_1, u_3) \leq M, \quad \rho(v_1, v_3) \leq M$$

and

$$(3.6) \quad \rho(u_2, u_4) \leq M, \quad \rho(v_2, v_4) \leq M.$$

From (3.3) and property (3) of the gauge C , one of the equalities in (3.5) and one in (3.6) must hold. To show that all four equalities must hold, suppose:

(3.7) $\rho(u_1, u_3) = M, \rho(v_1, v_3) < M$. Let R be the point $[\rho(u_1, u_3), \rho(v_1, v_3)]$. From $\|\rho(u_1, u_3), \rho(v_1, v_3)\| = \|(M, M)\|$ it follows, using similar triangles, that the gauge point R' on the line joining O and R lies on the line $x = \bar{x}$. Since the points \bar{P} and (M, O) are on C , the chord joining them must lie on or interior to C . Hence, R'' , the intersection of this chord with the radial line through R , must lie on or interior to C , which is impossible since $OR'' > OR'$. This contradiction shows that if $\rho(u_1, u_3) = M$ then $\rho(v_1, v_3) = M$. The four possibilities of one inequality holding in (3.5) and (3.6) lead to a similar contradiction, hence all four equalities hold. But from (1) that implies each u_i and each $v_i, i = 1, 2, 3, 4$, is either \bar{b} or b^* and establishes (3).

(3.8) *Case 2.* $\|(M, M)\| = 1$. Then (M, M) as well as (M, O) and (O, M) are gauge points. From the definition of convexity then there can be no gauge point interior to the square $(O, O), (M, O), (M, M), (O, M)$. But now from (3.2) and (3.3) all six distances in the set $\{T(P_i)\}, i = 1, 2, 3, 4$, are 1, therefore the metrizing point correspond-

ing to any pair of these elements must lie on C , and hence must have either its 1st or 2nd coordinate equal to M . Then either $\rho(u_i, u_j) = M$ or $\rho(v_i, v_j) = M$, $i, j = 1, 2, 3, 4$, $i \neq j$. Now assume $v_1 \neq \bar{b}$ or b^* . Then $\rho(v_1, v_2) \neq M$, $\rho(v_1, v_3) \neq M$, $\rho(v_1, v_4) \neq M$. Therefore, $\rho(u_1, u_2) = \rho(u_1, u_3) = \rho(u_1, u_4) = M$. Take $u_1 = \bar{b}$, then $u_2 = u_3 = u_4 = b^*$. Therefore $\rho(u_2, u_3) = \rho(u_2, u_4) = \rho(u_3, u_4) = 0$. Then $\rho(v_2, v_4) = \rho(v_2, v_3) = \rho(v_3, v_4) = M$. Let $v_2 = \bar{b}$ then $v_3 = v_4 = b^*$, and $\rho(v_3, v_4) = 0$ contradicting $\rho(v_3, v_4) = M$. Therefore v_1 must equal \bar{b} or b^* and by the same argument each u_i and each v_i , $i = 1, 2, 3, 4$, must be \bar{b} or b^* , which establishes (3).

(4) If we think of P_1, P_3 and P_2, P_4 as diagonal pairs in A^2 and Q_1, Q_3 and Q_2, Q_4 as diagonal pairs in B^2 it is clear, since there are six pairs in each set, that T must map at least one nondiagonal pair of the P_i 's onto a nondiagonal pair of the Q_i 's. The symmetry of C makes the argument the same in any case and we may suppose for definiteness that $T(P_1) = Q_1$, $T(P_2) = Q_2$.

We define a mapping, S , of A onto B as follows. To any element a_i in A there corresponds in A^2 the point $P_i(a_i, \bar{a})$. Let $T(P_i) = Q_i = (b_i, b_k)$, that is, label the first coordinate of the transformed point b_i . Define S by $S(a_i) = b_i$, $S(\bar{a}) = \bar{b}$, $S(a^*) = b^*$.

(5) S establishes $A \equiv B$.

(5.1) From the coordinates of P_1, P_i, P_2 we have $\rho(P_1, P_i) = \rho(\bar{a}, a_i)/M$, $\rho(P_i, P_2) = \rho(a_i, a^*)/M$, $\rho(P_1, P_2) = \rho(\bar{a}, a^*)/M$.

(5.2) Linearity in A gives $\rho(\bar{a}, a_i) + \rho(a_i, a^*) = \rho(\bar{a}, a^*)$.

(5.3) (5.1) and (5.2) imply $\rho(P_1, P_i) + \rho(P_i, P_2) = \rho(P_1, P_2)$.

(5.4) From (5.3) and $A^2 \equiv B^2$ we have $\rho(Q_1, Q_i) + \rho(Q_i, Q_2) = \rho(Q_1, Q_2)$.

(5.5) Consider the point $\bar{Q}_i(b_i, \bar{b})$. From the coordinates of Q_1, \bar{Q}_i, Q_2 and linearity in B it follows that $\rho(Q_1, \bar{Q}_i) + \rho(\bar{Q}_i, Q_2) = \rho(Q_1, Q_2)$.

(5.6) (5.4) and (5.5) give $\rho(Q_1, Q_i) + \rho(Q_i, Q_2) = \rho(Q_1, \bar{Q}_i) + \rho(\bar{Q}_i, Q_2)$.

(5.7) The metrizing point corresponding to the couple Q_1, Q_i has its coordinates respectively equal to or greater than those for the point corresponding to the couple Q_1, \bar{Q}_i . Hence, $\rho(Q_1, Q_i) \geq \rho(Q_1, \bar{Q}_i)$. Similarly $\rho(Q_i, Q_2) \geq \rho(\bar{Q}_i, Q_2)$. These relations together with (5.6) imply $\rho(Q_1, Q_i) = \rho(Q_1, \bar{Q}_i)$ and $\rho(Q_i, Q_2) = \rho(\bar{Q}_i, Q_2)$.

(5.8) Since $\rho(Q_1, \bar{Q}_i) = \rho(\bar{b}, b_i)/M$, from (5.1), $A^2 \equiv B^2$ and (5.7) we obtain $\rho(\bar{a}, a_i) = M\rho(P_1, P_i) = M\rho(Q_1, Q_i) = M\rho(Q_1, \bar{Q}_i) = \rho(\bar{b}, b_i)$.

(5.9) Let a_i, a_j be any two elements in A with b_i, b_j their transforms under S . From (5.8) we have $\rho(\bar{a}, a_i) = \rho(\bar{b}, b_i)$ and $\rho(\bar{a}, a_j) = \rho(\bar{b}, b_j)$, and from the linearity of A and B it follows that $\rho(a_i, a_j) = \rho(b_i, b_j)$. This last implies that if $a_i \neq a_j$ then $b_i \neq b_j$ so the mapping cannot be many-to-one. Also S is not a mapping of A onto a part of B

