

ON A PROPERTY OF POSITIVE-DEFINITE FUNCTIONS¹

BY R. G. LAHA

Communicated by S. Bochner, May 20, 1960

In the present note we prove the following theorem:

THEOREM. *Let $\{t_k\}$ ($k = \pm 1, \pm 2, \dots$) be a sequence of real numbers such that $t_k > 0$ and $t_{-k} = -t_k$ for any $k > 0$ and that $t_k \rightarrow 0$ as $k \rightarrow \infty$. Let $f(t)$ be a positive-definite function of the real variable t and let $\psi(z)$ be a function of the complex variable z ($z = t + iv$, t and v both real) which is regular in a circle about the point $z = 0$. Suppose that the function $f(t)$ coincides with $\psi(t)$ for every value of the sequence $\{t_k\}$. Then $f(t)$ coincides with $\psi(t)$ for all real t .*

PROOF. Let the function $\psi(z)$ be regular in the circle $|z| < R$ ($R > 0$) about the point $z = 0$. Then for every real t in the interval $|t| < R$ we have

$$\psi(t) = \sum_{j=0}^{\infty} (\alpha_j + i\beta_j) \frac{t^j}{j!}$$

where α_j and β_j are real numbers. Let $A(t)$ and $B(t)$ denote respectively the real and imaginary parts of $\psi(t)$ so that

$$A(t) = \sum_{j=0}^{\infty} \alpha_j \frac{t^j}{j!} \quad \text{and} \quad B(t) = \sum_{j=0}^{\infty} \beta_j \frac{t^j}{j!}.$$

We now make use of the hermitian property of $f(t)$ and the equation $f(t_k) = \psi(t_k)$ and obtain easily the relations $A(-t_k) = A(t_k)$ and $B(-t_k) = -B(t_k)$ holding for every value of the sequence $\{t_k\}$. Since the point $t = 0$ is the limit point of the sequence of real numbers $\{t_k\}$, we can verify easily that

$$A(t) = \sum_{j=0}^{\infty} \alpha_{2j} \frac{t^{2j}}{(2j)!} \quad \text{and} \quad B(t) = \sum_{j=0}^{\infty} \beta_{2j+1} \frac{t^{2j+1}}{(2j+1)!}$$

for all real t in $|t| < R$.

We next introduce the functions $g(t) = f(t)f(-t)$ and $\theta(z) = \psi(z)\psi(-z)$ ($z = t + iv$, t and v real) and note the following: The function $g(t)$ is a real-valued, even and positive-definite function of the real variable t ; the function $\theta(z)$ is regular in the same circle $|z| < R$ about the point $z = 0$ and for real t

¹ This work was supported by the National Science Foundation through grant NSF-G-4220 and NSF-G-9968.

$$\theta(t) = |\psi(t)|^2 = \{A(t)\}^2 + \{B(t)\}^2 = \sum_{j=0}^{\infty} \gamma_{2j} \frac{t^{2j}}{(2j)!}$$

where γ_{2j} is real and therefore $\theta(t)$ is a real-valued and even function of t . Before proceeding further we first prove the following lemma:

LEMMA. *The function $g(t)$ has finite derivatives of all orders for every real t .*

PROOF OF THE LEMMA. We can represent $g(t)$ in the form $g(t) = \int_{-\infty}^{\infty} \cos tx dG(x)$ where $G(x)$ is a nondecreasing function of bounded variation. We show first that the function $g(t)$ can be differentiated twice for any real t . We note that the relation $g(t_k) = \theta(t_k)$ holds for every value of the sequence $\{t_k\}$, $t_k \rightarrow 0$ so that we must have $g(0) = \theta(0)$ and therefore $\gamma_0 = \int_{-\infty}^{\infty} dG(x) > 0$. Thus for every value of the sequence $\{t_k\}$ $t_k \rightarrow 0$ we have

$$\frac{g(t_k) - g(0)}{t_k^2} = \frac{\theta(t_k) - \theta(0)}{t_k^2} = \frac{\gamma_2}{2} + O(t_k^2)$$

so that

$$\lim_{k \rightarrow \infty} \frac{g(t_k) - g(0)}{t_k^2} = \frac{\gamma_2}{2}$$

exists and is finite. But we note that

$$\frac{g(t_k) - g(0)}{t_k^2} = -2 \int_{-\infty}^{\infty} \frac{\sin^2 \frac{t_k x}{2}}{t_k^2} dG(x)$$

and therefore

$$\int_{-\infty}^{\infty} \frac{\sin^2 \frac{t_k x}{2}}{t_k^2} dG(x)$$

tends to a finite limit as $t_k \rightarrow 0$. Then we use Fatou's Lemma and deduce immediately that $\int_{-\infty}^{\infty} x^2 dG(x) < \infty$, that is, $|g''(0)| < \infty$. But it is well known that the existence of the $2p$ th derivative of a positive-definite function at the origin $t=0$ ensures that it is differentiable $2p$ times for any real t . Therefore $g''(t)$ exists and is finite for any real t . We now prove the lemma by induction. We suppose that the function $g(t)$ has finite derivatives up to an even order $2n$ and then show that the derivative of order $2n+2$ will also exist. We note that

the function $g(t) - \theta(t)$ is real-valued, even and can be differentiated $2n$ times and further vanishes for every value of the sequence $\{t_k\}$, $t_k \rightarrow 0$. We now apply the Theorem of Rolle: *Between two zeros of a differentiable function there exists at least one zero of its derivative.* Therefore, it follows that the function $g'(t) - \theta'(t)$ vanishes for every value of a sequence $\{t_{k_1}\}$ of real numbers, $t_{k_1} \rightarrow 0$ which lie between the numbers of the sequence $\{t_k\}$. We thus apply Rolle's Theorem successively $2n$ times and finally conclude that the function $g^{(2n)}(t)$ coincides with the function $\theta^{(2n)}(t)$ for every value of a sequence $\{t_{k_{2n}}\}$ of real numbers, $t_{k_{2n}} \rightarrow 0$. For simplicity in notation we denote this sequence by $\{t'_k\}$. The relation $g^{(2n)}(t'_k) = \theta^{(2n)}(t'_k)$ holds for every value t'_k , $t'_k \rightarrow 0$ so that we must have $g^{(2n)}(0) = \theta^{(2n)}(0)$ and therefore $(-1)^n \gamma_{2n} = \int_{-\infty}^{\infty} x^{2n} dG(x) > 0$. Thus for every value of the sequence $\{t'_k\}$, $t'_k \rightarrow 0$ we have

$$\frac{g^{(2n)}(t'_k) - g^{(2n)}(0)}{t'^k{}^2} = \frac{\theta^{(2n)}(t'_k) - \theta^{(2n)}(0)}{t'^k{}^2} = \frac{\gamma_{2n+2}}{2} + O(t'^k{}^2)$$

so that

$$\lim_{k \rightarrow \infty} \frac{g^{(2n)}(t'_k) - g^{(2n)}(0)}{t'^k{}^2}$$

exists and is finite. But we can verify easily that

$$\frac{g^{(2n)}(t'_k) - g^{(2n)}(0)}{t'^k{}^2} = (-1)^{n+1} \int_{-\infty}^{\infty} x^{2n} \frac{\sin^2 \frac{t'_k x}{2}}{t'^k{}^2} dG(x)$$

so that the integral on the right-hand side tends to a finite limit as $t'_k \rightarrow 0$. We apply again Fatou's Lemma and deduce that $\int_{-\infty}^{\infty} x^{2n+2} dG(x) < \infty$ that is, $g^{(2n+2)}(0)$ exists. Therefore $g^{(2n+2)}(t)$ exists and is finite for any real t , thus completing the induction.

Now we turn to the proof of the theorem. We note that the function $g(t)$ has finite derivatives of all orders and further $g^{(2n)}(0) = \theta^{(2n)}(0)$ for every n . Hence we have

$$\limsup_{n \rightarrow \infty} \left[\frac{|g^{(2n)}(0)|}{(2n)!} \right]^{1/2n} = \limsup_{n \rightarrow \infty} \left[\frac{|\theta^{(2n)}(0)|}{(2n)!} \right]^{1/2n} = \frac{1}{R}$$

so that the positive definite function $g(z)$ as a function of the complex variable z is also regular in the circle $|z| < R$ about the point $z=0$. Then it follows immediately from the theorem of Raikov [3] that the positive definite function $f(z)$ is also regular in the same circle

$|z| < R$ about the point $z=0$. Thus we conclude that both the functions $f(z)$ and $\psi(z)$ are regular in the circle $|z| < R$ and further note that they coincide for every value of a sequence $\{t_k\}$ of real numbers, $t_k \rightarrow 0$. Therefore the functions $f(z)$ and $\psi(z)$ coincide for every z in the circle $|z| < R$. It then follows at once from the theorem due to Boas [1] that the function $f(t)$ coincides with $\psi(t)$ for every real t and further the function $f(z)$ (z complex) is regular in the strip $|\operatorname{Im} z| < R$.

The following corollary is an immediate consequence of the above theorem:

COROLLARY. *Let $f(t)$ be a real-valued, even and positive-definite function of the real variable t and let $\psi(z)$ be a regular function of the complex variable z such that $\psi(t)$ is real-valued and even. Suppose that $f(t)$ coincides with $\psi(t)$ for every value of the sequence $\{t_k\}$ of real numbers, $t_k \rightarrow 0$. Then $f(t)$ coincides with $\psi(t)$ for every real t .*

This result has been proved by Linnik in [2] under the additional condition that the function $\psi(t)$ is also a positive-definite function.

REFERENCES

1. R. P. Boas, *Sur les series et intégrales de Fourier à coefficients positifs*, C. R. Acad. Sci. Paris vol. 228 (1949) pp. 1837-1838.
2. Yu. V. Linnik, *The problem of determining a general probability distribution by the distribution of a statistic*, (Russian), Teor. Veroyatnost. i Primenen. vol. 1 (1956) pp. 466-478.
3. D. A. Raikov, *On the decomposition of Gauss and Poisson laws*, (Russian), Izv. Akad. Nauk SSSR. Ser. Mat. vol. 1 (1938) pp. 91-124.

CATHOLIC UNIVERSITY OF AMERICA