

DIFFERENTIABLE OPEN MAPS¹

BY P. T. CHURCH

Communicated by Deane Montgomery, June 4, 1962

Let $f: M^n \rightarrow N^n$ be a continuous function, where M^n and N^n are n -manifolds (without boundary). It will be implicitly assumed that the manifolds share the differentiability properties of f , e.g., $f \in C'$ implies that M^n and N^n are C' manifolds. The map f is called *open* if, whenever U is open in M^n , $f(U)$ is open in N^n ; it is *light* if, for every $y \in N^n$, $\dim(f^{-1}(y)) \leq 0$.

For $n=2$, it is well-known that a nonconstant complex analytic function is open and light. Conversely, Stoilow proved that every light open map is locally, at each point, topologically equivalent [9, p. 198] to one of the canonical analytic maps g_d , defined by $g_d(z) = z^d$ ($d=1, 2, \dots$). If it is not assumed that f is light, however, f may be quite different from a g_d . R. D. Anderson in [1] (see also [2]) constructed an open map $f: S^2 \rightarrow S^2$ such that, for each $y \in S^2$, $f^{-1}(y)$ is a nondegenerate continuum.

For $n \geq 2$, let $F_{n,d}: E^n \rightarrow E^n$ be the canonical open map defined by: $F_{n,d}(x_1, x_2, \dots, x_n) = (u_1, u_2, \dots, u_n)$, where $u_1 + iu_2 = (x_1 + ix_2)^d$ ($i = \sqrt{-1}$) and $u_j = x_j$ ($j=3, 4, \dots, n; d=1, 2, \dots$). Since each $F_{n,d}$ is a generalization of g_d , it is natural to wonder (for $n \geq 3$) how much an arbitrary open map f , satisfying some additional condition, differs locally from one of them.

The *branch set* B_f is the set of points in M^n at which f fails to be a local homeomorphism (defined in [3]).

THEOREM. *Let $f: M^n \rightarrow N^n$ be C^n and open ($n \geq 2$), where M^n is compact or f is light. Then there exists a closed set E , $\dim E \leq n-3$, such that, for each x in $M^n - E$, there exists a neighborhood of x on which f is topologically equivalent to one of the canonical maps $F_{n,d}$ ($d=1, 2, \dots$). Moreover, E is nowhere dense in B_f unless f is a local homeomorphism.*

In particular, for $n=2$ we have the classical structure. In [4, p. 620, (4.3)] there is a 2-to-1 open map $f: S^5 \rightarrow S^5$ for which B_f is not locally a manifold at any point (it is necessarily [4, p. 620, (4.2)] a 3-gm mod 2); thus *some* differentiability assumption is required above. There is a C^∞ open map $f: E^2 \rightarrow E^2$ for which B_f is the y -axis; thus either compactness of the domain or lightness of the map is needed. An example $f: E^3 \rightarrow E^3$ (or $f: S^3 \rightarrow S^3$) given by E. Hemmingsen

¹ Research supported in part by National Science Foundation grant 18049.

and the author in [4, p. 620, (3.3)] indicates the extent of possible pathology. There B_f has a Cantor set of point components, so that the exceptional set E in the Theorem is necessary (f can be shown to be topologically equivalent to a C^∞ map).

The following corollary is a generalization of the inverse function theorem. Let Z be the set of zeros of the Jacobian determinant.

COROLLARY. *If $f: E^n \rightarrow E^n$, $n \geq 3$, $f \in C^n$, and $\dim Z \leq 0$, then f is a local homeomorphism.*

PROOF. The map f is light, and its Jacobian determinant is either non-negative or nonpositive everywhere. Thus f is open [8], and the result follows from the Theorem. More generally, the conclusion holds if $\dim(B_f) \leq 0$.

A basic lemma for the proof of the Theorem follows. The set of points in M^n at which the Jacobian matrix has rank at most q is denoted by R_q .

LEMMA. *Let $h: M^n \rightarrow N^p$, where $h \in C^n$ and M^n and N^p are n - and p -manifolds, respectively. Then $\dim(f(R_q)) \leq q$.*

In particular, $\dim(h(M^n)) \leq n$. The lemma is related to the theorem of A. P. Morse [6] on the image of the critical set of a real-valued function, and to Sard's Theorem [7]. If f is light, then [5, pp. 91–92] $\dim(R_q) \leq q$.

The proof of the Theorem employs Morse's Theorem, a uniform form of the implicit function theorem, and some results from [3]. Detailed proofs will appear elsewhere.

REFERENCES

1. R. D. Anderson, *On monotone interior mappings in the plane*, Trans. Amer. Math. Soc. **73** (1952), 211–222.
2. ———, *Open mappings of compact continua*, Proc. Nat. Acad. Sci. U.S.A. **42** (1956), 347–349.
3. P. T. Church and E. Hemmingsen, *Light open maps on n -manifolds*, Duke Math. J. **27** (1960), 527–536.
4. ———, *Light open maps on n -manifolds. II*, Duke Math. J. **28** (1961), 607–624.
5. W. Hurewicz and H. Wallman, *Dimension theory*, Princeton Mathematical Series, Vol. 4, Princeton Univ. Press, Princeton, N. J., 1941.
6. A. P. Morse, *The behavior of a function on its critical set of points*, Ann. of Math. **40** (1939), 62–70.
7. A. Sard, *The measure of the critical values of differentiable maps*, Bull. Amer. Math. Soc. **48** (1942), 883–890.
8. C. J. Titus and G. S. Young, *A Jacobian condition for interiority*, Michigan Math. J. **1** (1952), 89–94.
9. G. T. Whyburn, *Analytic topology*, Amer. Math. Soc. Colloq. Publ. Vol. 28, Amer. Math. Soc., New York, 1942.