

MINIMAL TOTAL ABSOLUTE CURVATURE FOR ORIENTABLE SURFACES WITH BOUNDARY

BY JAMES H. WHITE

Communicated by S. S. Chern, September 22, 1973

Let M be an orientable surface with single smooth boundary curve C which is C^2 imbedded in Euclidean three-space E^3 . (M may be thought of as a closed orientable surface with a single disc removed.) Let M_ϵ be the set of points of E^3 at a distance ϵ from M . M_ϵ is, of course, for small ϵ , an imbedded closed surface which is almost everywhere C^2 . Using N. Grossman's [1] adaptation of N. Kuiper's [2] definition, we say that M has minimal total absolute curvature if M_ϵ is tightly imbedded or has the two piece property, TPP [2].

We announce the following result:

THEOREM. *Let M be an orientable surface of genus g with a single smooth boundary curve which is C^2 imbedded in E^3 . Then M has minimal total absolute curvature if and only if M has $g=0$ and is a planar disc bounded by a convex curve.*

The proof uses a series of integral equations and geometric arguments. The outline is as follows. First, in his paper [1], N. Grossman shows that an orientable surface M of genus g with boundary curve C has minimal total absolute curvature only if the following integral equality holds:

$$(1) \quad \frac{1}{2\pi} \int_M |K| dA + \frac{1}{2\pi} \int_C \kappa ds = 1 + 2g,$$

where K is the Gauss curvature of M and κ is the Frenet curvature of the boundary curve C considered as a space curve in E^3 , where dA is the area element of M and ds is the arc element of C . Note that the right-hand side is the sum of the betti-numbers of M and compare with Kuiper [2] for closed surfaces.

Next, the theorem of Gauss-Bonnet yields

$$(2) \quad \frac{1}{2\pi} \int_M K dA + \frac{1}{2\pi} \int_C \kappa_g ds = 1 - 2g,$$

AMS (MOS) subject classifications (1970). Primary 53A05; Secondary 58E99.

Copyright © American Mathematical Society 1974

where κ_g is the geodesic curvature of C considered as a curve on the surface M .

Adding (1) and (2), we obtain that if M has minimal total absolute curvature,

$$(3) \quad \frac{2}{2\pi} \int_{M:\{K>0\}} K \, dA + \frac{1}{2\pi} \int_C (\kappa + \kappa_g) \, ds = 2,$$

where the first integral is taken over the points of M where $K > 0$.

LEMMA 1. *If M has minimal total absolute curvature, then M has TPP.*

In [3], L. Rodriguez shows that, if M has TPP,

$$(4) \quad \frac{1}{2\pi} \int_{M:\{K>0\}} K \, dA + \frac{1}{2\pi} \int_C (\kappa + \kappa_g) \, ds = 2.$$

Subtracting (4) from (3), we obtain $(1/2\pi) \int_{M:\{K>0\}} K \, dA = 0$, and hence $K \leq 0$ in the interior of M .

LEMMA 2. *$K \leq 0$ in the interior of M .*

LEMMA 3. *C is a plane convex curve.*

Lemma 3 is proved by using Morse theory and studying the convex hull of M_e .

LEMMA 4. *$K \equiv 0$ in the interior of M .*

This follows immediately from Lemmas 2 and 3.

Now Lemma 4 implies $\int_M |K| \, dA = 0$, and Lemma 3 implies $(1/2\pi) \int_C \kappa \, ds = 1$. Thus, in order for equation (1) to hold g must be zero and M must be a planar disc bounded by a convex curve.

REFERENCES

1. N. Grossman, *Relative Chern-Lashof theorems*, J. Differential Geometry 7 (1972), 611–618.
2. N. H. Kuiper, *Minimal total absolute curvature for immersions*, Invent. Math. 10 (1970), 209–238. MR 42 #2499.
3. L. Rodriguez, *The two-piece-property and relative tightness for surfaces with boundary* (xeroxed thesis), Brown University, Providence, R.I.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, CALIFORNIA 90024