

## CONTINUED FRACTALS AND THE SEIFERT CONJECTURE

BY JENNY HARRISON<sup>1</sup>

In 1950 Herbert Seifert posed a question today known as the Seifert Conjecture:

“Every  $C^r$  vector field on the three-sphere has either a zero or a closed integral curve.”

Paul Schweitzer published his celebrated  $C^1$  counterexample in 1971 [Sch]. We show how to obtain a  $C^{3-\varepsilon}$  counterexample  $X$  by using techniques from number theory, analysis, fractal geometry, and differential topology [H1 and H2].  $X$  is  $C^2$  and its second derivative satisfies a  $(1 - \varepsilon)$ -Hölder condition.

**1. Continued fractions and quasi-circles.** Any irrational number  $\alpha$ ,  $0 < \alpha < 1$ , can be expressed as a continued fraction

$$\alpha = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}$$

where the  $a_i$  are positive integers. One writes  $\alpha = [a_1, a_2, a_3, \dots]$ . The truncation  $[a_1, \dots, a_n] = p_n/q_n$  is the best approximation to  $\alpha$  among all rational numbers  $p/q$  with  $0 < q \leq q_n$ . The growth rate of the  $a_i$  tells “how irrational”  $\alpha = [a_i]$  is. At one extreme is the Golden Mean,  $\gamma = [1, 1, \dots]$ ; at the other are Liouville numbers such as  $\lambda = [1^{1!}, 2^{2!}, 3^{3!}, \dots]$ . The former is “very irrational” while the latter is “almost rational”.

To study  $\alpha$  dynamically it is standard to consider  $R_\alpha$ , the rigid rotation of the circle  $S^1$  of unit length through angle  $\alpha$ . Choose  $x \in S^1$  and consider its  $R_\alpha$ -orbit  $O_\alpha(x)$ . Since  $\alpha$  is irrational,  $O_\alpha(x)$  is dense in  $S^1$ . But how is it dense? For Liouville  $\lambda$ ,  $O_\lambda(x)$  contains long strings  $\{R_\lambda^n(x), R_\lambda^{n+1}(x), \dots, R_\lambda^m(x)\}$  that are poorly distributed. They “bunch up”. In contrast, the Golden Mean’s orbit distributes itself fairly evenly throughout  $S^1$ .

Unfortunately, it is hard to distinguish visually (and hence geometrically) between bunched-up dense orbits and well distributed ones. After many iterates, the orbit picture becomes blurred. This is due in fact to the picture’s being drawn on the circle. As a remedy, we “unfold”  $S^1$  onto a canonically constructed curve  $Q_\alpha$  in the 2-sphere  $S^2$  as follows.

Choose a “Denjoy” projection  $\rho: S^1 \rightarrow S^1$ ; that is,  $\rho$  is onto and continuous,  $\rho^{-1}(n\alpha)$  is an interval  $I_n$  for all  $n \in \mathbf{Z}$ , the  $I_n$  are disjoint, and  $\rho$  is 1-1

Received by the editors January 13, 1982 and, in revised form, April 25, 1985.

1980 *Mathematics Subject Classification*. Primary 58F25, 34C35, 10K10.

<sup>1</sup>Partially supported by NSF grant no. MCS-8302062.

away from  $\bigcup I_n$ . By  $\langle n\alpha \rangle$  we mean the fractional part of  $n\alpha$ . The *discrepancy* of  $\alpha$  is

$$D_n(\alpha) = \sup_I \left\{ \frac{1}{n} \left| \sum_{m=0}^{n-1} \chi_I \langle m\alpha \rangle - |I| \right| \right\}$$

where  $I$  is an interval in  $S^1 = \mathbf{R}/\mathbf{Z}$  and  $\chi_I$  is its characteristic function. Choose weights  $w_n > 0$  so that  $\sum w_n D_n(\alpha)$  converges and  $w_n D_n(\alpha)$  is monotone decreasing as  $|n| \rightarrow \infty$ . For any  $x \in S^1 \setminus \bigcup I_n$  define  $h_\alpha(x) = (h_1(x), h_2(x))$  in  $S^1 \times \mathbf{R}$  by

$$h_1(x) = \rho(x) + \sum_{|n|=0}^{\infty} w_n (\rho(x) - \chi_{[0, \rho(x))} \langle n\alpha \rangle)$$

$$h_2(x) = \sum_{|n|=0}^{\infty} \{ w_{2n+1} \chi_{[0, \rho(x))} \langle (2n+1)\alpha \rangle - w_{2n} \chi_{[0, \rho(x))} \langle 2n\alpha \rangle \}.$$

The mapping  $h_\alpha$  is uniformly continuous. Its extension to the closure of  $S^1 \setminus \bigcup I_n$  sends the endpoints of  $I_n$  onto points  $p_n, q_n$  in the cylinder  $S^1 \times \mathbf{R}^1$  joined by a line segment  $\Delta_n$  of slope  $\pm 1$ . Extend  $h_\alpha$  to  $S^1$  so it sends  $I_n$  onto  $\Delta_n$  homeomorphically.

The curve  $Q_\alpha = h_\alpha(S^1)$  is the *continued fractal* corresponding to  $\alpha$ . It depends only on  $\alpha$  and the choice of weights; different Denjoy projections just give it different parametrizations. When  $\alpha$  is of constant type (its  $a_i$  are bounded) we take  $w_n$  of the form  $1/(1 + |n|^\mu)$  with  $\frac{1}{2} < \mu < 1$ . In that case,  $Q_\alpha$  turns out to be an Ahlfors quasi-circle [Ah]. In any case,

**Think of the continued fractal  $Q_\alpha$  as a picture of  $\alpha$ .**

Its geometry embodies not only the early patterns apparent from the circle rotation but also much of its long-term behavior.

In Figure 1 are three examples drawn on the open cylinder with  $x = 0$  and  $x = 1$  identified. The quasi-circle  $Q_\alpha$  with  $\alpha = [4, 4, 4, \dots]$  leads to a  $C^{2+\delta}$  Seifert counterexample with  $\delta$  small. To raise the differentiability from  $C^{2+\delta}$  to  $C^{3-\epsilon}$  we take  $\alpha = [2N, 2N, \dots]$  with  $N$  large and prove a sharpened Denjoy-Koksma inequality for such numbers  $\alpha$  [H6]. The choice  $\alpha = \sqrt{21}$  also leads to a  $C^{2+\delta}$  example but  $(\sqrt{5} - 2)^{1/2}$  does not.

**2. Fractal geometry of  $Q$ .** When  $\alpha$  is of constant type, the continued fractal  $Q = Q_\alpha$  can be exhibited as the nested intersection of connected closed sets  $A^n$  called  *$\beta$ -diamond chains*, as in Figure 2. Formally,  $A^n = \Delta_1^n \cup T_1^n \cup \dots \cup \Delta_m^n \cup T_m^n$ , where  $m = m(n)$  and (suppressing the superscript  $n$  as appropriate)

(a)  $\Delta_i$  is a segment of slope  $\pm 1$  with respect to the cylinder's coordinates. The  $\Delta_i$  are *diagonals* of  $A^n$ ,  $1 \leq i \leq m$ .

(b)  $T_i$  is a parallelogram with interior, whose edges have slope  $\pm\beta$ . In practice,  $\beta \ll 1$ . The  $T_i$  are  *$\beta$ -diamonds* of  $A^n$ ,  $1 \leq i \leq m$ .

(c)  $\Delta_i$  slopes backwards and joins the right-hand vertex of  $T_{i-1}$  to the left-hand vertex of  $T_i$ ,  $1 \leq i \leq m$ . We call  $T_0 = T_m$  to take care of the case  $i = 1$ .




FIGURE 1. Continued Fractals.

(d)  $A^1 \supset A^2 \supset \dots \supset A^n \supset \dots$  and the diameters of the diamonds in  $A^n$  tend to zero as  $n \rightarrow \infty$ .

(e)  $\text{diam}(\Delta_1^n) \leq \text{diam}(T_j^n)$  if  $1 \leq i, j \leq m$  and  $\Delta_1^n$  is a diagonal of  $A^n$  but not of  $A^{n-1}$ .

The intersection  $C = \bigcap A^n$  is a Jordan curve consisting of all the diagonals  $\Delta_1^n$  of all the  $A^n$  plus a Cantor set  $\Gamma$ ,

$$C = \Delta \cup \Gamma, \quad \Delta = \bigcup \Delta_1^n.$$




FIGURE 2. A diamond circle.

Such a  $C$  is called a *diamond circle*. All continued fractals  $Q_\alpha$  with  $\alpha$  of constant type are diamond circles and all diamond circles are quasi-circles.

As a degenerate case, suppose all the diagonals of a diamond circle  $C$  are points. Then  $C$  is the graph of a Lipschitz function  $S^1 \rightarrow \mathbf{R}$  having Lipschitz constant  $\beta$  (and conversely). In the nondegenerate case and when  $\alpha$  is of constant type, its continued fractal  $Q = Q_\alpha$  turns out to have Hausdorff dimension  $> 1$ , so it cannot be a graph. Nevertheless,  $Q$  has the following

*Graph-like property.* There exist angles  $\eta', \eta$ ,  $0 < \eta' < \eta < \pi$ , a neighborhood  $U$  of  $Q$  in the cylinder, and a family of disjoint open sets  $D_i$ ,  $i \in \mathbf{Z}$ , such that

(a) Each  $D_i$  is a homothetic replica of a fixed hexagon and contains the interior of the diagonal  $\Delta_i$ . We denote  $D = \bigcup D_i$ .

(b) If  $x \in U \setminus D$  lies on the north side of  $Q$  then any point  $y \in Q$  nearest  $x$  lies in the downward pointing sector of angle  $\eta$  at  $x$ . If  $x$  lies near the bottom edge of  $D_i$  then  $y$  lies in the downward pointing sector of angle  $\eta'$  at  $x$ . Symmetric conditions prevail south of  $Q$ . See Figure 3.

If  $C$  is a Lipschitz graph with Lipschitz constant  $\beta$  this property is obvious;  $D$  is empty and  $\eta = 2 \arctan \beta$ . When  $C$  is a general diamond curve the proof is tricky.

**3. Denjoy homeomorphisms of  $Q$  and the Whitney extension theorem.** To introduce dynamics on  $Q_\alpha$  we consider any Denjoy homeomorphism  $\mathcal{D}$  of  $S^1$  satisfying  $\rho \mathcal{D} = R_{2\alpha} \rho$  (recall  $\rho$  from §1). Then we lift  $\mathcal{D}: S^1 \rightarrow S^1$  to  $f: Q_\alpha \rightarrow Q_\alpha$  via the embedding  $h_\alpha: S^1 \rightarrow Q_\alpha$ . Let  $\varepsilon > 0$  be given. Choose a large integer  $N$  and set  $\alpha = [2N, 2N, \dots]$ . The right choice of weights  $w_n$  in the definition of  $h_\alpha$  gives

$$(*) \quad \|f(x) - f(y) - (x - y)\| < C \|x - y\|^{3-\varepsilon}$$

for some constant  $C$  and all  $x, y$  in the Cantor set  $\Gamma$ . Using only (\*), the Whitney Extension Theorem [W, AR] and the fact that  $Q_\alpha$  is a quasi-circle,




FIGURE 3. Imposing semistability.

we find a  $C^{3-\epsilon}$  diffeomorphism  $F: S^2 \rightarrow S^2$  fixing the poles of  $S^2$  such that  $F|\Gamma = f|\Gamma$ ,  $F(Q_\alpha) = Q_\alpha$ ,  $DF|\Gamma = \text{Id}$ ,  $D^2F|\Gamma = 0$  and  $F(D_i) = D_{i+2}$ . In particular,  $F$  is a  $C^{3-\epsilon}$  Denjoy rotation of  $Q_\alpha$ , cf. [Ha, Kn].

Since no quasi-circles have  $C^3$  Denjoy rotations [H4], one is led to wonder if the differentiability class  $C^3$  separates Seifert counterexamples from a Seifert Theorem, much as happens in KAM theory [He, M]. It is also interesting to speculate about the relation between  $\epsilon$  and the Hausdorff dimension of  $\Gamma$ . Our  $F$  turns out to be of class  $C^{3-\epsilon}$  and our  $\Gamma$  has  $HD(\Gamma) = 2 - \epsilon$ , cf. [N]. Must  $HD(\Gamma)$  be large if the distortion of  $Df$  at  $\Gamma$  is small? Cf. [H4, H5].

**4. Semistability of  $g$  at  $Q$ .** We want a modification  $G$  of the Whitney extension  $F$  in §3 so that  $G = F$  on  $Q$ ,  $G$  is a  $C^{3-\epsilon}$  diffeomorphism of  $S^2$  and  $Q$  is  $G$ -semistable: under forward  $G$ -iterates  $Q$  attracts the north side of  $S^2 \setminus Q$  and the reverse holds south of  $Q$ .

North of  $Q$  we want to push  $F(x)$  closer to  $Q$  than  $x$  was. The crucial fact that lets us do so (in a  $C^2$  fashion near  $\Gamma$ ) is the  $C^2$ -flabby condition  $DF|\Gamma = \text{Id}$  and  $D^2F|\Gamma = 0$ . The Denjoy examples of Knill [Kn], Hall [Ha] and Herman [He] do not have this property and that is what prevents their use against the Seifert Conjecture.

In Figure 3 we indicate the directions in which we push  $F(x)$  toward  $Q$ . Since  $DF|\Gamma = \text{Id}$  and  $D^2F|\Gamma = 0$ , such pushing meets little resistance. At this stage of the construction we use the downward-pointing sector (shaded) from the graphlike property (§2), the fact that the quasi-slope  $\beta$  of  $Q$  is small, and the fact that the diagonals  $\Delta_i$  slope backward. Under the resulting diffeomorphism  $G$ ,  $Q$  is semistable. North of  $Q$ ,  $G(D_i) \subsetneq D_{i+2}$ , while south of  $Q$ ,  $G^{-1}(D_i) \subsetneq D_{i-2}$ .

**Seifert counterexamples and loxodromic diffeomorphisms.** The diffeomorphism  $G$  constructed in §4 sends some  $x_0 \in U \setminus D$  north of  $Q$  into  $D_0$ . Under  $G$  its  $\alpha$ -limit is the north pole  $N$  and its  $\omega$ -limit is  $\Gamma$ . Similarly, some  $y_0 \in D_0$  south of  $Q$  is sent into  $U \setminus D$  by  $G$ ; its  $\alpha$ -limit is  $\Gamma$  and its  $\omega$ -limit is the south pole  $S$ . Compose  $G$  with a  $C^\infty$  motion  $M$  of  $S^2$  such that  $M(G(x_0)) = y_0$  and  $M$  leaves all points of  $S^2 \setminus D_0$  fixed. The resulting  $C^{3-\varepsilon}$  diffeomorphism  $H = M \circ G: S^2 \rightarrow S^2$  has the following properties:

(a) The only periodic points of  $H$  are its fixed-point poles,  $N$  and  $S$ . They are a source and sink, respectively.

(b)  $\lim_{n \rightarrow -\infty} H^n(x_0) = N$  and  $\lim_{n \rightarrow \infty} H^n(x_0) = S$  for some  $x_0$ .

(c)  $H$  has a minimal set other than the poles.

(a) follows from disjointness of the  $G^n(D_0)$ ,  $n \in \mathbf{Z}$ ; (b) is by construction; (c) is clear— $\Gamma$  is the minimal set.

A suspension similar in spirit to Schweitzer's [Sch] lets us use  $H$  to construct a  $C^{3-\varepsilon}$  flow  $\phi$  on  $S^3$  with no compact orbits. By Hart's Smoothing Theorem [Ht],  $\phi$  is conjugate to a flow  $\psi$  whose generating vector field  $X$  is also of class  $C^{3-\varepsilon}$ .

*This vector field  $X$  is a  $C^{3-\varepsilon}$  counterexample to the Seifert conjecture. The same procedure applied to any  $C^r$  diffeomorphisms  $H: S^2 \rightarrow S^2$  obeying conditions (a), (b), (c) above would produce a counterexample to the  $C^r$  Seifert Conjecture.*

It is not known if  $X$  is  $C^2$  structurally stable. By Pugh's Closing Lemma it is not  $C^1$  structurally stable [P]. Any diffeomorphism of  $S^2$  obeying (a) and (b) but having no minimal set except the poles is topologically conjugate to the standard loxodromic diffeomorphism  $z \rightarrow \frac{1}{2}z$  of the closed complex plane  $C \cup \infty = S^2$ . Thus we put forward the

**CONJECTURE.** *Every  $C^3$  diffeomorphism of  $S^2$  satisfying conditions (a), (b) above is loxodromic.*

This is a dissipative analogue of Birkhoff's conjecture that any measure-preserving diffeomorphism of  $S^2$  whose only periodic points are the two fixed point poles must be topologically conjugate to a rigid irrational rotation of  $S^2$ .

## REFERENCES

- [AR] R. Abraham and J. Robbin, *Transversal mappings and flows*, Benjamin, New York, 1967.
- [Ah] L. V. Ahlfors, *Lectures on quasi-conformal mappings*, Van Nostrand, Princeton, N.J., 1966.
- [D] A. Denjoy, *Sur les courbes définies par les équations différentielles à la surface du tore*, J. Math. Pures Appl. **11** (1932), 333–375.
- [Ha] G. R. Hall, *A  $C^\infty$  Denjoy counterexample*, Ergodic Theory and Dynamical Systems, Vol. 1 (1981), 261–272.
- [H1] J. Harrison, *Denjoy fractals*, (preprint).
- [H2] —,  *$C^{2+\delta}$  Counterexamples to the Seifert conjecture*, (preprint).
- [H3] —, *Opening closed leaves of foliations*, Bulletin LMS, 1982.
- [H4] —, *Topological transitivity for Ahlfors quasi-circles*, (preprint).

- [H5] —, *Bounded distortion and an  $n$ -dimensional Denjoy Theorem* (in preparation).
- [H6] —, *Diophantine type and the Denjoy-Koksma inequality*, (in preparation).
- [HN1] J. Harrison and A. Norton, *An improved sampling procedure for the uniform distribution of  $\{na\}$* , (preprint).
- [HN2] —, *Denjoy quasi-circles in  $\mathbb{R}^n$  with Hausdorff dimension  $n - \varepsilon$*  (in preparation).
- [HY] J. Harrison and J. A. Yorke, *Vector fields on  $S^3$  and  $R^3$  without periodic-points*, Proc. Simposio Internac. de Sistemas Dinamicos, Springer Lecture Notes.
- [Ht] D. Hart, *On the smoothness of generators*, Topology **22** (1983), 357–363.
- [He] M. Herman, *Contre-exemples de Denjoy et contre-exemples de classe  $C^{3-\varepsilon}$  au théorème des courbes invariantes ayant un nombre de rotation fixe*, (preprint).
- [Kh] A. Khintchine, *Continued fractions*, translated by P. Wynn (1963) Groningen, Nordhoff.
- [Kn] R. J. Knill, *A  $C^\infty$  flow on  $S^3$  with a Denjoy minimal set*, J. Differential Geom. **16** (1981), 271–280.
- [KN] L. Kuipers and H. Niederreiter, *Uniform distribution of sequences*, Wiley, New York, 1974.
- [M] J. Mather, *Existence of quasi-periodic orbits for twist homeomorphisms of the annulus*, (preprint).
- [N] A. Norton, *A critical set with non-null image has large Hausdorff dimension*, (preprint).
- [P] C. Pugh, *The Closing Lemma*, Amer. J. Math. **89** (1967), 956–1021.
- [Sch] P. Schweitzer, *Counterexamples to the Seifert conjecture and opening closed leaves of foliations*, Ann. of Math. (2) **100** (1971), 386–400.
- [Sei] H. Seifert, *Closed integral curves in 3-space and isotopic two-dimensional deformations*, Proc. Amer. Math. Soc. **1** (1950), 287–302.
- [W] H. Whitney, *Analytic extensions of differentiable functions defined on closed sets*, Trans. Amer. Math. Soc. **36** (1934), 369–387.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, EVANS HALL,  
BERKELEY, CALIFORNIA 94720