

variational inequalities, and to applications in elasticity. This class of problems is approximated by Ritz and FEM methods. An associated approximation error estimate is given, and the stability of the methods is analyzed. In this context, stability results for the solution of general nonlinear recurrences and the Newton-Kantorovich method under perturbation and rounding errors are proved.

The monograph under review is basically a collection of work of Mikhlin, his students, and other Russian authors on numerical methods. The bibliography lists a great number of their publications since 1949. There are also references to work of non-Russian mathematicians, mostly in introductory remarks. It seems that this work had only a limited influence on the style and content of most of the book.

F. STUMMEL

JOHANN WOLFGANG GOETHE-UNIVERSITÄT  
E-mail address: stummel@math.uni.frankfurt.de

BULLETIN (New Series) OF THE  
AMERICAN MATHEMATICAL SOCIETY  
Volume 28, Number 1, January 1993  
©1993 American Mathematical Society  
0273-0979/93 \$1.00 + \$.25 per page

*Two-dimensional geometric variational problems*, by Jürgen Jost. Wiley, New York, 1991, x+234 pp. \$87.95, ISBN 0-471-92839-9

In the first chapter the author presents the most important and fascinating geometric variational problem, namely, to span a disc-type surface of minimal area into a given spatial Jordan curve  $\Gamma$ . This mathematical problem can be realized by soap films spanning into a given wire, initiated by experiments of the Belgian physicist Plateau. Then the general two-dimensional conformally invariant variational problem is considered, where the function  $u \in H^{1,2}(\Omega, \mathbb{R}^d)$  with  $d \geq 2$  renders the integral

$$(1) \quad I(u) := \frac{1}{2} \int_{\Omega} \{g_{ik}(u) \nabla u^i \nabla u^k + b_{ik}(u) \det(\nabla u^i, \nabla u^k)\} dx dy$$

stationary. Here  $(g_{ik})_{ik}$  is a positive definite and  $(b_{ik})_{ik}$  a skew-symmetric matrix. The Euler-Lagrange equations of  $I$  are given by

$$(2) \quad \Delta u^i + \Gamma_{kl}^i \nabla u^k \nabla u^l = g^{im} (b_{mk,l} + b_{kl,m} + b_{lm,k}) \det(\nabla u^k, \nabla u^l)$$

with  $(g^{ik})_{ik} = (g_{ik})_{ik}^{-1}$ ,  $g_{kl,m} = \partial g_{kl} / \partial u_m$ , and the Christoffel symbols  $\Gamma_{kl}^i$ . One can interpret (2) as the equation for a surface of prescribed mean curvature in a Riemannian manifold. The two-dimensional conformally invariant variational problems especially give rise to conformal maps between surfaces, parametric minimal surfaces in Riemannian manifolds, and harmonic maps from a surface into a Riemannian manifold.

For surfaces of prescribed mean curvature in  $\mathbb{R}^3$  one considers the following functional of Heinz and Hildebrandt

$$(3) \quad I(u) = \frac{1}{2} \int_{\Omega} \{|\nabla u|^2 + Q(u) \cdot u_x \wedge u_y\} dx dy$$

for functions  $u: \Omega \rightarrow \mathbb{R}^3$  with  $\Omega \subset \mathbb{R}^2$ , where the vector field  $Q$  satisfies  $\operatorname{div} Q(u) = 4H(u)$  with the mean curvature  $H$  and  $\wedge$  denotes the cross product.

The functional (3) has Rellich's *H-surface system*

$$(4) \quad \Delta u = 2H(u)u_x \wedge u_y \quad \text{in } \Omega$$

as its Euler-Lagrange equation.

For harmonic maps one considers the critical points of the energy functional

$$(5) \quad I(u) = \frac{1}{2} \int_{\Omega} g_{ik}(u_x^i u_x^k + u_y^i u_y^k) dx dy$$

generating the *harmonic functions*

$$(6) \quad \Delta u^i + \Gamma_{kl}^i(u_x^k u_x^l + u_y^k u_y^l) = 0, \quad i = 1, \dots, d.$$

The author discusses holomorphic quadratic functionals and gives a new proof of H. Hopf's celebrated theorem that a closed surface of genus 0 that is immersed with constant mean curvature into  $\mathbb{R}^3$  is necessarily a sphere.

Using the Darboux system of equations

$$(7) \quad \begin{aligned} \Delta u + \left( \Gamma_{11}^1 + \frac{1}{2} \frac{\partial}{\partial u} \log K \right) |\nabla u|^2 \\ + \left( 2\Gamma_{12}^1 + \frac{1}{2} \frac{\partial}{\partial v} \log K \right) (u_x v_x + u_y v_y) + \Gamma_{22}^1 |\nabla v|^2 = 0, \\ \Delta v + \Gamma_{11}^2 |\nabla u|^2 + \left( 2\Gamma_{12}^2 + \frac{1}{2} \frac{\partial}{\partial u} \log K \right) (u_x v_x + u_y v_y) \\ + \left( \Gamma_{22}^2 + \frac{1}{2} \frac{\partial}{\partial v} \log K \right) |\nabla v|^2 = 0, \end{aligned}$$

a beautiful proof of Liebmann's theorem is given.

The second chapter is devoted to regularity questions. It begins with the introduction of harmonic coordinates due to Karcher and Jost and continues with the fundamental uniqueness theorem of Jäger and Kaul for harmonic maps. Then Grüter's ingenious method is utilized to show continuity of weak solutions via the monotonicity formula: At first continuity of weak minimal surfaces (in the interior and up to the free boundary) is shown. Then weak solutions of the *H-surface system* (4) given in weakly conformal parameters

$$(8) \quad u_x \cdot u_y = 0, \quad |u_x| = |u_y| \quad \text{a.e. in } \Omega$$

are shown to be continuous. Finally, a general regularity result for two-dimensional geometric variational problems is presented. The celebrated theorem of Sacks and Uhlenbeck on the removability of isolated singularities is also derived.

Higher regularity, together with  $C^{2,\alpha}$ -estimates for harmonic maps, are then studied. Centrally important is the theorem of Hildebrandt, Kaul, and Widman: "Let  $N$  be a complete Riemannian manifold with a complete  $C^2$ -submanifold  $M$ . If  $u \in H^{1,2}(D, N)$  is a continuous, weakly harmonic map with free boundary  $M$ , then  $u \in C^{1,\alpha}(\bar{D}, N)$  holds true."

With the aid of the fundamental method of Heinz, the author derives gradient bounds for harmonic maps up to the free boundary. Also, the Hartman-Wintner

lemma for asymptotic expansions at singular points (in the interior and on the boundary) of the form

$$(9) \quad u_z = \frac{1}{2}(u_x - iu_y) = a(z - z_0)^m + o(|z - z_0|^m), \quad z \rightarrow z_0,$$

is explained.

The last section of Chapter 2 is devoted to estimates from below for the Jacobian of univalent harmonic mappings. The interior estimates are valid up to the boundary under convexity assumptions. Heinz discovered the fundamental importance of these kind of estimates for curvature bounds and  $C^{2,\alpha}$ -estimates for Monge-Ampère equations in connection with the system (7) and provided basic analytic tools to estimate the Jacobian from below.

Chapter 3 deals with conformal representation of surfaces homeomorphic to the sphere  $S^2$ , circular domains, and closed surfaces of higher genus. The proof is given by direct variational methods and not as usual by uniformization, completing a fragmentary proof of Morrey. The continuous and differentiable boundary correspondence is also directly attained.

Chapter 4 is devoted to existence results. Having established a local existence theorem, a harmonic map representing a saddle point of a certain functional is constructed, and many corollaries are presented. For this theorem originally due to Sacks and Uhlenbeck, the author invented a new interesting proof imitating the curve shortening process for the construction of unstable closed geodesics. Additionally, boundary conditions are imposed, especially of Plateau type. The author provides a new proof of the celebrated mountain pass lemma for minimal surfaces in Riemannian manifolds originally due to Ströhmer: "If  $u_1$  and  $u_2$  are two strict relative minima (w.r.t. the  $C^0$ - or  $H^{1,2}$ -topology), then there exists a third minimal surface  $u_3$  which is unstable." In this chapter also, the Plateau-Douglas problem in Riemannian manifolds is beautifully treated, namely, to span a minimal surface of higher topological type into a system of Jordan curves.

In the beginning of Chapter 5 a local existence result for disc-type harmonic diffeomorphisms between Riemannian manifolds is proved by a continuity method, which was invented by the author. This yields the basis for the following global result of Jost and Schoen: "Let  $\Sigma_1$  and  $\Sigma_2$  be compact surfaces without boundary and  $h: \Sigma_1 \rightarrow \Sigma_2$  a diffeomorphism. Then there exists a harmonic diffeomorphism  $u: \Sigma_1 \rightarrow \Sigma_2$  isotopic to  $h$ , which is energy-minimizing in this class."

Also, a global harmonic diffeomorphism with Dirichlet boundary data is constructed, and the following interesting result of Schoen, Yau, and Sampson is proved: "Let  $u: \Sigma_1 \rightarrow \Sigma_2$  be a harmonic map between closed oriented surfaces of the same genus with  $\deg(u) = \pm 1$ , and let the curvature  $K_2$  of  $\Sigma_2$  be nonpositive. Then  $u$  is a diffeomorphism."

With the aid of this well-developed theory of harmonic maps, the author provides a new approach to the Teichmüller theory in Chapter 6. Controlling the corresponding holomorphic quadratic differential, the basic structures of the Teichmüller space differential, the basic structures of the Teichmüller space are obtained, namely, the topological, differentiable, complex, metric, and Kählerian. Harmonic maps are better suited in this context than quasi-conformal maps, since the first class of functions can be better analytically controlled.

This excellent monograph, which is deeply rooted in the mathematical

areas of calculus of variations, nonlinear partial differential equations, differential geometry, complex analysis, and topology, covers a broad, central region of mathematics in great depth. In this treatise the author has reunited mathematical areas that had been a unit in the times of Riemann, and he has invested the mathematical strength now required. The book is well written and highly recommended to every mathematician with interests in one of the areas mentioned above.

FRIEDRICH SAUVIGNY  
 INSTITÜT FÜR MATHEMATIK DER  
 TECHNISCHEN UNIVERSITÄT COTTBÜS,  
 KARL-MARX STR. 17, D-0-7500 COTTBÜS, GERMANY

BULLETIN (New Series) OF THE  
 AMERICAN MATHEMATICAL SOCIETY  
 Volume 28, Number 1, January 1993  
 ©1993 American Mathematical Society  
 0273-0979/93 \$1.00 + \$.25 per page

*Additive number theory of polynomials over a finite field*, by Gove W. Effinger and David R. Hayes. Clarendon Press, Oxford, 157 pp., \$45.00. ISBN 0-19-853583-x

There cannot be many mathematicians who have not heard of the two most celebrated areas of additive number theory, the Goldbach (1742) conjectures and Waring's (1770) problem. The modern interpretation of the Goldbach conjectures is that every even number is the sum of at most two primes and that every odd number greater than one is the sum of at most three primes.

Waring's problem in its original form is the determination, for each  $n > 1$ , of  $g(n)$ , the smallest number  $s$  such that every number is the sum of at most  $s$  positive  $n$ th powers. It turns out that a few peculiar small numbers such as

$$2^n[(3/2)^n] - 1$$

need a relatively large number of  $n$ th powers to represent them, so the value of  $g(n)$  has been established for most values of  $n$ . In view of this, the modern form of Waring's problem has become the determination of  $G(n)$ , the smallest  $s$  such that every sufficiently large number can be written as the sum of at most  $s$  positive  $n$ th powers.

Over the last seventy years there has been a considerable amount of work on these problems, much of it through an analytic technique introduced in a series of seminal papers by Hardy and Littlewood [3-10] and then developed and refined by a number of researchers, most notably Vinogradov [13] and Davenport [2]. The Hardy-Littlewood method has many applications to other problems, many generalizations, and variants, for example, to algebraic number fields, and many of the associated technical devices have relevance to other areas of analytic number theory, such as the theory of the Riemann zeta function and the theory of diophantine approximation. There have been a number of accounts of the method over the years in its various forms and applications, e.g., by Landau [11], Vinogradov [13], Davenport [1], and Vaughan [12]. The basic principle of the method is that one can use the orthogonality of the additive characters