

NONABELIAN SYLOW SUBGROUPS OF FINITE GROUPS OF EVEN ORDER

NAOKI CHIGIRA, NOBUO HIYORI, AND HIROYOSHI YAMAKI

(Communicated by Efim Zelmanov)

ABSTRACT. We have been able to prove that every nonabelian Sylow subgroup of a finite group of even order contains a nontrivial element which commutes with an involution. The proof depends upon the consequences of the classification of finite simple groups.

The purpose of this note is to announce [4]:

Main Theorem. *Every nonabelian Sylow subgroup of a finite group of even order contains a nontrivial element which commutes with an involution.*

Let G be a finite group and $\Gamma(G)$ the prime graph of G . $\Gamma(G)$ is the graph such that the vertex set is the set of prime divisors of $|G|$, and two distinct vertices p and r are joined by an edge if and only if there exists an element of order pr in G . Let $n(\Gamma(G))$ be the number of connected components of $\Gamma(G)$ and $d_G(p, r)$ the distance between two vertices p and r of $\Gamma(G)$. It has been proved that $n(\Gamma(G)) \leq 6$ in [12], [10], [11], [9].

Theorem 1. *Let G be a finite group of even order and p a prime divisor of $|G|$. If $d_G(2, p) \geq 2$, then a Sylow p -subgroup of G is abelian.*

Theorem 1 is a restatement of the Main Theorem in terms of the prime graph $\Gamma(G)$.

Corollary 1. *Let G be a finite group of even order and p a prime divisor of $|G|$. If Δ is a connected component of $\Gamma(G) - \{p\}$ not containing 2, then a Sylow r -subgroup of G is abelian for any $r \in \Delta$.*

There is a certain relation between a subgraph $\Gamma(G) - \{p\}$ of $\Gamma(G)$ and Brauer characters of p -modular representations of G (see [3]).

Theorem 2. *Let G be a finite nonabelian simple group and p an odd prime divisor of $|G|$. Then $d_G(2, p) = 1$ or 2 provided $d_G(2, p) < \infty$.*

The significance of the prime graphs of finite groups can be found in [1], [3], [5], [6], [7], [8], [14], [15]. We apply the classification of finite simple groups (see [1], [2], [4], [7], [10], [13]). It has been proved that a minimal counterexample to Theorem 1

Received by the editors October 20, 1997.

1991 *Mathematics Subject Classification.* Primary 20D05, 20D06, 20D20.

Key words and phrases. Sylow subgroups, prime graphs, simple groups.

The third author was supported in part by Grant-in-Aid for Scientific Research (No. 8304003, No. 08640051), Ministry of Education, Science, Sports and Culture, Japan.

is a nonabelian simple group. We will give some examples of case by case analysis for finite simple groups. Theorem 1 holds true for the sporadic simple groups by Atlas of Finite Groups although we can find several typos in it. For a positive integer k let $\pi(k)$ be the set of all prime divisors of k . Let $\pi_0 = \{p \in \pi(G) \mid d_G(2, p) = 1\}$. Then we do not have to think about primes in π_0 in order to give the proof of Theorem 1.

Example. Let G be the alternating group on n -letters and $p \in \pi(G)$. It is trivial that Theorem 1 holds true for A_5 and A_6 . Assume that $n \geq 7$. If $p \leq n - 4$, then $d_G(2, p) = 1$. If $p \geq n - 3$, then Sylow p -subgroups of G are cyclic of order p . Thus Theorem 1 holds true for the alternating groups.

Example. Let $G = PSL(n, q)$, $q \equiv 0 \pmod{2}$. Then $|G| = q^{n(n-1)/2} \prod_{i=1}^{n-1} (q^{i+1} - 1)d^{-1}$, $d = (n, q - 1)$. Let I_j be the $j \times j$ identity matrix. Put

$$t'_k = \begin{pmatrix} I_k & 0 & 0 \\ 0 & I_{n-2k} & 0 \\ I_k & 0 & I_k \end{pmatrix}.$$

Then t'_k ($1 \leq k \leq r$), where $r = \lfloor n/2 \rfloor$, are representatives of the conjugacy classes of involutions in $SL(n, q)$. The centralizer of t'_k in $SL(n, q)$ is the set of all matrices of the form

$$\begin{pmatrix} A & 0 & 0 \\ H & B & 0 \\ K & L & A \end{pmatrix},$$

where $(\det A)^2 \det B = 1$ and A is a $k \times k$ nonsingular matrix. Denote t_k the homomorphic image of t'_k in $PSL(n, q)$. Then t_k ($1 \leq k \leq r$) are representatives of the conjugacy classes of involutions in $PSL(n, q)$. Let $C_k = C_G(t_k)$. Then

$$\pi(C_k) = \pi(2 \prod_{i=1}^{n-2k} (q^i - 1)/(q - 1)d)$$

and

$$\pi_0 = \pi\left(\prod_{k=1}^r |C_k|\right) = \pi\left(2 \prod_{i=1}^{n-2} (q^i - 1)\right).$$

Suppose $n \geq 4$. Then the only factors of $|G|$ to be considered are $(q^{n-1} - 1)(q^n - 1)$. There are maximal tori $T(A_{n-2})$ of order $(q^{n-1} - 1)d^{-1}$ and $T(A_{n-1})$ of order $(q^n - 1)/(q - 1)d$. Let $p \in \pi(T(X)) - \pi_0$, where $X = A_{n-1}$ or A_{n-2} . Let P be a Sylow p -subgroup of $T(X)$. Then $d_G(2, p) = 1$ or P is a Sylow p -subgroup of G . Since P is abelian, Theorem 1 holds true for $G = PSL(n, q)$, $n \geq 4$.

Suppose that $n = 3$. Then $|G| = q^3(q^2 - 1)(q^3 - 1)d^{-1}$ and there are three classes of maximal tori of orders

$$(q - 1)^2 d^{-1}, \quad (q^2 - 1)d^{-1}, \quad (q^2 + q + 1)d^{-1}.$$

We note that a torus of order $(q^2 + q + 1)d^{-1}$ is an isolated subgroup. If $q > 4$, then $d_G(2, r) = 2$ for $r \in \pi(q + 1)$. Let R be a Sylow r -subgroup of G . Then R is contained in a maximal torus of order $(q^2 - 1)d^{-1}$. If $q = 4$, then $G = PSL(3, 4)$ and $|G| = 2^6 \cdot 3^2 \cdot 5 \cdot 7$. If $q = 2$, then $G = PSL(3, 2)$ and $|G| = 2^3 \cdot 3 \cdot 7$. We have verified Theorem 1 for $n = 3$. It is trivial that Theorem 1 holds true for $PSL(2, q)$.

Theorem 3. *Let G be a simple group of Lie type and T a maximal torus. Let $p \in \pi(T) - \pi_0$. Then T contains a Sylow p -subgroup of G .*

Theorem 3 is a corollary of Theorem 1. Actually we prove Theorem 3 for specified tori when we verify Theorem 1 for the simple groups of Lie type.

REFERENCES

- [1] N. Chigira, Finite groups whose abelian subgroups have consecutive orders, *Osaka J. Math.* **35** (1998), 439–445.
- [2] N. Chigira, Number of Sylow subgroups and p -nilpotence of finite groups, *J. Algebra* **201** (1998), 71–85. CMP 98:09
- [3] N. Chigira and N. Iiyori, Prime graphs and Brauer characters, To appear in *J. Group Theory*.
- [4] N. Chigira, N. Iiyori and H. Yamaki, Nonabelian Sylow subgroups of finite groups of even order, in preparation.
- [5] N. Iiyori, Sharp characters and prime graphs of finite groups, *J. Algebra* **163** (1994), 1–8. MR **94m**:20021
- [6] N. Iiyori, A conjecture of Frobenius and the simple groups of Lie type IV, *J. Algebra* **154** (1993), 188–214. MR **94d**:20014
- [7] N. Iiyori and H. Yamaki, On a conjecture of Frobenius, *Bull. Amer. Math. Soc.* **25** (1991), 413–416. MR **92e**:20014
- [8] N. Iiyori and H. Yamaki, A conjecture of Frobenius and the simple groups of Lie type III, *J. Algebra* **145** (1992), 329–332. MR **93c**:20033
- [9] N. Iiyori and H. Yamaki, A conjecture of Frobenius, *Sugaku Expositions, Amer. Math. Soc.* **9** (1996), 69–85. MR **97a**:00046
- [10] N. Iiyori and H. Yamaki, Prime graph components of the simple groups of Lie type over the fields of even characteristic, *J. Algebra* **155** (1993), 335–343; *Corrigenda*, **181** (1996), 659. MR **94e**:05268
- [11] A. S. Kondrat'ev, Prime graph components of finite simple groups, *Math. USSR Sbornik* **67** (1990), 235–247. MR **90h**:20018
- [12] J. S. Williams, Prime graph components of finite groups, *J. Algebra* **69** (1981), 487–513. MR **82j**:20054
- [13] H. Yamaki, A characterization of the Suzuki simple groups of order 448, 345, 497, 600, *J. Algebra* **40** (1976), 229–244. MR **53**:13384
- [14] H. Yamaki, A conjecture of Frobenius and the sporadic simple groups I, *Comm. Algebra* **11** (1983), 2513–2518; II, *Math. Comp.* **46** (1986), 609–611; *Supplement. Math. Comp.* **46** (1986), S43–S46. MR **85k**:20049; MR **87i**:20033
- [15] H. Yamaki, A conjecture of Frobenius and the simple groups of Lie type I, *Arch. Math.* **42** (1984), 344–347; II, *J. Algebra* **96** (1985), 391–396. MR **85j**:20010; MR **87i**:20032

DEPARTMENT OF MATHEMATICAL SCIENCES, MURORAN INSTITUTE OF TECHNOLOGY, HOKKAIDO 050-8585, JAPAN

E-mail address: chigira@muroran-it.ac.jp

DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, YAMAGUCHI UNIVERSITY, YAMAGUCHI 753-8512, JAPAN

E-mail address: iiyori@po.yb.cc.yamaguchi-u.ac.jp

DEPARTMENT OF MATHEMATICS, KUMAMOTO UNIVERSITY, KUMAMOTO 860-8555, JAPAN

E-mail address: yamaki@gpo.kumamoto-u.ac.jp