

CONSTRUCTION OF ARBITRARY KAZHDAN-LUSZTIG POLYNOMIALS IN SYMMETRIC GROUPS

PATRICK POLO

ABSTRACT. To each polynomial P with integral nonnegative coefficients and constant term equal to 1, of degree d , we associate a certain pair of elements (y, w) in the symmetric group S_n , where $n = 1 + d + P(1)$, such that the Kazhdan-Lusztig polynomial $P_{y,w}$ equals P . This pair satisfies $\ell(w) - \ell(y) = 2d + P(1) - 1$, where $\ell(w)$ denotes the number of inversions of w .

INTRODUCTION

For every Coxeter group (W, S) , Kazhdan and Lusztig defined in [11] a polynomial $P_{y,w}(q) \in \mathbb{Z}[q]$ for each pair (y, w) of elements of W such that $y \leq w$ in the Bruhat-Chevalley order. It has constant term equal to 1 and, when $y \neq w$, degree at most equal to $(\ell(w) - \ell(y) - 1)/2$, where ℓ denotes the length function on (W, S) . When (W, S) is a finite or affine Weyl group, they showed in [12] that the $P_{y,w}(q)$ have nonnegative coefficients by relating them to the intersection cohomology of Schubert varieties.

In spite of their rather elementary definition in [11], the polynomials $P_{y,w}$ are quite difficult to compute explicitly. In view of the interpretation in terms of intersection cohomology of Schubert varieties, it seems that this difficulty reflects the rather complicated nature of the singularities of Schubert varieties in general. In fact, the only families of Kazhdan-Lusztig polynomials known so far correspond to situations where the geometry is simpler, namely Schubert varieties in Grassmannians ([13, 17]) and other minuscule G/Q ([3]), or the incidence variety in $\mathbb{P}^n \times (\mathbb{P}^n)^*$ ([4]).

In this paper we adopt the reverse approach and show that any polynomial $P \in 1 + q\mathbb{N}[q]$ is a Kazhdan-Lusztig polynomial. Namely, we prove the following. Let $d = \deg(P)$ and $n = 1 + d + P(1)$. We may and shall assume that $d \geq 1$.

Theorem. *There exists a pair of elements $y_P < w_P$ in the symmetric group S_n , explicitly defined, whose associated Kazhdan-Lusztig polynomial is P . Further, this pair satisfies $\ell(w_P) - \ell(y_P) = 2d + P(1) - 1$.*

Remarks. 1) In this construction, P is of degree $(\ell(w_P) - \ell(y_P) - 1)/2$ if and only if $P(1) = 2$, that is, if and only if P equals $1 + q^d$.

2) The special case where $P = 1 + q^d$ was known already: a simple geometric proof is given in [4, 4.6]. Also, A. Lascoux has informed me that he had obtained, jointly with M.P. Schützenberger, a combinatorial proof in that case.

Received by the editors December 11, 1998 and, in revised form, April 30, 1999.
1991 *Mathematics Subject Classification.* Primary 14M15; Secondary 20F55, 20G15.

The idea that this result should be true, as well as the precise conjectural form of the elements y_P, w_P , came from computations made in the symmetric group S_8 by using the computer program “Coxeter” written by Fokko Du Cloux ([7]). In order to prove that the postulated pair (y, w) actually has P as the Kazhdan-Lusztig polynomial, we use the interpretation in terms of intersection cohomology. Without loss, we shall assume that the ground field is \mathbb{C} , the field of complex numbers. Let $G = SL_n(\mathbb{C})$, let B be a Borel subgroup and $X_w \subset G/B$ the Schubert variety corresponding to w . We consider a certain resolution of singularities $\pi : Z_w \rightarrow X_w$ (of Bott-Samelson type), similar to the ones considered by Zelevinsky ([17]). At the difference of the situation in [loc. cit.], here π need not be a small map, nor even semi-small. Yet, thanks to the particular form of the permutations w and y , we are able to decompose explicitly the restriction to the open subset $\bigcup_{y \leq z \leq w} BzB/B$ of $R\pi_*(\mathbb{C})$ as a direct sum of shifted Deligne-Goresky-MacPherson complexes, using the decomposition theorem ([2]), ideas of Deodhar ([6]), and a monotonicity result due to Irving ([9]).

After introducing some notation in Section 1, we prove the theorem in Sections 2 and 3. More precisely, in Section 2 we associate to every nonconstant polynomial $P \in 1 + q\mathbb{N}[q]$ a certain pair $y_P < w_P$ and show that the proof of the equality $P_{y_P, w_P} = P$ can be reduced to the case where $P - 1$ is a monomial. We complete the proof in this case in Section 3. Finally, in Section 4 we establish, for any connected semi-simple group, an isomorphism theorem concerning certain locally closed subvarieties of Schubert varieties, which seems to have been overlooked in the literature, and which provides a geometric explanation (in the case of finite Weyl groups) for an equality of Kazhdan-Lusztig polynomials used in the reduction step in Section 2.

1. NOTATION AND RECOLLECTIONS

1.1. In Sections 1–3, the ground field is \mathbb{C} , the field of complex numbers, and algebraic varieties are equipped with the complex topology. For such a variety X , we denote by $IC(X)$ the Deligne-Goresky-MacPherson complex, and by $\mathcal{IH}^i(X)$ its (intersection) cohomology sheaves (see [2, 8, 12, 16]).

1.2. **The Ehresmann-Bruhat-Chevalley order.** Let S_n denote the symmetric group in n letters. Let us first recall that every permutation $w \in S_n$ is determined by its code $c(w)$, which is the n -tuple $(c_1(w), \dots, c_n(w))$, where $c_i(w)$ equals the number of $j > i$ such that $w(j) < w(i)$. Then $\ell(w)$, the number of inversions of w , equals $\sum_i c_i(w)$.

Let us also recall that the Ehresmann-Bruhat-Chevalley order on S_n , denoted by \leq , can be described in terms of keys of permutations. The r -th key of w , denoted by $K_r(w)$, is the array obtained by arranging the elements of $w([1, r])$ in increasing order: $i_1 < \dots < i_r$. The set of r -keys is endowed with the product order; that is, $(j_1 < \dots < j_r) \leq (i_1 < \dots < i_r)$ if and only if $j_s \leq i_s$ for all s . It is well known that $y \leq w$ if and only if $K_r(y) \leq K_r(w)$ for all r . One deduces, in particular, that if $c_i(w) \geq c_i(y)$ for all i , then $w \geq y$.

Most often, we shall denote a permutation $w \in S_n$ by the sequence (i_1, \dots, i_n) , where $i_r = w(r)$. Yet, the transposition which exchanges i and j is denoted, as usual, by (i, j) . Further, the transposition $(i, i + 1)$ will sometimes be denoted by s_i .

For future reference, let us also record here the following. Given $w \in S_n$ and $i < j$, it is well known and easy to check that $w(i, j) < w$ if and only if $w(i) > w(j)$, and in this case one has

$$(\dagger) \quad \ell(w(i, j)) = \ell(w) - 1 - 2 \#\{k \in]i, j[\mid w(i) > w(k) > w(j)\}.$$

For $y, w \in S_n$, we shall write $w \rightarrow y$ to mean that $w > y$ and $\ell(w) = \ell(y) + 1$.

1.3. Schubert varieties. For any $n \geq 2$, let $\mathcal{F}(n)$ denote the variety of complete flags $V^1 \subset \dots \subset V^{n-1} \subset \mathbb{C}^n$. Let $\{e_1, \dots, e_n\}$ be the standard basis of \mathbb{C}^n , and for every i let \mathbb{C}^i denote the span of e_1, \dots, e_i . The stabilizer of the standard flag $\mathbb{C}^1 \subset \dots \subset \mathbb{C}^{n-1}$ is the subgroup of upper triangular matrices, denoted by B , and $\mathcal{F}(n)$ identifies with $SL_n(\mathbb{C})/B$. For every $w \in S_n$, let \mathbf{V}_w denote the flag whose i -th term is the \mathbb{C} -span of $e_{w(1)}, \dots, e_{w(i)}$. Then the Schubert variety X_w is the closure (taken in the Zariski or, equivalently, complex topology) of the orbit $B\mathbf{V}_w$. It is well known that one has

$$X_w = \{V^1 \subset \dots \subset V^{n-1} \subset \mathbb{C}^n \mid \dim(V^p \cap \mathbb{C}^q) \geq \#\{i \leq p \mid w(i) \leq q\}\}.$$

1.4. The monomial case. The simpler case where $P - 1$ is a monomial $a q^d$ will play an important role in the sequel. In fact, the proof in the general case reduces, at a crucial step, to this case (see 2.5–2.7).

For the convenience of the reader, let us describe explicitly here the pair $w > y$ associated to $1 + a q^d$. Let $n = d + a + 2$ and consider the following elements of S_n :

$$\begin{aligned} w &= (d + 2, d + 3, \dots, n, d + 1, \dots, 3, 1, 2), \\ y &= (1, d + 2, \dots, n - 1, d + 1, \dots, 3, 2, n). \end{aligned}$$

Their codes are the n -tuples: $c(w) = ((d+1)^{a+1}, d, d-1, \dots, 2, 0, 0)$ and $c(y) = (0, d^a, d-1, \dots, 1, 0, 0)$, where m^b denotes a sequence of b terms equal to m . Thus $c_i(w) \geq c_i(y)$, for all i , and one has, therefore, $y < w$ and $\ell(w) - \ell(y) = 2d + a$. (Note that $\ell(w) - \ell(y) - 1 = 2d$ if and only if $a = 1$).

We will prove in Section 3 that $P_{y,w} = 1 + a q^d$. First, we begin with some results in the general case.

2. MAJORATION OF $P_{y,w}$ AND REDUCTION TO THE MONOMIAL CASE

2.1. Let (a_1, \dots, a_d) be a d -tuple of nonnegative integers, with $a_d \neq 0$, and let $P = 1 + a_1 q + \dots + a_d q^d$. Let $|a| = \sum_i a_i$ and $n = d + P(1) + 1 = d + |a| + 2$. For $i = 1, \dots, |a|$, let d_i be the smallest integer s such that $i \leq a_1 + \dots + a_s$. (In other words, the d_i 's are defined by the equality $\sum_{i=1}^{|a|} q^{d_i} = \sum_{s=1}^d a_s q^s$.) Note that $d_i \leq d_{i+1}$ for all i .

Consider now the elements w and y in S_n defined by the following conditions: $w(i) = i + d_i + 1$ for $i \leq |a| + 1$, $w(n) = 2$, and $w(i) > w(i+1)$ for $|a| + 2 \leq i \leq n - 2$, while $y(1) = 1$, $y(i) = w(i - 1)$ for $2 \leq i \leq |a| + 1$, $y(i) = w(i)$ for $|a| + 2 \leq i \leq n - 2$, $y(n - 1) = 2$ and $y(n) = n$. (More explicitly, one has $w(n - 1) = 1$ and $w(n - s) = a_1 + \dots + a_{s-1} + s + 1$, for $s = 2, \dots, d$.)

Therefore, the codes of w and y are the following n -tuples: (here, m^a denotes a sequence of a terms equal to m ; by convention this is the empty sequence if $a = 0$)

$$\begin{aligned} c(w) &= (2^{a_1}, \dots, (d + 1)^{a_d}, d + 1, d, \dots, 2, 0, 0), \\ c(y) &= (0, 1^{a_1}, \dots, d^{a_d}, d - 1, d - 2, \dots, 1, 0, 0). \end{aligned}$$

Thus, $c_i(w) \geq c_i(y)$ for all i . It follows that $w > y$ and that $\ell(w) - \ell(y) = |a| + 2d$. The main result of this paper is the following.

Theorem. *One has $P_{y,w} = 1 + a_1 q + \dots + a_d q^d$.*

The proof will be given in a number of steps.

2.2. First, one observes that the Schubert variety X_w is the closed subvariety of $\mathcal{F}(n)$ consisting of those flags $V^1 \subset \dots \subset V^{n-1} \subset \mathbb{C}^n$ such that $V^i \subset \mathbb{C}^{i+d_i+1}$, for $i = 1, \dots, |a|$, and $\mathbb{C}^1 \subset V^{n-1}$.

The union of the orbits $B\mathbf{V}_z$, for $z \in [y, w]$, is an open subset of X_w , that we shall denote by $C_{[y,w]}$. On the other hand, let Z_w be the variety consisting of pairs of flags $V^1 \subset \dots \subset V^{n-1}$ and $U^2 \subset \dots \subset U^{|a|+1}$ in \mathbb{C}^n which satisfy the relations of inclusion indicated in the diagram below:

$$\left\{ \begin{array}{cccccccc} & V^1 & \subset & V^2 & \subset & \dots & \subset & V^{|a|} & \subset \dots \subset & V^{n-2} \\ \mathbb{C}^1 \subset & \cap & & \cap & & & & \cap & & \cap \\ & U^2 & \subset & U^3 & \subset & \dots & \subset & U^{|a|+1} & \subset & V^{n-1} \\ & \cap & & \cap & & & & \cap & & \\ & \mathbb{C}^{d_1+2} & \subset & \mathbb{C}^{d_2+3} & \subset & \dots & \subset & \mathbb{C}^{n-1} & & \end{array} \right\},$$

and let π denote the natural projection $Z_w \rightarrow X_w$.

It is easily seen that Z_w is smooth, since it is obtained as an iteration of locally trivial fibrations with fibres isomorphic to projective spaces. Moreover, π is a resolution of singularities of X_w . In fact, π is proper, and one has the following lemma. Let X_w^1 denote the closed, B -stable, subvariety of X_w defined by the condition that $\mathbb{C}^1 \subseteq V^{|a|}$.

Lemma. *π is an isomorphism over the complement of X_w^1 .*

Proof. Set $\Omega = X_w \setminus X_w^1$. Observe that if $\pi(V^\bullet, U^\bullet) \in \Omega$, then $U^{i+1} = V^i + \mathbb{C}^1$ for all $i \leq |a|$. Thus, π is bijective over Ω . Since π is proper and X_w normal ([15, 14, 1]), it follows from Zariski's main theorem that π is an isomorphism over Ω .

We will see later that π is not a small resolution if $d \geq 2$, and is not semi-small if $d \geq 3$ (see 3.2). Yet, using the decomposition theorem ([2]), plus ideas of Deodhar ([6]) and a result of Irving ([9]), we will show that, thanks to the particular form of w and y , one can obtain enough information to decompose explicitly $j^* R\pi_*(\mathbb{C})$ as a direct sum of shifted Deligne-Goresky-MacPherson complexes, where j denotes the inclusion $C_{[y,w]} \hookrightarrow X_w$.

2.3. Set $q = t^2$ and $H_{z,\pi}(t) = \sum_{i \geq 0} \dim(R^i \pi_* \mathbb{C})_{\mathbf{V}_z} t^i$, for any $z \leq w$. On the one hand, since π is proper, one has $(R^i \pi_* \mathbb{C})_{\mathbf{V}_z} \cong H^i(\pi^{-1}(\mathbf{V}_z), \mathbb{C})$ (see e.g. [10, Th.III.6.2]). On the other hand, since π is B -equivariant and the B -orbits in X_w are simply connected, the decomposition theorem ([2]) implies that

$$R\pi_*(\mathbb{C}[\ell(w)]) \cong IC(X_w) \oplus \bigoplus_{v < w} E_v \otimes IC(X_v),$$

where the E_v are finite dimensional graded \mathbb{C} -vector spaces with the property that $\dim E_v^i = \dim E_v^{-i}$, for all i . Setting $E_v(t) = \sum_i (\dim E_v^i) t^i$, this is equivalent to

the fact that $E_v(t^{-1}) = E_v(t)$. Moreover, by [12, 16], one has, for all $z \leq w$,

$$\sum_i \dim \mathcal{I}\mathcal{H}^i(X_w)_{\mathbf{V}_z} t^i = t^{-\ell(w)} P_{z,w}(q).$$

One deduces that, for all $z \leq w$,

$$(1) \quad H_{z,\pi}(t) = P_{z,w}(q) + \sum_{z \leq v < w} t^{\ell(w)-\ell(v)} E_v(t) P_{z,v}(q).$$

2.4. For all $i \leq |a|$, let γ_i be the (d_i+1) -cycle which transforms $(i, n-d_i, \dots, n-1)$ into $(n-1, i, \dots, n-2)$ and let $v_i = w\gamma_i$.

- Proposition.** a) *The X_{v_i} , for $i = 1, \dots, |a|$, are exactly the irreducible components of X_w^1 . Further, one has $v_i \geq y$ and $\ell(v_i) = \ell(w) - d_i$ for all i .*
 b) *For each i , the fibre $\pi^{-1}(\mathbf{V}_{v_i})$ is isomorphic to projective space \mathbb{P}^{d_i-1} and, therefore, $t^{\ell(w)-\ell(v_i)} E_{v_i}(t) = q + \dots + q^{d_i-1}$ and \mathbf{V}_{v_i} is a smooth point of X_w .*
 c) *One has $H_{y,\pi}(t) = 1 + \sum_{i=1}^{|a|} (q + \dots + q^{d_i}) = 1 + \sum_{s=1}^d a_s (q + \dots + q^s)$.*
 d) *\mathbf{V}_y is a smooth point in each X_{v_i} .*

Proof. First, one checks that $v_i(i) = 1$ and $v_i(j) = w(j)$, for all $j \in [1, |a| + 1] \setminus \{i\}$. From this one deduces that $X_y \subseteq X_{v_i} \subseteq X_w^1$, and that $v_i \not\leq v_j$ if $i \neq j$ (by comparing the i -th and j -th keys). Further, let X_z be an irreducible component of X_w^1 and let $i = z^{-1}(1)$. Since $\mathbb{C}^1 \subseteq \mathbf{V}_z^{|a|}$, one has $i \leq |a|$. From the definition of v_i , again, this implies that $z \leq v_i$. It follows that the X_{v_i} are exactly the irreducible components of X_w^1 . (In fact, the orbit $B\mathbf{V}_{v_i}$ is defined inside X_w by the condition that $\mathbb{C}^1 \subseteq V^i$ but $\mathbb{C}^1 \not\subseteq V^{i-1}$.) Further, using 1.2(†) plus the fact that, for $j \in]i, n-d_i[$ one has $w(j) \geq w(n-d_i-1) > w(i)$, one obtains that

$$w \rightarrow ws_{n-2} \rightarrow \dots \rightarrow ws_{n-2} \cdots s_{n-d_i} \rightarrow ws_{n-2} \cdots s_{n-d_i}(i, n-d_i) = v_i.$$

Therefore, $\ell(v_i) = \ell(w) - d_i$. This proves assertion a).

b) Let us now describe the fibre $\pi^{-1}(\mathbf{V}_{v_i})$. First, note that $\mathbb{C}^{i+d_i+1} \cap \mathbf{V}_{v_i}^{n-1}$ has dimension $d_i + i$. Second, it is easily seen that the natural projection

$$p_i : \pi^{-1}(\mathbf{V}_{v_i}) \rightarrow \mathbb{P}((\mathbb{C}^{i+d_i+1} \cap \mathbf{V}_{v_i}^{n-1})/\mathbf{V}_{v_i}^i) \cong \mathbb{P}^{d_i-1}, \quad (U^2, \dots, U^{a+1}) \mapsto U^{i+1},$$

is surjective. But, clearly, for $j < i$, one has $U^{j+1} = \mathbb{C}^1 \oplus \mathbf{V}_{v_i}^j$ (since $\mathbb{C}^1 \not\subseteq \mathbf{V}_{v_i}^{i-1}$). Further, for every $j \in [i+1, |a|]$, one has $\mathbf{V}_{v_i}^j \not\subseteq \mathbb{C}^{d_{j-1}+j}$, whence $\mathbf{V}_{v_i}^j \neq U^j$, and therefore $U^{j+1} = U^j + \mathbf{V}_{v_i}^j$. This shows that p_i is bijective, whence an isomorphism by Zariski's main theorem. This proves the first part of assertion b). On the other hand, since $H_{z,\pi} = 1 = P_{z,w}$ for all $z \in]v_i, w]$, by Lemma 2.2 and assertion a), then (1) gives

$$H_{v_i,\pi}(t) = P_{v_i,w}(q) + t^{d_i} E_{v_i}(t).$$

The second part of b) is then an immediate consequence of this formula. Indeed, this is clear if $d_i = 1$, so we may suppose that $d_i \geq 2$. Then, since $H_{v_i,\pi} - P_{v_i,w}$ has nonnegative coefficients and $\deg P_{v_i,w} \leq (d_i - 1)/2 < d_i - 1$, it follows that $P_{v_i,w} = 1 + \dots + q^{s-1}$, for some $s \in [1, d_i - 1]$. Therefore,

$$E_{v_i}(t) = t^{-d_i}(q^s + \dots + q^{d_i-1}) = t^{2s-d_i} + \dots + t^{d_i-2},$$

and since $E_{v_i}(t^{-1}) = E_{v_i}(t)$, this implies that $s = 1$, whence $t^{d_i} E_{v_i}(t) = q + \dots + q^{d_i-1}$ and $P_{v_i,w} = 1$. Thus, by [5], \mathbf{V}_{v_i} is a smooth point of X_w .

Let us prove assertion c). The fibre $\pi^{-1}(\mathbf{V}_y)$ is the variety of flags $U^2 \subset \dots \subset U^{|a|+1}$ which satisfy the following relations of inclusion:

$$\left\{ \begin{array}{ccccccc} \mathbb{C}^1 & \subset \dots \subset & \mathbf{V}_y^i & \subset \dots \subset & \mathbf{V}_y^{|a|} & \subset \dots \subset & \mathbf{V}_y^{n-2} \\ \cap & & \cap & & \cap & & \cap \\ U^2 & \subset \dots \subset & U^{i+1} & \subset \dots \subset & U^{|a|+1} & & \\ \cap & & \cap & & \cap & & \\ \mathbb{C}^{d_1+2} & \subset \dots \subset & \mathbb{C}^{i+d_i+1} & \subset \dots \subset & \mathbb{C}^{n-1} & = & \mathbf{V}_y^{n-1} \end{array} \right\}.$$

Observe that, for $i = 2, \dots, |a|$, \mathbf{V}_y^i is the \mathbb{C} -span of $e_1, e_{d_1+2}, \dots, e_{d_{i-1}+i}$ and, therefore, one has $\mathbf{V}_y^i \subseteq \mathbb{C}^{d_{i-1}+i}$ but $\mathbf{V}_y^i \not\subseteq \mathbb{C}^{d_{i-1}+i-1}$.

We claim that $\pi^{-1}(\mathbf{V}_y)$ is the disjoint union of locally closed subvarieties $Y_1, \dots, Y_{|a|}$ (with Y_1 open and $Y_{|a|}$ closed), such that, for $i < |a|$, Y_i is isomorphic to $\mathbb{P}^{d_i} \setminus \{pt\}$, whereas $Y_{|a|} \cong \mathbb{P}^d$. Of course, this claim immediately implies assertion c).

Let Y_1 be the open subvariety of $\pi^{-1}(\mathbf{V}_y)$ consisting of those flags $(U^2, \dots, U^{|a|+1})$ such that $U^2 \neq \mathbf{V}_y^2$. Let us prove that $Y_1 \cong \mathbb{P}^{d_1} \setminus \{pt\}$. Let φ denote the restriction to Y_1 of the natural projection from $\pi^{-1}(\mathbf{V}_y)$ to $\mathbb{P}(\mathbb{C}^{d_1+2}/\mathbb{C}^1) \cong \mathbb{P}^{d_1}$, which sends $(U^2, \dots, U^{|a|+1})$ to U^2 . Clearly, φ maps Y_1 onto $\mathbb{P}^{d_1} \setminus \{pt\}$. Thus, by Zariski's main theorem, it suffices to prove that φ is injective. The hypothesis that $U^2 \neq \mathbf{V}_y^2$ implies that $U^3 = \mathbf{V}_y^2 + U^2$, whence $U^3 \subseteq \mathbb{C}^{d_1+2}$. Thus, since $\mathbf{V}_y^3 \not\subseteq \mathbb{C}^{d_2+2}$, it follows that $\mathbf{V}_y^3 \neq U^3$. Since $\mathbf{V}_y^i \subseteq \mathbb{C}^{d_{i-1}+i}$ but $\mathbf{V}_y^i \not\subseteq \mathbb{C}^{d_{i-1}+i-1}$, for all $i \in [2, |a|]$, one can repeat this argument to conclude that $U^i \neq \mathbf{V}_y^i$ and $U^{i+1} = \mathbf{V}_y^i + U^2$, for all $i \in [2, |a|]$. This proves that φ is bijective, and hence that $Y_1 \cong \mathbb{P}^{d_1} \setminus \{pt\}$.

Now, for $i = 2, \dots, |a|$, let Y_i denote the locally closed subvariety consisting of those flags $(U^2, \dots, U^{|a|+1})$ such that $U^{j+1} = \mathbf{V}_y^{j+1}$ for $j < i$ and $U^{i+1} \neq \mathbf{V}_y^{i+1}$ (this last condition being omitted if $i = |a|$). Using an argument similar to the above, one deduces that $Y_i \cong \mathbb{P}^{d_i} \setminus \{pt\}$, for $i < |a|$, whereas $Y_{|a|} \cong \mathbb{P}^d$, and that $\pi^{-1}(\mathbf{V}_y)$ is the disjoint union of the Y_i . It follows that

$$H_{y,\pi} = 1 + \sum_{i=1}^{|a|} (q + \dots + q^{d_i}) = 1 + \sum_{s=1}^d a_s (q + \dots + q^s).$$

This proves assertion c).

Finally, let us prove assertion d). From the definition of v_i , one derives easily that X_{v_i} admits a resolution of singularities $\phi : Z_{v_i} \rightarrow X_{v_i}$, where Z_{v_i} is the variety of pairs of flags $((V^k)_{1 \leq k < n}, (U^j)_{2 \leq j < i})$ satisfying the following relations of inclusion:

$$\left\{ \begin{array}{ccccccc} & V^1 & \subset \dots \subset & V^{i-2} & \subset & V^{i-1} & \\ \cap & & & \cap & & \cap & \\ \mathbb{C}^1 \subset & U^2 & \subset \dots \subset & U^{i-1} & \subset & V^i & \subset & V^{i+1} & \subset \dots \subset & V^{|a|} \\ \cap & & & \cap & & \cap & & \cap & & \\ & \mathbb{C}^{d_1+2} & \subset \dots \subset & \mathbb{C}^{d_{i-2}+i-1} & \subset & \mathbb{C}^{d_{i-1}+i} & \subset & \mathbb{C}^{d_{i+1}+i+2} & \subset \dots \subset & \mathbb{C}^{n-1} \end{array} \right\}.$$

It is easily seen that ϕ is bijective over the open subvariety Ω of X_{v_i} defined by the condition that $\dim(V^i \cap \mathbb{C}^{d_j+j+1}) \leq j+1$, for $j = 1, \dots, i-2$. In fact, this condition implies that $U^{j+1} = \mathbb{C}^{d_j+j+1} \cap V^i$ for all $j \in [1, i-2]$. Since ϕ is proper and X_{v_i} normal, ϕ is an isomorphism over Ω , by Zariski's main theorem. But, clearly, \mathbf{V}_y

belongs to Ω . Therefore, \mathbf{V}_y is a smooth point of X_{v_i} . This completes the proof of Proposition 2.4.

2.5. It follows from the previous proposition that

$$H_{y,\pi} - \sum_{i=1}^{|a|} t^{\ell(w)-\ell(v_i)} E_{v_i}(t) P_{y,v_i} = P.$$

Therefore, by (1), $P - P_{y,w}$ has nonnegative coefficients and the sought-for equality $P_{y,w} = P$ is equivalent to the vanishing of E_v for all $v \in [y, w[\setminus\{v_1, \dots, v_{|a|}\}]$.

On the other hand, by [9, Cor. 4], $P_{y,w} - P_{z,w}$ has nonnegative coefficients, for all $z \in [y, w]$. Therefore, in order to prove the equality $P_{y,w} = P$, it suffices to exhibit, for all $s \in [1, d]$, an element z_s in $[y, w]$ such that $P_{z_s,w} = 1 + a_s q^s$. We construct z_s as follows.

Let $f_s = \sum_{i < s} a_i$ and $z_s = w(n-1, n-s-1)(n-1, n)\sigma_s$, where σ_s is the (a_s+1) -cycle which transforms $(f_s+1, \dots, f_s+a_s, n-s-1)$ into $(n-s-1, f_s+1, \dots, f_s+a_s)$. We shall prove that

$$(2) \quad P_{z_s,w} = 1 + a_s q^s.$$

As observed above, this will complete the proof of the theorem.

2.6. Let $s \in [1, d]$. Set $h_s = w(n-s-1) = f_{s+1} + s + 2$. One observes that $z_s(i) = w(i)$ if $i \leq f_s$, and $z_s^{-1}(j) = w^{-1}(j)$ if $j > h_s$. Let $F_s = [f_s + 1, n] \cap w^{-1}[1, h_s] = [f_s + 1, n] \cap z_s^{-1}[1, h_s]$. From the definition of w , it follows that $F_s = [f_s + 1, f_s + a_s] \sqcup [n-s-1, n]$. In particular, $\#F_s = a_s + s + 2 := m_s$. Let ϕ (resp. ψ) be the unique increasing bijection from $[1, m_s]$ to F_s (resp. to $w(F_s) = z_s(F_s)$). Set $\bar{w} = \psi^{-1} \circ w|_{F_s} \circ \phi$, and define \bar{z}_s similarly. (Note that \bar{w} depends on s ; in order not to burden the notation we omit to indicate this.) Then one has the following proposition (see Section 4 for a geometric interpretation of this result in a more general setting).

Proposition. *One has $P_{z_s,w} = P_{\bar{z}_s,\bar{w}}$.*

Proof. Let us first recall the following known lemma, which is obtained easily by comparing the element C'_u defined in [11, (1.1.c)] with $C'_v C'_{x'}$.

Lemma. *Let W be a Coxeter group and W' a parabolic subgroup. Let $x \leq u$ in W and suppose that $xW' = uW'$ (resp. $W'x = W'u$). Let v be the unique minimal element in this coset and set $u' = v^{-1}u$, $x' = v^{-1}x$ (resp. $u' = uv^{-1}$ and $x' = xv^{-1}$). Then $P_{x,u} = P_{x',u'}$.*

Now, let us show that the proposition is a consequence of the lemma. Let us write, for simplicity, f, z, h, F, m instead of f_s, z_s, h_s, F_s, m_s . Let us denote by W_I the subgroup of permutations of $[f+1, n]$. Then $wW_I = zW_I$. Let v be the unique minimal element in this coset, and $w_1 = v^{-1}w$, $z_1 = v^{-1}z$. By the lemma, one has $P_{z,w} = P_{z_1,w_1}$.

Observe that the restriction of v to the interval $[f+1, n]$ is precisely the unique increasing bijection from $[f+1, n]$ to $w([f+1, n])$. Thus, recalling that F denotes $[f+1, n] \cap w^{-1}[1, h]$, one deduces that v^{-1} induces an increasing bijection from $w(F)$ to an interval $[f+1, g]$, for some $g \leq n$. Denoting by η the unique increasing bijection from $[1, m]$ to $[f+1, g]$, it follows that $\psi^{-1} = \eta^{-1} \circ v^{-1}$. Therefore, $\bar{w} = \eta^{-1} \circ v^{-1} \circ w|_F \circ \phi$.

Further, denoting by W_K the subgroup of permutations of $[f + 1, g]$, the hypotheses also imply that $W_K w_1 = W_K z_1$. Let θ be the unique minimal element in this coset, and let $w_2 = w_1\theta^{-1}$ and $z_2 = z_1\theta^{-1}$. On the one hand, by the lemma again, one has $P_{z_1, w_1} = P_{z_2, w_2}$ and, therefore, $P_{z, w} = P_{z_2, w_2}$. On the other hand, one observes that θ^{-1} is the unique increasing bijection from $[f + 1, g]$ to F . One deduces that $\phi = \theta^{-1} \circ \eta$, and hence that $\bar{w} = \eta^{-1} \circ v^{-1} \circ w|_F \circ \theta^{-1} \circ \eta$.

But $v^{-1} \circ w|_F \circ \theta^{-1}$ identifies with w_2 , and therefore one obtains that $\bar{w} = \eta^{-1} \circ w_2 \circ \eta$, and similarly for \bar{z} and z_2 . Since the map $x \mapsto \eta^{-1} \circ x \circ \eta$ induces an order-preserving isomorphism of Coxeter groups $S_{[1, m]} \cong S_{[f+1, g]}$, it follows that $P_{z_2, w_2} = P_{\bar{z}, \bar{w}}$. The proposition is proved.

2.7. Again, let $s \in [1, d]$ be fixed. It follows from the definitions that the codes of \bar{w} and \bar{z}_s are

$$c(\bar{w}) = ((s+1)^{a_s+1}, s, s-1, \dots, 2, 0, 0) \quad \text{and} \quad c(\bar{z}_s) = (0, s^{a_s}, s-1, \dots, 1, 0, 0).$$

Thus, $\bar{z}_s < \bar{w}$ is the pair associated to the polynomial $1 + a_s q^s$. Therefore, in order to complete the proof of the theorem in general, it suffices to do so in the special case where $P - 1$ is a monomial, since, by Proposition 2.6, this will imply the sought-for equality 2.5.(2).

3. COMPLETION OF THE PROOF IN THE MONOMIAL CASE

In this section, let a, d be positive integers, $n = a + d + 2$, and $P = 1 + a q^d$. Let $w > y$ be the elements of S_n associated to P in 2.1 (see also 1.4). In fact, one has $w(i) = i + 1 + d$ for $i \leq a + 1$, $w(i) = n + 1 - i$ for $i \in [a + 2, n - 2]$ (this condition being void if $d = 1$), $w(n - 1) = 1$ and $w(n) = 2$, whereas $y(1) = 1$, $y(i) = w(i - 1)$ for $i \in [2, a + 1]$, $y(i) = w(i)$ for $i \in [a + 2, n - 2]$, $y(n - 1) = 2$ and $y(n) = n$. Further, $\ell(w) - \ell(y) = 2d + a$.

3.1. The notation and results of the previous section apply in this special case. In particular, with notation as in Proposition 2.4, we know already that $H_{y, \pi}(t) = 1 + a(q + \dots + q^d)$ and that $t^d E_{v_i}(t) = q + \dots + q^{d-1}$, for $i = 1, \dots, a$. Therefore, one has

$$1 + a q^d = P_{y, w} + \sum_{\substack{v \in [y, w[\\ v \notin \{v_1, \dots, v_a\}}} t^{\ell(w) - \ell(v)} E_v(t) P_{y, v}.$$

Further, using the fact that the terms in the right-hand side have nonnegative coefficients and that $P_{y, w}$ and $P_{y, v}$ have constant term equal to 1, one deduces that if $E_v(t) \neq 0$ for some v in the above sum, then, necessarily, $P_{y, v} = 1$ and $t^{\ell(w) - \ell(v)} E_v(t) = c q^d$ for some $c \in [1, a]$. Since $E_v(t) = E_v(t^{-1})$, this implies that $\ell(w) - \ell(v) = 2d$. Therefore, the sought-for equality $P_{y, w} = 1 + a q^d$ is a consequence of the following proposition.

Proposition. *For all $v \in [y, w]$ such that $\ell(w) - \ell(v) = 2d$, one has $\deg_t H_{v, \pi} < 2d$, and hence $E_v = 0$. In fact, one has more precisely $H_{v, \pi} = 1 + b(q + \dots + q^{d-1})$, where $b = \#\{i \in [1, a] \mid v \leq v_i\}$.*

Proof. Recall first that, if $X_v \not\subseteq X_w^1$, that is, if $v^{-1}(1) > a$, then $H_{v, \pi} = 1$, by Lemma 2.2. Thus, we may assume that $v^{-1}(1) \leq a$. We will need the following lemma.

Lemma. *Let $v \in [y, w]$ be such that $v^{-1}(1) = i \leq a$. If $\mathbb{C}^{d+i+1} \subseteq \mathbf{V}_v^{n-1}$, then $\ell(w) - \ell(v) \geq 2d + 1$.*

Proof. Since $v \leq w$, then $v(j) \leq j + d + 1$, for $j = 1, \dots, a$. Further, we assume that $v^{-1}(1) = i \leq a$. On the other hand, the hypothesis $\mathbb{C}^{d+i+1} \subseteq \mathbf{V}_v^{n-1}$ is equivalent to $v(n) > d + i + 1$. Thus, one has $v(n) = d + j + 1$ for some $j \in [i + 1, a + 1]$.

One then deduces that v is smaller than the permutation $w_{i,j}$ defined by $w_{i,j}(k) = w(k)$ if $k \notin \{i, j, n - 1, n\}$, while $w_{i,j}(i) = 1$, $w_{i,j}(j) = d + i + 1$, $w_{i,j}(n - 1) = 2$ and $w_{i,j}(n) = d + j + 1$. Let $u_{i,j} = v_i(j, a + 2)$. Using 1.2(†), it is easily seen that

$$v_i \rightarrow u_{i,j} \rightarrow u_{i,j}s_{a+2} \rightarrow \cdots \rightarrow u_{i,j}s_{a+2} \cdots s_{n-1} = w_{i,j}.$$

Thus, $\ell(v_i) - \ell(w_{i,j}) = d + 1$ and hence $\ell(w) - \ell(w_{i,j}) = 2d + 1$. The lemma is proved.

Let us now prove the proposition. Let $v \in W$ be such that $y \leq v \leq w$ and $v^{-1}(1) := i_1 \leq a$. These hypotheses, together with the fact that $w([1, j]) = [d + 2, d + j + 1]$ and $y([1, j]) = \{1\} \cup [d + 2, d + j]$ for $j = 1, \dots, a$, imply that there exists a sequence $i_1 < \cdots < i_b$ such that $v(j) = j + d + 1$ for $j \in [1, a] \setminus \{i_1, \dots, i_b\}$ while $v(i_r) = i_{r-1} + d + 1$ for $r = 2, \dots, b$ (this condition being empty if $b = 1$). In particular, one has $v([1, a]) = \{1\} \cup [d + 2, n - 1] \setminus \{i_b + d + 1\}$. Set $J = \{i_1, \dots, i_b\}$. Then, clearly, $J = \{j \in [1, a] \mid \mathbf{V}_v^j \subseteq \mathbb{C}^{d+j}\}$ and one has, for every $j \in [1, a]$:

$$(*) \quad \dim(\mathbf{V}_v^a \cap \mathbb{C}^{d+j+1}) = \begin{cases} j + 1 & \text{if } j < i_b; \\ j & \text{if } j \geq i_b. \end{cases}$$

Further, it follows from the definition of the v_i 's (see 2.4) that $v \leq v_j$ if $j \in J$; conversely, since $\mathbf{V}_{v_i}^i \subseteq \mathbb{C}^{d+i}$, then $v \not\leq v_i$ if $i \notin J$. Therefore, one also has $J = \{j \in [1, a] \mid v \leq v_j\}$.

Suppose, moreover, that $\ell(w) - \ell(v) \leq 2d$. Then, by the previous lemma, we have $\mathbb{C}^{d+i+1} \not\subseteq \mathbf{V}_v^{n-1}$ and hence

$$(**) \quad \dim(\mathbb{C}^{d+j+1} \cap \mathbf{V}_v^{n-1}) = d + j, \quad \forall j \in [i_1, a].$$

Now, we claim that the fibre $\pi^{-1}(\mathbf{V}_v)$ is the disjoint union of locally closed subvarieties Y_{i_1}, \dots, Y_{i_b} (with Y_{i_1} open and Y_{i_b} closed), such that Y_{i_r} is isomorphic to $\mathbb{P}^{d-1} \setminus \{pt\}$ for $r < b$, whereas $Y_{i_b} \cong \mathbb{P}^{d-1}$.

In fact, set $U^1 = \mathbb{C}^1$ and for every $j = 2, \dots, a + 1$, let Y_j denote the locally closed subvariety of $\pi^{-1}(\mathbf{V}_v)$ consisting of those flags (U^\bullet) such that $U^{j-1} \subseteq \mathbf{V}_v^a$ and $U^j \not\subseteq \mathbf{V}_v^a$. Then the Y_j 's are clearly disjoint and, since $U^1 = \mathbb{C}^1 \subseteq \mathbf{V}_v^a$ but $U^{a+1} \not\subseteq \mathbf{V}_v^a$, they form a partition of $\pi^{-1}(\mathbf{V}_v)$.

Next, observe that if $j \notin J$, then $\mathbf{V}_v^j \neq U^j$ (since $\mathbf{V}_v^j \not\subseteq \mathbb{C}^{j+d}$) and hence $U^{j+1} = U^j + \mathbf{V}_v^j$. Therefore, $Y_j = \emptyset$ if $j \notin J$. Now, let us prove that the Y_{i_r} , for $r = 1, \dots, b$, are as we claimed above.

Let $r \in [1, b]$. Suppose that $(U^\bullet) \in Y_{i_r}$. Then, clearly, one has $U^{j+1} = U^{i_r+1} + \mathbf{V}_v^j$ for $j \in [i_r, a]$. Secondly, one has $U^{i_r} = \mathbf{V}_v^{i_r}$ (for otherwise one would have $U^{i_r+1} = U^{i_r} + \mathbf{V}_v^{i_r} \subseteq \mathbf{V}_v^a$) and hence, using (*), one obtains that $U^j = U^{i_r} \cap \mathbb{C}^{j+d}$ for $j \leq i_r$. Conversely, this shows that the projection $p_{i_r} : (U^\bullet) \mapsto U^{i_r}$, from Y_{i_r} to the projective space of $(\mathbb{C}^{i_r+1+d} \cap \mathbf{V}_v^{n-1})/\mathbf{V}_v^{i_r}$ is a bijection onto its image.

Further, since $\dim(\mathbb{C}^{i_r+1+d} \cap \mathbf{V}_v^{n-1})/\mathbf{V}_v^{i_r} = d$, by (**), whereas

$$\dim(\mathbb{C}^{i_r+1+d} \cap \mathbf{V}_v^a)/\mathbf{V}_v^{i_r} = \begin{cases} 1 & \text{for } r < b; \\ 0 & \text{for } r = b, \end{cases}$$

by (*), it follows that the image of p_{i_r} is $\mathbb{P}^{d-1} \setminus \{pt\}$ when $r < b$, and \mathbb{P}^{d-1} when $r = b$. The claim then follows, by Zariski's main theorem, and one deduces that $H_{z,\pi}(q) = 1 + b(q + \dots + q^{d-1})$.

This completes the proof of Proposition 3.1 and, therefore, of Theorem 2.1.

Remark. Along the same lines as above, one can prove, using [5], that the $w_{i,j}$ defined in the proof of Lemma 3.1 (for $1 \leq i < j \leq a+1$) are exactly the irreducible components of the singular locus of X_w , and that the corresponding Kazhdan-Lusztig polynomials are all equal to $1 + q^d$. We leave this to the interested reader.

3.2. Let us now return to the general case; that is, let $y < w$ be as in 2.1. Since Theorem 2.1 is proved, then, by the discussion at the beginning of 2.5, we have also shown that $E_v = 0$ for all $v \in [y, w[\setminus \{v_1, \dots, v_{|a|}\}]$. Therefore, denoting by j the inclusion of the open subset $C_{[y,w]}$ in X_w , one obtains that

$$j^* R\pi_*(\mathbb{C}[\ell(w)]) \cong j^* IC(X_w) \oplus \bigoplus_{i=1}^{|a|} H^\bullet(\mathbb{P}^{d_i-2})[d_i - 2] \otimes j^* IC(X_{v_i}).$$

Note that $H^\bullet(\mathbb{P}^{d-2})[d - 2]$ equals 0 if and only if $d = 1$, and equals 0 in degrees $\neq 0$ if and only if $d \leq 2$. Thus π is small (resp. semi-small) over $C_{[y,w]}$ if and only if $d = 1$ (resp. $d \leq 2$).

3.3. Let us conclude this section with the following remark. First, if W_1, W_2 are Coxeter groups, $W = W_1 \times W_2$, and $y_i \leq w_i$ in W_i , for $i = 1, 2$, then it is known (and easy to check) that $P_{y_1 y_2, w_1 w_2} = P_{y_1, w_1} P_{y_2, w_2}$.

Now, let $P_1, P_2 \in 1 + q\mathbb{N}[q]$ and let $P = P_1 P_2$. Set $a_i = P_i(1) - 1$, $a = a_1 + a_2$ and $d = \deg P$. Then the pair $y_P < w_P$ belongs to $S_{n(P)}$, where $n(P) = 1 + d + P(1) = 2 + d + a + a_1 a_2$, and satisfies $\ell(w_P) - \ell(y_P) = 2d + a + a_1 a_2$. On the other hand, let $n'(P) = n(P_1) + n(P_2) = 4 + d + a$. Then, the elements $y'_P := y_{P_1} y_{P_2}$ and $w'_P := w_{P_1} w_{P_2}$ belong to $S_{n(P_1)} \times S_{n(P_2)} \subset S_{n'(P)}$ and, by the above, one has $P_{y'_P, w'_P} = P$. This is an improvement on the general construction, in the sense that $\ell(w'_P) - \ell(y'_P) = 2d + a < \ell(w_P) - \ell(y_P)$.

4. A FOCALIZATION THEOREM

In this section, which is independent of the previous ones, we study certain open neighbourhoods in Schubert varieties and prove an isomorphism which provides, in a more general setting, a geometric explanation for the equality in Proposition 2.6.

4.1. Let k be an algebraically closed field of arbitrary characteristic, G a connected semi-simple algebraic group over k , T a maximal torus inside a Borel subgroup B , W the Weyl group, R the root system, and Δ (resp. R^+) the set of simple (resp. positive) roots associated with B . Let U (resp. U^-) be the unipotent radical of B (resp. of the opposite Borel subgroup). For $w \in W$, we denote by e_{wB} the point wB/B in G/B and by C_w its B -orbit (Bruhat cell). Then the Schubert variety X_w is the closure of C_w . Let \leq denote the Bruhat-Chevalley order on W .

Suppose that $y \leq w$. Set $\mathcal{V}_{y,w} = y(U^-)e_{yB} \cap X_w$. This is a T -stable, open affine neighbourhood of y in X_w . In fact, since e_{yB} is the unique closed T -orbit in $y(U^-)e_{yB}$, one can show that $\mathcal{V}_{y,w}$ is the unique T -stable, open affine neighbourhood of y in X_w , see [4, Lemma 2.4]. Set $\mathcal{N}_{y,w} = (y(U^-) \cap U^-)e_{yB} \cap X_w$. This is a closed, T -stable, subvariety of $\mathcal{V}_{y,w}$. Recall from [11, Lemma A4.(b)] the isomorphism $\mathcal{V}_{y,w} \cong C_y \times \mathcal{N}_{y,w}$.

We shall denote by $C_{[y,w]}$ (resp. $B[y,w]B$) the union of the Bruhat cells C_z (resp. of the double cosets BzB), for $z \in [y,w]$. This is an open subvariety of X_w (resp. of \overline{BwB} , the closure of BwB in G). Further, it is well known, and easy to verify, that $\mathcal{V}_{y,w} \subseteq C_{[y,w]}$.

4.2. If I is a subset of Δ , let R_I be the corresponding root system, and W_I and P_I the corresponding parabolic subgroups of W and G .

Let $w \in W$ and let v be the unique minimal element in the coset wW_I . It is well known that the map $z \mapsto vz$ induces an isomorphism of posets $W_I \cong wW_I$, and that $\ell(vz) = \ell(v) + \ell(z)$, for all $z \in W_I$. We shall need to describe similarly the intersection $wW_I \cap W_Jw$, where J is a second, arbitrary, subset of Δ .

Proposition. *Let I, J be subsets of Δ and let $w \in W$. Let v be the minimal element in the coset wW_I , let $K = I \cap v^{-1}R_J$ and $w_1 = v^{-1}w$, and let θ be the minimal element in the coset W_Kw_1 . Then the map $\psi : W_K \rightarrow wW_I \cap W_Jw$, $z \mapsto vz\theta$ is an isomorphism of posets. Further, one has $\ell(vz\theta) = \ell(v) + \ell(z) + \ell(\theta)$, for all $z \in W_K$.*

This is certainly known; yet since we are not aware of a reference, we give a proof, which begins with the following lemma. Set $v^{-1}W_Jv = v^{-1}(W_J)$.

Lemma. *One has $W_I \cap v^{-1}(W_J) = W_K$.*

Proof. It suffices to prove that $v(W_I) \cap W_J$ is generated by the reflections s_β , for $\beta \in v(I) \cap R_J$. Denote by H the subgroup generated by these reflections. Let $w \in v(W_I) \cap W_J$ and let $w = vs_{\alpha_1} \cdots s_{\alpha_r}v^{-1}$ be a reduced decomposition in $v(W_I)$ (that is, $\alpha_i \in I$). We prove by induction on r that $v\alpha_i \in R_J$ for all i , and hence that $w \in H$. This is immediate if $r = 1$, so assume that $r \geq 2$. Set $N(w) = \{\gamma \in R^+ \mid w\gamma \in R^-\}$. Since $w \in W_J$, then $N(w) \subseteq R_J$. On the other hand, by the assumption on v , one has $v(R_I^\pm) \subseteq R^\pm$, and it follows that $v\alpha_r \in R^+$ and $wv\alpha_r \in R^-$. One deduces that $v\alpha_r \in R_J$. Therefore, $ws_{v\alpha_r} \in H$ and we are done by induction.

Let us then derive the proposition. First, one sees immediately that ψ is an order isomorphism onto its image, and that the second assertion holds. Further, let $x \in wW_I \cap W_Jw$ and $x_1 = v^{-1}x$. Then $vx_1 \in W_Jv w_1$ and hence $x_1 w_1^{-1} \in v^{-1}(W_J) \cap W_I = W_K$. It follows that ψ is surjective, and the proposition is proved.

Remark. The statement of the proposition is not symmetric in I and J . But, of course, their roles can be reversed, as follows. Let u be the minimal element in the coset W_Jw , $K' = J \cap u(R_I)$, $w'_1 = wu^{-1}$, and γ the minimal element in the coset $w'_1W_{K'}$. Then one obtains similarly that the map $\psi' : W_{K'} \rightarrow wW_I \cap W_Jw$, $z' \mapsto \gamma z'u$ is an isomorphism of posets.

Therefore, the map $\psi^{-1}\psi' : z' \mapsto v^{-1}\gamma z'u\theta^{-1}$, from $W_{K'}$ to W_K , is an isomorphism of posets. Further, since the neutral element of W is the minimal element in both W_K and $W_{K'}$, one deduces that $u\theta^{-1} = (v^{-1}\gamma)^{-1}$ and, therefore, that $\psi^{-1}\psi'$ is an isomorphism of Coxeter groups.

4.3. Let $y < w$ in W and suppose that $y \in wW_I \cap W_Jw$. Let v be the unique minimal element in the coset wW_I and let $w_1 = v^{-1}w$ and $y_1 = v^{-1}y$. Let $K = I \cap v^{-1}R_J$ and let θ be the unique minimal element in the coset W_Kw_1 . Then, let $w_2 = w_1\theta^{-1}$ and $y_2 = y_1\theta^{-1}$.

Theorem. *There is a T -equivariant isomorphism*

$$C_{[y,w]} \cong C_v \times v(B[y_2, w_2]B \times^B C_\theta).$$

Further, it induces a T -equivariant isomorphism $\mathcal{N}_{y,w} \cong v\mathcal{N}_{y_2, w_2}$.

Proof. First, let us introduce some notation. For $x \in W$ and H a subgroup of G normalized by T , let $x(H) := \dot{x}H\dot{x}^{-1}$, where \dot{x} is any representative of x in $N(T)$, the normalizer of T in G . Then, let $U^x = U \cap x(U^-)$.

Set $Q = P_I$ and denote by U_Q^- the unipotent radical of the opposite parabolic subgroup. Let π be the projection $G/B \rightarrow G/Q$ and let $e_Q = \pi(e_B)$. Then, by Bruhat's decomposition, the multiplication map $v(U_Q^-) \times vQ/B \rightarrow G/B$, $(u, x) \mapsto ux$, induces an isomorphism $\phi_0 : v(U_Q^-) \times vQ/B \cong \pi^{-1}(vU_Q^-e_Q)$. Note that $v(U_Q^-) \cap U = U^v$ (since v is minimal in its coset vW_I) and that, by Bruhat's decomposition again, one has $U^v \cong C_{vQ}$ via $u \mapsto ue_{vQ}$. Therefore, ϕ_0 induces an isomorphism

$$\phi_1 : U^v \times vQ/B \rightarrow \pi^{-1}(C_{vQ}), \quad (u, x) \mapsto ux.$$

Further, the hypothesis that $y \in wW_I$ implies that $C_{[y,w]}$ is contained in $\pi^{-1}(C_{vQ})$. Since $C_{[y,w]}$ is B -stable, one deduces that ϕ_1 restricts to a TU^v -equivariant isomorphism

$$\phi : U^v \times (vQ/B \cap C_{[y,w]}) \xrightarrow{\cong} C_{[y,w]}, \quad (u, x) \mapsto ux.$$

We claim that $vQ/B \cap C_{[y,w]} = vC_{[y_1, w_1]}$. In fact, let $z \in [y, w]$. Since $[y, w] = v[y_1, w_1]$, then for any $z \in [y, w]$ one has $U^z = U^v v(U^{z_1})$, where $z_1 = v^{-1}z \in W_I$. This shows, first, that $vC_{[y_1, w_1]} \subseteq vQ/B \cap C_{[y,w]}$ and, second, that the restriction of ϕ to $U^v \times vC_{[y_1, w_1]}$ is surjective. Since ϕ is bijective, the claim follows. Therefore, we obtain a TU^v -equivariant isomorphism

$$(1) \quad \phi : U^v \times vC_{[y_1, w_1]} \xrightarrow{\cong} C_{[y,w]}, \quad (u, x) \mapsto ux.$$

Now, let us prove that $C_{[y_1, w_1]} \cong B[y_2, w_2]B \times^B C_\theta$. Let P_K be the parabolic subgroup corresponding to K , and let w_K denote the longest element in W_K . By hypothesis, $y_1, w_1 \in W_K\theta$ and hence, by Bruhat's decomposition, one has $B[y_1, w_1]B \subseteq P_K\theta B$. We claim that the morphism $\psi : P_K \times^B C_\theta \rightarrow P_K C_\theta = C_{[\theta, w_K\theta]}$ is an isomorphism.

Note first that the orbit C_θ (resp. $C_{w_K\theta}$) is closed (resp. open) in $C_{[\theta, w_K\theta]}$. Now, ψ is proper, since it factors as the composite of the closed immersion $P_K \times^B C_\theta \hookrightarrow P_K \times^B C_{[\theta, w_K\theta]}$ followed by the projection from $P_K \times^B C_{[\theta, w_K\theta]} \cong (P_K/B) \times C_{[\theta, w_K\theta]}$ onto $C_{[\theta, w_K\theta]}$. Further, one deduces from Bruhat's decomposition, again, that ψ maps bijectively $BzB \times^B C_\theta$ onto $C_{z\theta}$, for all $z \in W_K$, and that it induces an isomorphism $Bw_K B \times^B C_\theta \cong C_{w_K\theta}$ over the open orbit $C_{w_K\theta}$. Thus, ψ is birational and bijective. Since $C_{[\theta, w_K\theta]}$ is normal ([15, 14, 1]), it follows from Zariski's main theorem that ψ is an isomorphism.

Moreover, since $\psi(BzB \times^B C_\theta) = C_{z\theta}$ for all $z \in W_K$, and since multiplication by θ induces an isomorphism of Bruhat intervals $[x, z] \cong [x\theta, z\theta]$, one concludes that ψ induces a B -equivariant isomorphism $B[x, z]B \times^B C_\theta \cong C_{[x\theta, z\theta]}$, for all $x \leq z$ in W_K . Thus, one obtains, in particular, that

$$(2) \quad C_{[y_1, w_1]} \cong B[y_2, w_2]B \times^B C_\theta.$$

This completes the proof of the first assertion of the theorem.

Let us prove the second assertion. Let p denote the fibration $\overline{Bw_2B} \times^B C_\theta \rightarrow X_{w_2}$ and let $\Omega' = p^{-1}(\mathcal{V}_{y_2, w_2})$, a T -stable open subvariety of $C_{[y_1, w_1]}$. Since, by Bruhat's decomposition, the fibration $\overline{Bw_2B} \rightarrow X_{w_2}$ trivializes over \mathcal{V}_{y_2, w_2} , then so does p . Thus, there is a T -equivariant isomorphism $\Omega' \cong \mathcal{V}_{y_2, w_2} \times C_\theta$. It follows that $\phi(U^v \times v\Omega')$ is an open, T -stable, affine neighbourhood of e_{yB} in $C_{[y, w]}$. By [4, Lemma 2.4], this implies that $\phi(U^v \times v\Omega') = \mathcal{V}_{y, w}$.

Set $U(K)^- = U^- \cap P_K$ and $\mathcal{Z}_{y_2, w_2} = \{u \in y_2(U(K)^-) \cap U^- \mid ue_{y_2B} \in X_{w_2}\}$. Then, clearly, one has T -equivariant isomorphisms $\mathcal{Z}_{y_2, w_2} \cong \mathcal{N}_{y_2, w_2}$, $u \mapsto ue_{y_2}$ and

$$U^{y_2} \times \mathcal{Z}_{y_2, w_2} \times y_2(U^\theta) \cong \Omega', \quad (u_1, u_2, u_3) \mapsto u_1u_2u_3e_{y_1B}.$$

Thus, using (1), one obtains a T -equivariant isomorphism

$$\sigma : U^v \times v(U^{y_2}) \times v(\mathcal{Z}_{y_2, w_2}) \times vy_2(U^\theta) \xrightarrow{\cong} \mathcal{V}_{y, w}, \quad (u_1, u_2, u_3, u_4) \mapsto u_1u_2u_3u_4e_{yB}.$$

Therefore, it only remains to show that $\sigma(v(\mathcal{Z}_{y_2, w_2})) = \mathcal{N}_{y, w}$. Since both are closed subvarieties of $\mathcal{V}_{y, w}$ of dimension $\ell(w) - \ell(y)$, it is enough to prove that $\sigma(v(\mathcal{Z}_{y_2, w_2})) \subseteq \mathcal{N}_{y, w}$. Further, since $\sigma(v(\mathcal{Z}_{y_2, w_2})) \subseteq X_w$, it suffices to prove that $v(y_2(U(K)^-) \cap U^-) \subseteq U^- \cap y(U^-)$.

Thus, let $\alpha \in y_2(R_K^-) \cap R^-$. Note that $s_\alpha y_2 > y_2$ and $s_\alpha y_2 \in W_K$. Since v is minimal in its coset vW_I , then $v\alpha \in U^-$ and, therefore, the condition $v\alpha \in y(R^-)$ is equivalent to $s_{v\alpha}y > y$. But one has $s_{v\alpha}y = vs_\alpha y_2\theta$, and hence, since $s_\alpha y_2\theta \in W_K\theta \subseteq W_I$,

$$\ell(s_{v\alpha}y) = \ell(v) + \ell(s_\alpha y_2) + \ell(\theta) > \ell(v) + \ell(y_2) + \ell(\theta) = \ell(y).$$

This completes the proof of the equality $\sigma(v(\mathcal{Z}_{y_2, w_2})) = \mathcal{N}_{y, w}$ and, therefore, of the second assertion of the theorem.

4.4. Application to the symmetric group. Let $W = S_n$ and let $W_I = S_{[a, b]}$ and $W_J = S_{[c, d]}$, for some $a < b$ and $c < d$ in $[1, n]$. (We restrict ourselves to this case for simplicity, but, of course, the discussion below extends readily to the case of arbitrary Young subgroups.) Let $y < w$ in S_n and suppose that $y(i) = w(i)$ if $i < a$ or $i > b$, and that $y^{-1}(j) = w^{-1}(j)$ if $j < c$ or $j > d$.

On the one hand, let $v, w_1, y_1, \theta, w_2, y_2$ be as in 4.3. On the other hand, note that $w[a, b] = y[a, b]$ and $w^{-1}[c, d] = y^{-1}[c, d]$ and let $F = [a, b] \cap w^{-1}[c, d]$ and $E = [c, d] \cap w[a, b]$. Then w and y induce bijections \tilde{w}, \tilde{y} from F to E , and these give rise in a natural manner to elements \bar{w}, \bar{y} of S_m , where $m = \#F$. Explicitly, let ϕ (resp. ψ) be the unique increasing bijection from $[1, m]$ to F (resp. to E); then $\bar{w} = \psi^{-1} \circ \tilde{w} \circ \phi$, and similarly for \bar{y} (compare 2.6).

Further, as in 2.6, one observes that v^{-1} induces an increasing bijection from E to some interval $[f, g]$, where $a \leq f \leq g \leq b$, and that, denoting by η the unique increasing bijection from $[1, m]$ to $[f, g]$, one has $w_2 = \eta \circ \bar{w} \circ \eta^{-1}$, and similarly for y_2 and \bar{y} .


Therefore, the permutations w_2 and y_2 can be read off from the graphs of w and y by "focusing" on the intersections of the columns C_i , $a \leq i \leq b$, and the lines L_j , $c \leq j \leq d$, see the example below. This is the explanation for the title of this section.

4.5. Example. In S_{12} consider the permutations

$$\begin{aligned} w &= (6, 7, 2, 9, 12, 10, 11, 4, 1, 5, 3, 8); \\ y &= (6, 4, 2, 7, 12, 9, 11, 5, 1, 10, 3, 8). \end{aligned}$$

Note that $y(i) = w(i)$ if $i < 2$ or $i > 10$ and that $y^{-1}(j) = w^{-1}(j)$ if $j < 4$ or $j > 10$.

It is helpful to represent each permutation $\tau \in S_n$ by its graph $G(\tau)$, that is, the set of points $(i, \tau(i))$ in $[1, n] \times [1, n]$. In the picture below, the graph of w (resp. y) is represented by bullets and crosses (resp. bullets and stars).



Then, the associated \bar{w} and \bar{y} in S_5 are respectively $(3, 4, 5, 1, 2)$ and $(1, 3, 4, 2, 5)$; they correspond to the pair associated to the polynomial $1 + 2q$ in 1.4. Therefore, $P_{y,w} = 1 + 2q$.

ACKNOWLEDGEMENTS

This paper owes much to Fokko Du Cloux, who communicated to me several years ago his computer program “Coxeter” ([7]). It was on the ground of calculations made in the symmetric group S_8 that I was led to conjecture the theorem, as well as the form of the pair $y_P < w_P$. More recently, during a visit to the Institut Girard Desargues at Université Lyon I, discussions with Fokko du Cloux about this conjecture renewed my interest in the subject. It is a pleasure to thank him for sharing his program with me and for these discussions.

REFERENCES

1. H.H. Andersen, Schubert varieties and Demazure’s character formula, *Invent. Math.* **79** (1985), 611-618. MR **86h**:14042
2. A. Beilinson, J. Bernstein and P. Deligne, Faisceaux pervers, in “Analyse et topologie sur les espaces singuliers (I)”, Astérisque 100, Soc. Math. France, 1982, pp. 3-171. MR **86g**:32015
3. B.D. Boe, Kazhdan-Lusztig polynomials for hermitian symmetric spaces, *Trans. Amer. Math. Soc.* **309** (1988), 279-294. MR **89i**:22024
4. M. Brion and P. Polo, Generic singularities of certain Schubert varieties, *Math. Z.* **231** (1999), 301-324.
5. V.V. Deodhar, Local Poincaré duality and nonsingularity of Schubert varieties, *Comm. Algebra* **13** (1985), 1379-1388. MR **86i**:14015
6. V.V. Deodhar, A combinatorial setting for questions in Kazhdan-Lusztig theory, *Geom. Dedicata* **36** (1990), 95-119. MR **91h**:20075
7. F. Du Cloux, program “Coxeter” (available at www.desargues.univ-lyon1.fr/home/ducloux/coxeter.html).
8. M. Goresky and R. MacPherson, Intersection homology II, *Invent. Math.* **72** (1983), 77-129. MR **84i**:57012
9. R. Irving, The socle filtration of a Verma module, *Ann. Sci. École Norm. Sup.* **21** (1988), 47-65. MR **89h**:17015

10. B. Iversen, "Cohomology of sheaves", Springer-Verlag, Berlin Heidelberg 1986. MR **87m**:14013
11. D. Kazhdan and G. Lusztig, Representations of Coxeter groups and Hecke algebras, *Invent. Math.* **53** (1979), 165-184. MR **81j**:20066
12. D. Kazhdan and G. Lusztig, Schubert varieties and Poincaré duality, in Proc. Symposia Pure Math. Vol. 36, Amer. Math. Soc., 1980, pp. 185-203. MR **84g**:14054
13. A. Lascoux and M.P. Schützenberger, Polynômes de Kazhdan-Lusztig pour les grassmanniennes, in "Tableaux de Young et foncteurs de Schur en algèbre et géométrie", Astérisque 87-88, Soc. Math. France, 1981, pp. 249-266. MR **83i**:14045
14. S. Ramanan and A. Ramanathan, Projective normality of flag varieties and Schubert varieties, *Invent. Math.* **79** (1985), 217-224. MR **86j**:14051
15. C.S. Seshadri, Line bundles on Schubert varieties, in "Proceedings of the Bombay colloquium on vector bundles on algebraic varieties", 1984. MR **88i**:14047
16. T.A. Springer, Quelques applications de la cohomologie d'intersection, Séminaire Bourbaki, Vol. 1981/1982, Astérisque, 92-93, Soc. Math. France, Paris, 1982, pp. 249-273. MR **85i**:32016b
17. A. Zelevinsky, Small resolutions of singularities of Schubert varieties, *Funct. Anal. Appl.* **17** (1982), 142-144.

CNRS, UMR 7539, INSTITUT GALILÉE, DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ PARIS-NORD, 93430 VILLETANEUSE, FRANCE

E-mail address: `polo@math.univ-paris13.fr`