

CHARACTER SHEAVES ON DISCONNECTED GROUPS, VIII

G. LUSZTIG

ABSTRACT. In this paper we continue the study of character sheaves on a reductive group. To each subset of the set of simple reflections in the Weyl group we associate an algebra of the same kind as an Iwahori Hecke algebra with unequal parameters in terms of parabolic character sheaves. We also prove a Mackey type formula for character sheaves. We define a duality operation for character sheaves. We also prove a quasi-rationality property for character sheaves.

INTRODUCTION

Throughout this paper, G denotes a fixed, not necessarily connected, reductive algebraic group over an algebraically closed field \mathbf{k} . This paper is a part of a series [L9] which attempts to develop a theory of character sheaves on G .

In Section 36 we associate to any subset J of the set of simple reflections an algebra $\underline{\mathfrak{K}}^J$ over $\mathbf{Q}(v)$ (with v an indeterminate) defined using certain character sheaves on Z_{J,G^0} . When $J = \emptyset$, this is the standard Iwahori-Hecke algebra attached to \mathbf{W} . For general J we show that this algebra shares several basic features with an Iwahori-Hecke algebra with unequal parameters. This opens the possibility of studying Iwahori-Hecke algebras with unequal parameters in the framework of the theory of perverse sheaves.

In Section 37 we prove a Mackey type formula for character sheaves on $Z_{J,D}$ where $J \subset \mathbf{I}$ and D is a connected component of G . This is essentially an identity involving certain induction and restriction functors analogous to one in the representation theory of reductive groups over a finite field.

In Section 38 we study a duality operation for character sheaves on a connected component D of G generalizing the case $D = G^0$ considered in [L3, III, 15]. This duality operator is analogous to the known duality operator for representations of a reductive group over a finite field; see 38.12.

In Section 39 we prove a quasi-rationality property for representations of certain extensions of an (irreducible) Weyl group. This generalizes [L3, III, (12.9.3)] and is a step in the proof of a key property of character sheaves (see 39.8).

Notation. We write $\mathfrak{s}, \underline{\mathfrak{s}}$ instead of $\mathfrak{s}(\mathbf{T}), \underline{\mathfrak{s}}(\mathbf{T})$ (see 28.1, 28.3).

Errata to Part V. In 25.2, line 1, replace $\tilde{\mathcal{E}}'$ by $\pi_1' \tilde{\mathcal{E}}'$.

In 27.1 replace the diagram $N_D P/U_P \xleftarrow{a} V_1 \xrightarrow{a} V_2 \xrightarrow{a'} D$ by $N_D P/U_P \xleftarrow{a} V_1 \xrightarrow{a'} V_2 \xrightarrow{a''} D$.

Received by the editors November 9, 2006 and, in revised form, January 21, 2006.

2000 *Mathematics Subject Classification.* Primary 20G99.

Supported in part by the National Science Foundation.

©2006 American Mathematical Society
Reverts to public domain 28 years from publication

In the commutative diagram on p. 372 replace $\hat{Z}_1 \xrightarrow{a} \hat{Z}_2, Z_1 \xrightarrow{b} Z_2, V_1 \xrightarrow{c} V_2 \xrightarrow{a'}$
 D by $\hat{Z}_1 \xrightarrow{a'} \hat{Z}_2, Z_1 \xrightarrow{b'} Z_2, V_1 \xrightarrow{c'} V_2 \xrightarrow{a''} D$ respectively.

Errata to Part VI.

In 28.5 replace $t_0 : \text{Ad}(d^{-1}\dot{w}^{-1})(t_0)tt_0^{-1}$ by $t_0 : t \mapsto \text{Ad}(d^{-1}\dot{w}^{-1})(t_0)tt_0^{-1}$.

In 28.19(a) replace $\hat{Z}_{\emptyset, \epsilon_D(J), D^{-1}}^{\mathcal{L}'}$ by $\hat{Z}_{\epsilon_D(J), D^{-1}}^{\mathcal{L}'}$ (twice).

In 31.4, line 4 replace \mathfrak{a}_D^i by \mathfrak{a}_D .

Errata to Part VII.

In 32.5 replace $> b_2$ by b_2 .

In 32.15 replace C_u by C^u .

In 32.18(a) replace $T_{\epsilon'(a_{r+r'}^{-1})}$ by $T_{\epsilon'(a_{r+r'}^{-1})}$.

In 32.23 replace the first $=$ by $) =$.

After the statement of Corollary 32.23 insert: Take $D' = D^{-1}$ so that $\Delta = G^0$.

In the third line of Corollary 32.24 replace λ' by λ'' .

In the fourth line of Corollary 32.24 replace λ'' by \mathcal{L}'' .

In 32.26, line 13, replace tT_x by \tilde{T}_x .

In 35.6, line 8, after $h_{i-1}^{-1}h_i \in U^*n_iU^*$, add: ($i \in [1, r]$).

CONTENTS

- 36. The algebra $\underline{\mathfrak{K}}^J$.
- 37. A Mackey type formula.
- 38. Duality.
- 39. Quasi-rationality.

36. THE ALGEBRA $\underline{\mathfrak{K}}^J$

36.1. We fix a connected component D of G . We write ϵ instead of $\epsilon_D : \mathbf{W} \rightarrow \mathbf{W}$ (see 26.2). Occassionally we shall consider one (or two) other component(s), say D' (or D', D''); we write ϵ' instead of $\epsilon_{D'}$.

For $J \subset \mathbf{I}$ we identify $Z_{J,D}$ (see 26.2) with

$$\{(P, gU_P); P \in \mathcal{P}_J, gU_P \in D/U_P\}$$

by $(P, P', gU_P) \mapsto (P, gU_P)$. To any $(P, gU_P) \in Z_{J,D}$ we associate an element $w_{P, gU_P} \in \mathbf{W}$ by the following requirements. Let $z = \text{pos}(gPg^{-1}, P) \in {}^{\epsilon(J)}\mathbf{W}^J$; then

- (i) $w_{P, gU_P} = w_{P_1, gU_{P_1}}$ where $P_1 = (g^{-1}Pg \cap P)U_P \in \mathcal{P}_{J \cap \epsilon^{-1}(\text{Ad}(z)J)}$,
- (ii) $w_{P, gU_P} = z$ if $\epsilon^{-1}(\text{Ad}(z)J) = J$.

Note that (i), (ii) define uniquely w_{P, gU_P} by induction on $|J|$: if $|J| = 0$, then w_{P, gU_P} is given by (ii); if $|J| \geq 1$ and $\epsilon^{-1}(\text{Ad}(z)J) = J$, then w_{P, gU_P} is again given by (ii); if $|J| \geq 1$ and $\epsilon^{-1}(\text{Ad}(z)J) \neq J$, then $|J \cap \epsilon^{-1}(\text{Ad}(z)J)| < |J|$ and w_{P, gU_P} is determined by (i) since $w_{P_1, gU_{P_1}}$ is known from the induction hypothesis.

From definitions we see that the map $Z_{J,D} \rightarrow \mathbf{W}, (P, gU_P) \mapsto w_{P, gU_P}$ is the composition of $\beta' : Z_{J,D} \rightarrow \mathcal{T}(J, \epsilon)$ in 26.2 (see also [L10, 3.11]) with the bijection $\mathcal{T}(J, \epsilon) \rightarrow {}^{\epsilon(J)}\mathbf{W}$ given by [L10, 2.4, 2.5] and with the inclusion ${}^{\epsilon(J)}\mathbf{W} \rightarrow \mathbf{W}$. In particular, we have $w_{P, gU_P} \in {}^{\epsilon(J)}\mathbf{W}$ and $\text{pos}(gPg^{-1}, P) = \min(\mathbf{W}_{\epsilon(J)}w_{P, gU_P}\mathbf{W}_J)$ for any $(P, gU_P) \in Z_{J,D}$. (For any $\mathbf{W}_{\epsilon(J)}, \mathbf{W}_J$ double coset Ω in \mathbf{W} we denote by $\min(\Omega)$ the unique element of minimal length in Ω .)

For any $J \subset \mathbf{I}$, $w \in {}^{\epsilon(J)}\mathbf{W}$, we set

$${}^wZ_{J,D} = \{(P, gU_P) \in Z_{J,D}; w_{P,gU_P} = w\}.$$

Then $Z_{J,D} = \bigcup_{w \in {}^{\epsilon(J)}\mathbf{W}} {}^wZ_{J,D}$ is a partition; it is the same as the partition $Z_{J,D} = \bigcup_{\mathbf{t} \in \mathcal{T}(J,\epsilon)} {}^{\mathbf{t}}Z_{J,D}$ in 26.2 (see also [L10, 3.11]). In particular, each ${}^wZ_{J,D}$ is a locally closed, smooth subvariety of $Z_{J,D}$ stable under the G^0 -action $h : (P, gU_P) \mapsto (hPh^{-1}, hgh^{-1}U_{hPh^{-1}})$.

From definitions, for $J \subset \mathbf{I}$, $w \in {}^{\epsilon(J)}\mathbf{W}$ we have a map

$$\vartheta_{J,w} : {}^wZ_{J,D} \rightarrow {}^wZ_{J_1,D}, (P, gU_P) \mapsto (P_1, gU_{P_1})$$

with $J_1 = J \cap \epsilon^{-1}(\text{Ad}(w)J)$, $w = \min(\mathbf{W}_{\epsilon(J)} w \mathbf{W}_J)$, $P_1 = (g^{-1}Pg \cap P)U_P$. This map may be identified with a map ϑ as in [L10, 3.11]; in particular, it is an affine space bundle (see [L10, 3.12(b)]).

36.2. If $J \subset \mathbf{I}$ and $w \in {}^{\epsilon(J)}\mathbf{W}$ satisfies $\epsilon^{-1}(\text{Ad}(w)J) = J$ (hence $w \in {}^{\epsilon(J)}\mathbf{W}^J$), we have ${}^wZ_{J,D} = \{(P, gU_P) \in Z_{J,D}, \text{pos}(gPg^{-1}, P) = w\}$. In this case we pick $P \in \mathcal{P}_J, P' \in \mathcal{P}_{\epsilon(J)}$ such that $\text{pos}(P', P) = w$ and a common Levi L of P', P . Let $\mathbf{d} = \{g \in D; gLg^{-1} = L, gPg^{-1} = P'\}$, a connected component of the reductive group N_GL with identity component L . We have a diagram

$$(a) \quad \mathbf{d} \xleftarrow{pr_2} G^0/(U_P \cap U_{P'}) \times \mathbf{d} \xrightarrow{j} {}^wZ_{J,D}$$

where $j(h(U_P \cap U_{P'}), g) = (hPh^{-1}, hgh^{-1}U_{hPh})$ is a principal L -bundle for an L -action on $G^0/(U_P \cap U_{P'}) \times \mathbf{d}$ compatible under pr_2 with the L -action on \mathbf{d} (by conjugation in N_GL). If X is a character sheaf on \mathbf{d} , then $pr_2^{\star}X$ is therefore an L -equivariant simple perverse sheaf on $G^0/(U_P \cap U_{P'}) \times \mathbf{d}$ which must be of the form $j^{\star}X'$ for a well-defined simple perverse sheaf X' on ${}^wZ_{J,D}$.

The collection of simple perverse sheaves on ${}^wZ_{J,D}$ of the form X' with X as above is denoted by ${}^w\hat{Z}_{J,D}$. This collection is independent of the choice of P, P', L . Note that $X \mapsto X'$ defines a bijection between the set of isomorphism classes of character sheaves on \mathbf{d} and the set of isomorphism classes of objects in ${}^w\hat{Z}_{J,D}$.

36.3. More generally, for $J \subset \mathbf{I}$ and $w \in {}^{\epsilon(J)}\mathbf{W}$, we define by induction on $|J|$ a collection of simple G^0 -equivariant perverse sheaves ${}^w\hat{Z}_{J,D}$ on ${}^wZ_{J,D}$.

If $|J| = 0$, then ${}^w\hat{Z}_{J,D}$ is defined as in 36.2. If $|J| \geq 1$ and $\epsilon^{-1}(\text{Ad}(w)J) = J$, then ${}^w\hat{Z}_{J,D}$ is again defined as in 36.2. If $|J| \geq 1$ and $\epsilon^{-1}(\text{Ad}(w)J) \neq J$, then $\epsilon^{-1}(\text{Ad}(w)J) \neq J$ where $w = \min(\mathbf{W}_{\epsilon(J)} w \mathbf{W}_J)$. Thus, if $J_1 = J \cap \epsilon^{-1}(\text{Ad}(w)J)$, then $|J_1| < |J|$ and the class of perverse sheaves ${}^w\hat{Z}_{J_1,D}$ on ${}^wZ_{J_1,D}$ is defined from the induction hypothesis. By definition, ${}^w\hat{Z}_{J,D}$ consists of the simple perverse sheaves on ${}^wZ_{J,D}$ of the form $\vartheta_{J,w}^{\star}(X)$ for some $X \in {}^w\hat{Z}_{J_1,D}$ (with $\vartheta_{J,w}$ as in 36.1). This completes the inductive definition of ${}^w\hat{Z}_{J,D}$. The objects of ${}^w\hat{Z}_{J,D}$ are said to be *character sheaves on ${}^wZ_{J,D}$* . Let $\mathcal{I}_{J,w,D}$ be a set of representatives for the isomorphism classes of character sheaves on ${}^wZ_{J,D}$.

For $J \subset \mathbf{I}$ and $w \in {}^{\epsilon(J)}\mathbf{W}$, let $\mathcal{D}^{cs}({}^wZ_{J,D})$ be the subcategory of $\mathcal{D}({}^wZ_{J,D})$ whose objects are those $K \in \mathcal{D}({}^wZ_{J,D})$ such that for any j , any simple subquotient of ${}^pH^jK$ is in ${}^w\hat{Z}_{J,D}$. Let $i_{J,w} : {}^wZ_{J,D} \rightarrow Z_{J,D}$ be the inclusion.

$$(a) \text{ If } K \in \mathcal{D}^{cs}(Z_{J,D}), \text{ then } i_{J,w}^*(K) \in \mathcal{D}^{cs}({}^wZ_{J,D}).$$

It is enough to prove (a) for $K \in \hat{Z}_{J,D}$. In this case (a) follows from [L10, 4.12].

For $J \subset \mathbf{I}$, $w \in {}^{\epsilon(J)}\mathbf{W}$ and $K \in {}^w\hat{Z}_{J,D}$, let K^\sharp be the unique simple perverse sheaf on $Z_{J,D}$ such that $i_{J,w}^*(K^\sharp) = K$ and $\text{supp}(K^\sharp)$ is the closure in $Z_{J,D}$ of $\text{supp}(K)$.

(b) We have $K^\sharp \in \hat{Z}_{J,D}$.

This is proved in [L10, 4.17] based on the following statement:

(c) Let Y be a locally closed subvariety of an algebraic variety Y' and let $i : Y \rightarrow Y'$ be the inclusion. Let $C \in \mathcal{D}(Y)$ and let A be a simple perverse sheaf on Y such that $A \dashv C$. Let A^\sharp be the unique simple perverse sheaf on Y such that $i^*(A^\sharp) = A$ and $\text{supp}(A^\sharp)$ is the closure in Y' of $\text{supp}(A)$. Then $A^\sharp \dashv i_!C$.

(In *loc.cit.* this is applied with $Y = {}^wZ_{J,D}$, $Y' = Z_{J,D}$, $A = K$ and an $C \in \mathcal{D}^{cs}(Z_{J,D})$.) Since the proof of (c) is omitted in *loc.cit.* we give a proof here. Let Y_1 be the closure of Y in Y' . Let $Y \xrightarrow{i_1} Y_1 \xrightarrow{i_2} Y'$ be the inclusions. Clearly, if (c) holds for i_1 and i_2 instead of i , then it holds for i . Thus we may assume that $i = i_1$ or $i = i_2$, that is, that Y is open or closed in Y' . Assume first that Y is closed in Y' . Then $A^\sharp = i_!A$ and $i_!$ commutes with taking ${}^pH^j$; hence $i_!A \dashv i_!C$ as desired. Next, assume that Y is open in Y' . Then i^* commutes with taking ${}^pH^j$ and $i^*i_!C = C$. Hence for any j we have ${}^pH^j(C) = i^*({}^pH^j(i_!C))$ and for some j we have $A \dashv i^*({}^pH^j(i_!C))$. Let $0 = F_0 \subset F_1 \subset \dots \subset F_m = {}^pH^j(i_!C)$ be a sequence of perverse subsheaves of ${}^pH^j(i_!C)$ such that F_k/F_{k-1} is simple for $k \in [1, m]$. Then $0 = i^*F_0 \subset i^*F_1 \subset \dots \subset i^*F_m = i^*({}^pH^j(i_!C))$ is a sequence of perverse subsheaves of $i^*({}^pH^j(i_!C))$ such that $i^*F_k/i^*F_{k-1} = i^*(F_k/F_{k-1})$ is simple or 0 for $k \in [1, m]$. Hence $A \cong i^*F_k/i^*F_{k-1} = i^*(F_k/F_{k-1})$ for some $k \in [1, m]$. Then $A' = F_k/F_{k-1}$ is a simple perverse sheaf on Y' such that $A' \dashv i_!C$, $i^*A' \cong A$. We must have $A' \cong A^\sharp$ and (c) is proved.

(d) Let $K' \in \hat{Z}_{J,D}$. There exists a unique $w \in {}^{\epsilon(J)}\mathbf{W}$ and a unique $K \in {}^w\hat{Z}_{J,D}$ (up to isomorphism) such that $K' \cong K^\sharp$.

This is proved in [L10, 4.13]; it is an immediate consequence of (a).

Let $\mathcal{I}_{J,D} = \bigsqcup_{w \in {}^{\epsilon(J)}\mathbf{W}} \{K^\sharp; K \in \mathcal{I}_{J,w,D}\}$.

From (b), (d) we see that

(e) $\mathcal{I}_{J,D}$ is a set of representatives for the isomorphism classes of character sheaves on $Z_{J,D}$.

From (a), (b) we deduce:

(f) If $w \in {}^{\epsilon(J)}\mathbf{W}$ and $K \in \mathcal{D}^{cs}({}^wZ_{J,D})$, then $(i_{J,w})_!K \in \mathcal{D}^{cs}(Z_{J,D})$.

36.4. For $J \subset J' \subset \mathbf{I}$ and $P \in \mathcal{P}_J$ let $Q_{J',P}$ be the unique parabolic in $\mathcal{P}_{J'}$ that contains P . We have a diagram

$$Z_{J,D} \xleftarrow{\mathfrak{c}} Z_{J,J',D} \xrightarrow{\mathfrak{d}} Z_{J',D}$$

where

$$\begin{aligned} Z_{J,J',D} &= \{(P, gU_Q); P \in \mathcal{P}_J, Q = Q_{J',P}, gU_Q \in D/U_Q\}, \\ \mathfrak{c}(P, gU_Q) &= (P, gU_P), \mathfrak{d}(P, gU_Q) = (Q, gU_Q). \end{aligned}$$

Define functors

$$\mathfrak{f}_{J,J'} : \mathcal{D}(Z_{J,D}) \rightarrow \mathcal{D}(Z_{J',D}), \mathfrak{e}_{J,J'} : \mathcal{D}(Z_{J',D}) \rightarrow \mathcal{D}(Z_{J,D})$$

by $\mathfrak{f}_{J,J'}A = \mathfrak{d}!\mathfrak{c}^*A$, $\mathfrak{e}_{J,J'}A' = \mathfrak{c}!\mathfrak{d}^*A'$. Now \mathfrak{d} is proper and \mathfrak{c} is an affine space bundle with fibres of dimension $a = \dim \mathcal{P}_J - \dim \mathcal{P}_{J'}$. Hence $\mathfrak{f}_{J,J'}$ commutes with Verdier

duality up to a shift and a twist:

$$(a) \quad \mathfrak{D}f_{J,J'} = f_{J,J'}[[a]]\mathfrak{D} : \mathcal{D}(Z_{J,D}) \rightarrow \mathcal{D}(Z_{J',D}).$$

For $J \subset J' \subset J'' \subset \mathbf{I}$ we have (see [L10, 6.2]):

$$(b) \quad f_{J,J''} = f_{J',J''}f_{J,J'}, \quad \mathfrak{e}_{J,J''} = \mathfrak{e}_{J,J'}\mathfrak{e}_{J',J''}.$$

Clearly, $f_{J,J} = 1, \mathfrak{e}_{J,J} = 1$.

For $J \subset J' \subset \mathbf{I}$ we show that the convolution bifunctor

$$\mathcal{D}(Z_{J,D}) \times \mathcal{D}(Z_{\epsilon(J),D'}) \rightarrow \mathcal{D}(Z_{J,D'D}), \quad A, B \mapsto A * B,$$

(see 32.5) is compatible with the functors $\mathfrak{e}_{J,J'}$ in the following sense: for $A \in \mathcal{D}(Z_{J,D}), A' \in \mathcal{D}(Z_{\epsilon(J),D'})$, we have

$$(c) \quad (\mathfrak{e}_{J,J'}A) * (\mathfrak{e}_{J,J'}A') = \mathfrak{e}_{J,J'}(A * A').$$

We have a commutative diagram

$$\begin{array}{ccccc} Z_{J',D} \times Z_{\epsilon(J'),D'} & \xleftarrow{b'_1} & Z'_0 & \xrightarrow{b'_2} & Z_{J',D'D} \\ \mathfrak{d} \times \mathfrak{d}' \uparrow & & f_1 \uparrow & & \mathfrak{d}'' \uparrow \\ Z_{J,J',D} \times Z_{\epsilon(J),\epsilon(J'),D'} & \xleftarrow{f_2} & \underline{Z} & \xrightarrow{f_3} & Z_{J,J',D'D} \\ \mathfrak{c} \times \mathfrak{c}' \downarrow & & f_4 \downarrow & & \mathfrak{c}'' \downarrow \\ Z_{J,D} \times Z_{\epsilon(J),D'} & \xleftarrow{b_1} & Z_0 & \xrightarrow{b_2} & Z_{J,D'D} \end{array}$$

Here $\mathfrak{c}, \mathfrak{d}$ are as above,

$$Z_{\epsilon(J'),D'} \xleftarrow{\mathfrak{c}'} Z_{\epsilon(J),\epsilon(J'),D'} \xrightarrow{\mathfrak{d}'} Z_{\epsilon(J'),D'}, \quad Z_{J,D'D} \xleftarrow{\mathfrak{c}''} Z_{J,J',D'D} \xrightarrow{\mathfrak{d}''} Z_{J',D'D}$$

are the analogous maps when (J, J', D) is replaced by $(\epsilon(J), \epsilon(J'), D')$ or by $(J, J', D'D)$, b_1, b_2 are as in 32.5, b'_1, b'_2 are the analogous maps with J replaced by J' ,

$$\begin{aligned} \underline{Z} = \{ & (X, P, gU_P, g'U_{P'}); X \in \mathcal{P}_J, P = Q_{J',X}, gU_P \in D/U_P, \\ & P' = gPg^{-1}, g'U_{P'} \in D'/U_{P'} \}, \end{aligned}$$

$$f_1(X, gU_P, g'U_{P'}) = (P, gPg^{-1}, g'gPg^{-1}g'^{-1}, gU_P, g'U_{gPg^{-1}}),$$

$$f_2(X, gU_P, g'U_{P'}) = ((X, gU_P), (gXg^{-1}, g'U_{gPg^{-1}})),$$

$$f_3(X, gU_P, g'U_{P'}) = (X, g'gU_P),$$

$$f_4(X, gU_P, g'U_{P'}) = (X, gXg^{-1}, g'gXg^{-1}g'^{-1}, gU_X, g'U_{gXg^{-1}}).$$

It is enough to show that

$$b_{2!}b_1^*(\mathfrak{c} \times \mathfrak{c}')!(\mathfrak{d} \times \mathfrak{d}')^*(A \boxtimes A') = \mathfrak{c}_1''\mathfrak{d}''^*b_{2!}'b_1'^*(A \boxtimes A').$$

Since the upper right and lower left squares are cartesian, we have

$$b_1^*(\mathfrak{c} \times \mathfrak{c}')! = f_{4!}f_2^*, \quad \mathfrak{d}''^*b_{2!}' = f_{3!}f_1^*.$$

Hence it is enough to show that

$$b_{2!}f_{4!}f_2^*(\mathfrak{d} \times \mathfrak{d}')^*(A \boxtimes A') = \mathfrak{c}_1''f_{3!}f_1^*b_1'^*(A \boxtimes A').$$

This follows from the commutativity of the diagram above: we have $b_{2!}f_{4!} = \mathfrak{c}_1''f_{3!}$ and $f_2^*(\mathfrak{d} \times \mathfrak{d}')^* = f_1^*b_1'^*$. This proves (c).

If $A \in \mathcal{D}(Z_{J,D})$, $B \in \mathcal{D}(Z_{\epsilon(J),D'})$, $C \in \mathcal{D}(Z_{\epsilon'\epsilon(J),D''})$, then, from definitions, we have the associativity property

$$(d) \quad (A * B) * C = A * (B * C).$$

Consider the isomorphism $\partial : Z_{J,D} \xrightarrow{\sim} Z_{\epsilon(J),D^{-1}}$ as in 28.19. Note that the composition $Z_{J,D} \xrightarrow{\partial} Z_{\epsilon(J),D^{-1}} \xrightarrow{\partial} Z_{J,D}$ is the identity map. From definitions, we have for A, B as above,

$$(e) \quad \partial_!(A * B) = (\partial_!B) * (\partial_!A).$$

For $J \subset \mathbf{I}$ we define a functor $\tau : \mathcal{D}(Z_{J,G^0}) \rightarrow \mathcal{D}(\text{point})$ by $\tau(C) = p'_!i^*(C)$ where $i : \mathcal{P}_J \rightarrow Z_{J,G^0}$ is the imbedding $P \mapsto (P, U_P)$ and $p' : \mathcal{P}_J \rightarrow \text{point}$ is the obvious map. We define a bifunctor $(:) : \mathcal{D}(Z_{J,D}) \times \mathcal{D}(Z_{J,D}) \rightarrow \mathcal{D}(\text{point})$ by $(A : B) = p_!i'^*(A \boxtimes B)$ where $i' : Z_{J,D} \rightarrow Z_{J,D} \times Z_{J,D}$ is the diagonal and $p : Z_{J,D} \rightarrow \text{point}$ is the obvious map. As in 32.23(b) we have

$$(f) \quad (A : B) = \tau(A * (\partial_!B)).$$

From definitions we have

$$(g) \quad (A : B) = (\partial_!A : \partial_!B).$$

If $A \in \mathcal{D}(Z_{J,D})$, $B \in \mathcal{D}(Z_{\epsilon(J),D'})$, $C \in \mathcal{D}(Z_{\epsilon'\epsilon(J),D^{-1}D'^{-1}})$, then from (d) we have $\tau((A * B) * C) = \tau(A * (B * C))$. Using (f) we rewrite this as $(A * B : \partial_!C) = (A : \partial_!(B * C))$ or, using (g), as

$$(h) \quad (A * B : \partial_!C) = (\partial_!A : B * C).$$

For $J \subset J'$, $\mathfrak{c}, \mathfrak{d}$ as above and $A \in \mathcal{D}(Z_{J,D})$, $B \in \mathcal{D}(Z_{J',D})$ we show that

$$(i) \quad (B : f_{J,J'}A) = (e_{J,J'}B : A) \in \mathcal{D}(\text{point}).$$

Let p_1, p_2, p_3 be the obvious maps from $Z_{J',D}, Z_{J,D}, Z_{J,J',D}$ to the point. We must show that $p_{1!}(B \otimes \mathfrak{d}_!c^*A) = p_{2!}(c_!\mathfrak{d}^*B \otimes A)$. Since $p_{1!}\mathfrak{d}_! = p_{3!} = p_{2!}c_!$, both sides are equal to $p_{3!}(\mathfrak{d}^*B \otimes c^*A)$. This proves (i).

36.5. For $J \subset J' \subset \mathbf{I}$ and $w \in {}^{\epsilon(J)}\mathbf{W}$, $w' \in {}^{\epsilon(J')}\mathbf{W}$ we have a diagram

$${}^w Z_{J,D} \xleftarrow{c_{w,w'}} {}^{w,w'} Z_{J,J',D} \xrightarrow{\mathfrak{d}_{w,w'}} {}^{w'} Z_{J',D}$$

where

$${}^{w,w'} Z_{J,J',D} = \{(P, gU_Q) \in Z_{J,J',D}; (P, gU_P) \in {}^w Z_{J,D}, (Q, gU_Q) \in {}^{w'} Z_{J',D}\}$$

and $c_{w,w'}, \mathfrak{d}_{w,w'}$ are the restrictions of c, \mathfrak{d} in 36.4. Consider the functor

$$f_{J,w,J',w'} : \mathcal{D}({}^w Z_{J,D}) \rightarrow \mathcal{D}({}^{w'} Z_{J',D}), A \mapsto \mathfrak{d}_{w,w'}c_{w,w'}^*A.$$

From definitions we see that, for $A \in \mathcal{D}({}^w Z_{J,D})$, we have

$$(a) \quad f_{J,w,J',w'}(A) = i_{J',w'}^*f_{J,J'}(i_{J,w}A).$$

36.6. Let $J \subset \mathbf{I}$ and let $w \in {}^{\epsilon(J)}\mathbf{W}$. We have a sequence of affine space bundles

$$(a) \quad {}^wZ_{J,D} \xrightarrow{\vartheta_{J,w}} {}^wZ_{J_1,D} \xrightarrow{\vartheta_{J_1,w}} {}^wZ_{J_2,D} \xrightarrow{\vartheta_{J_2,w}} \dots$$

where $\vartheta_{J,w}$ is as in 36.1, $\vartheta_{J_1,w}$ is the analogous map with J replaced by J_1 (we set $(J_1)_1 = J_2$), $\vartheta_{J_2,w}$ is the analogous map with J replaced by J_2 (we set $(J_2)_1 = J_3$), etc. We have $J \supset J_1 \supset J_2 \supset \dots$. Let $J_\infty = J_r$ for large r . Since $\vartheta_{J_r,w} = 1$ for large r , the composition of the maps (a) is a well-defined map $\underline{\vartheta} : {}^wZ_{J,D} \rightarrow {}^wZ_{J_\infty,D}$. For any $(P, gU_P) \in {}^wZ_{J,D}$ we have

$$\vartheta_{J,w}(P, gU_P) = (P_1, gU_{P_1}), \vartheta_{J_1,w}(P_1, gU_{P_1}) = (P_2, gU_{P_2}), \dots$$

where $P \supset P_1 \supset P_2 \supset \dots$. Let $P_\infty = P_r$ for large r . We show:

(b) *The map $\alpha : {}^{w,w}Z_{J_\infty,J,D} \rightarrow {}^wZ_{J,D}$, $(R, gU_P) \mapsto (P, gU_P)$, is an isomorphism.*

We show only that α is a bijection. Let $(P, gU_P) \in {}^wZ_{J,D}$. By [L10, 4.14(b)] there exists $(B, h) \in \mathcal{B} \times gU_P$ such that $B \subset P$ and $\text{pos}(hBh^{-1}, B) = w$. We have also $\text{pos}(gBg^{-1}, B) = w$. Let $R = Q_{J_\infty,B}$. By [L10, 4.14(a)] we have $(R, gU_R) \in {}^wZ_{J_\infty,D}$. We have $(R, gU_P) \in {}^{w,w}Z_{J_\infty,J,D}$ and $\alpha(R, gU_P) = (P, gU_P)$. Thus α is surjective.

We show that α is injective. Let $(R, gU_P) \in {}^{w,w}Z_{J_\infty,J,D}$. Define $P_1, P_2, \dots, P_\infty$ in terms of (P, gU_P) as above. It is enough to show that $R = P_\infty$. Let $P' = gPg^{-1}, R' = gRg^{-1}$. Since $(R, gU_R) \in {}^wZ_{J_\infty,D}$ and $\epsilon^{-1}(\text{Ad}(w)J_\infty) = J_\infty$, we see that R, R' have a common Levi and $\text{pos}(R', R) = w$. Hence if B' is any Borel of R' , there exists a Borel B of R such that $\text{pos}(B', B) = w$. If $w = \text{pos}(P', P)$, we have $w = w.x$ where $x \in \mathbf{W}_J$ (see [L10, 2.1(b)]) and $l(w.) + l(x) = l(w)$. Hence we can find a Borel B_1 of G^0 such that $\text{pos}(B', B_1) = w., \text{pos}(B_1, B) = x$. Since $B \subset R \subset P$, we have $B_1 \subset P$. Since $\text{pos}(B', B_1) = \text{pos}(P', P) = w.,$ we have $B' \subset (P' \cap P)U_{P'}$ (see [L10, 2.7]). Since this holds for any Borel B' of R' and R' is the union of its Borels, it follows that $R' \subset (P' \cap P)U_{P'}$. Hence $R = g^{-1}R'g \subset (g^{-1}Pg \cap P)U_P = P_1$. Hence $(R, gU_{P_1}) \in {}^{w,w}Z_{J_\infty,J_1,D}$ is well defined. Repeating the previous argument for (R, gU_{P_1}) instead of (R, gU_P) we see that $R \subset P_2$. Continuing in this way we obtain $R \subset P_r$ for any $r \geq 0$. In particular, $R \subset P_\infty$. Since R, P_∞ are parabolics of the same type, we see that $R = P_\infty$. This proves (b).

Next we show that

(c) *if $y \in {}^{\epsilon(J)}\mathbf{W}$, $y \neq w$, then ${}^{w,y}Z_{J_\infty,J,D} = \emptyset$.*

Assume that $(R, gU_P) \in {}^{w,y}Z_{J_\infty,J,D}$. We have $(R, gU_R) \in {}^wZ_{J_\infty,D}$ and by [L10, 4.14(b)] there exists $(B, h) \in \mathcal{B} \times gU_R$ such that $B \subset R$ and $\text{pos}(hBh^{-1}, B) = w$. We have also $\text{pos}(gBg^{-1}, B) = w$. Since $B \subset P$ and $w \in {}^{\epsilon(J)}\mathbf{W}$, we see, using [L10, 4.14(a)], that $(P, gU_P) \in {}^wZ_{J,D}$. Since $(P, gU_P) \in {}^yZ_{J,D}$ and ${}^yZ_{J,D}, {}^wZ_{J,D}$ are disjoint for $y \neq w$, we have a contradiction. This proves (c).

Let $A \in {}^w\hat{Z}_{J,D}$. From definitions we have $A = \underline{\vartheta}^\star(A')$ with $A' \in {}^w\hat{Z}_{J_\infty,D}$. We show:

(d) $f_{J_\infty,w,J,w}(A') \cong A[-\delta]$ for some integer δ .

(e) $f_{J_\infty,w,J,y}(A') = 0$ for any $y \in {}^{\epsilon(J)}\mathbf{W}$, $y \neq w$.

Now (e) follows immediately from (c). To prove (d) we recall that $f_{J_\infty,w,J,w}(A') = \alpha_1 \tilde{a}^\star(A')$ where $\tilde{a} : {}^{w,w}Z_{J_\infty,J,D} \rightarrow {}^wZ_{J_\infty,D}$ is given by $(R, gU_P) \mapsto (R, gU_R)$. Define

$e : {}^w Z_{J,D} \rightarrow {}^{w,w} Z_{J_\infty,J,D}$ by $e(P, gU_P) = (P_\infty, gU_P)$ with P_∞ as above. Clearly, $\tilde{\alpha}e = \underline{\vartheta}$, $\alpha e = 1$. Since α is an isomorphism, we see that

$$\alpha_! \tilde{\alpha}^*(A') = e^* \tilde{\alpha}^*(A') = \underline{\vartheta}^*(A') = \underline{\vartheta}^\star(A')[-\delta] = A[-\delta]$$

where δ is the dimension of any fibre of $\underline{\vartheta}$. This proves (d).

36.7. Let $L \subset J \subset \mathbf{I}$ and let $w \in {}^{\epsilon(J)}\mathbf{W}$, $y \in {}^{\epsilon(L)}\mathbf{W}$. Assume that $\epsilon^{-1}(\text{Ad}(w)J) = J$, $\epsilon^{-1}(\text{Ad}(y)L) = L$. From definitions we have

$$\begin{aligned} {}^{y,w} Z_{L,J,D} &= \{(R, gU_P); R \in \mathcal{P}_L, P = Q_{J,R}, gU_P \in D/U_P, \\ \text{pos}(gRg^{-1}, R) &= y, \text{pos}(gPg^{-1}, P) = w\}. \end{aligned}$$

We show:

(a) ${}^{y,w} Z_{L,J,D} = \emptyset$ unless $y \in \mathbf{W}_{\epsilon(J)} w = w \mathbf{W}_J$, that is, $w = \min(\mathbf{W}_{\epsilon(J)} y \mathbf{W}_J)$.

Assume that $(R, gU_P) \in {}^{y,w} Z_{L,J,D}$. Let B be a Borel of gRg^{-1} . Then $B \subset gPg^{-1}$. In our case gPg^{-1}, P have a common Levi; hence there exists a Borel B' of P such that $\text{pos}(B, B') = \text{pos}(gPg^{-1}, P) = w$. Similarly, since gRg^{-1}, R have a common Levi, there exists a Borel B'' of R such that $\text{pos}(B, B'') = \text{pos}(gRg^{-1}, R) = y$. Let $z = \text{pos}(B', B'')$. Since $B' \subset P, B'' \subset P$, we have $z \in \mathbf{W}_J$. Since $w \in \mathbf{W}^J$ we have $l(wz) = l(w) + l(z)$ hence from $\text{pos}(B, B') = w, \text{pos}(B', B'') = z$ we deduce $\text{pos}(B, B'') = wz$. Thus $y = wz$ and (a) follows.

Now for $J \subset \mathbf{I}$, $w \in {}^{\epsilon(J)}\mathbf{W}$, $w' \in {}^{\epsilon' \epsilon(J)}\mathbf{W}$, $y \in {}^{\epsilon' \epsilon(J)}\mathbf{W}$, the diagram

$$Z_{J,D} \times Z_{\epsilon(J),D'} \xleftarrow{b_1} Z_0 \xrightarrow{b_2} Z_{J,D'D}$$

(see 32.5) restricts to a diagram

$${}^w Z_{J,D} \times {}^{w'} Z_{\epsilon(J),D'} \xleftarrow{b_{w,w',1}} {}^{w,w',y} Z_0 \xrightarrow{b_{w,w',2}} {}^y Z_{J,D'D}$$

where ${}^{w,w',y} Z_0 = b_1^{-1}({}^w Z_{J,D} \times {}^{w'} Z_{\epsilon(J),D'}) \cap b_2^{-1}({}^y Z_{J,D'D})$. Define a bifunctor $\mathcal{D}({}^w Z_{J,D}) \times \mathcal{D}({}^{w'} Z_{\epsilon(J),D'}) \rightarrow \mathcal{D}({}^y Z_{J,D'D})$ by

$$A, B \mapsto A *_y B = b_{w,w',2}! b_{w,w',1}^*(A \boxtimes B).$$

Let $K \in \mathcal{D}({}^w Z_{J,D}), K' \in \mathcal{D}({}^{w'} Z_{\epsilon(J),D'})$. From definitions we see that

(b)
$$i_{J,y}^*((i_{J,w}! K) * (i_{\epsilon(J),w'}! K')) = K *_y K'.$$

36.8. In the remainder of this section we assume that \mathbf{k} is an algebraic closure of a finite field. As in 31.2 let $\mathcal{A} = \mathbf{Z}[v, v^{-1}]$, v an indeterminate.

Let V be an algebraic variety with a given family of simple perverse sheaves (called “character sheaves”) with the following property: any character sheaf on V comes from a mixed complex on V relative to a rational structure of V over a finite subfield of \mathbf{k} . Let $\mathcal{D}^{cs}(V)$ be the subcategory of $\mathcal{D}(V)$ whose objects are complexes K such that for any j , any composition factor of ${}^p H^j(K)$ is a “character sheaf”. Let $\mathfrak{K}(V)$ be the free \mathcal{A} -module with basis given by the isomorphism class of “character sheaves” on V . Let K be an object of $\mathcal{D}^{cs}(V)$ with a given mixed structure relative to a rational structure of V over a finite subfield of \mathbf{k} . We set

$$gr(K) = \sum_A \sum_{j,h \in \mathbf{Z}} (-1)^j (\text{multiplicity of } A \text{ in } {}^p H^j(K)_h) v^h A \in \mathfrak{K}(V),$$

where A runs over a set of representatives for the isomorphism classes of “character sheaves” on V and the subscript h denotes the subquotient of pure weight h of a mixed perverse sheaf.

Now let V' be another algebraic variety with a given family of simple perverse sheaves (called “character sheaves”) like that of V . Then $\mathcal{D}^{cs}(V')$ is defined. Assume that we are given a functor $\Theta : \mathcal{D}(V) \rightarrow \mathcal{D}(V')$ which restricts to a functor $\mathcal{D}^{cs}(V) \rightarrow \mathcal{D}^{cs}(V')$. We also assume that Θ is a composition of functors of the form $a_!, a^*$ induced by various maps a between algebraic varieties. In particular, Θ preserves the triangulated structures and makes sense also on the mixed level. We define an \mathcal{A} -linear map $gr(\Theta) : \mathfrak{K}(V) \rightarrow \mathfrak{K}(V')$ by the following requirement: If A is a “character sheaf” on V regarded as a mixed complex of pure weight 0, then $gr(\Theta)(A) = gr(\Theta(A))$ where $\Theta(A)$ is regarded as a mixed complex on V' (with mixed structure defined by that of A). Note that $gr(\Theta)(A)$ does not depend on the choice of mixed structure on A . If $\Theta' : \mathcal{D}(V') \rightarrow \mathcal{D}(V'')$ is another functor like Θ , then so is $\Theta'\Theta$ and we have $gr(\Theta'\Theta) = gr(\Theta')gr(\Theta)$.

The previous discussion applies, in particular, to $V = Z_{J,D}$ or ${}^wZ_{J,D}$ where $J \subset \mathbf{I}$ and $w \in {}^{\epsilon(J)}\mathbf{W}$ and to the functors

$$i_{J,w}^* : \mathcal{D}(Z_{J,D}) \rightarrow \mathcal{D}({}^wZ_{J,D}), \quad i_{J,w!} : \mathcal{D}({}^wZ_{J,D}) \rightarrow \mathcal{D}(Z_{J,D})$$

(see 36.3(a), 36.3(f)); the character sheaves on ${}^wZ_{J,D}$ are by definition the simple perverse sheaves isomorphic to ones in ${}^w\hat{Z}_{J,D}$; see 36.3. Hence $\mathfrak{K}(Z_{J,D}), \mathfrak{K}({}^wZ_{J,D})$ are defined and the \mathcal{A} -linear maps

$$gr(i_{J,w}^*) : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}({}^wZ_{J,D}), \quad gr(i_{J,w!}) : \mathfrak{K}({}^wZ_{J,D}) \rightarrow \mathfrak{K}(Z_{J,D})$$

are defined. Since $i_{J,w}^*i_{J,w!} = 1$, we have $gr(i_{J,w}^*)gr(i_{J,w!}) = 1$.

Let $\mathfrak{K}'(Z_{J,D}) = \bigoplus_{w \in {}^{\epsilon(J)}\mathbf{W}} \mathfrak{K}({}^wZ_{J,D})$. Define an \mathcal{A} -linear map $\phi : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}'(Z_{J,D})$ by $\phi = \bigoplus_{w \in {}^{\epsilon(J)}\mathbf{W}} gr(i_{J,w}^*)$.

From 36.3(e) we see that the matrix of ϕ is indexed by $\mathcal{I}_{J,D} \times \mathcal{I}_{J,D}$ and from 36.3(b),(d) we see that this matrix is square and upper triangular (with 1 on diagonal) with respect to a suitable order on $\mathcal{I}_{J,D}$. In particular,

(a) $\phi : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}'(Z_{J,D})$ is an isomorphism.

Note that the inverse isomorphism restricted to $\mathfrak{K}({}^wZ_{J,D})$ is just $gr(i_{J,w!})$. Clearly,

(b) $\mathfrak{K}(Z_{J,D}) = \bigoplus_{w \in {}^{\epsilon(J)}\mathbf{W}} {}^w\mathfrak{K}(Z_{J,D})$,

where ${}^w\mathfrak{K}(Z_{J,D}) = \phi^{-1}(\mathfrak{K}({}^wZ_{J,D}))$.

We have a partition $\mathcal{I}_{J,D} = \bigsqcup_{\mathfrak{k} \in \mathbf{W} \backslash \mathfrak{s}} \mathcal{I}_{J,D}^{\mathfrak{k}}$ where $\mathbf{W} \backslash \mathfrak{s}$ is the set of \mathbf{W} -orbits on \mathfrak{s} and $\mathcal{I}_{J,D}^{\mathfrak{k}}$ consists of those $A \in \mathcal{I}_{J,D}$ such that $A \in \hat{Z}_{J,D}^{\mathcal{L}}$ for some $\mathcal{L} \in \mathfrak{s}$ whose isomorphism class is in \mathfrak{k} . We have

(c) $\mathfrak{K}(Z_{J,D}) = \bigoplus_{\mathfrak{k} \in \mathbf{W} \backslash \mathfrak{s}} \mathfrak{K}^{\mathfrak{k}}(Z_{J,D})$

where $\mathfrak{K}^{\mathfrak{k}}(Z_{J,D})$ is the \mathcal{A} -submodule of $\mathfrak{K}(Z_{J,D})$ generated by $\mathcal{I}_{J,D}^{\mathfrak{k}}$. From definitions we see that each $\mathfrak{K}({}^wZ_{J,D})$ and each ${}^w\mathfrak{K}(Z_{J,D})$ has a natural direct sum decomposition indexed by $\mathbf{W} \backslash \mathfrak{s}$ analogous to (c). It follows that the decompositions (b),(c) are compatible in the sense that

(d) $\mathfrak{K}(Z_{J,D}) = \bigoplus_{w, \mathfrak{k}} ({}^w\mathfrak{K}(Z_{J,D}) \cap \mathfrak{K}^{\mathfrak{k}}(Z_{J,D}))$.

36.9. Let $J \subset J' \subset \mathbf{I}$. As in [L10, 6.4, 6.7(b)] we see that $f_{J,J'}, e_{J,J'}$ restrict to functors $\mathcal{D}^{cs}(Z_{J,D}) \rightarrow \mathcal{D}^{cs}(Z_{J',D}), \mathcal{D}^{cs}(Z_{J',D}) \rightarrow \mathcal{D}^{cs}(Z_{J,D})$. Hence the \mathcal{A} -linear maps $gr(f_{J,J'}) : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J',D}), gr(e_{J,J'}) : \mathfrak{K}(Z_{J',D}) \rightarrow \mathfrak{K}(Z_{J,D})$ are defined as in 36.8; we denote them again by $f_{J,J'}, e_{J,J'}$. The identities 36.4(b) continue to hold for these linear maps. We define a group homomorphism $\mathfrak{D} : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J,D})$ by

$\mathfrak{D}(v^m A) = v^{-m} A^*$ where $A^* \in \mathcal{I}_{J,D}$ is isomorphic to the Verdier dual of $A \in \mathcal{I}_{J,D}$. From 36.4(a) we deduce

$$(a) \quad \mathfrak{f}_{J,J'} = v^{-2a} \mathfrak{f}_{J,J'} \mathfrak{D} : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J',D})$$

with a as in 36.4(a).

If $J \subset J'$, $\mathfrak{c}, \mathfrak{d}, a$ are as in 36.4, then for $A \in \mathcal{I}_{J,D}$ we have

$$\mathfrak{f}_{J,J'} A = \sum_{A_1 \in \mathcal{I}_{J',D}} x_{A,A_1} A_1,$$

$$(b) \quad x_{A,A_1} = \sum_j (-v)^j (\text{multiplicity of } A_1 \text{ in } {}^p H^{j-a}(\mathfrak{d}_! \mathfrak{c}^* A[a])) \in \mathcal{A}.$$

(Indeed, if we regard A as a pure perverse sheaf of weight 0, then $\mathfrak{c}^* A[a]$ is a pure perverse sheaf of weight a (since \mathfrak{c} is an affine space bundle with fibres of dimension a); by [De, 6.2.6] applied to the proper morphism \mathfrak{d} , $\mathfrak{d}_! \mathfrak{c}^* A[a]$ is a pure complex of weight a . From [BBD, 5.4.1], we see that ${}^p H^j(\mathfrak{d}_! \mathfrak{c}^* A) = {}^p H^{j-a}(\mathfrak{d}_! \mathfrak{c}^* A[a])$ is pure of weight $(j - a) + a = j$.)

Let $\xi \mapsto \tilde{\xi}$ be the group homomorphism $\mathcal{A} \rightarrow \mathcal{A}$ given by $v^m \mapsto v^{-m}$ for $m \in \mathbf{Z}$. Since $\mathfrak{c}^* A[a]$ is a pure perverse sheaf, we have (using the relative hard Lefschetz theorem [BBD, 5.4.10]):

$$\begin{aligned} \bar{x}_{A,A_1} &= \sum_j (-v)^{-j} (\text{multiplicity of } A_1 \text{ in } {}^p H^{j-a}(\mathfrak{d}_! \mathfrak{c}^* A[a])) \\ &= \sum_j (-v)^{-j} (\text{multiplicity of } A_1 \text{ in } {}^p H^{-j+a}(\mathfrak{d}_! \mathfrak{c}^* A[a])) \\ &= \sum_{j'} (-v)^{j'-2a} (\text{multiplicity of } A_1 \text{ in } {}^p H^{j'-a}(\mathfrak{d}_! \mathfrak{c}^* A[a])) = v^{-2a} x_{A,A_1}. \end{aligned}$$

Define a group homomorphism $\beta_J : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J,D})$ by $\beta_J(v^m A) = v^{-m} A$ for $A \in \mathcal{I}_{J,D}, m \in \mathbf{Z}$. We see that $\beta_{J'}(\mathfrak{f}_{J,J'} A) = v^{-2a} \mathfrak{f}_{J,J'} A$ for any $A \in \mathcal{I}_{J,D}$. Hence for any $\xi \in \mathfrak{K}(Z_{J,D})$ we have

$$(c) \quad \beta_{J'}(\mathfrak{f}_{J,J'} \xi) = v^{-2a} \mathfrak{f}_{J,J'}(\beta_J(\xi)).$$

Define $\tilde{\mathfrak{D}} : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J,D})$ by $\tilde{\mathfrak{D}}(\xi) = \mathfrak{D}\beta_J(\xi) = \beta_J \mathfrak{D}(\xi)$. Clearly, $\tilde{\mathfrak{D}}$ is \mathcal{A} -linear. Note that

(d) *the maps $\mathfrak{f}_{J,J'}, \mathfrak{e}_{J,J'}, \beta_J$ are compatible with the decompositions of type 36.8(c) while $\mathfrak{D}, \tilde{\mathfrak{D}}$ map the summand corresponding to \mathfrak{k} in 36.8(c) onto the summand corresponding to $\tilde{\mathfrak{k}}$. (Here $\tilde{\mathfrak{k}}$ is the image of \mathfrak{k} under $\mathcal{L} \mapsto \tilde{\mathcal{L}}$; see 28.18.)*

Let $J, D, D', b_1, b_2, \epsilon$ be as in 32.5. By 32.21, $b_2! b_1^* : \mathcal{D}(Z_{J,D} \times Z_{\epsilon(J),D'}) \rightarrow \mathcal{D}(Z_{J,D'D})$ restricts to a functor $\mathcal{D}^{cs}(Z_{J,D} \times Z_{\epsilon(J),D'}) \rightarrow \mathcal{D}^{cs}(Z_{J,D'D})$, where the character sheaves on $Z_{J,D} \times Z_{\epsilon(J),D'}$ are by definition complexes of the form $A \boxtimes A'$ with $A \in \hat{Z}_{J,D}, A' \in \hat{Z}_{\epsilon(J),D'}$. Hence the \mathcal{A} -linear map $gr(b_2! b_1^*) : \mathfrak{K}(Z_{J,D} \times Z_{\epsilon(J),D'}) \rightarrow \mathfrak{K}(Z_{J,D'D})$ or equivalently $\mathfrak{K}(Z_{J,D}) \otimes_{\mathcal{A}} \mathfrak{K}(Z_{\epsilon(J),D'}) \rightarrow \mathfrak{K}(Z_{J,D'D})$ is well defined. (We have canonically $\mathfrak{K}(Z_{J,D} \times Z_{\epsilon(J),D'}) = \mathfrak{K}(Z_{J,D}) \otimes_{\mathcal{A}} \mathfrak{K}(Z_{\epsilon(J),D'})$.) We write $\xi * \xi'$ instead of $gr(b_2! b_1^*)(\xi \boxtimes \xi')$ where $\xi \in \mathfrak{K}(Z_{J,D}), \xi' \in \mathfrak{K}(Z_{\epsilon(J),D'})$. For $\mathfrak{k}, \mathfrak{k}' \in \mathbf{W} \setminus \underline{\mathfrak{s}}$ and $\xi \in \mathfrak{K}^{\mathfrak{k}}(Z_{J,D}), \xi' \in \mathfrak{K}^{\mathfrak{k}'}(Z_{\epsilon(J),D'})$, we have $\xi * \xi' \in \mathfrak{K}^{\mathfrak{k}}(Z_{J,D'D})$ if $\mathfrak{k}' = \underline{D}(\mathfrak{k})$ and $\xi * \xi' = 0$ if $\mathfrak{k}' \neq \underline{D}(\mathfrak{k})$ (see 32.6(a),(b)).

From 28.19 we see that $\partial_! : \mathcal{D}(Z_{J,D}) \rightarrow \mathcal{D}(Z_{\epsilon(J),D-1})$ restricts to an equivalence of categories $\mathcal{D}^{cs}(Z_{J,D}) \rightarrow \mathcal{D}^{cs}(Z_{\epsilon(J),D-1})$. Hence the \mathcal{A} -linear isomorphism

$gr(\partial_1) : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{\epsilon(J),D-1})$ is well defined; we denote it again by ∂ . Now the composition $\mathfrak{K}(Z_{J,D}) \xrightarrow{\partial} \mathfrak{K}(Z_{\epsilon(J),D-1}) \xrightarrow{\partial} \mathfrak{K}(Z_{J,D})$ is the identity map.

In the setup of 36.5, $f_{J,w,J',w'}$ restricts to a functor $\mathcal{D}^{cs}({}^w Z_{J,D}) \rightarrow \mathcal{D}^{cs}({}^{w'} Z_{J',D})$. (We use the analogous statement for $f_{J,J'}$, as above, together with 36.5(a), 36.3(a), 36.3(f).) Hence the \mathcal{A} -linear map $gr(f_{J,w,J',w'}) : \mathfrak{K}({}^w Z_{J,D}) \rightarrow \mathfrak{K}({}^{w'} Z_{J',D})$ is well defined. We denote it again by $f_{J,w,J',w'}$. Let $f'_{J,J'} : \mathfrak{K}'(Z_{J,D}) \rightarrow \mathfrak{K}'(Z_{J',D})$ be the unique \mathcal{A} -linear map such that $pr_{w'} f'_{J,J'} pr_w = f_{J,w,J',w'}$ for any w, w' as above. (Here pr_w is the projection of $\mathfrak{K}'(Z_{J',D})$ onto the direct summand $\mathfrak{K}({}^w Z_{J',D})$.) From 36.5(a) we see that for $x \in \mathfrak{K}({}^w Z_{J',D})$ we have

$$(e) \quad f'_{J,J'} = \phi f_{J,J'} \phi^{-1}.$$

We also have

$$(f) \quad \partial f_{J,J'} = f_{\epsilon(J),\epsilon(J')} \partial, \quad \partial \mathfrak{e}_{J,J'} = \mathfrak{e}_{\epsilon(J),\epsilon(J')} \partial.$$

Now the functor $\tau : \mathcal{D}(Z_{J,G^0}) \rightarrow \mathcal{D}(\text{point})$ (see 36.4) restricts to a functor $\mathcal{D}^{cs}(Z_{J,G^0}) \rightarrow \mathcal{D}^{cs}(\text{point})$ where the point is regarded as having exactly one character sheaf, $\bar{\mathbf{Q}}_l$. Hence the \mathcal{A} -linear map $gr(\tau) : \mathfrak{K}(Z_{J,G^0}) \rightarrow \mathfrak{K}(\text{point}) = \mathcal{A}$ is well defined; we denote it again by τ .

Let p, i' be as in 36.4. Then $p_i i'^* : \mathcal{D}(Z_{J,D} \times Z_{J,D}) \rightarrow \mathcal{D}(\text{point})$ restricts to a functor $\mathcal{D}^{cs}(Z_{J,D} \times Z_{J,D}) \rightarrow \mathcal{D}^{cs}(\text{point})$ where the character sheaves on $Z_{J,D} \times Z_{J,D}$ are by definition complexes of the form $A \boxtimes A'$ with $A, A' \in \hat{Z}_{J,D}$. Hence the \mathcal{A} -linear map $gr(p_i i'^*) : \mathfrak{K}(Z_{J,D} \times Z_{J,D}) \rightarrow \mathfrak{K}(\text{point}) = \mathcal{A}$ or equivalently $\mathfrak{K}(Z_{J,D}) \otimes_{\mathcal{A}} \mathfrak{K}(Z_{J,D}) \rightarrow \mathcal{A}$ is well defined. (We have canonically $\mathfrak{K}(Z_{J,D} \times Z_{J,D}) = \mathfrak{K}(Z_{J,D}) \otimes_{\mathcal{A}} \mathfrak{K}(Z_{J,D})$.) We write $(\xi : \xi')$ instead of $gr(p_i i'^*)(\xi \boxtimes \xi')$ where $\xi, \xi' \in \mathfrak{K}(Z_{J,D})$. From [BBD, 5.1.14] we see that for $A, B \in \mathcal{I}_{J,D}$ we have $(A : B) \in \mathbf{Z}[v^{-1}]$ and from [L3, II,7.4] and its proof we see that the constant term of $(A : B)$ is δ_{A,B^*} . Thus,

$$(g) \quad (A : B) \in \delta_{A,B^*} + v^{-1} \mathbf{Z}[v^{-1}].$$

We show that

(h) if $\xi \in \mathfrak{K}(Z_{J,D})$, $(\xi : \tilde{\mathfrak{D}}(\xi)) = 0$, then $\xi = 0$.

Assume that $\xi \neq 0$. We have $\xi = \sum_{A \in \mathcal{I}_{J,D}, m \in \mathbf{Z}} g_{A,m} v^m A$ where $g_{A,m} \in \mathbf{Z}$ is zero for all but finitely many A, m . We can find $e \in \mathbf{Z}$ such that $g_{A,e} \neq 0$ for some $A \in \mathcal{I}_{J,D}$ and $g_{A,m} = 0$ for all $m > e$ and all $A \in \mathcal{I}_{J,D}$. Then

$$(\xi : \tilde{\mathfrak{D}}(\xi)) = \sum_{A_1, A, m_1, m} g_{A_1, m_1} g_{A, m} v^{m_1+m} (A_1 : A^*).$$

By (g) this equals $\sum_A g_{A,e}^2 v^{2e} +$ an element in $v^{2e-1} \mathbf{Z}[v^{-1}]$. This is nonzero since $\sum_A g_{A,e}^2 \in \mathbf{Z}_{>0}$. This proves (h).

We show that

(i) if $\mathfrak{k}, \mathfrak{k}' \in \mathbf{W} \setminus \underline{\mathfrak{s}}$ and $\mathfrak{k}' \neq \mathfrak{k}$, then $(\mathfrak{K}^{\mathfrak{k}}(Z_{J,D}) : \mathfrak{K}^{\mathfrak{k}'}(Z_{J,D})) = 0$.

Let $A \in \hat{Z}_{J,D}^{\mathcal{L}}, B \in \hat{Z}_{J,D}^{\mathcal{L}_1}$ where $\mathcal{L}, \mathcal{L}_1 \in \mathfrak{s}$. It is enough to show that

if $H_c^j(Z_{J,D}, A \otimes B) \neq 0$ for some j , then the isomorphism class of \mathcal{L}_1 is in the \mathbf{W} -orbit of the isomorphism class of $\check{\mathcal{L}}$.

From our assumption we have $A * \partial_1(B) \neq 0$. We have $A \dashv \bar{K}_{J,D}^{\bar{\mathfrak{s}}, \mathcal{L}}$ as in 28.13(v) and similarly $B \dashv \bar{K}_{J,D}^{\bar{\mathfrak{s}}', \mathcal{L}_1}$. Hence $\partial_1 B \dashv \bar{K}_{\epsilon(J), D-1}^{\bar{\mathfrak{s}}', \mathcal{L}''}$ where $\mathcal{L}'' = (\underline{D}^{-1})^* \check{\mathcal{L}}$, see 28.19. We

then have $\bar{K}_{J,D}^{\bar{s},\mathcal{L}} * \bar{K}_{\epsilon^{(J)},D^{-1}}^{\bar{s}',\mathcal{L}''} \neq 0$. Using 32.6(a) we see that $\mathcal{L}, \mathcal{L}_1$ have the required property. This proves (i).

We show that (j) if $w, w' \in \mathbf{W}$ and $w \neq w'$, then $({}^w\mathfrak{R}(Z_{J,D}) : {}^{w'}\mathfrak{R}(Z_{J,D})) = 0$.

It is enough to show that if $A \in \mathcal{D}^{cs}({}^wZ_{J,D}), A' \in \mathcal{D}^{cs}({}^{w'}Z_{J,D})$, then

$$p!i'^*(i_{J,w!}A \boxtimes i_{J,w'!}A') = 0$$

with p, i' as in 36.4. It is enough to show that $(i_{J,w} \times i_{J,w'})!i'^*(A \boxtimes A') = 0$. This follows from the fact that ${}^wZ_{J,D} \cap {}^{w'}Z_{J,D} = \emptyset$.

We show that

(k) if $w \in \epsilon^{(J)}\mathbf{W}$, $w \neq 1$, then $\tau({}^w\mathfrak{R}(Z_{J,G^0})) = 0$.

Since ${}^wZ_{J,G^0}, {}^1Z_{J,G^0}$ are disjoint, we have $i_{J,1}^*i_{J,w!} = 0$, hence $i^*i_{J,w!} = 0$ with i as in 36.4; (k) follows.

From the definitions we have (l) $\partial({}^w\mathfrak{R}(Z_{J,D})) = w^{-1}\mathfrak{R}(Z_{\epsilon^{(J)},D^{-1}})$

for any $w \in \epsilon^{(J)}\mathbf{W}$ such that $\epsilon^{-1}(\text{Ad}(w)J) = J$. (For such w we have $w^{-1} \in {}^J\mathbf{W}$.)

36.10. Let

$$\tilde{\mathfrak{R}}(Z_{J,D}) = \sum_{L; L \subsetneq J} f_{L,J}(\mathfrak{R}(Z_{L,D})), \mathfrak{R}_0 = \bigoplus_{w \in \epsilon^{(J)}\mathbf{W}; \epsilon^{-1}(\text{Ad}(w)J) \neq J} {}^w\mathfrak{R}(Z_{J,D}).$$

We show that

$$(a) \quad \tilde{\mathfrak{R}}(Z_{J,D}) = \mathfrak{R}_0 \oplus \bigoplus_{\substack{w \in \epsilon^{(J)}\mathbf{W} \\ \epsilon^{-1}(\text{Ad}(w)J) = J}} (\tilde{\mathfrak{R}}(Z_{J,D}) \cap {}^w\mathfrak{R}(Z_{J,D})).$$

Setting

$$\tilde{\mathfrak{R}}'(Z_{J,D}) = \sum_{L; L \subsetneq J} f'_{L,J}(\mathfrak{R}'(Z_{L,D})), \mathfrak{R}'_0 = \bigoplus_{w \in \epsilon^{(J)}\mathbf{W}; \epsilon^{-1}(\text{Ad}(w)J) \neq J} \mathfrak{R}'({}^wZ_{J,D})$$

and using 36.9(e), we see that it is enough to show that

$$(b) \quad \tilde{\mathfrak{R}}'(Z_{J,D}) = \mathfrak{R}'_0 \oplus \bigoplus_{\substack{w \in \epsilon^{(J)}\mathbf{W} \\ \epsilon^{-1}(\text{Ad}(w)J) = J}} (\tilde{\mathfrak{R}}'(Z_{J,D}) \cap \mathfrak{R}'({}^wZ_{J,D})).$$

From 36.6(d), 36.6(e) we see that, if $w \in \epsilon^{(J)}\mathbf{W}, \epsilon^{-1}(\text{Ad}(w)J) \neq J$, then $A = f'_{J_\infty, J}A'$ for some $A' \in \mathfrak{R}'({}^yZ_{L',D})$ and $J_\infty \neq J$. Hence

$$(c) \quad \mathfrak{R}'_0 \subset \tilde{\mathfrak{R}}'(Z_{J,D}).$$

Using again 36.6(d), 36.6(e) and also 36.4(b) (or rather its analogue for $f'_{J,J'}$) and (c), we see that

$$\tilde{\mathfrak{R}}'(Z_{J,D}) = \sum_{\substack{L; L \subsetneq J \\ y \in \epsilon^{(L)}\mathbf{W} \\ \epsilon^{-1}(\text{Ad}(y)L) = L}} f'_{L,J}(\mathfrak{R}'({}^yZ_{L,D})) = \mathfrak{R}'_0 + \sum_{\substack{L; L \subsetneq J \\ y \in \epsilon^{(L)}\mathbf{W} \\ \epsilon^{-1}(\text{Ad}(y)L) = L}} f'_{L,J}(\mathfrak{R}'({}^yZ_{L,D})).$$

Thus, (b) holds and (a) holds.

Let

$$\bar{\mathfrak{R}}(Z_{J,D}) = \mathfrak{R}(Z_{J,D}) / \tilde{\mathfrak{R}}(Z_{J,D}).$$

For any $w \in {}^{\epsilon(J)}\mathbf{W}$ such that $\epsilon^{-1}(\text{Ad}(w)J) = J$, let ${}^w\bar{\mathfrak{K}}(Z_{J,D})$ be the image of ${}^w\mathfrak{K}(Z_{J,D})$ under the obvious map $\mathfrak{K}(Z_{J,D}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D})$. From (a) we see that

$$(d) \quad \bar{\mathfrak{K}}(Z_{J,D}) = \bigoplus_{\substack{w \in {}^{\epsilon(J)}\mathbf{W} \\ \epsilon^{-1}(\text{Ad}(w)J) = J}} {}^w\bar{\mathfrak{K}}(Z_{J,D}).$$

For any $\mathfrak{k} \in \mathbf{W} \setminus \underline{\mathfrak{g}}$ let $\bar{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D})$ be the image of $\mathfrak{K}^{\mathfrak{k}}(Z_{J,D})$ under the obvious map $\mathfrak{K}(Z_{J,D}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D})$. From 36.9(d) we see that $\bar{\mathfrak{K}}(Z_{J,D})$ is compatible with the decomposition 36.8(c). It follows that

$$(e) \quad \bar{\mathfrak{K}}(Z_{J,D}) = \bigoplus_{\mathfrak{k} \in \mathbf{W} \setminus \underline{\mathfrak{g}}} \bar{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D}).$$

Using 36.8(d) we see that that the decompositions (d),(e) are compatible in the sense that

$$(f) \quad \bar{\mathfrak{K}}(Z_{J,D}) = \bigoplus_{w, \mathfrak{k}} ({}^w\bar{\mathfrak{K}}(Z_{J,D}) \cap \bar{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D})).$$

Using 36.9(a), 36.9(c), we see that

(g) $\mathfrak{D}, \beta_J, \tilde{\mathfrak{D}} : \mathfrak{K}(Z_{J,D}) \rightarrow \mathfrak{K}(Z_{J,D})$ map $\tilde{\mathfrak{K}}(Z_{J,D})$ into itself hence induces group homomorphisms $\bar{\mathfrak{K}}(Z_{J,D}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D})$ denoted again by $\mathfrak{D}, \beta_J, \tilde{\mathfrak{D}}$.

36.11. For $J' \subset J \subset \mathbf{I}$ and $x \in \mathfrak{K}(Z_{J',D}), x' \in \mathfrak{K}(Z_{\epsilon(J),D'})$, we show that

$$(a) \quad (\mathfrak{f}_{J',J}x) * x' = \mathfrak{f}_{J',J}(x * \mathfrak{e}_{\epsilon(J'),\epsilon(J)}x').$$

Using successively 36.4(h), 36.9(f), 36.4(i), 36.4(c), 36.9(f), 36.4(h), 36.4(i), we see that, for any $x'' \in \mathfrak{K}(Z_{J,D'D})$, we have

$$\begin{aligned} ((\mathfrak{f}_{J',J}x) * x' : x'') &= ((\partial \mathfrak{f}_{J',J}x) : x' * \partial x'') \\ &= (\mathfrak{f}_{\epsilon(J'),\epsilon(J)}\partial(x) : x' * (\partial x'')) = (\partial x : \mathfrak{e}_{\epsilon(J'),\epsilon(J)}(x' * (\partial x''))) \\ &= (\partial x : \mathfrak{e}_{\epsilon(J'),\epsilon(J)}(x') * \mathfrak{e}_{\epsilon(J'),\epsilon(J)}(\partial x'')) = (\partial x : \mathfrak{e}_{\epsilon(J'),\epsilon(J)}x' * \partial \mathfrak{e}_{J',J}x'') \\ &= (x * \mathfrak{e}_{\epsilon(J'),\epsilon(J)}x' : \mathfrak{e}_{J',J}x'') = (\mathfrak{f}_{J',J}(x * \mathfrak{e}_{\epsilon(J'),\epsilon(J)}x') : x''). \end{aligned}$$

Thus, if $\xi = (\mathfrak{f}_{J',J}x) * x' - \mathfrak{f}_{J',J}(x * \mathfrak{e}_{\epsilon(J'),\epsilon(J)}x')$, then $(\xi : x'') = 0$ for any $x'' \in \mathfrak{K}(Z_{J,D'D})$. In particular, $(\xi : \tilde{\mathfrak{D}}(\xi)) = 0$. Using 36.9(h) we see that $\xi = 0$. This proves (a).

A similar argument shows that, if $x \in \mathfrak{K}(Z_{J,D}), x' \in \mathfrak{K}(Z_{\epsilon(J'),D'})$, then

$$(b) \quad x * (\mathfrak{f}_{\epsilon(J'),\epsilon(J)}x') = \mathfrak{f}_{J',J}(\mathfrak{e}_{J',J}x * x').$$

From (a),(b) we see that

$$\tilde{\mathfrak{K}}(Z_{J,D}) * \mathfrak{K}(Z_{\epsilon(J),D'}) \subset \tilde{\mathfrak{K}}(Z_{J,D'D}), \quad \mathfrak{K}(Z_{J,D}) * \tilde{\mathfrak{K}}(Z_{\epsilon(J),D'}) \subset \tilde{\mathfrak{K}}(Z_{J,D'D}).$$

It follows that $\mathfrak{K}(Z_{J,D}) \times \mathfrak{K}(Z_{\epsilon(J),D'}) \rightarrow \mathfrak{K}(Z_{J,D'D}), A, B \mapsto A * B$, induces an \mathcal{A} -bilinear pairing $\tilde{\mathfrak{K}}(Z_{J,D}) \times \tilde{\mathfrak{K}}(Z_{\epsilon(J),D'}) \rightarrow \tilde{\mathfrak{K}}(Z_{J,D'D})$. We denote it again by $A, B \mapsto A * B$. The following result relates this bilinear pairing to the decompositions of type 36.10(d).

For $w \in {}^{\epsilon(J)}\mathbf{W}, w' \in {}^{\epsilon'\epsilon(J)}\mathbf{W}$ such that $\epsilon^{-1}(\text{Ad}(w)J) = J, \epsilon'^{-1}(\text{Ad}(w')\epsilon(J)) = \epsilon(J)$, let $X_{w,w'}$ be the set of all $y \in {}^{\epsilon'\epsilon(J)}\mathbf{W}$ such that $\epsilon^{-1}\epsilon'^{-1}(\text{Ad}(y)J) = J$ and such that for some $Q \in \mathcal{P}_J, Q' \in \mathcal{P}_{\epsilon(J)}, Q'' \in \mathcal{P}_{\epsilon'\epsilon(J)}$ we have

$$\text{pos}(Q', Q) = w, \text{pos}(Q'', Q') = w', \text{pos}(Q'', Q) = y.$$

Then

$$(c) \quad {}^w \bar{\mathfrak{K}}(Z_{J,D}) * {}^{w'} \bar{\mathfrak{K}}(Z_{\epsilon(J),D'}) \subset \bigoplus_{y \in X_{w,w'}} {}^y \bar{\mathfrak{K}}(Z_{J,D'}).$$

Indeed, using 36.7(a) we see that it is enough to show that

if ${}^{w,w'} {}^y Z_0 \neq \emptyset$ (notation of 36.7), then $y \in X_{w,w'}$.

This is immediate from definitions.

36.12. As a special case of the pairing $A, B \mapsto A * B$ in 36.11 we have an \mathcal{A} -bilinear pairing $\bar{\mathfrak{K}}(Z_{J,G^0}) \times \bar{\mathfrak{K}}(Z_{J,G^0}) \rightarrow \bar{\mathfrak{K}}(Z_{J,G^0})$. This defines an associative algebra structure on $\bar{\mathfrak{K}}(Z_{J,G^0})$ (not necessarily with 1).

Also as special cases of the pairing $A, B \mapsto A * B$ in 36.11 we have \mathcal{A} -bilinear pairings

$$\bar{\mathfrak{K}}(Z_{J,G^0}) \times \bar{\mathfrak{K}}(Z_{J,D}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D}), \quad \bar{\mathfrak{K}}(Z_{J,D}) \times \bar{\mathfrak{K}}(Z_{\epsilon(J),G^0}) \rightarrow \bar{\mathfrak{K}}(Z_{J,D}),$$

which make $\bar{\mathfrak{K}}(Z_{J,D})$ into a (not necessarily unital) $(\bar{\mathfrak{K}}(Z_{J,G^0}), \bar{\mathfrak{K}}(Z_{\epsilon(J),G^0}))$ -bimodule.

Let \heartsuit be the \mathbf{W} -orbit $\{\mathbf{Q}_I\}$ in $\underline{\mathfrak{g}}$. By 36.10(f) we have

$$\bar{\mathfrak{K}}^\heartsuit(Z_{J,G^0}) = \bigoplus_{\substack{w \in \epsilon^{(J)} \mathbf{W} \\ \epsilon^{-1}(\text{Ad}(w)J) = J}} ({}^w \bar{\mathfrak{K}}(Z_{J,G^0}) \cap \bar{\mathfrak{K}}^\heartsuit(Z_{J,G^0})).$$

From 36.10(g) we see that $\mathfrak{D}, \beta_J, \tilde{\mathfrak{D}}$ may be regarded as group homomorphisms $\bar{\mathfrak{K}}^\heartsuit(Z_{J,G^0}) \rightarrow \bar{\mathfrak{K}}^\heartsuit(Z_{J,G^0})$.

36.13. For any \mathcal{A} -module V we set $\underline{V} = \mathbf{Q}(v) \otimes_{\mathcal{A}} V$. If V, V' are \mathcal{A} -modules and $f : V \rightarrow V'$ is \mathcal{A} -linear we denote again by f the $\mathbf{Q}(v)$ -linear map $\underline{V} \rightarrow \underline{V}'$ such that $1 \otimes x \mapsto 1 \otimes f(x)$ for $x \in V$.

In particular, the $\mathbf{Q}(v)$ -vector spaces $\bar{\mathfrak{K}}(Z_{J,D}), \tilde{\mathfrak{K}}(Z_{J,D})$ and $\underline{\mathfrak{K}}(Z_{J,D})$ are defined. Note that $\tilde{\mathfrak{K}}(Z_{J,D}) = \sum_{L; L \subsetneq J} f_{L,J}(\bar{\mathfrak{K}}(Z_L,D))$, $\underline{\mathfrak{K}}(Z_{J,D}) = \bigoplus_{\mathfrak{k} \in \mathbf{W} \setminus \underline{\mathfrak{g}}} \bar{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D})$; see 36.8(c).

The symmetric bilinear form $(:): \bar{\mathfrak{K}}(Z_{J,D}) \times \bar{\mathfrak{K}}(Z_{J,D}) \rightarrow \mathcal{A}$ extends uniquely to a symmetric $\mathbf{Q}(v)$ -bilinear form $\underline{\mathfrak{K}}(Z_{J,D}) \times \underline{\mathfrak{K}}(Z_{J,D}) \rightarrow \mathbf{Q}(v)$ denoted again by $(:)$.

A vector subspace E of $\underline{\mathfrak{K}}(Z_{J,D})$ is said to be *homogeneous* if $E = \sum_{\mathfrak{k} \in \mathbf{W} \setminus \underline{\mathfrak{g}}} E^{\mathfrak{k}}$ where $E^{\mathfrak{k}} = E \cap \bar{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D})$. For such E we set $E^\perp = \{x \in \underline{\mathfrak{K}}(Z_{J,D}); (x : E) = 0\}$. Then E^\perp is homogeneous. We show that if, in addition, E is stable under $\tilde{\mathfrak{D}}$, then:
(a) $\underline{\mathfrak{K}}(Z_{J,D}) = E \oplus E^\perp$.

We first show that $E \cap E^\perp = 0$. Assume that $x \in E \cap E^\perp$. Then $f\tilde{\mathfrak{D}}(x) \in E$ and $(x : \tilde{\mathfrak{D}}(x)) = 0$. We can find $\lambda \in \mathcal{A} - \{0\}$ such that $\lambda x \in \bar{\mathfrak{K}}(Z_{J,D})$. Then $(\lambda x, \tilde{\mathfrak{D}}(\lambda x)) = 0$. By 36.9(h) we have $\lambda x = 0$. Hence $x = 0$ so that $E \cap E^\perp = 0$. It remains to show that $\dim(E^\perp)^{\mathfrak{k}} + \dim E^{\mathfrak{k}} = \dim \bar{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D})$ for any \mathfrak{k} . This is a consequence of the following statement:

(b) *For any \mathfrak{k} , the form $(:)$ on the finite dimensional vector space $\bar{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D}) + \tilde{\mathfrak{K}}^{\mathfrak{k}}(Z_{J,D})$ is nonsingular.*

More generally, we show that for any finite dimensional subspace E' of $\underline{\mathfrak{K}}(Z_{J,D})$ which is stable under $\tilde{\mathfrak{D}}$, the form $(:)$ on E' is nonsingular. Let $x \in E'$ be such that $(x : x') = 0$ for any $x' \in E'$. In particular, we have $(x, \tilde{\mathfrak{D}}(x)) = 0$. As above, this implies that $x = 0$. This proves (b), hence also (a).

We set

$$\underline{\mathfrak{K}}(Z_{J,D})^J = \{\xi \in \underline{\mathfrak{K}}(Z_{J,D}); \mathfrak{e}_{H,J}\xi = 0 \quad \forall H \subsetneq J\}.$$

We show:

(c)
$$\tilde{\mathfrak{K}}(Z_{J,D})^\perp = \underline{\mathfrak{K}}(Z_{J,D})^J.$$

We have

$$\begin{aligned} \tilde{\mathfrak{K}}(Z_{J,D}) &= \{x \in \underline{\mathfrak{K}}(Z_{J,D}); (x : \mathfrak{f}_{H,J}\underline{\mathfrak{K}}(Z_{H,D})) = 0 \quad \forall H \subsetneq J\} \\ &= \{\xi \in \underline{\mathfrak{K}}(Z_{J,D}); (\mathfrak{e}_{H,J}x : \underline{\mathfrak{K}}(Z_{H,D})) = 0 \quad \forall H \subsetneq J\} \\ &= \{\xi \in \underline{\mathfrak{K}}(Z_{J,D}); \mathfrak{e}_{H,J}x = 0 \quad \forall H \subsetneq J\}, \end{aligned}$$

as required. (We have used that $\underline{\mathfrak{K}}(Z_{H,D})^\perp = 0$ which follows from (a), applied to H instead of J .) This proves (c).

We show that

(d)
$$\underline{\mathfrak{K}}(Z_{J,D}) = \tilde{\mathfrak{K}}(Z_{J,D}) \oplus \underline{\mathfrak{K}}(Z_{J,D})^J.$$

This follows from (a),(c) using the fact that $\tilde{\mathfrak{K}}(Z_{J,D})$ is stable under $\tilde{\mathfrak{D}}$; see 36.10(g).

From (d) we see that the second projection $\underline{\mathfrak{K}}(Z_{J,D}) \rightarrow \underline{\mathfrak{K}}(Z_{J,D})^J$ induces an isomorphism $\underline{\mathfrak{K}}(Z_{J,D})/\tilde{\mathfrak{K}}(Z_{J,D}) \xrightarrow{\sim} \underline{\mathfrak{K}}(Z_{J,D})^J$, that is, an isomorphism

(e)
$$\tilde{\mathfrak{K}}(Z_{J,D}) \xrightarrow{\sim} \underline{\mathfrak{K}}(Z_{J,D})^J.$$

36.14. The \mathcal{A} -algebra structure on $\mathfrak{K}(Z_{J,G^0})$ (resp. $\tilde{\mathfrak{K}}(Z_{J,G^0})$) given by $*$ extends to a $\mathbf{Q}(v)$ -algebra structure on $\underline{\mathfrak{K}}(Z_{J,G^0})$ (resp. $\tilde{\underline{\mathfrak{K}}}(Z_{J,G^0})$) denoted again by $*$. Moreover, $\tilde{\underline{\mathfrak{K}}}(Z_{J,G^0})$ is a two-sided ideal of $\underline{\mathfrak{K}}(Z_{J,G^0})$ (see 36.11). Note that $\underline{\mathfrak{K}}(Z_{J,D})^J$ is also a two-sided ideal of $\underline{\mathfrak{K}}(Z_{J,G^0})$, since $\mathfrak{e}_{H,J} : \underline{\mathfrak{K}}(Z_{J,G^0}) \rightarrow \underline{\mathfrak{K}}(Z_{H,G^0})$ is an algebra homomorphism for any $H \subset J$ (see 36.4(c)). Since the two summands in the right-hand side of 36.13(d) are two-sided ideals, they annihilate each other under the product $*$. We see also that the isomorphism 36.13(e) respects the algebra structures.

From 36.9(f) we see that the two summands in the right-hand side of 36.13(d) are stable under $\partial : \underline{\mathfrak{K}}(Z_{J,G^0}) \rightarrow \underline{\mathfrak{K}}(Z_{J,G^0})$ and from 36.4(h) we have

(a)
$$(x * y : \partial(z)) = (\partial(x) : y * z)$$

for $x, y, z \in \underline{\mathfrak{K}}(Z_{J,G^0})^J$.

Let $\mathcal{W} := \{w \in {}^J\mathbf{W}; \text{Ad}(w)J = J\}$. We show that

(b)
$$\underline{\mathfrak{K}}(Z_{J,G^0})^J = \bigoplus_{w \in \mathcal{W}} {}^w\underline{\mathfrak{K}}(Z_{J,G^0})^J$$

where ${}^w\underline{\mathfrak{K}}(Z_{J,G^0})^J = {}^w\underline{\mathfrak{K}}(Z_{J,G^0}) \cap \underline{\mathfrak{K}}(Z_{J,G^0})^J$. Let $x \in \underline{\mathfrak{K}}(Z_{J,G^0})^J$. By 36.8(b) we can write uniquely $x = \sum_{w \in {}^J\mathbf{W}} x_w$ where $x_w \in {}^w\underline{\mathfrak{K}}(Z_{J,G^0})$. It is enough to show that $x_w \in \underline{\mathfrak{K}}(Z_{J,G^0})^J$ (that is, $(y : x_w) = 0$ for any $y \in \tilde{\underline{\mathfrak{K}}}(Z_{J,G^0})$) for all w and $x_w = 0$ (that is, $(y' : x_w) = 0$ for any $y' \in \underline{\mathfrak{K}}(Z_{J,G^0})$) if $\text{Ad}(w)J \neq J$. For x_w, y as above we have $y = \sum_{w' \in {}^J\mathbf{W}} y_{w'}$ with $y_{w'} \in {}^{w'}\underline{\mathfrak{K}}(Z_{J,G^0}) \cap \tilde{\underline{\mathfrak{K}}}(Z_{J,G^0})$; see 36.10(a). Using 36.9(j) twice, we have $(y : x_w) = (y_w : x_w) = (y_w : x) = 0$, as required. Now assume that $w \in {}^J\mathbf{W}$, $\text{Ad}(w)J \neq J$ and $y' \in \underline{\mathfrak{K}}(Z_{J,G^0})$. By 36.8(b) we have $y' = \sum_{w' \in {}^J\mathbf{W}} y'_{w'}$ with $y'_{w'} \in {}^{w'}\underline{\mathfrak{K}}(Z_{J,G^0})$; moreover, by 36.10(a) we have $y'_{w'} \in \tilde{\underline{\mathfrak{K}}}(Z_{J,G^0})$. Using 36.9(j) twice, we have $(y' : x_w) = (y'_{w'} : x_w) = (y'_{w'} : x) = 0$, as required. This proves (b).

From definitions we see that

(c) the decomposition (b) corresponds under 36.13(e) to the decomposition $\bar{\mathfrak{K}}(Z_{J,G^0}) = \bigoplus_{w \in \mathcal{W}} {}^w \bar{\mathfrak{K}}(Z_{J,G^0})$ (see 36.10(d)).

Under the isomorphism 36.13(e), the involution $\tilde{\mathfrak{D}} : \bar{\mathfrak{K}}(Z_{J,G^0}) \rightarrow \bar{\mathfrak{K}}(Z_{J,G^0})$ corresponds to an involution $\tilde{\mathfrak{D}}' : \mathfrak{K}(Z_{J,G^0})^J \rightarrow \mathfrak{K}(Z_{J,G^0})^J$; this is related to the involution $\tilde{\mathfrak{D}} : \mathfrak{K}(Z_{J,G^0}) \rightarrow \mathfrak{K}(Z_{J,G^0})$ by $\tilde{\mathfrak{D}}'(x) = \tilde{\mathfrak{D}}(x) \bmod \tilde{\mathfrak{K}}(Z_{J,G^0})$ for $x \in \mathfrak{K}(Z_{J,G^0})^J$. Hence for $x, x' \in \mathfrak{K}(Z_{J,G^0})^J$ we have $(\tilde{\mathfrak{D}}'(x) : x') = (\tilde{\mathfrak{D}}(x) : x')$. Using this and 36.9(h) we see that

(d) if $x \in \mathfrak{K}(Z_{J,G^0})^J$ satisfies $(x : \tilde{\mathfrak{D}}(x)) = 0$, then $x = 0$.

Note that $\mathfrak{K}^\heartsuit(Z_{J,G^0})$ is a subalgebra of $\mathfrak{K}(Z_{J,G^0})$. Hence

$$\mathfrak{K}^J := \mathfrak{K}^\heartsuit(Z_{J,G^0}) \cap \mathfrak{K}(Z_{J,G^0})^J$$

is a subalgebra of $\mathfrak{K}(Z_{J,G^0})^J$. From (b) and 36.8(d) we see that

(e)
$$\mathfrak{K}^J = \bigoplus_{w \in \mathcal{W}} {}^w \mathfrak{K}^J$$

where ${}^w \mathfrak{K}^J = {}^w \mathfrak{K}^\heartsuit(Z_{J,G^0}) \cap \mathfrak{K}^J$. Now $\partial : \mathfrak{K}(Z_{J,G^0})^J \rightarrow \mathfrak{K}(Z_{J,G^0})^J$ leaves \mathfrak{K}^J stable. Moreover, for $w \in \mathcal{W}$ we have

(f)
$$\partial({}^w \mathfrak{K}^J) = {}^{w^{-1}} \mathfrak{K}^J.$$

(see 36.9(l)).

Note that \mathcal{W} is the same as the set of all $w \in \mathbf{W}$ such that the corresponding permutation of the set of roots leaves stable the set of simple roots corresponding to elements of J . Hence \mathcal{W} is a subgroup of \mathbf{W} .

36.15. By 36.4(f) we have $(x : x') = (x' : x) = \tau(x * \partial(x'))$ for $x, x' \in \mathfrak{K}^J$. By 36.13(b), the bilinear form $(:)$ is nondegenerate on (the finite dimensional vector space) \mathfrak{K}^J . Hence there is a unique vector $x_0 \in \mathfrak{K}^J$ such that $(x_0 : x) = \tau(x)$ for all $x \in \mathfrak{K}^J$. Hence for $x, x' \in \mathfrak{K}^J$ we have $(x_0 : x * \partial(x')) = \tau(x * \partial(x')) = (x : x')$. Using 36.14(a) we rewrite this as $(\partial(x_0) * x : x') = (x : x')$. (We use also that $\partial^2 = 1$ on \mathfrak{K}^J .) Using the nondegeneracy of $(:)$ on \mathfrak{K}^J we deduce $\partial(x_0) * x = x$ for all $x \in \mathfrak{K}^J$. For $x, x' \in \mathfrak{K}^J$ we have also $(\partial(x) * x' : x_0) = \tau(\partial(x) * x') = (\partial(x) : \partial(x')) = (x : x')$ (we use 36.4(g)). Using 36.14(a) we rewrite this as $(x : x' * \partial(x_0)) = (x : x')$. Using the nondegeneracy of $(:)$ on \mathfrak{K}^J we deduce $x' * \partial(x_0) = x'$ for all $x' \in \mathfrak{K}^J$. We see that the algebra \mathfrak{K}^J has a unit element, namely $\mathbf{1} = \partial(x_0)$. By 36.14(e) we have $x_0 = \sum_{w \in \mathcal{W}} x_0^w$ where $x_0^w \in {}^w \mathfrak{K}^J$. For any $x \in \mathfrak{K}^J$ we have similarly $x = \sum_{w \in \mathcal{W}} x^w$ where $x^w \in {}^w \mathfrak{K}^J$. From 36.9(k) we see that $\tau(x) = \tau(x^1)$. Hence $(x_0 : x^1) = \tau(x^1) = \tau(x) = (x_0 : x)$. Using 36.9(j) we see that $(x_0 : x^1) = (x_0^1 : x^1) = (x_0^1 : x)$ hence $(x_0^1 : x) = (x_0 : x)$. Using the nondegeneracy of $(:)$ on \mathfrak{K}^J we deduce $x_0 = x_0^1$, that is, $x_0 \in {}^1 \mathfrak{K}^J$. Using 36.14(f) we deduce that $\mathbf{1} \in {}^1 \mathfrak{K}^J$.

36.16. We preserve the setup of 36.14, 36.15. In this subsection we assume that $G = G^0$ is a symplectic group $Sp_{2n}(\mathbf{k})$ ($n \geq 1$) and we describe in this case the structure of the algebra \mathfrak{K}^J . (The proofs, which depend on results in this and future sections, will be given elsewhere.) We have $\mathbf{I} = \{s_1, s_2, \dots, s_n\}$ where $s_i s_{i+1}$ has order 3 if $i = 1, 2, \dots, n-2$ and order 4 if $i = n-1$; we have $s_i s_j = s_j s_i$ if $|i-j| \geq 2$.

(i) We have $\mathfrak{K}^J = 0$ unless

(*) $J = \{s_{k+1}, s_{k+2}, \dots, s_n\}$ with $0 \leq k \leq n$ such that $n - k = a^2 + a$ for some $a \in \mathbf{N}$.

Now assume that (*) holds. Then \mathcal{W} is a Weyl group of type B_k with standard generators $\sigma_1, \sigma_2, \dots, \sigma_k$ where $\sigma_i = s_i$ for $1 \leq i < k$ and σ_k is the unique element in the subgroup of \mathbf{W} generated by s_k, s_{k+1}, \dots, s_n such that $\sigma_k \in \mathcal{W} - \{1\}$ (if $k \geq 1$). Let $\tilde{l} : \mathcal{W} \rightarrow \mathbf{N}$ be the length function of the Weyl group \mathcal{W} .

(ii) For any $w \in \mathcal{W}$ we have $\dim({}^w \underline{\mathfrak{K}}^J) = 1$.

(iii) For any $i \in [1, k]$ there is a unique element $x \in \sigma_i \underline{\mathfrak{K}}^J - \{0\}$ such that $(x + \mathbf{1}) * (x - c\mathbf{1}) = 0$ for some $c \in v\mathbf{Z}[v]$; in fact, we have $c = v^2$ if $1 \leq i < k$ and $c = v^{4a+2}$ if $i = k$. We set $x = t_i$.

(iv) For any $w \in \mathcal{W}$ there is a unique element $t_w \in {}^w \underline{\mathfrak{K}}^J - \{0\}$ such that the following hold: $t_{\sigma_i} = t_i$ for $i \in [1, k]$; $t_w * t_{w'} = t_{w'w}$ if $w, w' \in \mathcal{W}$, $\tilde{l}(w'w) = \tilde{l}(w) + \tilde{l}(w')$.

We see that $\underline{\mathfrak{K}}^J$ is an Iwahori-Hecke algebra with not necessarily equal parameters. Similar results hold for other classical groups.

37. A MACKEY TYPE FORMULA

37.1. We fix a connected component D of G . With notation in 26.1, if $J, J' \subset \mathbf{I}, P \in \mathcal{P}_J, Q \in \mathcal{P}_{J'}, u = \text{pos}(P, Q) \in \mathbf{W}$, then $u \in {}^J \mathbf{W}^{J'}$. Setting $P^Q = (P \cap Q)U_P$, we have $P^Q \in \mathcal{P}_{J \cap \text{Ad}(u)J'}$.

For $K, K' \subset \mathbf{I}$ and $u \in {}^K \mathbf{W}^{K'}$ let

$$\Upsilon_u = \{(X, Y, g(U_X \cap U_Y)); X \in \mathcal{P}_{K \cap \text{Ad}(u)K'}, Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}, \\ g(U_X \cap U_Y) \in D / (U_X \cap U_Y), \text{pos}(X, Y) = u\}.$$

We have a diagram

$$Z_{K \cap \text{Ad}(u)K', D} \xleftarrow{j} \Upsilon_u \xrightarrow{h} Z_{K' \cap \text{Ad}(u^{-1})K, D}$$

where $j(X, Y, g(U_X \cap U_Y)) = (X, gU_X), h(X, Y, g(U_X \cap U_Y)) = (Y, gU_Y)$. Set

$$\Phi_u = h!j^* : \mathcal{D}(Z_{K \cap \text{Ad}(u)K', D}) \rightarrow \mathcal{D}(Z_{K' \cap \text{Ad}(u^{-1})K, D}).$$

Proposition 37.2. Let $K, K', J \subset \mathbf{I}$ be such that $K \subset J, K' \subset J$. Let $A' \in \mathcal{D}(Z_{K, D})$. We set $\mathfrak{B} = \mathfrak{e}_{K', J} \mathfrak{f}_{K, J} A' \in \mathcal{D}(Z_{K', D})$. For any $u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J$ we set $\mathfrak{B}_u = \mathfrak{f}_{K' \cap \text{Ad}(u^{-1})K, K'} \Phi_u \mathfrak{e}_{K \cap \text{Ad}(u)K', K} A' \in \mathcal{D}(Z_{K', D})$ and $m_u = \dim(U_P \cap U_R) / U_Q$ where $P \in \mathcal{P}_K, R \in \mathcal{P}_{K'}, \text{pos}(P, R) = u$ and $Q = Q_{J, P} = Q_{J, R} \in \mathcal{P}_J$ (notation of 36.4). We have

$$\mathfrak{B} \simeq \{\mathfrak{B}_u[[-m_u]]; u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J\},$$

with \simeq as in 32.15.

We have a commutative diagram with a cartesian square

$$\begin{array}{ccccc} \mathfrak{E} & \xrightarrow{b} & Z_{K', J, D} & \xrightarrow{c'} & Z_{K', D} \\ a \downarrow & & & & \mathfrak{d}' \downarrow \\ Z_{K, D} & \xleftarrow{c} & Z_{K, J, D} & \xrightarrow{d} & Z_{J, D} \end{array}$$

Here

$$\mathfrak{E} = \{(P, R, gU_Q); P \in \mathcal{P}_K, R \in \mathcal{P}_{K'}, gU_Q \in D / U_Q, Q = Q_{J, P} = Q_{J, R}\}, \\ \mathfrak{c}(P, gU_Q) = (P, gU_P), \mathfrak{d}(P, gU_Q) = (Q, gU_Q) \text{ with } Q = Q_{J, P},$$

$$\begin{aligned} \mathfrak{c}'(R, gU_Q) &= (R, gU_R), \mathfrak{d}'(R, gU_Q) = (Q, gU_Q) \text{ with } Q = Q_{J,R}, \\ \mathfrak{a}(P, R, gU_Q) &= (P, gU_Q), \mathfrak{b}(P, R, gU_Q) = (R, gU_Q). \end{aligned}$$

We have

$$\mathfrak{B} = \mathfrak{c}'_! \mathfrak{d}'^* \mathfrak{d}_! \mathfrak{c}^* A' = \mathfrak{c}'_! \mathfrak{b}_! \mathfrak{a}^* \mathfrak{c}^* A' = (\mathfrak{c}'\mathfrak{b})_! (\mathfrak{c}\mathfrak{a})^* A' = \mathfrak{q}_! \mathfrak{p}^* A'$$

where $\mathfrak{q} = \mathfrak{c}'\mathfrak{b} : \mathfrak{E} \rightarrow Z_{K',D}, \mathfrak{p} = \mathfrak{c}\mathfrak{a} : \mathfrak{E} \rightarrow Z_{K,D}$ are given by $\mathfrak{q}(P, R, gU_Q) = (R, gU_R), \mathfrak{p}(P, R, gU_Q) = (P, gU_P)$. We have a partition

$$\mathfrak{E} = \bigsqcup_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J} \mathfrak{E}_u$$

where $\mathfrak{E}_u = \{(P, R, gU_Q) \in \mathfrak{E}; \text{pos}(P, R) = u\}$ is locally closed in \mathfrak{E} . Let $\mathfrak{p}_u = \mathfrak{p}|_{\mathfrak{E}_u} : \mathfrak{E}_u \rightarrow Z_{K,D}, \mathfrak{q}_u = \mathfrak{q}|_{\mathfrak{E}_u} : \mathfrak{E}_u \rightarrow Z_{K'}$. By 32.15, we have

$$\mathfrak{q}_! \mathfrak{p}^* A' \simeq \{\mathfrak{q}_u! \mathfrak{p}_u^* A'; u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J\}.$$

It remains to show that, for u as above, we have

$$\mathfrak{q}_u! \mathfrak{p}_u^* A' = \mathfrak{B}_u[[-m_u]].$$

We have a commutative diagram

$$\begin{array}{ccccc} Z_{K,D} & \xleftarrow{\tilde{\mathfrak{d}}} & Z_{K \cap \text{Ad}(u)K', K, D} & \xrightarrow{\tilde{\mathfrak{e}}} & Z_{K \cap \text{Ad}(u)K', D} \\ & & \tilde{\mathfrak{a}} \uparrow & & \uparrow \tilde{\mathfrak{j}} \\ & & \mathfrak{S}'_u & \xrightarrow{\tilde{\mathfrak{b}}} & \Upsilon_u & \xrightarrow{\mathfrak{h}} & Z_{K' \cap \text{Ad}(u^{-1})K, D} \\ & & \mathfrak{r} \uparrow & & & & \uparrow \tilde{\mathfrak{c}}' \\ & & \mathfrak{S}_u & \xrightarrow{\mathfrak{q}} & & \longrightarrow & Z_{K' \cap \text{Ad}(u^{-1})K, K', D} \\ & & & & & & \downarrow \tilde{\mathfrak{d}}' \\ & & & & & & Z_{K', D} \end{array}$$

Here

$$\begin{aligned} \tilde{\mathfrak{c}}(X, gU_P) &= (X, gU_X), \tilde{\mathfrak{d}}(X, gU_P) = (P, gU_P), \\ \tilde{\mathfrak{c}}'(Y, gU_R) &= (Y, gU_Y), \tilde{\mathfrak{d}}'(Y, gU_R) = (R, gU_R), \\ \mathfrak{S}'_u, \tilde{\mathfrak{a}}, \tilde{\mathfrak{b}} &\text{ are defined so that the square } (\tilde{\mathfrak{a}}, \tilde{\mathfrak{b}}, \tilde{\mathfrak{c}}, \tilde{\mathfrak{j}}) \text{ is cartesian;} \\ \mathfrak{S}_u, \mathfrak{r}, \mathfrak{q} &\text{ are defined so that the square } (\mathfrak{r}, \mathfrak{q}, \mathfrak{h}\tilde{\mathfrak{b}}, \tilde{\mathfrak{c}}') \text{ is cartesian.} \end{aligned}$$

Then

$$\begin{aligned} \mathfrak{S}_u &= \{(X, Y, g(U_X \cap U_Y), g'U_P, g''U_R); X \in \mathcal{P}_{K \cap \text{Ad}(u)K'}, Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}, \\ &g(U_X \cap U_Y) \in D/(U_X \cap U_Y), g'U_P \in D/U_P, g''U_R \in D/U_R, \text{pos}(X, Y) = u, \\ &P = Q_{K,X}, R = Q_{K',Y}, g'U_X = gU_X, g''U_Y = gU_Y\}. \end{aligned}$$

Set $\mathfrak{r} = \tilde{\mathfrak{d}}'\mathfrak{q} : \mathfrak{S}_u \rightarrow Z_{K',D}, \mathfrak{s} = \tilde{\mathfrak{d}}\mathfrak{r} : \mathfrak{S}_u \rightarrow Z_{K,D}$. Then

$$\begin{aligned} \mathfrak{r}(X, Y, g(U_X \cap U_Y), g'U_P, g''U_R) &= (P, g'U_P), \\ \mathfrak{s}(X, Y, g(U_X \cap U_Y), g'U_P, g''U_R) &= (R, g''U_R). \end{aligned}$$

We have

$$\begin{aligned} \mathfrak{B}_u &= \tilde{\mathfrak{d}}'_! \tilde{\mathfrak{c}}'^* \mathfrak{h}_! \tilde{\mathfrak{j}}^* \tilde{\mathfrak{c}}_! \tilde{\mathfrak{d}}^* A' = \tilde{\mathfrak{d}}'_! \tilde{\mathfrak{c}}'^* \mathfrak{h}_! \tilde{\mathfrak{b}}_! \tilde{\mathfrak{a}}^* \tilde{\mathfrak{d}}^* A' \\ &= \tilde{\mathfrak{d}}'_! \tilde{\mathfrak{c}}'^* (\mathfrak{h}\tilde{\mathfrak{b}})_! (\tilde{\mathfrak{d}}\tilde{\mathfrak{a}})^* A' = \tilde{\mathfrak{d}}'_! \mathfrak{r}_! \mathfrak{s}^* (\tilde{\mathfrak{d}}\tilde{\mathfrak{a}})^* A' = (\tilde{\mathfrak{d}}'\mathfrak{r})_! (\tilde{\mathfrak{d}}\tilde{\mathfrak{a}})^* A' = \mathfrak{r}_! \mathfrak{s}^* A'. \end{aligned}$$

We show that $\mathfrak{q}_u! \mathfrak{p}_u^* A' = \mathfrak{r}_! \mathfrak{s}^* A' [[-m_u]]$. We have a commutative diagram

$$\begin{CD} Z_{K,D} @<\mathfrak{p}_u<< \mathfrak{E}_u @>\mathfrak{q}_u>> Z_{K',D} \\ @V1VV @VV\mathfrak{t}V @VV1V \\ Z_{K,D} @<\mathfrak{s}<< \mathfrak{S}_u @>\mathfrak{r}>> Z_{K',D} \end{CD}$$

where $\mathfrak{t}(P, R, gU_Q) = (P^R, R^P, g(U_{P^R} \cap U_{R^P}), gU_P, gU_R)$ is well defined since $U_Q \subset U_P, U_Q \subset U_R, U_Q \subset U_{P^R}, U_Q \subset U_{R^P}$. We continue the proof assuming that

(a) \mathfrak{t} is an affine space bundle with fibres of dimension m_u .

For any $\tilde{A} \in \mathcal{D}(\mathfrak{S}_u)$ we have $\mathfrak{t}_! \mathfrak{t}^*(\tilde{A}) = \tilde{A} [[-m_u]]$. Hence

$$\mathfrak{r}_! \mathfrak{s}^* A' [[-m_u]] = \mathfrak{r}_! \mathfrak{t}_! \mathfrak{t}^* \mathfrak{s}^* A' = (\mathfrak{r}\mathfrak{t})_! (\mathfrak{s}\mathfrak{t})^* A' = \mathfrak{q}_u! \mathfrak{p}_u^* A'$$

as required.

We prove (a). We only show that each fibre of \mathfrak{t} is an affine space of dimension m_u . First we show that $\mathfrak{t} : \mathfrak{E}_u \rightarrow \mathfrak{S}_u$ is surjective. Assume that we are given $(X, Y, g_0(U_X \cap U_Y), g'_0 U_P, g''_0 U_R) \in \mathfrak{S}_u$. Then $g_0, g'_0, g''_0 \in D, v' = g_0^{-1} g'_0 \in U_X, v'' = g_0^{-1} g''_0 \in U_Y$. We must show that there exists $g \in D$ such that $g_0 \in g(U_X \cap U_Y), g'_0 \in gU_P, g''_0 \in gU_R$. Setting $y = g^{-1} g_0$, we must show that there exists $y \in U_X \cap U_Y$ such that $yv' \in U_P, yv'' \in U_R$. We have $v' = v'_1 v'_2$ where $v'_1 \in U_R \cap P, v'_2 \in U_P$ and $v'' = v''_1 v''_2$ where $v''_1 \in U_P \cap R, v''_2 \in U_R$. Then $v'_1 \in U_X \cap U_Y, v''_1 \in U_X \cap U_Y$. Setting $y = (v'_1 v''_1)^{-1} \in U_X \cap U_Y$ we have $yv' = v''_1^{-1} v'_2 \in U_P$ and $yv'' = v''_1^{-1} v'_1^{-1} v''_2 \in v''_1^{-1} U_R v''_2 = U_R$, as desired.

It remains to show that, if $(P, R, gU_Q) \in \mathfrak{E}_u$, then

(b) $F = \{(P', R', g'U_{Q'}) \in E_u; \mathfrak{t}(P', R', g'U_{Q'}) = \mathfrak{t}(P, R, gU_Q)\}$ is an affine space of dimension m_u .

For $(P', R', g'U_{Q'}) \in F$, both P, P' contain $P^R = (P')^{R'}$ and have the same type, hence $P = P'$. Similarly, $R = R', Q = Q'$. Hence

$$\begin{aligned} F &\cong \{g'U_Q; gU_P = g'U_P, gU_R = g'U_R, g(U_{P^R} \cap U_{R^P}) = g'(U_{P^R} \cap U_{R^P})\} \\ &= \{g'U_Q; g^{-1}g' \in U_P \cap U_R \cap U_{P^R} \cap U_{R^P}\} \\ &= \{g'U_Q; g^{-1}g' \in U_P \cap U_R\} \cong (U_P \cap U_R)/U_Q, \end{aligned}$$

and (b) follows. This completes the proof.

37.3. For any $n \in \mathbf{N}_k^*$ we have a $G^0 \times \mathbf{T}$ -action on $Z_{\emptyset, D}$ given by

$$(a) (g_1, t) : (B, gU_B) \mapsto (g_1 B g_1^{-1}, g_1 g b^{-n} g_1^{-1} U_{g_1 B g_1^{-1}})$$

where $b \in B$ is such that $f_B(t) = bU_B$ (f_B as in 28.3). Let $u \in \mathbf{W}$. We consider another $G^0 \times \mathbf{T}$ -action on $Z_{\emptyset, D}$ given by

$$(b) (g_1, t) : (B, gU_B) \mapsto (g_1 B g_1^{-1}, g_1 g b_1^{-n} g_1^{-1} U_{g_1 B g_1^{-1}})$$

where $b_1 \in B$ is such that $f_B(u^{-1}(t)) = b_1 U_B$ (f_B as in 28.3).

Using 36.3 we see that a simple perverse sheaf on $Z_{\emptyset, D}$ is a character sheaf if and only if it is $G^0 \times \mathbf{T}$ -equivariant for the action (a) for some n or equivalently if it is $G^0 \times \mathbf{T}$ -equivariant for the action (b) for some n .

Let $\Upsilon_{u,0}$ be like Υ_u in 37.1 but with K, K' replaced by \emptyset, \emptyset . We have a diagram $Z_{\emptyset, D} \xrightarrow{j_0} \Upsilon_{u,0} \xrightarrow{h_0} Z_{\emptyset, D}$ (a special case of the diagram in 37.1 with $K = K' = \emptyset$). Set $\Phi_u^\emptyset = h_0! j_0^* : \mathcal{D}(Z_{\emptyset, D}) \rightarrow \mathcal{D}(Z_{\emptyset, D})$. We show that

$$(c) \Phi_u^\emptyset \text{ restricts to a functor } \mathcal{D}^{cs}(Z_{\emptyset, D}) \rightarrow \mathcal{D}^{cs}(Z_{\emptyset, D}).$$

It is enough to show that if A is a character sheaf on $Z_{\emptyset,D}$, then $\Phi_u^\emptyset(A) \in \mathcal{D}^{cs}(Z_{\emptyset,D})$. Now $G^0 \times \mathbf{T}$ acts on $\Upsilon_{u,0}$ by

$$(g_1, t) : (B, B', g(U_B \cap U_{B'})) \mapsto (g_1 B g_1^{-1}, g_1 B' g_1^{-1}, g_1 g b^{-n} g_1^{-1} (U_{g_1 B g_1^{-1}} \cap U_{g_1 B' g_1^{-1}}))$$

where $b \in B \cap B'$ is such that $f_B(t) = bU_B$ (f_B as in 28.3). Note that j_0 (resp. \mathfrak{h}_0) is $G^0 \times \mathbf{T}$ -equivariant where $G^0 \times \mathbf{T}$ acts on $Z_{\emptyset,D}$ by (a) (resp. by (b)). Since A is equivariant for the action (a) for some n , it follows that ${}^p H^j(\Phi_u^\emptyset(A))$ is equivariant for the action (b). Hence any simple subquotient of ${}^p H^j(\Phi_u^\emptyset(A))$ is a character sheaf. This proves (c).

Let $K, K' \subset \mathbf{I}$ and let $u \in {}^K \mathbf{W}^{K'}$. Let $H = K \cap \text{Ad}(u)K'$, $H' = K' \cap \text{Ad}(u^{-1})K = \text{Ad}(u^{-1})H$. Let $\Phi_u : \mathcal{D}(Z_{H,D}) \rightarrow \mathcal{D}(Z_{H',D})$ be as in 37.1. We show that

(d) $\Phi_u \mathfrak{f}_{\emptyset,H}(A) = \mathfrak{f}_{\emptyset,H'} \Phi_u^\emptyset(A)$ for any $A \in \mathcal{D}(Z_{\emptyset,D})$.

We have a diagram

$$\begin{array}{ccccc} Z_{\emptyset,D} & \xleftarrow{a} & \mathfrak{Z} & \xrightarrow{b} & Z_{H',D} \\ \downarrow = & & \downarrow c & & \downarrow = \\ Z_{\emptyset,D} & \xleftarrow{a'} & \mathfrak{Z}' & \xrightarrow{b'} & Z_{H',D} \end{array}$$

where

$$\mathfrak{Z} = \{(B, P, P', g(U_P \cap U_{P'}); B \in \mathcal{B}, P \in \mathcal{P}_H, P' \in \mathcal{P}_{H'}, g \in D, B \subset P, \text{pos}(P, P') = u)\},$$

$$\mathfrak{Z}' = \{(B, B', P', g(U_B \cap U_{B'}), g'U_{P'}); B, B' \in \mathcal{B}, P' \in \mathcal{P}_{H'}, g \in D; g' \in D, gU_{B'} = g'U_{B'}, B' \subset P', \text{pos}(B, B') = u\},$$

$$a \text{ is } (B, P, P', g(U_P \cap U_{P'})) \mapsto (B, gU_B),$$

$$b \text{ is } (B, P, P', g(U_P \cap U_{P'})) \mapsto (P', gU_{P'}),$$

$$a' \text{ is } (B, B', P', g(U_B \cap U_{B'}), g'U_{P'}) \mapsto (B, gU_B),$$

$$b' \text{ is } (B, B', P', g(U_B \cap U_{B'}), g'U_{P'}) \mapsto (P', g'U_{P'}),$$

$$c \text{ is } (B, P, P', g(U_P \cap U_{P'})) \mapsto ((B, B', P', g(U_B \cap U_{B'}), g'U_{P'})) \text{ with } B' = (B \cap P')U_{P'}.$$

From the definition and the change of base theorem we see that $\Phi_u \mathfrak{f}_{\emptyset,H}(A) = b_! a^* A$, $\mathfrak{f}_{\emptyset,H'} \Phi_u^\emptyset(A) = b'_! a'^* A$. Since c is an isomorphism, the diagram above shows that $b_! a^* A = b'_! a'^* A$. This proves (d).

We show that

(e) Φ_u restricts to a functor $\mathcal{D}^{cs}(Z_{H,D}) \rightarrow \mathcal{D}^{cs}(Z_{H',D})$.

It is enough to show that if A is a character sheaf on $Z_{H,D}$, then $\Phi_u(A) \in \mathcal{D}^{cs}(Z_{H',D})$. We can find a character sheaf A' on $Z_{\emptyset,D}$ and $m \in \mathbf{Z}$ such that $A[m]$ is a direct summand of $\mathfrak{f}_{\emptyset,H} A'$. Hence $\Phi_u(A)[n]$ is a direct summand of $\Phi_u \mathfrak{f}_{\emptyset,H} A'$. By (d), $\Phi_u(A)[n]$ is a direct summand of $\mathfrak{f}_{\emptyset,H'} \Phi_u^\emptyset(A')$. It is enough to show that $\mathfrak{f}_{\emptyset,H'} \Phi_u^\emptyset(A') \in \mathcal{D}^{cs}(Z_{H',D})$. By (c), we have $\Phi_u^\emptyset(A') \in \mathcal{D}^{cs}(Z_{\emptyset,D})$. Since $\mathfrak{f}_{\emptyset,H'}$ carries $\mathcal{D}^{cs}(Z_{\emptyset,D})$ into $\mathcal{D}^{cs}(Z_{\emptyset,H'})$, we see that (e) holds.

In the remainder of this section we assume that we are in the setup of 36.8. From 37.2 we deduce that for $K, K', J \subset \mathbf{I}$ such that $K \subset J, K' \subset J$ we have

$$(f) \quad \mathfrak{e}_{K',J} \mathfrak{f}_{K,J} = \sum_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J} v^{2m_u} \mathfrak{f}_{K' \cap \text{Ad}(u^{-1})K, K'} \Phi_u \mathfrak{e}_{K \cap \text{Ad}(u)K', K}$$

as \mathcal{A} -linear maps $\mathfrak{K}(Z_{K,D}) \rightarrow \mathfrak{K}(Z_{K',D})$, where the \mathcal{A} -linear map

$$gr(\Phi_u) : \mathfrak{K}(Z_{K \cap \text{Ad}(u)K',D}) \rightarrow \mathfrak{K}(Z_{K' \cap \text{Ad}(u^{-1})K,D})$$

(see (e) and 36.8) is denoted again by Φ_u .

37.4. Let $K, K' \subset J$ and let $x \in \mathfrak{K}(Z_{K,D})^K, x' \in \mathfrak{K}(Z_{K',D})^{K'}$. We show that

$$(a) \quad (f_{K,J}x : f_{K',J}x') = \sum_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J; K = \text{Ad}(u)K'} v^{2m_u} (\Phi_u x : x').$$

Here m_u, Φ_u are as in 37.3; in our case, $\Phi_u : \mathfrak{K}(Z_{K,D}) \rightarrow \mathfrak{K}(Z_{K',D})$. Using 37.3 we have

$$\begin{aligned} (f_{K,J}x : f_{K',J}x') &= (\mathbf{e}_{K',J} f_{K,J}x : x') \\ &= \left(\sum_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J} v^{2m_u} f_{K' \cap \text{Ad}(u^{-1})K, K'} \Phi_u \mathbf{e}_{K \cap \text{Ad}(u)K', K} x : x' \right) \\ &= \sum_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J} v^{2m_u} (\Phi_u \mathbf{e}_{K \cap \text{Ad}(u)K', K} x : \mathbf{e}_{K' \cap \text{Ad}(u^{-1})K, K'} x') \\ &= \sum_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J; K \cap \text{Ad}(u)K' = K, K' \cap \text{Ad}(u^{-1})K = K'} v^{2m_u} (\Phi_u x : x'). \end{aligned}$$

The condition that $K \cap \text{Ad}(u)K' = K, K' \cap \text{Ad}(u^{-1})K = K'$ is equivalent to $K \subset \text{Ad}(u)K', K' \subset \text{Ad}(u^{-1})K$, that is, to $K = \text{Ad}(u)K'$. This proves (a).

Consider the equivalence relation on the set of subsets of J given by $K_1 \sim K_2$ if $\text{Ad}(u)K_1 = K_2$ for some $u \in \mathbf{W}_J$. For any equivalence class \mathfrak{o} under \sim we set

$$\underline{\mathfrak{K}}(Z_{J,D})^\mathfrak{o} = \sum_{H \subset J; H \in \mathfrak{o}} f_{H,J}(\underline{\mathfrak{K}}(Z_{H,D})^H) \subset \underline{\mathfrak{K}}(Z_{J,D}).$$

Note that $\underline{\mathfrak{K}}(Z_{J,D})^\mathfrak{o}$ is homogeneous.

Proposition 37.5. *We have*

$$(a) \quad \underline{\mathfrak{K}}(Z_{J,D}) = \bigoplus_{\mathfrak{o}} \underline{\mathfrak{K}}(Z_{J,D})^\mathfrak{o}$$

where \mathfrak{o} runs over the equivalence classes for \sim .

If $J = \emptyset$ we have $\underline{\mathfrak{K}}(Z_{J,D}) = \underline{\mathfrak{K}}(Z_{J,D})^J$ and the result is obvious. We may assume that $J \neq \emptyset$ and that the result is true when J is replaced by a strictly smaller subset. Using 36.13(d) and the induction hypothesis we have

$$\underline{\mathfrak{K}}(Z_{J,D}) = \underline{\mathfrak{K}}(Z_{J,D})^J + \sum_{L \subsetneq J} f_{L,J} \sum_{J'; J' \subset L} f_{J',L} \underline{\mathfrak{K}}(Z_{J',D})^{J'} \subset \sum_{J'; J' \subset J} f_{J',J} \underline{\mathfrak{K}}(Z_{J',D})^{J'}.$$

Thus, $\underline{\mathfrak{K}}(Z_{J,D}) = \sum_{\mathfrak{o}} \underline{\mathfrak{K}}(Z_{J,D})^\mathfrak{o}$. Next we show that $(\underline{\mathfrak{K}}(Z_{J,D})^\mathfrak{o} : \underline{\mathfrak{K}}(Z_{J,D})^{\mathfrak{o}'}) = 0$ if $\mathfrak{o} \neq \mathfrak{o}'$. It is enough to show that $(f_{H,J}(\underline{\mathfrak{K}}(Z_{H,D})^H) : f_{H',J}(\underline{\mathfrak{K}}(Z_{H',D})^{H'})) = 0$ if $H, H' \subset J, H \not\sim H'$. This follows from 37.4(a). It remains to use the following (easily verified) statement: If V is a finite dimensional vector space with a nonsingular symmetric bilinear form $(:)$ and V_1, V_2, \dots, V_k are subspaces such that $(V_i : V_j) = 0$ for $i \neq j$ and $\sum_i V_i = V$, then $V = \bigoplus_i V_i$.

38. DUALITY

38.1. We fix a connected component D of G that generates G . In this section we study an involution of the set of isomorphism classes of character sheaves on D called *duality*.

We write ϵ instead of $\epsilon_D : \mathbf{W} \rightarrow \mathbf{W}$ (see 26.2). For $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ we set, as in 30.3,

$$V_{J,D} = \{(P, gU_P); P \in \mathcal{P}_J, gU_P \in N_D P/U_P\}.$$

This is the same as ${}^1Z_{J,D}$; see 36.2. We also set $\mathbf{W}_J^\epsilon = \{w \in \mathbf{W}_J; \epsilon(w) = w\}$ where \mathbf{W}_J is as in 26.1. Let J_ϵ be the set of orbits of the restriction of ϵ to J .

For any $P_0 \in \mathcal{P}_J$ we have a functor $A_0 \mapsto A_0^b$ (see 30.3) from the category of P_0/U_{P_0} -equivariant perverse sheaves on the connected component $N_D P_0/U_{P_0}$ of $N_G P_0/U_{P_0}$ to the category of perverse sheaves on $V_{J,D}$.

Let $CS(V_{J,D})$ be the full subcategory of the category of perverse sheaves on $V_{J,D}$ whose objects are isomorphic to objects of the form A_0^b where A_0 is a direct sum of character sheaves on $N_D P_0/U_{P_0}$. Equivalently, $CS(V_{J,D})$ is the category of perverse sheaves on $V_{J,D}$ that are direct sums of perverse sheaves in ${}^1\hat{Z}_{J,D}$; see 36.2. We write also $CS(D)$ instead of $CS(V_{\mathbf{I},D})$. Note that $A_0 \mapsto A_0^b$ is an equivalence of categories $CS(N_D P_0/U_{P_0}) \xrightarrow{\sim} CS(V_{J,D})$.

For $J \subset J' \subset \mathbf{I}$ such that $\epsilon(J) = J, \epsilon(J') = J'$ we have functors $f_{J,J'} : \mathcal{D}(V_{J,D}) \rightarrow \mathcal{D}(V_{J',D})$ and $e_{J,J'} : \mathcal{D}(V_{J',D}) \rightarrow \mathcal{D}(V_{J,D})$; see 30.4. From definitions, for $J \subset J' \subset J'' \subset \mathbf{I}$ such that $\epsilon(J) = J, \epsilon(J') = J', \epsilon(J'') = J''$, we have

$$(a) \quad f_{J,J''} = f_{J',J''} f_{J,J'}, \quad e_{J,J''} = e_{J',J''} e_{J,J'}.$$

Clearly, $f_{J,J} = 1, e_{J,J} = 1$.

38.2. We show that

(a) for $J \subset J'$ as above, $e_{J,J'}$ restricts to a functor $CS(V_{J',D}) \rightarrow CS(V_{J,D})$ denoted again by $e_{J,J'}$.

Let $P \in \mathcal{P}_J, P' \in \mathcal{P}_{J'}$ be such that $P \subset P'$. Let $D_0 = N_D P/U_P, D'_0 = N_D P'/U_{P'}$. Let $C_0 \in CS(D'_0)$ and let $C = C_0^b$ be the corresponding object of $CS(V_{J',D})$. From 31.14 we see that $\text{res}_{D'_0}^{D_0}(C_0) \in CS(D_0)$ and from 30.4(b) we see that $e_{J,J'} C$ is the perverse sheaf $(\text{res}_{D'_0}^{D_0}(C_0))^b$ on $V_{J,D}$ hence $e_{J,J'} C \in CS(V_{J,D})$. This proves (a).

38.3. We show that

(a) for $J \subset J'$ as above, $f_{J,J'}$ restricts to a functor $CS(V_{J,D}) \rightarrow CS(V_{J',D})$ denoted again by $f_{J,J'}$.

Let P, P', D_0, D'_0 be as in 38.2. Let $A_0 \in CS(D_0)$ and let $A = A_0^b$ be the corresponding object of $CS(V_{J,D})$. By 30.4(a), $f_{J,J'} A = A'_0{}^b$ where $A'_0 = \text{ind}_{D_0}^{D'_0} A_0$ is a direct sum of simple admissible perverse sheaves on D'_0 . It remains to show that $A'_0 \in CS(D'_0)$. To do this we may assume that $J' = \mathbf{I}$, hence $P' = G^0, D'_0 = D$. Let $\alpha = \dim U_P$. Let L be a Levi of P . We can identify naturally $N_G P/U_P$ with $H = N_G P \cap N_G L$, a reductive group with $H^0 = L$. Then D_0 becomes $H \cap D$. We identify the canonical torus of L with the canonical torus \mathbf{T} of G^0 , and the Weyl group of L with the subgroup \mathbf{W}_J of \mathbf{W} as in 29.1. Let $\mathcal{L} \in \mathfrak{s}$. Let $\mathbf{s} = (s_1, s_2, \dots, s_r)$ be a sequence in $J \cup \{1\}$ such that $s_1 s_2 \dots s_r \underline{D} \in \mathbf{W}_{\mathcal{L}}^\bullet$ (see 28.3). Let $\bar{Z}_{\emptyset, \mathbf{I}, D}^{\mathbf{s}}, \bar{K}_D^{\mathbf{s}, \mathcal{L}} \in \mathcal{D}(D)$ be as in 28.12, $\bar{\mathcal{L}}$ as in 28.9, and let $\bar{Z}_{\emptyset, J, D_0}^{\mathbf{s}}, \bar{K}_{D_0}^{\mathbf{s}, \mathcal{L}} \in \mathcal{D}(D_0)$,

$\bar{\mathcal{L}}_0$, be the analogous objects defined in terms of H, D_0 instead of G, D . Consider the commutative diagram

$$\begin{array}{ccccc}
 \bar{Z}_{\emptyset, J, D_0}^s & \xleftarrow{pr_2} & V_1 \times_{D_0} \bar{Z}_{\emptyset, J, D_0}^s & \xrightarrow{f_0} & \bar{Z}_{\emptyset, \mathbf{I}, D}^s \\
 f_1 \downarrow & & pr_1 \downarrow & & f_2 \downarrow \\
 D_0 & \xleftarrow{a_1} & V_1 & \xrightarrow{a'} & V_2 \xrightarrow{a''} D
 \end{array}$$

where

$$\begin{aligned}
 V_1 &= \{(g, x) \in D \times G^0; x^{-1}gx \in N_D P\}, \\
 V_2 &= \{(g, xP) \in D \times G^0/P; x^{-1}gx \in N_D P\}, \\
 a_1(g, x) &= g' \text{ where } g_0 \in D_0 \text{ is such that } x^{-1}gx \in g_0 U_P, \\
 a'(g, x) &= (g, xP), a''(g, xP) = g, f_1(\beta_0, \beta_1, \dots, \beta_r, g_0) = g_0, \\
 f_2(B_0, B_1, \dots, B_r, g) &= (g, x_0 P) \text{ where } x_0 \in G^0 \text{ is such that } x_0^{-1}B_0 x_0 \subset P, \\
 f_0((g, x), (\beta_0, \beta_1, \dots, \beta_r, g_0)) &= (x\beta_0 U_P x^{-1}, x\beta_1 U_P x^{-1}, \dots, x\beta_r U_P x^{-1}, g).
 \end{aligned}$$

Both squares in the diagram are cartesian and the maps a', f_0 (resp. a_1, pr_2) are smooth with connected fibres of dimension $\dim G - \alpha$ (resp. $\dim G + \alpha$). From definitions we have $pr_2^* \bar{\mathcal{L}}' = f_0^* \bar{\mathcal{L}}$. Hence

$$a_1^*(\bar{K}_{D_0}^{s, \mathcal{L}}) = a_1^* f_1^* \bar{\mathcal{L}}_0 = pr_{1!} pr_2^* \bar{\mathcal{L}}_0 = pr_{1!} f_0^* \bar{\mathcal{L}} = a'^* f_2^* \bar{\mathcal{L}}.$$

We see that

$$a_1^\star(\bar{K}_{D_0}^{s, \mathcal{L}})[- \dim G - \alpha] = a'^\star \tilde{K}[- \dim G + \alpha]$$

where $\tilde{K} = f_{2!} \bar{\mathcal{L}}$, that is, $a_1^\star(\bar{K}_{D_0}^{s, \mathcal{L}}) = a'^\star \tilde{K}[2\alpha]$. Hence

$$\begin{aligned}
 a_1^\star({}^p H^i(\bar{K}_{D_0}^{s, \mathcal{L}})) &= {}^p H^i(a_1^\star(\bar{K}_{D_0}^{s, \mathcal{L}})) = {}^p H^i(a'^\star \tilde{K}[2\alpha]) = a'^\star({}^p H^i(\tilde{K}[2\alpha])) \\
 &= a'^\star({}^p H^{i+2\alpha} \tilde{K}).
 \end{aligned}$$

From this and definition (27.1) we see that

$$\text{ind}_{D_0}^D({}^p H^i(\bar{K}_{D_0}^{s, \mathcal{L}})) = a_1''({}^p H^{i+2\alpha} \tilde{K}).$$

We have

$$(b) \quad \bigoplus_i \text{ind}_{D_0}^D({}^p H^i(\bar{K}_{D_0}^{s, \mathcal{L}}))[-i] = \bigoplus_i {}^p H^{i+2\alpha}(\bar{K}_D^{s, \mathcal{L}})[-i] \in \mathcal{D}(D).$$

Indeed, the left-hand side is

$$\begin{aligned}
 \bigoplus_i a_1''({}^p H^{i+2\alpha} \tilde{K})[-i] &= a_1''(\tilde{K}[2\alpha]) = a_1'' f_{2!} \bar{\mathcal{L}}[2\alpha] = \bar{K}_D^{s, \mathcal{L}}[2\alpha] \\
 &= \bigoplus_i {}^p H^{i+2\alpha}(\bar{K}_D^{s, \mathcal{L}})[-i] \in \mathcal{D}(D),
 \end{aligned}$$

where we use that \tilde{K} and $\bar{K}_D^{s, \mathcal{L}}$ are semisimple complexes (a consequence of the decomposition theorem [BBD]). Since ${}^p H^i(\bar{K}_{D_0}^{s, \mathcal{L}})$ is a direct sum of character sheaves on D_0 , we see, using 30.6(a), that $\text{ind}_{D_0}^D({}^p H^i(\bar{K}_{D_0}^{s, \mathcal{L}}))$ is a perverse sheaf on D . Taking ${}^p H^i$ for both sides of (b) we therefore find

$$(c) \quad \text{ind}_{D_0}^D({}^p H^i(\bar{K}_{D_0}^{s, \mathcal{L}})) = {}^p H^{i+2\alpha}(\bar{K}_D^{s, \mathcal{L}})$$

for any $i \in \mathbf{Z}$. To prove (a) it is enough to verify the following statement.

(d) *If $A_1 \in \hat{D}_0^{\mathcal{L}}$, then $\text{ind}_{D_0}^D(A_1)$ is a direct sum of character sheaves in $\hat{D}^{\mathcal{L}}$.*

We may assume that A_1 is a direct summand of ${}^p H^i(\bar{K}_{D_0}^{s,\mathcal{L}})$. Then $\text{ind}_{D_0}^D(A_1)$ is a direct summand of $\text{ind}_{D_0}^D({}^p H^i(\bar{K}_{D_0}^{s,\mathcal{L}}))$. From (c) we see that $\text{ind}_{D_0}^D(A_1)$ is a direct summand of ${}^p H^{i+2\alpha}(\bar{K}_D^{s,\mathcal{L}})$ which is a direct sum of character sheaves in $\hat{D}^{\mathcal{L}}$. This proves (d) hence also (a).

38.4. For $J \subset J' \subset \mathbf{I}$ such that $\epsilon(J) = J, \epsilon(J') = J'$ we have functors $\tilde{f}_{J,J'} : \mathcal{D}(V_{J,D}) \rightarrow \mathcal{D}(V_{J',D}), \tilde{e}_{J,J'} : \mathcal{D}(V_{J',D}) \rightarrow \mathcal{D}(V_{J,D})$; see 30.4.

Let $K, K', J \subset \mathbf{I}$ be such that $K \subset J, K' \subset J, \epsilon(J) = J, \epsilon(K) = K, \epsilon(K') = K'$. For any $u \in {}^K \mathbf{W}^{K'} \in \mathbf{W}_J^\epsilon$ let

$$\Xi_u = \{(X, Y, g(U_X \cap U_Y)); X \in \mathcal{P}_{K \cap \text{Ad}(u)K'}, Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}, \\ g(U_X \cap U_Y) \in (N_D X \cap N_D Y)/(U_X \cap U_Y), \text{pos}(X, Y) = u\}.$$

We have a diagram

$$V_{K \cap \text{Ad}(u)K', D} \xleftarrow{j} \Xi_u \xrightarrow{h} V_{K' \cap \text{Ad}(u^{-1})K, D}$$

where $j(X, Y, g(U_X \cap U_Y)) = (X, gU_X), h(X, Y, g(U_X \cap U_Y)) = (Y, gU_Y)$. Set $\Psi_u = h_! j^* : \mathcal{D}(V_{K \cap \text{Ad}(u)K', D}) \rightarrow \mathcal{D}(V_{K' \cap \text{Ad}(u^{-1})K, D})$.

Lemma 38.5. *Let $A' \in \mathcal{D}(V_{K,D})$. We set $\tilde{\mathcal{C}} = \tilde{e}_{K',J} \tilde{f}_{K,J} A' \in \mathcal{D}(V_{K',D})$. For any $u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon$ we set $\tilde{\mathcal{C}}_u = \tilde{f}_{K' \cap \text{Ad}(u^{-1})K, K'} \Psi_u \tilde{e}_{K \cap \text{Ad}(u)K', K} A' \in \mathcal{D}(V_{K',D})$ and $m_u = \dim(U_P \cap U_R)/U_Q$ where $P \in \mathcal{P}_K, R \in \mathcal{P}_{K'}, \text{pos}(P, R) = u$ and $Q = Q_{J,P} = Q_{J,R} \in \mathcal{P}_J$ (notation of 36.4). We have*

$$\tilde{\mathcal{C}} \simeq \{\tilde{\mathcal{C}}_u[[-m_u]]; u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon\}$$

where \simeq is as in 32.15.

The proof is very similar to that of Proposition 37.2. We have a commutative diagram with a cartesian square

$$\begin{array}{ccccc} E & \xrightarrow{b} & V_{K',J,D} & \xrightarrow{c'} & V_{K',D} \\ a \downarrow & & d' \downarrow & & \\ V_{K,D} & \xleftarrow{c} & V_{K,J,D} & \xrightarrow{d} & V_{J,D} \end{array}$$

Here

$$E = \{(P, R, gU_Q); P \in \mathcal{P}_K, R \in \mathcal{P}_{K'}, gU_Q \\ \in (N_D P \cap N_D R)/U_Q, Q = Q_{J,P} = Q_{J,R}\}, \\ c(P, gU_Q) = (P, gU_P), d(P, gU_Q) = (Q, gU_Q) \text{ with } Q = Q_{J,P}, \\ c'(R, gU_Q) = (R, gU_R), d'(R, gU_Q) = (Q, gU_Q), \text{ with } Q = Q_{J,R}, \\ a(P, R, gU_Q) = (P, gU_Q), b(P, R, gU_Q) = (R, gU_Q).$$

We have

$$\tilde{\mathcal{C}} = c'_! d'^* d_! c^* A' = c'_! b_! a^* c^* A' = (c'b)_! (ca)^* A' = q_! p^* A'$$

where $q = c'b : E \rightarrow V_{K',D}, p = ca : E \rightarrow V_{K,D}$ are given by $q(P, R, gU_Q) = (R, gU_R), p(P, R, gU_Q) = (P, gU_P)$. We have a partition

$$E = \bigsqcup_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon} E_u$$

where $E_u = \{(P, R, gU_Q) \in E; \text{pos}(P, R) = u\}$ is locally closed in E . Let $p_u = p|_{E_u} : E_u \rightarrow V_{K,D}$, $q_u = q|_{E_u} : E_u \rightarrow V_{K',D}$. By 32.15, we have

$$q_! p^* A' \simeq \{q_u! p_u^* A'; u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon\}.$$

It remains to show that, for u as above, we have $q_u! p_u^* A' = \tilde{\mathcal{C}}_u[[-m_u]]$. We have a commutative diagram

$$\begin{array}{ccccc} V_{K,D} & \xleftarrow{\tilde{d}} & V_{K \cap \text{Ad}(u)K', K, D} & \xrightarrow{\tilde{c}} & V_{K \cap \text{Ad}(u)K', D} \\ & & \tilde{a} \uparrow & & j \uparrow \\ & & S'_u & \xrightarrow{\tilde{b}} & \Xi_u & \xrightarrow{h} & V_{K' \cap \text{Ad}(u^{-1})K, D} \\ & & x \uparrow & & & & \tilde{c}' \uparrow \\ & & S_u & \xrightarrow{y} & & \longrightarrow & V_{K' \cap \text{Ad}(u^{-1})K, K', D} \\ & & & & & & \tilde{d}' \downarrow \\ & & & & & & V_{K', D} \end{array}$$

Here

$$\begin{aligned} \tilde{c}(X, gU_P) &= (X, gU_X), \tilde{d}(X, gU_P) = (P, gU_P), \\ \tilde{c}'(Y, gU_R) &= (Y, gU_Y), \tilde{d}'(Y, gU_R) = (R, gU_R), \\ S'_u, \tilde{a}, \tilde{b} &\text{ are defined so that the square } (\tilde{a}, \tilde{b}, \tilde{c}, j) \text{ is cartesian;} \\ S_u, x, y &\text{ are defined so that the square } (x, y, h\tilde{b}, \tilde{c}') \text{ is cartesian.} \end{aligned}$$

Then

$$\begin{aligned} S_u &= \{(X, Y, g(U_X \cap U_Y), g'U_P, g''U_R); X \in \mathcal{P}_{K \cap \text{Ad}(u)K'}, Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}, \\ &g(U_X \cap U_Y) \in (N_D X \cap N_D Y)/(U_X \cap U_Y), g'U_P \in N_D X/U_P, \\ &g''U_R \in N_D Y/U_R, \text{pos}(X, Y) = u, P = Q_{K,X}, R = Q_{K',Y}, \\ &g'U_X = gU_X, g''U_Y = gU_Y\}. \end{aligned}$$

Set $r = \tilde{d}'y : S_u \rightarrow V_{K',D}$, $s : \tilde{d}\tilde{a}x : S_u \rightarrow V_{K,D}$. Then

$$\begin{aligned} r(X, Y, g(U_X \cap U_Y), g'U_P, g''U_R) &= (P, g'U_P), \\ s(X, Y, g(U_X \cap U_Y), g'U_P, g''U_R) &= (R, g''U_R). \end{aligned}$$

We have

$$\begin{aligned} \tilde{\mathcal{C}}_u &= \tilde{d}'_! \tilde{c}'^* h_! j^* \tilde{c}_! \tilde{d}^* A' = \tilde{d}'_! \tilde{c}'^* h_! \tilde{b}_! \tilde{a}^* \tilde{d}^* A' = \tilde{d}'_! \tilde{c}'^* (h\tilde{b})_! (\tilde{d}\tilde{a})^* A' \\ &= \tilde{d}'_! y_! x^* (\tilde{d}\tilde{a})^* A' = (\tilde{d}'y)_! (\tilde{d}\tilde{a}x)^* A' = r_! s^* A'. \end{aligned}$$

We show that $q_u! p_u^* A' = r_! s^* A'[[-m_u]]$. We have a commutative diagram

$$\begin{array}{ccccc} V_{K,D} & \xleftarrow{p_u} & E_u & \xrightarrow{q_u} & V_{K',D} \\ 1 \downarrow & & t \downarrow & & 1 \downarrow \\ V_{K,D} & \xleftarrow{s} & S_u & \xrightarrow{r} & V_{K',D} \end{array}$$

where $t(P, R, gU_Q) = (P^R, R^P, g(U_{P^R} \cap U_{R^P}), gU_P, gU_R)$ is well defined since $U_Q \subset U_P, U_Q \subset U_R, U_Q \subset U_{P^R}, U_Q \subset U_{R^P}$. We continue the proof assuming that

- (a) t is an affine space bundle with fibres of dimension m_u .

For any $\tilde{A} \in \mathcal{D}(S_u)$ we have $t_!t^*(\tilde{A}) = \tilde{A}[[-m_u]]$. Hence

$$r_!s^*A'[[-m_u]] = r_!t_!t^*s^*A' = (rt)_!(st)^*A' = q_u!p_u^*A',$$

as required.

We prove (a). Consider the commutative diagram

$$\begin{array}{ccc} E_u & \longrightarrow & \mathfrak{E}_u \\ t \downarrow & & \downarrow \mathfrak{t} \\ S_u & \longrightarrow & \mathfrak{S}_u \end{array}$$

where $\mathfrak{E}_u \xrightarrow{\mathfrak{t}} \mathfrak{S}_u$ is as in the proof of 37.2 and the horizontal maps are the obvious imbeddings. Clearly, this diagram is cartesian. Hence (a) is a consequence of the analogous statement 37.2(a) for $\mathfrak{E}_u \xrightarrow{\mathfrak{t}} \mathfrak{S}_u$. This completes the proof.

38.6. In the setup of 38.4, let $u \in {}^K\mathbf{W}^{K'} \cap \mathbf{W}^\epsilon_j$. Let $\delta = \dim(U_{X_u}/(U_{X_u} \cap U_{Y_u})) = \dim(U_{Y_u}/(U_{X_u} \cap U_{Y_u}))$ where $(X_u, Y_u, g(U_{X_u} \cap U_{Y_u})) \in \Xi_u$. Let $\alpha_u = \dim U_{X_u} = \dim U_{Y_u}$, $\alpha = \dim U_P, \alpha' = \dim U_R$ where $P \in \mathcal{P}_K, R \in \mathcal{P}_{K'}$. We show that

(a) *h and j in 38.4 are affine space bundles with fibres of dimension δ .*

It is enough to prove the statements on j (the statement on h is entirely similar). We show that j is surjective. Let X, g be such that $(X, gU_X) \in V_{K \cap \text{Ad}(u)K', D}$. We must show that there exist $Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}, g' \in N_D X \cap N_D Y$ such that $\text{pos}(X, Y) = u$ and $g'U_X = gU_X$. Setting $g^{-1}g' = v$, it is enough to show that for any $Y \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}$ with $\text{pos}(X, Y) = u$ there exists $v \in U_X$ such that $gv \in N_G Y$. Now X, Y contain a common Levi M . Since $g \in N_G X$, we can find $v \in U_X$ such that $g' = gv \in N_G X \cap N_G M$. There is a unique parabolic Y' of the same type as Y such that Y' has Levi M and $\text{pos}(X, Y') = u$. Then $\text{pos}(g'Xg'^{-1}, g'Yg'^{-1}) = u, g'Yg'^{-1}$ has Levi $g'Mg'^{-1} = M$. By uniqueness, we have $Y' = Y$. Thus, $g' \in N_G Y$.

We show that the fibres of j are affine spaces of dimension δ . Let $(X, Y, g(U_X \cap U_Y)) \in \Xi_u$. We must show that $F = \{(X, Y', g'(U_X \cap U_{Y'})) \in \Xi_u; gU_X = g'U_X\}$ is an affine space. Fix $Y' \in \mathcal{P}_{K' \cap \text{Ad}(u^{-1})K}$ such that $\text{pos}(X, Y') = u$. (The set of such Y' is a homogeneous space $U_X/(U_X \cap Y)$, hence is an affine space of dimension δ .) It is enough to show that $\{g'(U_X \cap U_{Y'}) \in (N_D X \cap N_D Y)/(U_X \cap U_Y); g' \in gU_X\}$ is a point. Now $g = g_0v_0$ where $g_0 \in N_D X \cap N_D Y, v_0 \in U_X$. It is enough to show that $\{v \in U_X; v_0v \in N_G X \cap N_G Y\}/(U_X \cap U_Y)$ is a point or that $(U_X \cap N_G X \cap N_G Y)/(U_X \cap U_Y)$ is a point or that $U_X \cap Y = U_X \cap U_Y$. This is clear.

38.7. Let $u \in {}^K\mathbf{W}^{K'} \cap \mathbf{W}^\epsilon_j$. We set $H = K \cap \text{Ad}(u)K', H' = K' \cap \text{Ad}(u^{-1})K = \text{Ad}(u^{-1})H$. Let $\Psi'_u = \Psi_u[[\delta]] : \mathcal{D}(V_{H,D}) \rightarrow \mathcal{D}(V_{H',D})$. We show that

(a) *If $A \in CS(V_{H,D})$, then $\Psi'_u(A) \in CS(V_{H',D})$.*

We have a commutative diagram in which the upper squares are cartesian and the

left and right vertical arrows are smooth with connected fibres:

$$\begin{array}{ccccc}
 V_{H,D} & \xleftarrow{j} & \Xi_u & \xrightarrow{h} & V_{H',D} \\
 f'_3 \uparrow & & f'_4 \uparrow & & f'_5 \uparrow \\
 V'_H & \xleftarrow{j'} & \Xi''_u & \xrightarrow{h'} & V'_{H'} \\
 f_3 \downarrow & & f_4 \downarrow & & f_5 \downarrow \\
 V_H^u & \xleftarrow{j^u} & \Xi'_u & \xrightarrow{h^u} & V_{H'}^u
 \end{array}$$

Here

$$\begin{aligned}
 V_H^u &= \{(X, gU_X) \in V_{H,D}; X = X_u\}, V_{H'}^u = \{(Y, gU_Y) \in V_{H',D}; Y = Y_u\}, \\
 \Xi'_u &= \{(X, Y, g(U_X \cap U_Y)) \in \Xi_u; X = X_u, Y = Y_u\}, \\
 V'_H &= G^0/U_{X_u} \times V_H^u, V'_{H'} = G^0/U_{Y_u} \times V_{H'}^u, \\
 \Xi''_u &= G^0/(U_{X_u} \cap U_{Y_u}) \times \Xi'_u, \\
 j^u, h^u &\text{ are the restrictions of } j, h, \\
 j'(x(U_{X_u} \cap U_{Y_u}), (X_u, Y_u, g(U_{X_u} \cap U_{Y_u}))) &= (xU_{X_u}, (X_u, gU_{X_u})), \\
 h'(x(U_{X_u} \cap U_{Y_u}), (X_u, Y_u, g(U_{X_u} \cap U_{Y_u}))) &= (xU_{Y_u}, (Y_u, gU_{Y_u})), \\
 f'_3(xU_{X_u}, (X_u, gU_{X_u})) &= (xX_u x^{-1}, xg x^{-1} U_{xX_u x^{-1}}), \\
 f_3(xU_{X_u}, (X_u, gU_{X_u})) &= (X_u, gU_{X_u}), \\
 f'_4(x(U_{X_u} \cap U_{Y_u}), (X_u, Y_u, g(U_{X_u} \cap U_{Y_u}))) &= (xX_u x^{-1}, xY_u x^{-1}, xg x^{-1} (U_{xX_u x^{-1}} \cap \\
 U_{xY_u x^{-1}})), \\
 f'_5(xU_{Y_u}, (Y_u, gU_{Y_u})) &= (xY_u x^{-1}, xg x^{-1} U_{xY_u x^{-1}}), \\
 f_4(x(U_{X_u} \cap U_{Y_u}), (X_u, Y_u, g(U_{X_u} \cap U_{Y_u}))) &= (X_u, Y_u, g(U_{X_u} \cap U_{Y_u})), \\
 f_5(xU_{Y_u}, (Y_u, gU_{Y_u})) &= (Y_u, gU_{Y_u}).
 \end{aligned}$$

We may identify $V_H^u = N_D X_u / U_{X_u}$, $V_{H'}^u = N_D Y_u / U_{Y_u}$ in an obvious way. We can find $C \in CS(N_D X_u / U_{X_u})$ such that $f_3^{\star} A = f_3^{\star} C$. Hence $f_3^{\star} A[\dim X_u / U_{X_u}] = f_3^{\star} C[\dim G / U_{X_u}]$ and $f_3^{\star} A = f_3^{\star} A_1$ where $A_1 = C[\alpha_u]$. Let $A_2 = h_1^u j^{u*} A_1, A_2 = h_1 j^* A$. Using 38.6(a) and the fact that h^u is an isomorphism, we have

$$\begin{aligned}
 f_5^* A_2 &= f_5^* h_1 j^* A = h_1' f_4^* j^* A = h_1' j'^* f_3^* A = h_1' j'^* f_3^* A_1 = h_1' f_4^* j^{u*} A_1 \\
 &= h_1' f_4^* h^{u*} h_1^u j^{u*} A_1 = h_1' h'^* f_5^* h_1^u j^{u*} A_1 = f_5^* h_1^u j^{u*} A_1[[-\delta]] = f_5^* A_2[[-\delta]].
 \end{aligned}$$

Thus, $f_5^* A_2[[-\delta]] = f_5^* A_2'$. Hence

$$f_5^{\star} A_2[-\dim G^0 / U_{Y_u}][[-\delta]] = f_5^{\star} A_2'[-\dim Y_u / U_{Y_u}], f_5^{\star} A_2[-\alpha_u] = f_5^{\star} A_2'[[\delta]].$$

Since h^u, j^u are isomorphisms we see that $A_2[-\alpha_u]$ is a perverse sheaf. Hence so is $f_5^{\star} A_2[-\alpha_u]$. Hence $f_5^{\star} A_2'[[\delta]]$ is perverse. Hence $A_2'[[\delta]]$ is perverse and $h_1 j^* A[[\delta]]$ is perverse. To show that $h_1 j^* A[[\delta]] = A_2'[[\delta]] \in CS(V_{H',D})$, it is enough to show that $A_2[-\alpha_u] \in CS(N_D Y_u / U_{Y_u})$ or that $h_1^u j^{u*} C \in CS(N_D Y_u / U_{Y_u})$.

Let M be a common Levi subgroup of X_u, Y_u . Let $\tilde{M} = N_G X_u \cap N_G Y_u \cap N_G M$. Then \tilde{M} is a reductive group with $\tilde{M}^0 = M$ and $\tilde{M}^1 = N_D X_u \cap N_D Y_u \cap N_D M$ is a connected component of \tilde{M} . Moreover, the obvious maps $\tilde{M}^1 \rightarrow N_D X_u / U_{X_u}, \tilde{M}^1 \rightarrow N_D Y_u / U_{Y_u}, \tilde{M}^1 \rightarrow (N_D X_u \cap N_D Y_u) / (U_{X_u} \cap U_{Y_u})$ are isomorphisms (see 1.25). Hence the bottom row $V_H^u \xleftarrow{j^u} \Xi'_u \xrightarrow{h^u} V_{H'}^u$ of the commutative diagram above may be identified with $\tilde{M}^1 \leftarrow \tilde{M}^1 \rightarrow \tilde{M}^1$ where both maps are the identity. Thus $h_1^u j^{u*} C \in CS(N_D Y_u / U_{Y_u})$ follows immediately from $C \in CS(N_D X_u / U_{X_u})$. This proves (a).

We now show that (b) for A as above we have $f_{H',J} \Psi'_u(A) = f_{H,J}(A)$.

Let $Q \in \mathcal{P}_J$ be such that $X_u \subset Q \supset Y_u$. Let M_1 be the unique Levi of Q such that $M \subset M_1$. Let $\tilde{M}_1 = N_G Q \cap N_G M_1$. Then \tilde{M}_1 is a reductive group with $\tilde{M}_1^0 = M_1$ and $\tilde{M}_1^1 = N_D Q \cap N_D M_1$ is a connected component of \tilde{M}_1 . Moreover, $X_u \cap \tilde{M}_1, Y_u \cap \tilde{M}_1$ are parabolic subgroups of M_1 with a common Levi, M . Let $C \in CS(N_D X_u / U_{X_u}) = CS(\tilde{M}_1^1)$ be as above. We may assume that A is simple so that C is also simple. Using the proof of (a) and that of 38.2 we see that it is enough to verify that $\text{ind}_{\tilde{M}_1^1}^{\tilde{M}_1}(C)$ (defined in terms of the parabolic $X_u \cap \tilde{M}_1$) is isomorphic to $\text{ind}_{\tilde{M}_1^1}^{\tilde{M}_1}(C)$ (defined in terms of the parabolic $Y_u \cap \tilde{M}_1$). Since C is an admissible complex on \tilde{M}_1^1 (see 30.12), this follows from 27.2(d) which shows (for G instead of \tilde{M}_1) that $\text{ind}_{\tilde{M}_1^1}^{\tilde{M}_1}(C)$ can be defined without reference to a choice of parabolic. This proves (b).

Proposition 38.8. *Let K, K', J be as in 38.4. Let $A \in CS(V_{K,D})$. We have*

$$e_{K',J} f_{K,J} A \cong \bigoplus_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon} f_{K' \cap \text{Ad}(u^{-1})K, K'} \Psi'_u e_{K \cap \text{Ad}(u)K', K} A$$

in $CS(V_{K',D})$.

We set $\mathfrak{C} = e_{K',J} f_{K,J} A \in \mathcal{D}(V_{K,D})$. For any $u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon$ we set $\mathfrak{C}_u = f_{K' \cap \text{Ad}(u^{-1})K, K'} \Psi'_u e_{K \cap \text{Ad}(u)K', K} A \in \mathcal{D}(V_{K',D})$. Assume that we can show that

$$\mathfrak{C} \simeq \{\mathfrak{C}_u; u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon\}.$$

From the definition of \simeq (see 32.15) it would then follow that

$$\sum_i (-1)^i ({}^p H^i(\mathfrak{C})) = \sum_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon} \sum_i (-1)^i ({}^p H^i(\mathfrak{C}_u))$$

in the Grothendieck group of the category of perverse sheaves on $V_{K',D}$. From 38.2(a), 38.3(a), 38.7(a) we see that $\mathfrak{C}, \mathfrak{C}_u \in CS(V_{K',D})$ (hence are perverse sheaves); hence the previous equality implies that $\mathfrak{C} = \sum_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon} \mathfrak{C}_u$ in the Grothendieck group of the category of perverse sheaves on $V_{K',D}$. Since $\mathfrak{C}, \mathfrak{C}_u$ are semisimple perverse sheaves (being in $CS(V_{K',D})$) it follows that $\mathfrak{C} \cong \bigoplus_{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon} \mathfrak{C}_u$, as desired.

Assume that $P \in \mathcal{P}_K$ contains X_u and $R \in \mathcal{P}_{K'}$ contains Y_u so that $\text{pos}(P, R) = u$. (X_u, Y_u as in 38.6.) Let $Q \in \mathcal{P}_J$ be such that $P \subset Q \supset R$. Let $\beta = \dim U_Q$. We have

$$\begin{aligned} \mathfrak{C} &= \tilde{\mathfrak{C}}[\alpha + \alpha' - 2\beta](\alpha' - \beta), \\ \mathfrak{C}_u &= \tilde{\mathfrak{C}}_u\alpha_u - \alpha[\alpha_u - \alpha'] [2\delta](\delta). \end{aligned}$$

(Notation of 38.5.) Hence it is enough to show

$$\tilde{\mathfrak{C}}[\alpha + \alpha' - 2\beta](\alpha' - \beta) \simeq \{\tilde{\mathfrak{C}}_u[2\alpha_u - \alpha - \alpha' + 2\delta](\alpha_u - \alpha + \delta); u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon\}$$

or that

$$\tilde{\mathfrak{C}} \simeq \{\tilde{\mathfrak{C}}_u[2\alpha_u - 2\alpha - 2\alpha' + 2\delta - 2\beta](\alpha_u - \alpha - \alpha' + \delta + \beta); u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon\}.$$

By 38.5, it is enough to show that for any u we have $\alpha_u - \alpha - \alpha' + \delta + \beta = -m_u$ or that

$$\dim U_P + \dim U_R - \dim(U_P \cap U_R) = \dim U_{X_u} + \dim U_{Y_u} - \dim(U_{X_u} \cap U_{Y_u}).$$

It is enough to show that

$$\begin{aligned} & \dim \text{Lie } U_P + \dim \text{Lie } U_R - \dim(\text{Lie } U_P \cap \text{Lie } U_R) \\ &= \dim \text{Lie } U_{X_u} + \dim \text{Lie } U_{Y_u} - \dim(\text{Lie } U_{X_u} \cap \text{Lie } U_{Y_u}) \end{aligned}$$

or that $\dim(\text{Lie } U_P + \text{Lie } U_R) = \dim(\text{Lie } U_{X_u} + \text{Lie } U_{Y_u})$. We have

$$\text{Lie } U_{X_u} = \text{Lie } U_P + (\text{Lie } P \cap \text{Lie } U_R), \text{Lie } U_{Y_u} = \text{Lie } U_R + (\text{Lie } R \cap \text{Lie } U_P),$$

hence $\text{Lie } U_{X_u} + \text{Lie } U_{Y_u} \subset \text{Lie } U_P + \text{Lie } U_R$. The opposite inclusion is clear since $\text{Lie } U_P \subset \text{Lie } U_{X_u}, \text{Lie } U_R \subset \text{Lie } U_{Y_u}$. Thus we have

$$\text{Lie } U_{X_u} + \text{Lie } U_{Y_u} = \text{Lie } U_P + \text{Lie } U_R.$$

This completes the proof.

38.9. For $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ let $\mathcal{K}(V_{J,D})$ be the Grothendieck group of $CS(V_{J,D})$. Similarly let $\mathcal{K}(D)$ be the Grothendieck group of $CS(D)$. For $A, A' \in CS(V_{J,D})$ we set $(A, A') = \dim \text{Hom}(A, A')$. This induces a symmetric bilinear pairing $(\cdot, \cdot)_J : \mathcal{K}(V_{J,D}) \times \mathcal{K}(V_{J,D}) \rightarrow \mathbf{Z}$.

For $J \subset J' \subset \mathbf{I}$ such that $\epsilon(J) = J, \epsilon(J') = J'$, the functors $f_{J,J'} : CS(V_{J,D}) \rightarrow CS(V_{J',D})$ and $e_{J,J'} : CS(V_{J',D}) \rightarrow CS(V_{J,D})$ are compatible with direct sums hence they induce homomorphisms $\mathcal{K}(V_{J,D}) \rightarrow \mathcal{K}(V_{J',D}), \mathcal{K}(V_{J',D}) \rightarrow \mathcal{K}(V_{J,D})$ denoted again by $f_{J,J'}, e_{J,J'}$. From 30.5 we see that

$$(a) \quad (e_{J,J'} A', A)_J = (A', f_{J,J'} A)_{J'}$$

for $A \in \mathcal{K}(V_{J,D}), A' \in \mathcal{K}(V_{J',D})$.

In the setup of 38.7, for $u \in {}^K \mathbf{W}^{K'} \in \mathbf{W}_J^\epsilon$, the functor

$$\Psi'_u : CS(V_{K \cap \text{Ad}(u)K', D}) \rightarrow CS(V_{K' \cap \text{Ad}(u^{-1})K, D})$$

is compatible with direct sums hence induces a homomorphism

$$\mathcal{K}(V_{K \cap \text{Ad}(u)K', D}) \rightarrow \mathcal{K}(V_{K' \cap \text{Ad}(u^{-1})K, D})$$

denoted again by Ψ'_u . Below we shall need the following identity:

$$(b) \quad \sum_{\substack{K'; K' \subset J \\ \epsilon(K')=K}} \sum_{\substack{u \in {}^K \mathbf{W}^{K'} \cap \mathbf{W}_J^\epsilon \\ K \cap \text{Ad}(u)K'=H}} (-1)^{|K'|} = (-1)^{|H|}$$

for any $H \subset K \subset J$ such that $\epsilon(H) = H, \epsilon(K) = K$. In the case where $J = \mathbf{I}, \epsilon = 1$ this is proved in [Cu, 2.5]; the general case can be reduced to this special case by replacing \mathbf{W} by \mathbf{W}_J^ϵ which is itself a Weyl group with simple reflections in bijection with J_ϵ .

38.10. For $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ we define a homomorphism $\mathbf{d}_J : \mathcal{K}(V_{J,D}) \rightarrow \mathcal{K}(V_{J,D})$ by

$$(a) \quad \mathbf{d}_J A = \sum_{K; K \subset J, \epsilon(K)=K} (-1)^{|K|} f_{K,J} e_{K,J} A.$$

Using 38.9(a) we see that for $A, A' \in \mathcal{K}(V_{J,D})$ we have

$$(b) \quad (\mathbf{d}_J(A), A')_J = (A, \mathbf{d}_J(A'))_J.$$

We show that, for $K \subset J \subset \mathbf{I}$ such that $\epsilon(K) = K, \epsilon(J) = J$ and $A \in \mathcal{K}(V_{K,D})$, we have

$$(c) \quad \mathbf{d}_J f_{K,J} A = f_{K,J} \mathbf{d}_K A.$$

Using 38.8, 38.7(b), 38.9(b), 38.1(a) we have

$$\begin{aligned}
 \mathbf{d}_J f_{K,J} A &= \sum_{K'; K' \subset J, \epsilon(K')=K'} (-1)^{|K'_\epsilon|} f_{K',J} e_{K',J} f_{K,J} A \\
 &= \sum (-1)^{|K'_\epsilon|} f_{K',J} f_{K' \cap \text{Ad}(u^{-1})K, K'} \Psi'_u e_{K \cap \text{Ad}(u)K', K} A \\
 &= \sum (-1)^{|K'_\epsilon|} f_{\text{Ad}(u^{-1})H, J} \Psi'_u e_{H, K} A \\
 &= \sum (-1)^{|K'_\epsilon|} f_{H, J} e_{H, K} A \\
 &= \sum (-1)^{|K'_\epsilon|} f_{H, J} e_{H, K} A = \sum_{H; H \subset K, \epsilon(H)=H} (-1)^{|H_\epsilon|} f_{H, J} e_{H, K} A \\
 &= \sum_{H; H \subset K, \epsilon(H)=H} (-1)^{|H_\epsilon|} f_{K, J} f_{H, K} e_{H, K} A = f_{K, J} \mathbf{d}_K A
 \end{aligned}$$

and (c) is proved.

We show that, for $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ and $A \in \mathcal{K}(V_{J,D})$, we have

$$(d) \quad \mathbf{d}_J \mathbf{d}_J A = A.$$

Using (c), 38.1(a), we have

$$\begin{aligned}
 \mathbf{d}_J \mathbf{d}_J A &= \sum_{K; K \subset J, \epsilon(K)=K} (-1)^{|K_\epsilon|} \mathbf{d}_J f_{K, J} e_{K, J} A \\
 &= \sum_{K; K \subset J, \epsilon(K)=K} (-1)^{|K_\epsilon|} f_{K, J} \mathbf{d}_K e_{K, J} A \\
 &= \sum_{K, K'; K' \subset K \subset J, \epsilon(K)=K, \epsilon(K')=K'} (-1)^{|K_\epsilon|} (-1)^{|K'_\epsilon|} f_{K, J} f_{K', K} e_{K', K} e_{K, J} A \\
 &= \sum_{K, K'; K' \subset K \subset J, \epsilon(K)=K, \epsilon(K')=K'} (-1)^{|K_\epsilon|} (-1)^{|K'_\epsilon|} f_{K', J} e_{K', J} A \\
 &= \sum_{K'; K' \subset J, \epsilon(K')=K'} (-1)^{|K'_\epsilon|} \sum_{K; K' \subset K \subset J, \epsilon(K)=K} (-1)^{|K_\epsilon|} f_{K', J} e_{K', J} A \\
 &= \sum_{K'; K' \subset J, \epsilon(K')=K'} (-1)^{|K'_\epsilon|} \delta_{K', J} (-1)^{|K'_\epsilon|} f_{K', J} e_{K', J} A = f_{J, J} e_{J, J} A = A
 \end{aligned}$$

and (d) is proved.

We show that, for $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ and $A, A' \in \mathcal{K}(V_{J,D})$, we have

$$(e) \quad (\mathbf{d}_J A, \mathbf{d}_J A')_J = (A, A')_J.$$

Using (b),(d) we have $(\mathbf{d}_J A, \mathbf{d}_J A')_J = (A, \mathbf{d}_J \mathbf{d}_J A')_J = (A, A')_J$ as desired.

38.11. We write \mathbf{d} instead of $\mathbf{d}_{\mathbf{I}}$ and $(,)$ instead of $(,)_{\mathbf{I}}$. We call \mathbf{d} the duality operator on character sheaves. If A is a character sheaf on $D = V_{\mathbf{I},D}$, then $(A, A) = 1$ (where A is regarded as an element of $\mathcal{K}(D)$) hence, by 38.10(e), we have $(\mathbf{d}(A), \mathbf{d}(A)) = 1$. Since $\mathbf{d}(A)$ is a \mathbf{Z} -linear combination of isomorphism classes of character sheaves (which form an orthonormal basis of $\mathcal{K}(D)$ for $(,)$) it follows that $\mathbf{d}(A) = \pm A'$ where A' is a well-defined character sheaf on D (up to isomorphism). The sign can be described as follows. By 30.6(d) we can find a parabolic P_0 of G^0 such that $N_D P_0 \neq \emptyset$ and a cuspidal character sheaf A_0 on $D_0 := N_D P_0 / U_{P_0}$ such

that A is a direct summand of $\text{ind}_{D_0}^D(A_0)$. We have $P_0 \in \mathcal{P}_J$ where $J \subset \mathbf{I}, \epsilon(J) = J$. Then

$$(a) \quad \mathbf{d}(A) = (-1)^{|J^\epsilon|} A'.$$

Indeed, let $A_0^b \in CS(V_{D,J})$ be the perverse sheaf corresponding to A_0 as in 30.3. Then $e_{J',J} A_0^b = 0$ for any $J' \subsetneq J$ such that $\epsilon(J') = J'$; see 38.2. Hence $\mathbf{d}_J(A_0^b) = (-1)^{|J^\epsilon|} A_0^b$. Now A is a direct summand of $f_{J,\mathbf{I}} A_0^b$ and by 38.10(c) we have

$$\mathbf{d}f_{J,\mathbf{I}} A_0^b = f_{J,\mathbf{I}} \mathbf{d}_J A_0^b = (-1)^{|J^\epsilon|} f_{J,\mathbf{I}} A_0^b.$$

In $\mathcal{K}(D)$ we have $f_{J,\mathbf{I}} A_0^b = \sum_{k=1}^{k_0} n_k A_k$ where A_k are distinct character sheaves on D , $n_k \in \mathbf{Z}_{>0}$ and $A_1 = A$. We have $\mathbf{d}(A_k) = \iota_k A'_k$ where $\iota_k = \pm 1$ and A'_k are distinct character sheaves on D , $n_k \in \mathbf{Z}_{>0}$ and $A_1 = A$. We see that $\sum_k n_k \iota_k A'_k = (-1)^{|J^\epsilon|} \sum_k n_k A_k$. Since $\{A_k\}$ and $\{A'_k\}$ are parts of the same basis of $\mathcal{K}(D)$ we see that $\iota_k / (-1)^{|J^\epsilon|} > 0$ for any k . Hence $\iota_k = (-1)^{|J^\epsilon|}$ for any k . In particular, this holds for $k = 1$ and (a) follows.

Note that, by 38.10(d), $A \mapsto A'$ is an involution of the set of isomorphism classes of character sheaves on D .

38.12. The definition of the duality operator for character sheaves in 38.11 is entirely similar to that of a duality operator for representations of a reductive group over a finite field (given again by an alternating sum of compositions of a parabolic restriction and a parabolic induction) which was found by the author in 1977, who conjectured that it takes irreducibles to \pm irreducibles and is involutive. In 1977 I communicated this conjecture to C.W. Curtis and N. Kawanaka (see [Ka, p.412]); the conjecture was proved in [Cu], [Al] and in [Ka].

39. QUASI-RATIONALITY

39.1. The main result of this section is Proposition 39.7 which gives a quasi-rationality property of representations of certain extensions of a Weyl group. This is needed to prove a key property of character sheaves (Corollary 39.8).

Let W, I be a Weyl group (I is the set of simple reflections). We have canonically $W = \prod_{j \in J} W_j$ where W_j is an irreducible Weyl group (with set of simple reflections $I_j = I \cap W_j$). We identify W_j with a subgroup of W . Let $R(W) = \bigcup_{w \in W} wIw^{-1}$. We have $R(W) = \bigsqcup_{j \in J} R(W_j)$. For any $j \in J$ the set $R(W_j)$ is a single W_j -conjugacy class if W_j is of type A, D or E ; it is a union of two W_j -conjugacy classes, otherwise. A subset \mathcal{X} of $R(W)$ is said to be *special* if $\mathcal{X} = \bigsqcup_{j \in J} \mathcal{X}_j$ where \mathcal{X}_j is a W_j -conjugacy class in $R(W_j)$. Clearly, a special subset of $R(W)$ exists; we assume that a special subset \mathcal{X} of $R(W)$ is given. Let $\text{Aut}(W, I, \mathcal{X})$ be the group of automorphisms of W that preserve I and \mathcal{X} .

Lemma 39.2. *Let Γ be a finite group. Let $\gamma \mapsto \rho_\gamma$ be a homomorphism $\Gamma \rightarrow \text{Aut}(W, I, \mathcal{X})$. Let E be a simple $\mathbf{Q}[W]$ -module such that $\text{tr}(\rho_\gamma(w), E) = \text{tr}(w, E)$ for any $\gamma \in \Gamma, w \in W$. Assume that either*

- (i) $|J| = 1$, or
- (ii) Γ is an extension of a cyclic group by a cyclic group.

Then there exists a homomorphism $\Gamma \rightarrow \text{Aut}_{\mathbf{Q}}(E)$, $\gamma \mapsto t_\gamma$ such that $t_\gamma(w(e)) = \rho_\gamma(w)t_\gamma(e)$ for any $\gamma \in \Gamma, w \in W, e \in E$.

Γ acts on J by $\gamma : j \mapsto \gamma(j)$ where $W_{\gamma(j)} = \rho_\gamma(W_j)$. We may identify $E = \bigotimes_{j \in J} E_j$ as $\mathbf{Q}[W]$ -modules where E_j is a simple $\mathbf{Q}[W_j]$ -module for any $j \in J$. From our assumption we see that for any $j \in J, \gamma \in \Gamma$ there exists a \mathbf{Q} -linear isomorphism

(a) $H_j^\gamma : E_j \xrightarrow{\sim} E_{\gamma(j)}$ with $H_j^\gamma(w_j e) = \rho_\gamma(w_j) H_j^\gamma(e)$ for all $e \in E_j, w_j \in W_j$;

moreover, H_j^γ is unique up to multiplication by an element of \mathbf{Q}^* . It follows that

(b) $H_{\gamma(j)}^{\gamma'} H_j^\gamma \in \mathbf{Q}^* H_j^{\gamma' \gamma}$ for $j \in J$ and $\gamma, \gamma' \in \Gamma$.

For $\gamma \in \Gamma$ we define $\tilde{t}_\gamma \in \text{Aut}_{\mathbf{Q}}(E)$ by

(c) $\tilde{t}_\gamma(\bigotimes_j e_j) = \bigotimes_j e'_j$ where $e_j \in E_j$ and $e'_j = H_{\gamma^{-1}(j)}^\gamma(e_{\gamma^{-1}(j)}) \in E_j$.

From definitions we have $\tilde{t}_\gamma(w(e)) = \rho_\gamma(w) \tilde{t}_\gamma(e)$ for any $w \in W, e \in E$; moreover, for $\gamma, \gamma' \in \Gamma$, the maps $\tilde{t}_{\gamma' \gamma}, \tilde{t}_{\gamma'} \tilde{t}_\gamma$ are equal up to a factor in \mathbf{Q}^* . Thus the maps \tilde{t}_γ provide a homomorphism $\Gamma \rightarrow PGL(E)$ rather than a homomorphism $\Gamma \rightarrow GL(E)$.

We prove the lemma in the setup of (i). Replacing Γ by its image under $\gamma \mapsto \rho_\gamma$ we may assume that $\Gamma \subset \text{Aut}(W, I, \mathcal{X})$ and $\gamma \mapsto \rho_\gamma$ is the inclusion. We form the semidirect product $W\Gamma$ with W normal. It is enough to show that E extends to a $W\Gamma$ -module. We may assume that $\Gamma \neq \{1\}$. Then W is of type $A_n (n \geq 2), D_n$ or E_6 . If $\Gamma = \text{Aut}(W, I, \mathcal{X})$, then $W\Gamma$ is itself a Weyl group, of type $A_n \times A_1, D_{2n+1} \times A_1, B_{2n}, F_4, E_6 \times A_1$ for W of type $A_n (n \geq 2), D_{2n+1} (n \geq 2), D_{2n} (n \geq 3), D_4, E_6$ respectively and the desired result follows easily from the known properties of representations of such Weyl groups (in particular, from their rationality); the same applies if W is of type D_4 and $|\Gamma| = 2$ (in this case, $W\Gamma$ is a Weyl group of type B_4). In the only remaining case (W of type $D_4, |\Gamma| = 3$), the result follows by an argument in [L14, 3.2].

Next we prove the lemma in the setup of (ii). Now Γ has two generators a, c and relations $a^M = 1, c^N = a^u, cac^{-1} = a^k$ where M, N, k are integers ≥ 1 such that $k^N = 1 \pmod M$ and $u \in \mathbf{N}$ satisfies $uk = k \pmod M$.

We consider separately 3 cases in increasing order of generality.

Case 1. Assume that J is a single orbit of $a : J \rightarrow J$. We may identify $J = \mathbf{Z}/m$ so that $a(i) = i + 1, c(i) = ki - r$ for $i \in \mathbf{Z}/m$; here $r \in \mathbf{N}$ is independent of i . Since $a^M = 1 : J \rightarrow J$ we see that m divides M . For any i we have $i + u = c^N(i) = k^N i - (1 + k + k^2 + \dots + k^{N-1})r$ in \mathbf{Z}/m . Since $k^N = 1 \pmod M$ (hence $k^N = 1 \pmod m$) we have $u + (1 + k + k^2 + \dots + k^{N-1})r = fm$ for some $f \in \mathbf{N}$. We set $c' = a^r c$. Then $c'(i) = ki$ for $i \in \mathbf{Z}/m$. Let Γ' be the subgroup of Γ generated by a^m, c' . In Γ' we have

$$(a^m)^{M/m} = 1, c'^N = (a^m)^f, c' a^m c'^{-1} = (a^m)^k.$$

Since $a^m(0) = 0, c'(0) = 0$, the action of Γ on W restricts to an action of Γ' on W_0 . Using the lemma (setup of (i)) for W_0, E_0, Γ' instead of W, E, Γ we obtain a homomorphism $\Gamma' \rightarrow \text{Aut}_{\mathbf{Q}}(E_0)$ such that, denoting by s_0, s_1 the images of a^m, c' under this homomorphism, we have $s_0^{M/m} = 1, s_1^N = s_0^f, s_1 s_0 s_1^{-1} = s_0^k, s_0(w(e)) = \rho_a^m(w) s_0(e), s_1(w(e)) = \rho_{c'}(w) s_1(e)$ for any $w \in W_0, e \in E_0$.

For $i \in \mathbf{Z}/m$, let $H_i^a : E_i \xrightarrow{\sim} E_{i+1}$ be as in (a). Then $s_2 := H_{m-1}^a \dots H_1^a H_0^a : E_0 \xrightarrow{\sim} E_0$ satisfies $s_2(w(e)) = \rho_a^m(w) s_2(e)$ for any $w \in W_0, e \in E_0$. Hence $s_2^{-1} s_0(w(e)) = w(s_2^{-1} s_0(e))$ for any $w \in W_0, e \in E_0$. By the absolute irreducibility of the W_0 -module E_0 we see that $s_2^{-1} s_0 : E_0 \rightarrow E_0$ is a \mathbf{Q}^* -multiple of the identity

map. Hence, replacing H_0^a by a \mathbf{Q}^* -multiple, we may assume that $s_2 = s_0$, that is, $H_{m-1}^a \dots H_1^a H_0^a = s_0$. For any $h \in \mathbf{N}$ we set

$$b_h = H_{hk-1}^a H_{hk-2}^a \dots H_1^a H_0^a s_1 (H_{h-1}^a \dots H_1^a H_0^a)^{-1} : E_h \xrightarrow{\sim} E_{kh}$$

(there are hk factors to the left of s_1 and h factors to the right of s_1). For $h \geq 1$ we have

$$(d) \quad b_h = H_{hk-1}^a H_{hk-2}^a \dots H_{hk-k}^a b_{h-1} (H_{h-1}^a)^{-1}.$$

We show that $b_{h+m} = b_h$ for $h \in \mathbf{N}$. We argue by induction on h . Assume first that $h = 0$. We must show that $H_{mk-1}^a H_{mk-2}^a \dots H_1^a H_0^a s_1 (H_{m-1}^a \dots H_1^a H_0^a)^{-1} = s_1$; this is a reformulation of the already known equality $s_0^k s_1 s_0^{-1} = s_1$. Assume next that $h \geq 1$. Using (d) twice and the induction hypothesis we have

$$\begin{aligned} b_{h+m} &= H_{hk+mk-1}^a H_{hk+mk-2}^a \dots H_{hk+mk-k}^a b_{h+m-1} (H_{h+m-1}^a)^{-1} \\ &= H_{hk-1}^a H_{hk-2}^a \dots H_{hk-k}^a b_{h-1} (H_{h-1}^a)^{-1} = b_h, \end{aligned}$$

as desired. We see that b_h depends only on the image of h in \mathbf{Z}/m .

We set $k = k^{N-1}$. Then $kk' = 1 \pmod{M}$. Define $t_a, t_{c'} \in \text{Aut}_{\mathbf{Q}}(E)$ by $t_a(\bigotimes_{i \in \mathbf{Z}/m} e_i) = \bigotimes_{i \in \mathbf{Z}/m} e'_i$ where $e_i \in E_i$ and $e'_i = H_{i-1}^a(e_{i-1}) \in E_i$, $t_{c'}(\bigotimes_{i \in \mathbf{Z}/m} e_i) = \bigotimes_{i \in \mathbf{Z}/m} e''_i$ where $e_i \in E_i$ and $e''_i = b_{ik'}(e_{ik'}) \in E_{ik'k} = E_i$. From definitions we have $t_a(w(e)) = \rho_a(w)t_a(e)$, $t_{c'}(w(e)) = \rho_{c'}(w)t_{c'}(e)$ for any $w \in W, e \in E$.

We have $t_{c'} t_a t_{c'}^{-1} = t_a^k$. This follows from the identity

$$b_{hk'} H_{hk'-1}^a b_{hk'-1}^{-1} = H_{h-1}^a H_{h-2}^a \dots H_{h-k}^a : E_{h-k} \rightarrow E_h$$

for $h = 0, 1, \dots, m-1$ (here b_{-1} is taken to be b_{m-1}); an equivalent identity is

$$b_i H_{i-1}^a b_{i-1}^{-1} = H_{ik-1}^a H_{ik-2}^a \dots H_{ik-k}^a : E_{ik-k} \rightarrow E_{ik}$$

for $i = 0, 1, \dots, m-1$, which is the same as (d).

We have $t_a^M = 1$. This follows from the identity $H_{i+M-1}^a \dots H_{i+1}^a H_i^a = 1 : E_i \rightarrow E_i$ for $i = 0, 1, \dots, m-1$ which is equivalent to the known equality $s_0^{M/m} = 1$.

We show that

$$(e) \quad t_{c'}^N = t_a^{mf}.$$

This follows from the identity

$$b_{ik'} b_{ik'^2} \dots b_{ik'^N} = H_{i-1}^a H_{i-2}^a \dots H_{i-mf}^a \quad \text{for } i = 0, 1, \dots, m-1$$

or equivalently

$$b_{ik^{N-1}} b_{ik^{N-2}} \dots b_i = H_{i-1}^a H_{i-2}^a \dots H_{i-mf}^a \quad \text{for } i = 0, 1, \dots, m-1.$$

This is the same as

$$H_{ik^{N-1}}^a H_{ik^{N-2}}^a \dots H_1^a H_0^a s_1^N (H_{i-1}^a \dots H_1^a H_0^a)^{-1} = H_{i-1}^a H_{i-2}^a \dots H_{i-mf}^a$$

(with ik^N factors to the left of s_1^N). We have $ik^N - i = iMl$ for some $l \in \mathbf{N}$. From $s_0^{M/m} = 1$ we deduce $s_0^{iMl/m} = 1$ hence $H_{ik^{N-1}}^a H_{ik^{N-2}}^a \dots H_i^a = 1$ (with $ik^N - i$ factors). Since $s_1^N = s_0^f$, it remains to show that

$$H_{i-1}^a \dots H_1^a H_0^a s_0^f (H_{i-1}^a \dots H_1^a H_0^a)^{-1} = H_{i-1}^a H_{i-2}^a \dots H_{i-mf}^a.$$

If $f = 0$, this is obvious. Assume now that $f \geq 1$. Then $i - mf < 0$ and we see that it is enough to show $s_0^f (H_{i-1}^a \dots H_1^a H_0^a)^{-1} = H_{-1}^a H_{-2}^a \dots H_{i-mf}^a$, that is, $s_0^f = H_{-1}^a H_{-2}^a \dots H_{i-mf}^a (H_{i-1}^a \dots H_1^a H_0^a)$. This can be rewritten in the form $s_0^f =$

$H_{-1}^a H_{-2}^a \dots H_{i-mf}^a H_{i-mf-1}^a \dots H_{1-mf}^a H_{-mf}^a$ which follows from $s_0 = H_{m-1}^a \dots H_1^a H_0^a$. This proves (e).

We set $t_c = t_a^{-r} t_{c'} \in \text{Aut}_{\mathbf{Q}}(E)$. From definitions we have $t_c(w(e)) = \rho_c(w) t_c(e)$ for any $w \in W, e \in E$.

We have $t_c t_a t_c^{-1} = t_a^k$. This follows from the identity $t_{c'} t_a t_{c'}^{-1} = t_a^k$.

We have $t_c^N = t_a^u$. Indeed,

$$(t_a^{-r} t_{c'})^N = t_a^{-r(1+k+k^2+\dots+k^{N-1})} t_{c'}^N = t_a^{-mf+u} t_a^{mf} = t_a^u.$$

We see that $t_a, t_c \in \text{Aut}_{\mathbf{Q}}(E)$ satisfy the relations of Γ , hence they define a homomorphism $\Gamma \rightarrow \text{Aut}_{\mathbf{Q}}(E)$. This has the required properties.

Case 2. Assume that J is a single Γ -orbit. Let $\langle a \rangle$ be the subgroup of Γ generated by a . If X is an $\langle a \rangle$ -orbit in J , then cX is again an $\langle a \rangle$ -orbit. (We must show that, if $j \in X$ and $i \geq 1$, then $ca^i j, cj$ are in the same $\langle a \rangle$ -orbit. But $ca^i j = a^{ik} c j$.) Hence $c^h X$ is an $\langle a \rangle$ -orbit in J for any $h \in \mathbf{N}$. We can find an integer $z \geq 1$ such that $X, cX, \dots, c^{z-1}X$ are distinct and $c^z X = X$. (Clearly, z is a divisor of N .) Hence the notation $X_h = c^h X$ for $h \in \mathbf{Z}/z$ is meaningful. Now X_0, X_1, \dots, X_{z-1} are distinct. The union $\bigcup_{h \in [0, z-1]} X_h$ is c -stable and a -stable, hence it is equal to J (by our assumption on J). We see that $|J| = |X|z$. For $h \in \mathbf{Z}/z$ we set $W^h = \prod_{j \in X_h} W_j$, $E^h = \otimes_{j \in X_h} E_j$. We have naturally $W = \prod_{h \in [0, z-1]} W^h$, $E = \otimes_{h \in [0, z-1]} E^h$. Let Γ'' be the subgroup of Γ generated by a, c^z . In Γ'' we have

$$a^M = 1, (c^z)^{N/z} = a^u, c^z a c^{-z} = a^{k^z}.$$

Since $a(X_0) = X_0, c^z(X_0) = X_0$, the action of Γ on W restricts to an action of Γ'' on W^0 . Using case 1 for W^0, E^0, Γ'' instead of W, E, Γ we obtain a homomorphism $\Gamma'' \rightarrow \text{Aut}_{\mathbf{Q}}(E^0)$ such that, denoting by S_0, S_1 the images of a, c^z under this homomorphism, we have $S_0^M = 1, S_1^{N/z} = S_0^u, S_1 S_0 S_1^{-1} = S_0^{k^z}, S_0(w(e)) = \rho_a(w) S_0(e), S_1(w(e)) = \rho_{c^z}(w) S_1(e)$ for any $w \in W^0, e \in E^0$.

For $h \in \mathbf{Z}/z$ there exists a \mathbf{Q} -linear isomorphism $K_h : E^h \xrightarrow{\sim} E^{h+1}$ with $K_h(w e) = \rho_c(w) K_h(e)$ for all $e \in E^h, w \in W^h$ (note that $\rho_c(w) \in W^{h+1}$); moreover, K_h is unique up to multiplication by an element of \mathbf{Q}^* . (For example we can take K_h of the form $\otimes_{j \in X_h} H_j^c$ where $H_j^c : E_j \xrightarrow{\sim} E_{cj}$ are as in (a).) Then $S_2 := K_{z-1} \dots K_1 K_0 : E^0 \xrightarrow{\sim} E^0$ satisfies $S_2(w(e)) = \rho_{c^z}(w) S_2(e)$ for any $w \in W^0, e \in E^0$. Hence $S_2^{-1} S_1(w(e)) = w(S_2^{-1} S_1(e))$ for any $w \in W^0, e \in E^0$. By the absolute irreducibility of the W^0 -module E^0 we see that $S_2^{-1} S_1 : E^0 \rightarrow E^0$ is a \mathbf{Q}^* -multiple of the identity map. Hence, replacing K_0 by a \mathbf{Q}^* -multiple, we may assume that $S_2 = S_1$ that is, $K_{z-1} \dots K_1 K_0 = S_1$.

For any $h \in \mathbf{N}$ we define $\beta_{-h} \in \text{Aut}_{\mathbf{Q}}(E^{-h})$ by

$$\beta_{-h} = K_{-h}^{-1} \dots K_{-2}^{-1} K_{-1}^{-1} S_0^{k^h} K_{-1} K_{-2} \dots K_{-h}$$

(for $h = 0$ this is interpreted as $\beta_0 = S_0$). We show that $\beta_{-h-z} = \beta_{-h}$. We argue by induction on h . For $h = 0$ we must verify that

$$K_{-z}^{-1} \dots K_{-2}^{-1} K_{-1}^{-1} S_0^{k^z} K_{-1} K_{-2} \dots K_{-z} = S_0;$$

this follows from the known equality $S_1^{-1} S_0^{k^z} S_1 = S_0$. For $h \geq 1$ we have, using the induction hypothesis,

$$\beta_{-h-z} = K_{-h-z}^{-1} \beta_{-h-z+1} K_{-h-z} = K_{-h}^{-1} \beta_{-h+1} K_{-h} = \beta_{-z}.$$

We see that β_{-h} depends only on the image of h in \mathbf{Z}/z . Define $t_a, t_c \in \text{Aut}_{\mathbf{Q}}(E)$ by

$$\begin{aligned} \tau_a(\otimes_{h \in \mathbf{Z}/z} e_h) &= \otimes_{h \in \mathbf{Z}/z} e'_h \text{ where } e_h \in E^h \text{ and } e'_h = \beta_h(e_h) \in E^h, \\ \tau_c(\otimes_{h \in \mathbf{Z}/z} e_h) &= \otimes_{h \in \mathbf{Z}/z} e''_h \text{ where } e_h \in E^h \text{ and } e''_h = K_{h-1}(e_{h-1}) \in E^h. \end{aligned}$$

From definitions we have $t_a(w(e)) = \rho_a(w)t_a(e), t_c(w(e)) = \rho_c(w)t_c(e)$ for any $w \in W, e \in E$.

We have $t_c t_a t_c^{-1} = t_a^k$. This follows from the identity $\beta_{h-1} = K_{h-1}^{-1} \beta_h^k K_{h-1}$ for $h \in \mathbf{Z}/z$.

We have $t_a^M = 1$. This follows from the identity $\beta_{-h}^M = 1$ for any $h \in \mathbf{N}$. An equivalent statement is $K_{-h}^{-1} \dots K_{-2}^{-1} K_{-1}^{-1} S_0^{k^h M} K_{-1} K_{-2} \dots K_{-h} = 1$ which follows from $S_0^M = 1$.

We show that $t_c^N = t_a^u$. It is enough to show that $K_{-h-1} K_{-h-2} \dots K_{-h-N} = b_{-h}^u : E_{-h} \rightarrow E_{-h}$ for $h = 0, 1, \dots, z-1$. An equivalent statement is

$$(K_{-h}^{-1} \dots K_{-1}^{-1})(K_{-1} K_{-2} \dots K_{-N})(K_{-1} \dots K_{-h}) = K_{-h}^{-1} \dots K_{-1}^{-1} S_0^{k^h u} K_{-1} \dots K_{-h}.$$

The left-hand side is $(K_{-h}^{-1} \dots K_{-1}^{-1}) S_1^{N/z} (K_{-1} \dots K_{-h})$. It is enough to show that $S_1^{N/z} = S_0^{k^h u}$ or that $S_0^u = S_0^{k^h u}$ for $h = 0, 1, \dots, z-1$. Since $S_0^M = 1$, it is enough to show that $u = k^h u \pmod M$. This follows from $uk = u \pmod M$.

We see that $t_a, t_c \in \text{Aut}_{\mathbf{Q}}(E)$ satisfy the relations of Γ , hence they define a homomorphism $\Gamma \rightarrow \text{Aut}_{\mathbf{Q}}(E)$. This has the required properties.

Case 3. We now consider the general case. For any Γ -orbit Y on J we set $W^Y = \prod_{j \in Y} W_j, E^Y = \otimes_{j \in Y} E_j$. We have naturally $W = \prod_Y W^Y, E = \otimes_Y E^Y$ where Y runs over the Γ -orbits in J . Now the Γ -action on W restricts to a Γ -action on W^Y for each Y . Using case 2 for W^Y, E^Y, Γ instead of W, E, Γ we obtain for each Y a homomorphism $\Gamma \rightarrow \text{Aut}_{\mathbf{Q}}(E^Y)$. We define a homomorphism $\Gamma \rightarrow \text{Aut}_{\mathbf{Q}}(E)$ by $\gamma : \otimes_Y e_Y \mapsto \otimes_Y (\gamma(e_Y))$; here $e_Y \in E^Y$. This has the required properties.

This completes the proof in the setup of (ii). The lemma is proved.

39.3. Let Γ be a finite group. Assume that Γ is a semidirect product of a normal subgroup Γ' with a cyclic group of order n with generator b . Let $\bar{\mathfrak{U}}$ be an algebraic closed field of characteristic 0. Let E be a simple $\bar{\mathfrak{U}}[\Gamma]$ -module which is isotypical as a $\bar{\mathfrak{U}}[\Gamma']$ -module. We show that (a) E is simple as a $\bar{\mathfrak{U}}[\Gamma']$ -module.

There exists a simple $\bar{\mathfrak{U}}[\Gamma']$ -module E' such that, setting $V = \text{Hom}_{\bar{\mathfrak{U}}[\Gamma']}(E', E)$ we have $V \otimes E' \xrightarrow{\sim} E, f \otimes e' \mapsto f(e')$. For any $\gamma' \in \Gamma'$ we have $\text{tr}(b\gamma'b^{-1}, E') = \text{tr}(\gamma', E')$. Hence there exists $\xi \in \text{Aut}_{\bar{\mathfrak{U}}}(E')$ such that $\xi(\gamma'e') = (b\gamma'b^{-1})\xi(e')$ for $e' \in E', w' \in \Gamma'$. Then $\xi^n \in \text{Aut}_{\bar{\mathfrak{U}}}(E')$ commutes with the Γ' -action hence it is a scalar on E' . Replacing ξ by an $\bar{\mathfrak{U}}^*$ -multiple we can assume that $\xi^n = 1$. Then E' becomes a $\bar{\mathfrak{U}}[\Gamma]$ -module with b acting as ξ . Define $\eta : V \rightarrow V$ by $f \mapsto \eta(f)$ where $\eta(f)(e') = bf(\xi^{-1}(e'))$ for $e' \in E'$. We regard V as a $\bar{\mathfrak{U}}[\Gamma]$ -module in which Γ' acts trivially and b acts as η . Then the isomorphism $V \otimes E' \xrightarrow{\sim} E$ considered above is an isomorphism of $\bar{\mathfrak{U}}[\Gamma]$ -modules. Since E is simple it follows that V is a simple $\bar{\mathfrak{U}}[\Gamma]$ -module. Since Γ acts on V through a cyclic quotient, we see that $\dim V = 1$. It follows that $E \cong E'$ as a $\bar{\mathfrak{U}}[\Gamma']$ -module; (a) follows.

39.4. Let W, I, \mathcal{X} be as in 39.1. Let Γ be a finite group with generators a, c and relations $a^M = 1, c^N = 1, cac^{-1} = a^k$ where M, N, k are integers ≥ 1 such that $k^N = 1 \pmod M$. Let $\langle a \rangle$ (resp. $\langle c \rangle$) be the subgroup of Γ generated by a (resp.

c). Let $\gamma \mapsto \rho_\gamma$ be a homomorphism $\Gamma \rightarrow \text{Aut}(W, I, \mathcal{X})$. We form the semidirect product $W\Gamma, W\langle a \rangle$ with W normal. Note that $W\langle a \rangle$ is a subgroup of $W\Gamma$.

Lemma 39.5. *In the setup of 39.4 let E be a simple $\bar{\mathfrak{U}}[W\Gamma]$ -module. Assume that either $cxc^{-1} = x$ for any $x \in \langle a \rangle$, or $cxc^{-1} = x^{-1}$ for any $x \in \langle a \rangle$. Let $x \in \langle a \rangle$. There exists ζ , a root of 1 in $\bar{\mathfrak{U}}$, such that $\text{tr}(wxc, E) \in \zeta\mathbf{Z}$ for any $w \in W$.*

Let $W' = W\langle a \rangle$. We can write canonically $E = \bigoplus_{t \in T} E^t$ where E^t are isotypical $\bar{\mathfrak{U}}[W']$ -modules. Now Γ acts on T by $\gamma E^t = E_{\gamma(t)}$. This action is transitive since E is simple as a $\bar{\mathfrak{U}}[W\Gamma]$ -module. If $c(t) \neq t$ for any $t \in T$, then for any $w \in W$, $wxc : E \rightarrow E$ permutes the summands E^t and no summand is stable, hence $\text{tr}(wxc, E) = 0$. In this case the lemma is clear. Thus we may assume that $c(t) = t$ for some $t \in T$. Then $cE^t = E^t$, hence E^t is a $\bar{\mathfrak{U}}[W\Gamma]$ -submodule of E . Since E is simple, we have $E = E^t$. Thus, E is isotypical as a $\bar{\mathfrak{U}}[W']$ -module. Using 39.3(a) for $W\Gamma, W', \langle c \rangle$ instead of Γ, Γ', C we see that E is simple as a $\bar{\mathfrak{U}}[W']$ -module.

We can write canonically $E = \bigoplus_{h \in H} E_h$ where E_h are isotypical $\bar{\mathfrak{U}}[W]$ -modules. Now Γ acts on H by $\gamma E_h = E_{\gamma(h)}$. This action is transitive since E is simple as a $\bar{\mathfrak{U}}[W\Gamma]$ -module. The restriction of this action to $\langle a \rangle$ is also transitive since E is simple as a $\bar{\mathfrak{U}}[W']$ -module. For $h \in H$ let $\langle a \rangle_h$ be the stabilizer of h in $\langle a \rangle$. Let $W'_h = W\langle a \rangle_h \subset W'$. Then E_h is a W'_h -submodule of E and the W' -module E is induced by the W'_h -module E_h . Since E is simple as a W' -module, it follows that E_h is simple as a W'_h -module. Using 39.3(a) for W'_h, W, E_h instead of Γ, Γ', E , we see that E_h is simple as a W -module.

For $h \in H$ let $\Gamma_h = \{\gamma \in \Gamma; \gamma(h) = h\}$. Then E_h is a $W\Gamma_h$ -submodule of E . Note that Γ_h is an extension of a cyclic group (the image of Γ_h under $\Gamma \rightarrow \Gamma/\langle a \rangle$) by a cyclic group (the intersection $\Gamma_h \cap \langle a \rangle$). Let $H_0 = \{h \in H; xc(h) = h\}$. For any $w \in W$ we have $\text{tr}(wxc, E) = \sum_{h \in H_0} \text{tr}(wxc, E_h)$. In particular, if $H_0 = \emptyset$, then $\text{tr}(wxc, E) = 0$ so that the lemma is clear in this case. Thus we may assume that $H_0 \neq \emptyset$. Let $h \in H_0$. We can find a simple $\mathbf{Q}[W]$ -submodule E_h^0 of E_h such that $E_h = \bar{\mathfrak{U}} \otimes_{\mathbf{Q}} E_h^0$ as $\bar{\mathfrak{U}}[W]$ -modules. By Lemma 39.2 applied to W, Γ_h, E_h^0 instead of W, Γ, E we see that the $\mathbf{Q}[W]$ -structure on E_h^0 extends to a $\mathbf{Q}[W\Gamma_h]$ -module structure. (The hypotheses of that lemma are satisfied since the $\bar{\mathfrak{U}}[W]$ -module structure on E_h extends to a $\bar{\mathfrak{U}}$ -module structure.) For $\gamma \in \Gamma_h$ let $t_\gamma \in \text{Aut}_{\mathbf{Q}}(E_h^0)$ be the action of γ in this $\mathbf{Q}[W\Gamma_h]$ -module. By extension of scalars, t_γ defines an element $\tilde{t}_\gamma \in \text{Aut}_{\mathbf{Q}}(E_h)$. For $e \in E_h, w \in W$ we have $\tilde{t}_\gamma(we) = (\gamma w \gamma^{-1})(\tilde{t}_\gamma(e))$. Hence $\gamma^{-1}(\tilde{t}_\gamma(we)) = w \gamma^{-1} \tilde{t}_\gamma(e)$. Thus $\gamma^{-1} \tilde{t}_\gamma : E_h \rightarrow E_h$ commutes with the action of W . Since E_h is simple we see that $\gamma^{-1} \tilde{t}_\gamma : E_h \rightarrow E_h$ is a scalar $\lambda_h(\gamma) \in \bar{\mathfrak{U}}^*$. Thus $\tilde{t}_\gamma = \lambda_h(\gamma) \gamma : E_h \rightarrow E_h$. Clearly, $\gamma \mapsto \lambda_h(\gamma)$ is a homomorphism $\Gamma_h \rightarrow \bar{\mathfrak{U}}^*$. For $w \in W$ we have $wxc = \lambda_h(xc)^{-1} w \tilde{t}_{xc} : E_h \rightarrow E_h$, hence

$$\text{tr}(wxc, E_h) = \lambda_h(xc)^{-1} \text{tr}(w \tilde{t}_{xc}, E_h) = \lambda_{xc}^{-1} \text{tr}(w t_{xc}, E_h^0).$$

Note that $\text{tr}(w t_{xc}, E_h^0) \in \mathbf{Z}$ since E_h^0 is a $\mathbf{Q}[W\Gamma_h]$ -module. We deduce

$$\text{tr}(wxc, E) = \sum_{h \in H_0} \lambda_h(xc)^{-1} \text{tr}(w t_{xc}, E_h^0).$$

Let $H_1 = \{h \in H_0; \text{tr}(w' t_{xc}, E_h^0) \neq 0 \text{ for some } w' \in W\}$. Then, clearly,

$$\text{tr}(wxc, E) = \sum_{h \in H_1} \lambda_h(xc)^{-1} \text{tr}(w t_{xc}, E_h^0).$$

Since $\lambda_h(xc)$ is a root of 1, it is enough to verify the following statement: If $h, h' \in H_1$, then $\lambda_h(xc) = \pm\lambda_{h'}(xc)$. Since $\langle a \rangle$ acts transitively on H , we can find $y \in \langle a \rangle$ such that $h = yh'$. We have $yE_{h'} = E_h, y\Gamma_{h'}y^{-1} = \Gamma_h$. Also, for any $w' \in W$ we have

$$\text{tr}(w'xc, E_{h'}) = \text{tr}(yw'xcy^{-1}, E_h) = \text{tr}(yw'y^{-1}w'xcy^{-1}, E_h),$$

hence

$$\lambda_{h'}(xc)\text{tr}(w'xc, E_{h'}^0) = \lambda_h(yxcy^{-1})\text{tr}(yw'y^{-1}w'xcy^{-1}, E_h^0).$$

We take here $w' \in W$ such that $\text{tr}(w'xc, E_{h'}^0) \neq 0$. Dividing by $\text{tr}(w'xc, E_{h'}^0)$ we deduce $\lambda_{h'}(xc) \in \mathbf{Q}\lambda_h(yxcy^{-1})$. Since $\lambda_{h'}(xc), \lambda_h(yxcy^{-1})$ are roots of 1, it follows that $\lambda_{h'}(xc) = \pm\lambda_h(yxcy^{-1})$. Thus it is enough to show that $\lambda_h(xc) = \pm\lambda_h(yxcy^{-1})$ or that $\lambda_h(c^{-1}x^{-1}yxcy^{-1}) = \pm 1$ or that $\lambda_h(c^{-1}ycy^{-1}) = \pm 1$. More generally, we will verify the following statement: If $h \in H_0, y \in \langle a \rangle, c^{-1}ycy^{-1} \in \langle a \rangle_h$, then $\lambda_h(c^{-1}ycy^{-1}) = \pm 1$. If $cy = yc = 1$, this is obvious. Therefore, we may assume that $cy' = y'^{-1}c$ for any $y' \in \langle a \rangle$.

Let $u \in \langle a \rangle_h$. Since u, xc belong to Γ_h we have $\lambda_h((xc)^{-1}uxcu^{-1}) = 1$, that is, $\lambda_h(c^{-1}ucu^{-1}) = 1$ (we use $ux = xu$). But $c^{-1}uc = u^{-1}$. Hence $\lambda_h(u^{-2}) = 1$, that is, $\lambda_h(u)^{-2} = 1$ and $\lambda_h(u) = \pm 1$. We apply this with $u = c^{-1}y_1cy_1^{-1} \in \langle a \rangle_h$. We see that $\lambda_h(c^{-1}y_1cy_1^{-1}) = \pm 1$. The lemma is proved.

39.6. Let W, I, \mathcal{X} are as in 39.1. We assume that (W, I) is irreducible. Let $\tilde{I} = I \sqcup \{\omega\}$ where ω is a symbol. We define a map $\pi : \tilde{I} \rightarrow W$ as follows: $\pi(s) = s$ if $s \in I$ and $\pi(\omega)$ is the unique reflection of maximal length in \mathcal{X} . The restriction of π to \tilde{I} is injective if $|I| \geq 2$; if $|I| = 1$, it maps both elements of \tilde{I} to the unique element of I . Let Ω be the group of all permutations $\sigma : \tilde{I} \xrightarrow{\sim} \tilde{I}$ such that there exist $w \in W$ with $w\pi(\sigma)w^{-1} = \pi(\sigma(x))$ for all $x \in \tilde{I}$. Then $\sigma \mapsto [w_1 \mapsto ww_1w^{-1}]$ is a homomorphism of Ω into $\text{Inn}(W)$, the group of inner automorphisms of W . Let K be a subset of \tilde{I} such that $K \neq \tilde{I}$. Then π restricts to a bijection $K \xrightarrow{\sim} \pi(K)$. Let $\Omega^K = \{\sigma \in \Omega; \sigma(K) = K\}$. Let $W^{(K)}$ be the subgroup of W generated by $\pi(K)$. From the theory of affine Weyl groups we see that $W^{(K)}$ is a (finite) Coxeter group on the generators $\pi(K)$. We have canonically $W^{(K)} = \prod_{z \in Z} W_z^{(K)}$ where $W_z^{(K)}$ is an irreducible Weyl group with set of simple reflections $K_z = K \cap W_z^{(K)}$. For $z \in Z$ we set $\mathcal{X}_z^K = R(W_z^{(K)})$ if $R(W_z^{(K)})$ is a single $W_z^{(K)}$ -conjugacy class and $\mathcal{X}_z^K = \mathcal{X} \cap W_z^{(K)}$ if $R(W_z^{(K)})$ is a union of two $W_z^{(K)}$ -conjugacy classes; in any case, \mathcal{X}_z^K is a single $W_z^{(K)}$ -conjugacy class in $R(W_z^{(K)})$. (We use the following fact: If $s \in I$ is such that $s\pi(\omega)$ has order ≥ 4 , then $s, \pi(\omega)$ are not conjugate under W .) Then $\mathcal{X}^K = \bigsqcup_{z \in Z} \mathcal{X}_z^K$ is a special subset of $R(W^{(K)})$ (see 39.1). Now the image of Ω^K under $\Omega \rightarrow \text{Inn}(W)$ (as above) is contained in the group $\text{Aut}(W^{(K)}, K, \mathcal{X}^K)$ of automorphisms of $W^{(K)}$ which preserve K and \mathcal{X}^K . Thus we have a homomorphism $\Omega^K \rightarrow \text{Aut}(W^{(K)}, K, \mathcal{X}^K)$. Restricting this to a subgroup C of Ω^K we obtain a homomorphism $C \rightarrow \text{Aut}(W^{(K)}, K, \mathcal{X}^K)$. Let $c \in \text{Aut}(W, I, \mathcal{X})$ be such that $c(K) = K$. We have $c^N = 1$ for some $N \geq 1$. We extend the bijection $c : I \xrightarrow{\sim} I$ (restriction of $c : W \rightarrow W$) to a bijection $c : \tilde{I} \xrightarrow{\sim} \tilde{I}$ by $c(\omega) = \omega$. We have $c(W^{(K)}) = W^{(K)}, c(\mathcal{X}^K) = \mathcal{X}^K$, hence c restricts to an element of $\text{Aut}(W^{(K)}, K, \mathcal{X}^K)$. For any $\sigma \in \Omega$ we define an element $c(\sigma) \in \Omega$ by $c(\sigma)(x) = \sigma(c^{-1}x)$ for $x \in \tilde{I}$. Then $c : \Omega \xrightarrow{\sim} \Omega$ preserves Ω^K . Assume that $c(C) = C$. For $\sigma \in C, w \in W^{(K)}$ we have $c(\sigma(w)) = c(\sigma)(c(w))$. Let $\langle c \rangle$ be a cyclic group of order N with generator c . On the set $W^{(K)} \times C \times \langle c \rangle$ we have

a group structure $(w, \sigma, c^n)(w', \sigma', c^{n'}) = (w\sigma(c^n(w')), \sigma c^n(\sigma'), c^{n+n'})$. This is the semidirect product $W^{(K)}\Gamma$ of $W^{(K)}$ with the group $\Gamma = C \times \langle c \rangle$ with group structure $(\sigma, c^n)(\sigma', c^{n'}) = (\sigma c^n(\sigma'), c^{n+n'})$.

Proposition 39.7. *In the setup of 39.6 let E be a simple $\bar{\mathfrak{U}}[W^{(K)}\Gamma]$ -module. Let $\gamma \in \Gamma$. There exists ζ , a root of 1 in $\bar{\mathfrak{U}}$, such that $\text{tr}(w\gamma, E) \in \zeta\mathbf{Z}$ for any $w \in W^{(K)}$.*

From the theory of affine Weyl groups it is known that one of the following holds:

- (i) Ω is cyclic and $c(\sigma) = \sigma$ for all $\sigma \in \Omega$,
- (ii) Ω is cyclic and $c(\sigma) = \sigma^{-1}$ for all $\sigma \in \Omega$,
- (iii) $\Omega \cong \mathbf{Z}/2 \times \mathbf{Z}/2$.

Moreover, $C \subset \Omega$ (compatibly with the action of c). In cases (i),(ii), C is cyclic and the assumptions of Lemma 39.5 are satisfied (with $W^{(K)}$ instead of W). Hence the result follows from 39.5.

In the remainder of the proof we assume that we are in case (iii). If C is cyclic, then $c(\sigma) = \sigma = \sigma^{-1}$ for all $\sigma \in C$. The result follows again from Lemma 39.5. Hence we may assume that C is not cyclic so that $C = \Omega$. Let n be the order of $c : C \rightarrow C$. Then $n \in \{1, 2, 3\}$ and n divides N . Note that $c^{N/n}$ is in the center of Γ . Tensoring E by a suitable one-dimensional representation (which is trivial on $W^{(K)}C$) we may assume that $c^{N/n}$ acts trivially on E , hence E factors through the quotient of $W^{(K)}\Gamma$ by the subgroup generated by $c^{N/n}$. Hence we may assume that $N = n$. If $n = 1$, then $\Gamma \cong \mathbf{Z}/2 \times \mathbf{Z}/2$ is a group as in 39.5 and the result follows from 39.5. If $n = 2$, then Γ is a dihedral group of order 8 which is again a group as in 39.5 (an extension of $\mathbf{Z}/2$ by $\mathbf{Z}/4$) and the result follows from 39.5. If $n = 3$ then W must be of type D_4 , $W^{(K)}$ is an elementary abelian 2-group, E is one-dimensional and there exists a homomorphism $\mu : W^{(K)}\Gamma \rightarrow \bar{\mathfrak{U}}^*$ such that $\text{tr}(w\gamma', E) = \mu(w\gamma')$ for all $w \in W^{(K)}$, $\gamma' \in \Gamma$. Then $\zeta = \mu(\gamma)$ is a root of 1. For any $w \in W^{(K)}$ we have $\mu(w\gamma) = \zeta\mu(w)$ and $\mu(w) = \pm 1$, since $w^2 = 1$. Thus, $\text{tr}(w\gamma, E) = \pm\zeta$. The proposition is proved.

Corollary 39.8. *Assume that G, D, A, u are as in 35.22 and that G^0/Z_{G^0} is simple. Then $b_{A,u}^v \in \eta\mathbf{Q}$ for some η , a root of 1 (with $b_{A,u}^v$ as in 34.19.)*

This follows from 35.22 and 39.7.

REFERENCES

- [Al] D. Alvis, *The duality operation in the character ring of a finite Chevalley group*, Bull. Amer. Math. Soc. **1** (1979), 907-911. MR0546315 (81e:20012)
- [BBD] A. Beilinson, J. Bernstein, P. Deligne, *Faisceaux pervers*, Astérisque **100** (1982). MR0751966 (86g:32015)
- [Cu] C.W. Curtis, *Truncation and duality in the character ring of a finite group of Lie type*, J. Algebra **62** (1980), 320-332. MR0563231 (81e:20011)
- [De] P. Deligne, *La conjecture de Weil, II*, Publ. Math. I.H.E.S. **52** (1980), 137-252. MR0601520 (83c:14017)
- [Ka] N. Kawanaka, *Fourier transforms of nilpotently supported invariant functions on a simple Lie algebra over a finite field*, Invent. Math. **69** (1982), 411-435. MR0679766 (84c:20053)
- [L14] G. Lusztig, *Characters of reductive groups over a finite field*, Ann. Math. Studies 107, Princeton Univ. Press, 1984. MR0742472 (86j:20038)
- [L3] G. Lusztig, *Character sheaves, I*, Adv. Math. **56** (1985), 193-237; II, vol. 57, 1985, pp. 226-265; III, vol. 57, 1985, pp. 266-315; IV, vol. 59, 1986, pp. 1-63; V, vol. 61, 1986, pp. 103-155. MR0792706 (87b:20055); MR0806210 (87m:20118a); MR0825086 (87m:20118b); MR0849848 (87m:20118c)

- [L9] G. Lusztig, *Character sheaves on disconnected groups, I*, Represent. Theory (electronic) **7** (2003), 374-403; II, vol. 8, 2004, pp. 72-124; III, vol. 8, 2004, pp. 125-144; IV, vol. 8, 2004, pp. 145-178; Errata , vol. 8, 2004, pp. 179-179; V, vol. 8, 2004, pp. 346-376; VI, vol. 8, 2004, pp. 377-413; VII, vol. 9, 2005, pp. 209-266. MR2017063 (2006d:20090a); MR2048588 (2006d:20090b); MR2048589 (2006d:20090c); MR2048590 (2006d:20090d); MR2058725; MR2077486 (2005h:20111); MR2084488 (2005h:20112); MR2133758 (2006e:20089)
- [L10] G. Lusztig, *Parabolic character sheaves, I*, Moscow Math. J. **4** (2004), 153-179. MR2074987 (2006d:20091a)

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE,
MASSACHUSETTS 02139