

CHARACTER SHEAVES ON DISCONNECTED GROUPS, IX

G. LUSZTIG

ABSTRACT. We associate a two-sided cell to any (parabolic) character sheaf. We study the interaction between the duality operator for character sheaves and the operation of “twisted induction”.

INTRODUCTION

Throughout this paper, G denotes a fixed, not necessarily connected, reductive algebraic group over an algebraically closed field \mathbf{k} . This paper is a part of a series [L9] which attempts to develop a theory of character sheaves on G .

One of the main constructions in [L3] (going back to [L14]) was a procedure which to any character sheaf on G^0 associates a certain two-sided cell in an (extended) Coxeter group. A variant of this construction (restricted to “unipotent” character sheaves) was later given by Grojnowski [Gr]. Here we give a construction which generalizes that in [L3] (and takes into account the approach in [Gr]) which to any (parabolic) character sheaf on $Z_{J,D}$ associates a certain type of two-sided cell.

The paper is organized as follows. In Section 40 we study certain equivariant sheaves on $G^0/U^* \times G^0/U^*$ (where U^* is the unipotent radical of a Borel in G^0) under the convolution operation. Some results in this section are implicit in [L14, Ch.1]. In Section 41 we study the character sheaves on $Z_{\emptyset,D}$ (where D is a connected component of G) by connecting them with sheaves on $G^0/U^* \times G^0/U^*$. We use this study to attach a two-sided cell to any character sheaf on $Z_{J,D}$. (See 41.4.) In Section 42 we study the interaction between the duality operation \mathbf{d} (see 38.10, 38.11) and the functor $f_{\emptyset,\mathbf{I}}$ (see 36.4). The main result in this section is Proposition 42.9 which contains [L3, III, Cor. 15.8(b)] as a special case (with $G = G^0, v = 1$).

Notation. We fix a 1-dimensional $\bar{\mathbf{Q}}_l$ -vector space V with a given isomorphism $V^{\otimes 2} \xrightarrow{\sim} \bar{\mathbf{Q}}_l(1)$ (Tate twist of $\bar{\mathbf{Q}}_l$). For $n \in \mathbf{N}$ we set $\bar{\mathbf{Q}}_l(n/2) = V^{\otimes n}$. For $n \in \mathbf{Z}, n < 0$ let $\bar{\mathbf{Q}}_l(n/2)$ be the dual space of $\bar{\mathbf{Q}}_l(-n/2)$. If X is an algebraic variety and $A \in \mathcal{D}(X), n \in \mathbf{Z}$, we write $A(n/2)$ instead of $A \otimes \bar{\mathbf{Q}}_l(n/2)$ and $A[[n/2]]$ instead of $A[n](n/2)$. (When n is even, this agrees with the notation in [L9, II, p. 73].)

CONTENTS

- 40. Sheaves on $G^0/U^* \times G^0/U^*$.
- 41. Character sheaves and two-sided cells.
- 42. Duality and the functor $f_{\emptyset,\mathbf{I}}$.

Received by the editors February 8, 2006.
2000 *Mathematics Subject Classification.* Primary 20G99.
Supported in part by the National Science Foundation.

©2006 American Mathematical Society
Reverts to public domain 28 years from publication

40. SHEAVES ON $G^0/U^* \times G^0/U^*$

40.1. Let $\mathcal{A} = \mathbf{Z}[v, v^{-1}]$. Let \hat{H} (resp. H) be the \mathcal{A} -module consisting of all formal (resp. finite) linear combinations $\sum_{w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}} a_{w,\lambda} \tilde{T}_w 1_\lambda$ with $a_{w,\lambda} \in \mathcal{A}$. Note that H is naturally an \mathcal{A} -submodule of \hat{H} with \mathcal{A} -basis $\{\tilde{T}_w 1_\lambda; w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}\}$. For any $n \in \mathbf{N}_k^*$, the \mathcal{A} -submodule of H spanned by $\{\tilde{T}_w 1_\lambda; w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}_n\}$ may be naturally identified with H_n (see 31.2(a)). There is a unique \mathcal{A} -algebra structure on \hat{H} in which the product of two elements

$$h = \sum_{w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}} a_{w,\lambda} \tilde{T}_w 1_\lambda, \quad h' = \sum_{w' \in \mathbf{W}, \lambda' \in \underline{\mathfrak{s}}} a'_{w',\lambda'} \tilde{T}_{w'} 1_{\lambda'}$$

as above is $hh' = \sum_{y \in \mathbf{W}, \nu \in \underline{\mathfrak{s}}} b_{y,\nu} \tilde{T}_y 1_\nu$ where for any $\nu \in \underline{\mathfrak{s}}$,

$$\sum_{w, w' \in \mathbf{W}} a_{w, w'^{-1}\nu} a'_{w', \nu} \tilde{T}_w \tilde{T}_{w'} 1_\nu = \sum_{y \in \mathbf{W}} b_{y,\nu} \tilde{T}_y 1_\nu$$

is computed in the algebra structure of H_n for any n such that $\nu \in \underline{\mathfrak{s}}_n$. Thus \hat{H} becomes an associative algebra with 1; H is a subalgebra (without 1) and, for $n \in \mathbf{N}_k^*$, H_n is a subalgebra (with a different 1) with the algebra structure as in 31.2.

Now in the definition of \hat{H} given above, although $\tilde{T}_w 1_\lambda$ is defined, the elements $\tilde{T}_w, 1_\lambda$ are not defined separately (as was the case in H_n). To remedy this we set $\tilde{T}_w = \sum_{\lambda \in \underline{\mathfrak{s}}} \tilde{T}_w 1_\lambda \in \hat{H}$ (for $w \in \mathbf{W}$) and $1_\lambda = \tilde{T}_1 1_\lambda \in H$ (for $\lambda \in \underline{\mathfrak{s}}$). Then $\tilde{T}_w 1_\lambda$ is the product of $\tilde{T}_w, 1_\lambda$ in the algebra \hat{H} . Note that \tilde{T}_1 is the unit element of \hat{H} and the following equalities hold in \hat{H} :

$$\begin{aligned} 1_\lambda 1_{\lambda'} &= 1_\lambda \text{ for } \lambda \in \underline{\mathfrak{s}}, 1_\lambda 1_{\lambda'} = 0 \text{ for } \lambda \neq \lambda' \text{ in } \underline{\mathfrak{s}}; \\ \tilde{T}_w \tilde{T}_{w'} &= \tilde{T}_{ww'} \text{ for } w, w' \in \mathbf{W} \text{ such that } l(ww') = l(w) + l(w'); \\ \tilde{T}_w 1_\lambda &= 1_{w\lambda} \tilde{T}_w \text{ for } w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}; \\ \tilde{T}_s^2 &= \tilde{T}_1 + (v - v^{-1}) \sum_{\lambda \in \underline{\mathfrak{s}}; s \in \mathbf{W}_\lambda} \tilde{T}_s 1_\lambda \text{ for } s \in \mathbf{I}. \end{aligned}$$

By a standard argument we see that

(a) H is exactly the \mathcal{A} -algebra defined by the generators $\tilde{T}_w 1_l$ ($w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}$) and the relations:

$$\begin{aligned} (\tilde{T}_w 1_\lambda)(\tilde{T}_{w'} 1_{\lambda'}) &= 0 \text{ if } w, w' \in \mathbf{W}, \lambda, \lambda' \in \underline{\mathfrak{s}}, w'\lambda' \neq \lambda, \\ (\tilde{T}_w 1_{w'\lambda'})(\tilde{T}_{w'} 1_{\lambda'}) &= \tilde{T}_{ww'} 1_{\lambda'} \text{ if } w, w' \in \mathbf{W}, \lambda, \lambda' \in \underline{\mathfrak{s}}, l(ww') = l(w) + l(w'), \\ (\tilde{T}_s 1_{s\lambda'})(\tilde{T}_s 1_{\lambda'}) &= \tilde{T}_1 1_{\lambda'} + (v - v^{-1})c \tilde{T}_s 1_{\lambda'} \text{ if } s \in \mathbf{I}, \lambda' \in \underline{\mathfrak{s}} \text{ where } c = 1 \text{ for } s \in \mathbf{W}_{\lambda'} \\ &\text{and } c = 0 \text{ for } s \notin \mathbf{W}_{\lambda'}. \end{aligned}$$

40.2. Let R, R^+ be as in 28.3. The following result is well known:

(a) *If $w \in \mathbf{W}, \alpha \in R^+$ and s_α is as in 28.3, then we have $l(ws_\alpha) > l(w)$ if and only if $w(\alpha) \in R^+$.*

Let $\lambda \in \underline{\mathfrak{s}}$. Let $R_\lambda, R_\lambda^+, \mathbf{W}_\lambda, H_\lambda$ be as in 34.2. We write \vee_λ instead of \vee_λ^D (as in 34.4 with $D = G^0$). We show that

(b) *if $w \in \mathbf{W}$, then $w\mathbf{W}_\lambda$ contains a unique element w_1 of minimal length; it is characterized by the property $w_1(R_\lambda^+) \subset R^+$.*

Let w_1 be an element of minimal length in $w\mathbf{W}_\lambda$. Let $\alpha \in R_\lambda^+$. Then $l(w_1 s_\alpha) \geq l(w_1)$. Since $l(w_1 s_\alpha) = l(w_1) + 1 \pmod 2$ we see that $l(w_1 s_\alpha) > l(w_1)$. By (a) we have $w_1(\alpha) \in R^+$. Thus, $w_1(R_\lambda^+) \subset R^+$. Now let $u \in \mathbf{W}_\lambda - \{1\}$. We pick $\alpha \in R_\lambda^+$ such that $u(\alpha)^{-1} \in R_\lambda^+$; then $w_1 u(\alpha)^{-1} \in R^+$. If $w_1 u$ has minimal length in $w\mathbf{W}_\lambda$,

then by an earlier part of the argument applied to w_1u instead of w_1 we have $w_1u(\alpha) \in R^+$, a contradiction. We see that w_1 is the unique element of minimal length in $w\mathbf{W}_\lambda$. It remains to show that if $u \in \mathbf{W}_\lambda$ satisfies $w_1u(R_\lambda^+) \subset R^+$, then $u = 1$. If $u \neq 1$, then by an earlier part of the argument we have $w_1u(\alpha)^{-1} \in R^+$ for some $\alpha \in R_\lambda^+$, a contradiction. This proves (b).

We show that

(c) if $s \in \mathbf{I}$ and $w \in \mathbf{W}$ has minimal length in $w\mathbf{W}_\lambda$, then either (i) sw has minimal length in $sw\mathbf{W}_\lambda$ or (ii) $w^{-1}sw \in \mathbf{W}_\lambda$.

There is a unique $\beta \in R^+$ such that $s(\beta)^{-1} \in R^+$. Assume that (i) does not hold. By (b) there exists $\alpha \in R_\lambda^+$ such that $sw(\alpha)^{-1} \in R^+$; moreover, $w(\alpha) \in R^+$. Hence $w(\alpha) = \beta$. We have $w^{-1}(\beta) = \alpha \in R_\lambda$, hence $w^{-1}sw \in \mathbf{W}_\lambda$ and (ii) holds. This proves (c).

For $z \in \mathbf{W}_\lambda$ let $\tilde{T}_z^\lambda, c_z^\lambda \in H_\lambda$ be as in 34.2. Then $c_z^\lambda = \sum_{z' \in \mathbf{W}_\lambda} p_{z',z}^\lambda \tilde{T}_{z'}^\lambda$ where $p_{z',z}^\lambda \in \mathbf{Z}[v^{-1}]$ are uniquely defined.

For any $w \in \mathbf{W}$, $\lambda \in \underline{\mathfrak{g}}$ there is a unique element of H which is equal to $c_{w,\lambda} \in H_n$ (see 34.4) for any n such that $\lambda \in \underline{\mathfrak{g}}_n$; we denote this element again by $c_{w,\lambda}$. We have $c_{w,\lambda} = \sum_{w' \in \mathbf{W}} \pi_{w',w,\lambda} \tilde{T}_{w'}^\lambda$ where $\pi_{w',w,\lambda} \in \mathbf{Z}[v^{-1}]$ are uniquely defined. Note that $\{c_{w,\lambda}; w \in \mathbf{W}, \lambda \in \underline{\mathfrak{g}}\}$ is an \mathcal{A} -basis of H .

Now:

(d) Let $w, w' \in \mathbf{W}$. We write $w = w_1z, w' = w'_1z'$ where w_1 has minimal length in $w\mathbf{W}_\lambda$, w'_1 has minimal length in $w'\mathbf{W}_\lambda$ and $z, z' \in \mathbf{W}_\lambda$. If $w_1 \neq w'_1$, then $\pi_{w',w,\lambda} = 0$. If $w_1 = w'_1$, then $\pi_{w',w,\lambda} = p_{z',z}^\lambda$.

From the definitions we see that if $w\lambda \neq w'\lambda$, then $\pi_{w',w,\lambda} = 0$. Thus we may assume that $w\lambda = w'\lambda$. We choose a sequence s_1, s_2, \dots, s_r in \mathbf{I} such that $w\lambda = w'\lambda = s_r s_{r-1} \dots s_1 \lambda \neq s_{r-1} \dots s_1 \lambda \neq \dots \neq s_1 \lambda \neq \lambda$.

We show that for $k \in [0, r]$, $s_k s_{k-1} \dots s_1$ has minimal length in $s_k s_{k-1} \dots s_1 \mathbf{W}_\lambda$. We argue by induction. For $k = 0$ the result is obvious. Assume now that $k \in [1, r]$. Since $s_{k-1} \dots s_1$ has minimal length in $s_{k-1} \dots s_1 \mathbf{W}_\lambda$ and $s_k s_{k-1} \dots s_1 \lambda \neq s_{k-1} \dots s_1 \lambda$, we see from (c) that $s_k s_{k-1} \dots s_1$ has minimal length in $s_k s_{k-1} \dots s_1 \mathbf{W}_\lambda$ as required.

In particular, $s_r s_{r-1} \dots s_1$ has minimal length in $s_r s_{r-1} \dots s_1 \mathbf{W}_\lambda$. Since $w\lambda = s_r s_{r-1} \dots s_1 \lambda$ we have $w = s_r s_{r-1} \dots s_1 h_1 h_2$ where $h_1 \in \vee_\lambda, h_2 \in \mathbf{W}_\lambda$. Then both w_1 and $s_r s_{r-1} \dots s_1 h_1$ have minimal length in $s_r s_{r-1} \dots s_1 h_1 \mathbf{W}_\lambda = w\mathbf{W}_\lambda = w_1 \mathbf{W}_\lambda$; using (b) we deduce that $s_r s_{r-1} \dots s_1 h_1 = w_1$. Hence $s_1 \dots s_r w = s_1 \dots s_r w_1 z = h_1 z$. Similarly, $s_1 \dots s_r w' = h'_1 z'$ where $h'_1 \in \vee_\lambda$.

From the results in 34.7–34.10 we see that $\pi_{w',w,\lambda} = p_{s_1 \dots s_r w', s_1 \dots s_r w}^\lambda = p_{h'_1 z', h_1 z}^\lambda$. Using $h_1, h'_1 \in \vee_\lambda$ and the definitions (34.2) we see that $p_{h'_1 z', h_1 z}^\lambda = 0$ if $h_1 \neq h'_1$ and $p_{h'_1 z', h_1 z}^\lambda = p_{z',z}^\lambda$ if $h_1 = h'_1$.

It remains to show that we have $w_1 = w'_1$ if and only if $h_1 = h'_1$. We have $s_r s_{r-1} \dots s_1 = h_1^{-1} w_1$ and similarly $s_r s_{r-1} \dots s_1 = (h'_1)^{-1} w'_1$. Hence $h_1^{-1} w_1 = (h'_1)^{-1} w'_1$. We see that $w_1 = w'_1$ if and only if $h_1 = h'_1$. This proves (d).

For $w' \leq w$ in \mathbf{W} , $\lambda \in \underline{\mathfrak{g}}$ and $i \in \mathbf{Z}$ we define $N_{i,w',w,\lambda} \in \mathbf{Z}$ by

(e) $\pi_{w',w,\lambda} = v^{l(w')-l(w)} \sum_{i \in \mathbf{Z}} N_{i,w',w,\lambda} v^i$, that is,

$$p_{z',z}^\lambda = v^{l(w')-l(w)} \sum_{i \in \mathbf{Z}} N_{i,w',w,\lambda} v^i$$

if $w'\mathbf{W}_\lambda = w\mathbf{W}_\lambda$ and z, z' are as in (d), then

$$N_{i,w',w,\lambda} = 0 \text{ if } w'\mathbf{W}_\lambda \neq w\mathbf{W}_\lambda.$$

Note that $N_{i,w',w,\lambda}$ is 0 unless i is even.

40.3. Let $B^* \in \mathcal{B}$. Let $U^* = U_{B^*}$ and let T be a maximal torus of B^* . Let $r = \dim \mathbf{T}$. Let $W_T = N_{G^0}T/T$. We identify $T = \mathbf{T}, W_T = \mathbf{W}$ as in 28.5. For any $w \in \mathbf{W}$ we denote by \dot{w} a representative of w in $N_{G^0}T$.

Let $C = G^0/U^* \times G^0/U^*$. We have a partition $C = \bigcup_{w \in \mathbf{W}} C_w$ where

$$C_w = \{(hU^*, h'U^*) \in C; h^{-1}h' \in B^*\dot{w}B^*\}.$$

For $w \in \mathbf{W}$ let $d_w = \dim C_w$ and let

$$\bar{C}_w = \{(hU^*, h'U^*) \in C; h^{-1}h' \in \overline{B^*\dot{w}B^*}\}$$

(closure in G^0). Now \bar{C}_w is an irreducible variety and we have a partition $\bar{C}_w = \bigcup_{w', w' \leq w} C_{w'}$ with C_w smooth, open dense in \bar{C}_w .

Define $\gamma_{\dot{w}} : B^*\dot{w}B^* \rightarrow T$ by $\gamma_{\dot{w}}(g) = t$ where $g \in U^*\dot{w}tU^*$ with $t \in T$. Define $\psi : C_w \rightarrow T$ by $\psi(hU^*, h'U^*) = \gamma_{\dot{w}}(h^{-1}h')$.

For $\mathcal{L} \in \mathfrak{s}$ we set $\mathcal{L}_w = \psi^*\mathcal{L}$, a local system on C_w . (Using 28.1(c) we see that the isomorphism class of $\psi^*\mathcal{L}$ is independent of the choice of \dot{w} .) Let $\mathcal{L}_w^\sharp = IC(\bar{C}_w, \mathcal{L}_w) \in \mathcal{D}(\bar{C}_w)$.

40.4. For $w \in \mathbf{W}, \mathcal{L} \in \mathfrak{s}$ let $\underline{\mathcal{L}}_w = j_{w!}\mathcal{L}_w, \underline{\mathcal{L}}_w^\sharp = \bar{j}_{w!}\mathcal{L}_w^\sharp$ where $j_w : C_w \rightarrow C, \bar{j}_w : \bar{C}_w \rightarrow C$ are the inclusions. Let \hat{C} be the full subcategory of $\mathcal{D}(C)$ whose objects are the simple perverse sheaves on C which are equivariant for the $G^0 \times T \times T$ action

$$(a) (x, t, t') : (hU^*, h'U^*) \mapsto (xht^nU^*, xh't'^nU^*)$$

on C (for some $n \in \mathbf{N}_k^*$) or equivalently, are isomorphic to $\underline{\mathcal{L}}_w^\sharp[d_w]$ for some $\mathcal{L} \in \mathfrak{s}$ and some $w \in \mathbf{W}$. Let $\mathcal{D}^{cs}(C)$ be the subcategory of $\mathcal{D}(C)$ whose objects are those $K \in \mathcal{D}(C)$ such that for any j , any simple subquotient of ${}^pH^jK$ is in \hat{C} .

If w, \mathcal{L} are as above, then $\underline{\mathcal{L}}_w \in \mathcal{D}^{cs}(C)$. Indeed this constructible sheaf is equivariant for the action (a) (for some n), hence so is each ${}^pH^j(\underline{\mathcal{L}}_w)$.

We have a diagram $C \times C \xleftarrow{r} (G^0/U^*)^3 \xrightarrow{s} C$ where

$$\begin{aligned} r(h_1U^*, h_2U^*, h_3U^*) &= ((h_1U^*, h_2U^*), (h_2U^*, h_3U^*)), \\ s(h_1U^*, h_2U^*, h_3U^*) &= (h_1U^*, h_3U^*). \end{aligned}$$

We define a bi-functor $\mathcal{D}(C) \times \mathcal{D}(C) \rightarrow \mathcal{D}(C)$ by $A, A' \mapsto A * A' = s_!r^*(A \boxtimes A')$. The “product” $A * A'$ is associative in an obvious sense. We show that

(b) $A, A' \mapsto A * A'$ restricts to a bi-functor $\mathcal{D}^{cs}(C) \times \mathcal{D}^{cs}(C) \rightarrow \mathcal{D}^{cs}(C)$.

Let $A, A' \in \mathcal{D}^{cs}(C)$. To show that $A * A' \in \mathcal{D}^{cs}(C)$ we may assume that $A, A' \in \hat{C}$. Then each ${}^pH^j(A * A')$ is equivariant for the action (a) (for some n). This proves (b).

40.5. For $w' \leq w$ in $\mathbf{W}, \lambda \in \underline{\mathfrak{s}}, \mathcal{L} \in \lambda$ and $i \in \mathbf{Z}$ we show that

$$(a) \mathcal{H}^i(\mathcal{L}_w^\sharp)|_{C_{w'}} \cong (\mathcal{L}_{w'}(-i/2))^{\oplus N_{i,w',w,\lambda}}.$$

(Both sides are 0 unless i is even.)

Let

$$\begin{aligned} \tilde{C}_w &= \{(h, h') \in G^0 \times G^0; h^{-1}h' \in B^* \dot{w} B^*\} \times \mathbf{k}^*, \\ \bar{C}_w &= \{(h, h') \in G^0 \times G^0; h^{-1}h' \in \overline{B^* \dot{w} B^*}\} \times \mathbf{k}^*. \end{aligned}$$

Now \bar{C}_w is an irreducible variety and we have a partition $\bar{C}_w = \bigcup_{w', w' \leq w} \bar{C}_{w'}$ with \tilde{C}_w smooth, open dense in \bar{C}_w . Define $\bar{d} : \bar{C}_w \rightarrow \bar{C}_w$, $d : \tilde{C}_w \rightarrow \bar{C}_w$ by $(h, h', z) \mapsto (hU^*, h'U^*)$. Let $\tilde{\mathcal{L}}_w = d^* \mathcal{L}_w$, a local system on \tilde{C}_w . Let $\tilde{\mathcal{L}}_w^\sharp = IC(\tilde{C}_w, \tilde{\mathcal{L}}_w)$. Since d, \bar{d} are principal $U^* \times \mathbf{k}^*$ -bundles, it is enough to show that

$$(b) \mathcal{H}^i(\tilde{\mathcal{L}}_w^\sharp)|_{\tilde{C}_{w'}} \cong (\tilde{\mathcal{L}}_{w'}(-i/2))^{\oplus N_{i, w', w, \lambda}}.$$

(Both sides are 0 unless i is even.)

We choose $\kappa \in \text{Hom}(T, \mathbf{k}^*), \mathcal{E} \in \mathfrak{s}(\mathbf{k}^*)$ such that $\mathcal{L} \cong \kappa^* \mathcal{E}$; see 28.1(c).

Now B^* acts on $(B^* \dot{w} B^*) \times \mathbf{k}^*$ and on $(\overline{B^* \dot{w} B^*}) \times \mathbf{k}^*$ by

$$t_1 u : (g, z) \mapsto (g(t_1 u)^{-1}, \kappa(t_1)z)$$

where $t_1 \in T, u \in U^*$. Let $\bar{\mathbf{P}}_w^\kappa = ((\overline{B^* \dot{w} B^*}) \times \mathbf{k}^*)/B^*$, $PP_w^\kappa = ((B^* \dot{w} B^*) \times \mathbf{k}^*)/B^*$. Now \mathbf{P}_w^κ is a smooth open dense subvariety of the irreducible variety $\bar{\mathbf{P}}_w^\kappa$ and $\bar{\mathbf{P}}_w^\kappa = \bigcup_{w', w' \leq w} \bar{\mathbf{P}}_{w'}^\kappa$ is a partition. The morphism $(B^* \dot{w} B^*) \times \mathbf{k}^* \rightarrow \mathbf{k}^*$ given by $(g, z) \mapsto \kappa(\gamma_{\dot{w}}(g))z$ factors through a morphism $\phi : \mathbf{P}_w^\kappa \rightarrow \mathbf{k}^*$. Let $\mathcal{E}_w^\kappa = \phi^* \mathcal{E}$, a local system of rank 1 on \mathbf{P}_w^κ . Let $\mathcal{E}_w^{\kappa \sharp} = IC(\bar{\mathbf{P}}_w^\kappa, \mathcal{E}_w^\kappa) \in \mathcal{D}(\bar{\mathbf{P}}_w^\kappa)$. From [L14, 1.24] we see that

$$(c) \mathcal{H}^i(\mathcal{E}_w^{\kappa \sharp})|_{\bar{\mathbf{P}}_{w'}} \cong (\mathcal{E}_{w'}^\kappa(-i/2))^{\oplus N_{i, w', w, \lambda}}.$$

(Both sides are 0 unless i is even.)

We can find $n \in \mathbf{N}_{\mathbf{k}^*}$ such that $\mathcal{E} \in \mathfrak{s}_n(\mathbf{k}^*)$. Define $\bar{c} : \bar{C}_w \rightarrow \bar{\mathbf{P}}_w, c : \tilde{C}_w \rightarrow \bar{\mathbf{P}}_w$ by $(h, h', z) \mapsto B^* - \text{orbit of } (h^{-1}h', z^n)$. Now \bar{c}, c are locally trivial fibrations with smooth fibres of pure dimension. Hence (b) follows from (c) provided that we can show that $c^* \mathcal{E}_{w'}^\kappa = \tilde{\mathcal{L}}_{w'}$ for $w' \leq w$. We may assume that $w = w'$. We have a commutative diagram

$$\begin{array}{ccccc} \mathbf{P}_w^\kappa & \xleftarrow{c} & \tilde{C}_w \times \mathbf{k}^* & \xrightarrow{d} & C_w \\ \phi \downarrow & & \phi \downarrow & & \kappa \psi \downarrow \\ \mathbf{k}^* & \xleftarrow{c'} & \mathbf{k}^* \times \mathbf{k}^* & \xrightarrow{d'} & \mathbf{k}^* \end{array}$$

with ϕ, ψ, c, d as above, $\phi'(h, h', z) = (\kappa(\gamma_{\dot{w}}(h^{-1}h')), z)$, $c'(z', z) = z'z^n$, $d'(z', z) = z'$. Using this and the definitions we have $\tilde{\mathcal{L}}_w = \phi'^* d'^* \mathcal{E}, c^* \mathcal{E}_w = \phi'^* c'^* \mathcal{E}$. It remains to show that $d'^* \mathcal{E} = c'^* \mathcal{E}$. This expresses the fact that \mathcal{E} is equivariant for the \mathbf{k}^* -action $z_1 : z \mapsto z_1^n z$ on \mathbf{k}^* which follows from $\mathcal{E} \in \mathfrak{s}_n(\mathbf{k}^*)$. This proves (b), hence (a).

40.6. Let $w, w' \in \mathbf{W}, \mathcal{L}, \mathcal{L}' \in \mathfrak{s}$. We set $L = \underline{\mathcal{L}}_w * \underline{\mathcal{L}}_{w'} \in \mathcal{D}^{cs}(C)$. Let

$$\begin{aligned} X &= \{(h_1 U^*, h_2 U^*, h_3 U^*) \in (G^0/U^*)^3; h_1^{-1}h_2 \in B^* \dot{w} B^*, h_2^{-1}h_3 \in B^* \dot{w}' B^*\}, \\ \bar{X} &= \{(h_1 U^*, h_2 B^*, h_3 U^*) \in G^0/U^* \times G^0/B^* \times G^0/U^*; \\ &\quad h_1^{-1}h_2 \in B^* \dot{w} B^*, h_2^{-1}h_3 \in B^* \dot{w}' B^*\}. \end{aligned}$$

We have a commutative diagram with a cartesian square

$$\begin{array}{ccccc} X & \xrightarrow{f} & \bar{X} & \xrightarrow{\bar{\sigma}} & C \\ \tau \downarrow & & \bar{\tau} \downarrow & & \\ T \times T & \xrightarrow{f'} & T & & \end{array}$$

where f is given by $(h_1U^*, h_2U^*, h_3U^*) \mapsto (h_1U^*, h_2B^*, h_3U^*)$,
 f' is $(t, t') \mapsto \text{Ad}(\dot{w}')^{-1}(t)t'$,
 τ is $(h_1U^*, h_2U^*, h_3U^*) \mapsto (t, t')$ with $h_1^{-1}h_2 \in U^*\dot{w}tU^*$, $h_2^{-1}h_3 \in U^*\dot{w}'t'U^*$,
 $\bar{\tau}$ is $(h_1U^*, h_2B^*, h_3U^*) \mapsto \text{Ad}(\dot{w}')^{-1}(t)t'$ with t, t' as in the definition of τ ,
 $\bar{\sigma}$ is $(h_1U^*, h_2B^*, h_3U^*) \mapsto (h_1U^*, h_3U^*)$.

From the definitions we have $L = \bar{\sigma}_! f_! \tau^*(\mathcal{L} \boxtimes \mathcal{L}')$. Using the diagram above, we have $L = \bar{\sigma}_! \bar{\tau}^* f'_!(\mathcal{L} \boxtimes \mathcal{L}')$. From the definitions we see that either (i) or (ii) below holds:

- (i) $\mathcal{L} \not\cong (\text{Ad}(\dot{w}')^{-1})^* \mathcal{L}'$ and $f'_!(\mathcal{L} \boxtimes \mathcal{L}') = 0$;
- (ii) $\mathcal{L} \cong (\text{Ad}(\dot{w}')^{-1})^* \mathcal{L}'$ and $\mathcal{L} \boxtimes \mathcal{L}' = f'^* \mathcal{L}'$.

If (i) holds, then $K = 0$. If (ii) holds, then, as in 32.16, we have

$$\begin{aligned} f'_!(\mathcal{L} \boxtimes \mathcal{L}') &= f'_! f'^* \mathcal{L}' = \mathcal{L}' \otimes f'_! \bar{\mathbf{Q}}_l \simeq \{\mathcal{L}' \otimes \mathcal{H}^e(f'_! \bar{\mathbf{Q}}_l)[-e], e \in \mathbf{Z}\}, \\ \mathcal{L}' \otimes \mathcal{H}^e(f'_! \bar{\mathbf{Q}}_l)[-e] &\simeq \{\mathcal{L}'(\mathbf{r} - e), \dots, \mathcal{L}'(\mathbf{r} - e), (\binom{\mathbf{r}}{2\mathbf{r} - e} \text{ copies})\}. \end{aligned}$$

Setting $\bar{L} = \bar{\sigma}_! \bar{\tau}^*(\mathcal{L}')$, it follows that

$$L \simeq \{\bar{L}(\mathbf{r} - e)[-e], \dots, \bar{L}(\mathbf{r} - e)[-e], (\binom{\mathbf{r}}{2\mathbf{r} - e} \text{ copies}), e \in \mathbf{Z}\}.$$

We now consider \bar{L} for certain choices of w, w' .

If w, w' satisfy $l(ww') = l(w) + l(w')$, then $\bar{\sigma}$ restricts to an isomorphism $\bar{X} \rightarrow C_{ww'}$ and $\bar{L} = \underline{\mathcal{L}}'_{ww'}$.

Now assume that $\alpha, \check{\alpha}, s_\alpha$ are as in 28.3 and that $w = w' = s_\alpha \in \mathbf{I}$. We have

$$\bar{L} \simeq \{j_{u!} \bar{L}_u; u \in \mathbf{W}\}$$

where $j_u : C_u \rightarrow C$ is the inclusion and $\bar{L}_u = j_u^* \bar{L}$. Let $\bar{X}_u = \bar{\sigma}^{-1}(C_u)$. Then $\bar{L}_u = \bar{\sigma}_{u!} \bar{\tau}_u^*(\mathcal{L}')$ where $\bar{\sigma}_u : \bar{X}_u \rightarrow C_u$, $\bar{\tau}_u : \bar{X}_u \rightarrow T$ are the restrictions of $\bar{\sigma}, \bar{\tau}$.

If $u \notin \{1, s_\alpha\}$, then $\bar{X}_u = \emptyset$ and $\bar{L}_u = 0$. If $u = 1$, then $\bar{\sigma}_u : \bar{X}_u \rightarrow C_u$ is an affine line bundle and $\bar{\tau}_u^*(\mathcal{L}') = \bar{\sigma}_u^* \mathcal{L}'_u$; hence $\bar{\sigma}_{u!} \bar{\tau}_u^*(\mathcal{L}') = \bar{\sigma}_{u!} \bar{\sigma}_u^* \mathcal{L}'_u = \mathcal{L}'_u[[-1]]$. If $u = s_\alpha$, then $\bar{\sigma}_u : \bar{X}_u \rightarrow C_u$ is a principal \mathbf{k}^* -bundle and either (iii) or (iv) below holds:

- (iii) $\check{\alpha}^* \mathcal{L}' \not\cong \bar{\mathbf{Q}}_l$ and $\bar{\sigma}_{u!} \bar{\tau}_u^*(\mathcal{L}') = 0$,
- (iv) $\check{\alpha}^* \mathcal{L}' \cong \bar{\mathbf{Q}}_l$ and $\bar{\tau}_u^*(\mathcal{L}') = \bar{\sigma}_u^* \mathcal{L}'_u$.

If (iv) holds, then, as in case (ii) above, we have

$$\begin{aligned} \bar{\sigma}_{u!} \bar{\tau}_u^*(\mathcal{L}') &= \bar{\sigma}_{u!} \bar{\sigma}_u^* \mathcal{L}'_u = \mathcal{L}'_u \otimes \bar{\sigma}_{u!} \bar{\mathbf{Q}}_l \simeq \{\mathcal{L}'_u \otimes \mathcal{H}^e(\bar{\sigma}_{u!} \bar{\mathbf{Q}}_l)[-e], e \in \mathbf{Z}\}, \\ \mathcal{L}'_u \otimes \mathcal{H}^e(\bar{\sigma}_{u!} \bar{\mathbf{Q}}_l)[-e] &\simeq \{\mathcal{L}'_u(1 - e), \dots, \mathcal{L}'_u(1 - e), (\binom{1}{2 - e} \text{ copies})\}. \end{aligned}$$

40.7. In this subsection we assume that \mathbf{k} is an algebraic closure of a finite field. Now the \mathcal{A} -module $\mathfrak{K}(C)$ is defined as in 36.8 (the character sheaves on C are taken to be the objects in \hat{C}).

For $(w, \lambda) \in \mathbf{W} \times \underline{\mathfrak{s}}$, let $[w; \lambda]$ be the basis element of $\mathfrak{K}(C)$ given by $\underline{\mathcal{L}}_w^\sharp[[d_w/2]]$; we choose $\mathcal{L} \in \lambda$ and we regard $\underline{\mathcal{L}}_w, \underline{\mathcal{L}}_w^\sharp$ as mixed complexes on C whose restriction to C_w is pure of weight 0; then $gr(\underline{\mathcal{L}}_w), gr(\underline{\mathcal{L}}_w^\sharp)$ are defined in $\mathfrak{K}(C)$ as in 36.8. We denote these elements of $\mathfrak{K}(C)$ by $[w; \lambda]', [w; \lambda]'^\sharp$ respectively. From 40.5(a) we see that

$$(a) \quad (-v)^{d_w} [w; \lambda] = [w; \lambda]'^\sharp = \sum_{w' \in \mathbf{W}} \sum_{i \in 2\mathbf{Z}} N_{i, w', w, \lambda} v^i [w'; \lambda]' \text{ in } \mathfrak{K}(C)$$

where $N_{i, w', w, \lambda}$ is as in 40.2(e).

Let r, s be as in 40.4. By 40.4(b), $s!r^* : \mathcal{D}(C \times C) \rightarrow \mathcal{D}(C)$ restricts to a functor $\mathcal{D}^{cs}(C \times C) \rightarrow \mathcal{D}^{cs}(C)$ where the character sheaves on $C \times C$ are by definition complexes of the form $A \boxtimes A'$ with $A \in \hat{C}, A' \in \hat{C}$. Hence the \mathcal{A} -linear map $gr(s!r^*) : \mathfrak{K}(C \times C) \rightarrow \mathfrak{K}(C)$ or equivalently $\mathfrak{K}(C) \otimes_{\mathcal{A}} \mathfrak{K}(C) \rightarrow \mathfrak{K}(C)$ is well defined. (We have canonically $\mathfrak{K}(C \times C) = \mathfrak{K}(C) \otimes_{\mathcal{A}} \mathfrak{K}(C)$.) We write $\xi * \xi'$ instead of $gr(s!r^*)(\xi \boxtimes \xi')$ where $\xi, \xi' \in \mathfrak{K}(C)$. Note that $\xi, \xi' \mapsto \xi * \xi'$ defines an associative \mathcal{A} -algebra structure on $\mathfrak{K}(C)$.

Let $w, w' \in \mathbf{W}, \lambda, \lambda' \in \underline{\mathfrak{s}}$. From 40.6 we see that

if $w' \lambda' \neq \lambda$, then $[w; \lambda]' * [w'; \lambda']' = 0$ in $\mathfrak{K}(C)$;

if $w' \lambda' = \lambda$ and $l(ww') = l(w) + l(w')$, then $[w; \lambda]' * [w'; \lambda']' = (v^2 - 1)^r [ww'; \lambda]'$ in $\mathfrak{K}(C)$;

if $s \in \mathbf{I}$ and $s\lambda' = \lambda$, then $[s; \lambda]' * [s; \lambda']' = (v^2 - 1)^r (v^2 [1; \lambda']' + (v^2 - 1)c[s; \lambda']')$

where $c = 1$ for $s \in \mathbf{W}_{\lambda'}$ and $c = 0$ for $s \notin \mathbf{W}_{\lambda'}$.

Using this and (a), 40.1(a), 40.2(e), we see that

(b) the unique \mathcal{A} -linear isomorphism $\omega : \mathfrak{K}(C) \rightarrow H$ (H as in 40.1) given by $[w; \lambda]' \mapsto v^{l(w)} \tilde{T}_w 1_\lambda$ for $w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}$, satisfies $\omega([w; \lambda]) = (-v)^{-d_w} v^{l(w)} c_{w, \lambda}$ for $w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}$ and $\omega(x * x') = (v^2 - 1)^r \omega(x) \omega(x')$ for any $x, x' \in \mathfrak{K}(C)$.

40.8. For $w, w' \in \mathbf{W}$ and $\lambda, \lambda' \in \underline{\mathfrak{s}}$ we have $c_{w, \lambda} c_{w', \lambda'} = \sum_{y \in \mathbf{W}, \nu \in \underline{\mathfrak{s}}} \gamma_{y, \nu}^{w, \lambda; w', \lambda'} c_{y, \nu}$ in the algebra H . Here $\gamma_{y, \nu}^{w, \lambda; w', \lambda'} \in \mathcal{A}$. We have:

$$(a) \quad \gamma_{y, \nu}^{w, \lambda; w', \lambda'} \in \mathbf{N}[v, v^{-1}].$$

By the arguments in 34.4–34.10 (with $D = G^0$) this is reduced to the analogous (well-known) statement for the structure constants of the algebra H_λ^D with its basis (c_w^λ) (see 34.2).

40.9. For any $J \subset \mathbf{I}$ let H_J be the \mathcal{A} -submodule of H spanned by $\{c_{w, \lambda}; w \in \mathbf{W}_J, \lambda \in \underline{\mathfrak{s}}\}$ or equivalently by $\{\tilde{T}_w 1_\lambda; w \in \mathbf{W}_J, \lambda \in \underline{\mathfrak{s}}\}$. From the definitions we see that H_J is a subalgebra of H . For any $J \subset \mathbf{I}, J' \subset \mathbf{I}$ we define a relation $\preceq_{J, J'}$ on $\mathbf{W} \times \underline{\mathfrak{s}}$ as follows. We say that $(y, \nu) \preceq_{J, J'} (w, \lambda)$ if there exist $w_1 \in \mathbf{W}_J, w_2 \in \mathbf{W}_{J'}, \lambda_1, \lambda_2 \in \underline{\mathfrak{s}}$ such that in the expansion (in the algebra H)

$$c_{w_1, \lambda_1} c_{w, \lambda} c_{w_2, \lambda_2} = \sum_{y' \in \mathbf{W}, \nu' \in \underline{\mathfrak{s}}} a_{y', \nu'} c_{y', \nu'}$$

(with $a_{y', \nu'} \in \mathcal{A}$) we have $a_{y, \nu} \neq 0$.

Using the associativity of the product in H , the fact that $H_J, H_{J'}$ are subalgebras of H and 40.8(a), we see that $\preceq_{J, J'}$ is transitive. Using the formula $c_{1, w\lambda} c_{w, \lambda} c_{1, \lambda} = c_{w, \lambda}$ we see that it is reflexive. Thus, it is a preorder. Let $\sim_{J, J'}$ be the equivalence relation attached to $\preceq_{J, J'}$; thus, $(y, \nu) \sim_{J, J'} (w, \lambda)$ if

$(y, \nu) \preceq_{J, J'} (w, \lambda)$ and $(w, \lambda) \preceq_{J, J'} (y, \nu)$. The equivalence classes for $\sim_{J, J'}$ are called (J, J') -two-sided cells. The (\mathbf{I}, \mathbf{I}) -two-sided cells in $\mathbf{W} \times \underline{\mathfrak{s}}$ are also called two-sided cells.

40.10. Let $w, w', w'' \in \mathbf{W}$, $\mathcal{L}, \mathcal{L}', \mathcal{L}'' \in \mathfrak{s}$. We set $K = \underline{\mathcal{L}}_w * \underline{\mathcal{L}}_{w'}^\sharp * \underline{\mathcal{L}}_{w''} \in \mathcal{D}^{cs}(C)$. Let

$$X = \{(h_1U^*, h_2U^*, h_3U^*, h_4U^*) \in (G^0/U^*)^4; \\ h_1^{-1}h_2 \in B^*\dot{w}B^*, h_2^{-1}h_3 \in \overline{B^*\dot{w}'B^*}, h_3^{-1}h_4 \in B^*\dot{w}''B^*\},$$

an irreducible variety. Let X_0 be the smooth open dense subset of X defined by the condition $h_2^{-1}h_3 \in B^*\dot{w}'B^*$. Define $\sigma : X \rightarrow C$ by

$$(h_1U^*, h_2U^*, h_3U^*, h_4U^*) \mapsto (h_1U^*, h_4U^*).$$

Define $\tau : X_0 \rightarrow T \times T \times T$ by

$$(h_1U^*, h_2U^*, h_3U^*, h_4U^*) \mapsto (t, t', t'')$$

with $h_1^{-1}h_2 \in U^*\dot{w}tU^*$, $h_2^{-1}h_3 \in U^*\dot{w}'t'U^*$, $h_3^{-1}h_4 \in U^*\dot{w}''t''U^*$.

Let $\mathcal{F} = \tau^*(\mathcal{L} \boxtimes \mathcal{L}' \boxtimes \mathcal{L}'')$, a local system on X_0 . Then $\mathcal{F}^\sharp := IC(X, \mathcal{F}) \in \mathcal{D}(X)$ is defined and we have $K = \sigma_*\mathcal{F}^\sharp$.

Let \bar{X} (resp. \bar{X}_0) be the the variety of all $(h_1U^*, h_2B^*, h_3B^*, h_4U^*) \in G^0/U^* \times G^0/B^* \times G^0/B^* \times G^0/U^*$ that satisfy the same equations as those defining X (resp. X_0). Note that \bar{X} is irreducible and \bar{X}_0 is an open dense smooth subset of \bar{X} . We have a cartesian diagram

$$\begin{array}{ccccc} X & \xrightarrow{f} & \bar{X} & \xrightarrow{\bar{\sigma}} & C \\ \uparrow & & \uparrow & & \\ X_0 & \xrightarrow{f_0} & \bar{X}_0 & & \\ \tau \downarrow & & \bar{\tau} \downarrow & & \\ T \times T \times T & \xrightarrow{f'} & T & & \end{array}$$

where $X_0 \rightarrow X, \bar{X}_0 \rightarrow \bar{X}$ are the obvious imbeddings,

f, f_0 are given by $(h_1U^*, h_2U^*, h_3U^*, h_4U^*) \mapsto (h_1U^*, h_2B^*, h_3B^*, h_4U^*)$,

f' is $(t, t', t'') \mapsto \text{Ad}(\dot{w}'\dot{w}'')^{-1}(t)\text{Ad}(\dot{w}'')^{-1}(t')t''$,

$\bar{\tau}$ is $(h_1U^*, h_2B^*, h_3B^*, h_4U^*) \mapsto \text{Ad}(\dot{w}'\dot{w}'')^{-1}(t)\text{Ad}(\dot{w}'')^{-1}(t')t''$ with t, t', t'' as in the definition of τ , $\bar{\sigma}$ is $(h_1U^*, h_2B^*, h_3B^*, h_4U^*) \mapsto (h_1U^*, h_4U^*)$. Assume that $\mathcal{L} \cong (\text{Ad}(\dot{w}')^{-1})^*\mathcal{L}'$ and $\mathcal{L}'' \cong (\text{Ad}(\dot{w}'')^{-1})^*\mathcal{L}''$. Then $\mathcal{L} \boxtimes \mathcal{L}' \boxtimes \mathcal{L}'' = f'^*\mathcal{L}''$. We have $\mathcal{F} = \tau^*f'^*\mathcal{L}'' = f_0^*\bar{\tau}^*\mathcal{L}''$. Since f is a principal $T \times T$ -bundle and $X_0 = f^{-1}(\bar{X}_0)$ it follows that $\mathcal{F}^\sharp = f^*IC(\bar{X}, \bar{\tau}^*\mathcal{L}'')$. Note that $f_!\bar{\mathbf{Q}}_l \simeq \{\mathcal{H}^e(f_!\bar{\mathbf{Q}}_l)[-e], 2\mathbf{r} \leq e \leq 4\mathbf{r}\}$,

$$\mathcal{H}^e(f_!\bar{\mathbf{Q}}_l) \simeq \{\bar{\mathbf{Q}}_l(2\mathbf{r} - e), \dots, \bar{\mathbf{Q}}_l(2\mathbf{r} - e), \left(\binom{2\mathbf{r}}{4\mathbf{r} - e} \text{ copies}\right)\}.$$

Hence setting $\bar{K} = \bar{\sigma}_!(IC(\bar{X}, \bar{\tau}^*\mathcal{L}''))$ we have

$$K = \sigma_*f^*IC(\bar{X}, \bar{\tau}^*\mathcal{L}'') = \bar{\sigma}_!f_!f^*IC(\bar{X}, \bar{\tau}^*\mathcal{L}'') = \bar{\sigma}_!(IC(\bar{X}, \bar{\tau}^*\mathcal{L}'') \otimes f_!\bar{\mathbf{Q}}_l),$$

(a) $K \simeq \{\bar{K}(2\mathbf{r} - e)[-e], \dots, \bar{K}(2\mathbf{r} - e)[-e], \left(\binom{2\mathbf{r}}{4\mathbf{r} - e} \text{ copies}\right), 2\mathbf{r} \leq e \leq 4\mathbf{r}\}.$

We now show that

(b) if $A \in \hat{C}$ is such that $A \dashv \bar{K}$, then $A \dashv K$.

We may regard $\mathcal{L}, \mathcal{L}', \mathcal{L}''$ as mixed local systems (with respect to a rational structure over a sufficiently large finite subfield of \mathbf{k}) which are pure of weight 0. Then K, \bar{K} are naturally mixed complexes and (a) is compatible with the mixed structures. For any mixed perverse sheaf P , let P_h be the subquotient of P of pure weight h . We can find $h \in \mathbf{Z}$ such that $A \dashv {}^p H^j(\bar{K})_h$ for some $j \in \mathbf{Z}$; moreover, we may assume that h is maximum possible. Note that $A \dashv {}^p H^{j+4\mathbf{r}}(\bar{K}[-4\mathbf{r}](-2\mathbf{r}))_{h+2\mathbf{r}}$ and $A \dashv {}^p H^{j'}(\bar{K}[-e](2\mathbf{r}-e))_{h+2\mathbf{r}}$ for $2\mathbf{r} \leq e < 4\mathbf{r}$ and any j' ; hence from (a) we see that $A \dashv {}^p H^{j+4\mathbf{r}}(K)_{h+2\mathbf{r}}$. In particular, $A \dashv K$, and (b) is proved.

40.11. Let $w, w', \mathcal{L}, \mathcal{L}', X, \bar{X}, \tau$ be as in 40.6. We set $\mathbf{L} = \underline{\mathcal{L}}_w^\# * \underline{\mathcal{L}}_{w'}^\# \in \mathcal{D}^{cs}(C)$. Let $A = \underline{\mathcal{L}}_{w''}^\#[d_{w''}]$. We show that

(a) if $A \dashv \mathbf{L}$, then $[w'', \lambda'']$ appears with nonzero coefficient in the expansion of the product $[w, \lambda] * [w', \lambda']$ in terms of the basis $([y, \nu])$ of $\mathfrak{K}(C)$.

Let

$$\begin{aligned} \mathbf{X} &= \{(h_1 U^*, h_2 U^*, h_3 U^*) \in (G^0/U^*)^3; h_1^{-1} h_2 \in \overline{B^* \dot{w} B^*}, h_2^{-1} h_3 \in \overline{B^* \dot{w}' B^*}\}, \\ \bar{\mathbf{X}} &= \{(h_1 U^*, h_2 B^*, h_3 U^*) \in G^0/U^* \times G^0/B^* \times G^0/U^*; \\ &\quad h_1^{-1} h_2 \in \overline{B^* \dot{w} B^*}, h_2^{-1} h_3 \in \overline{B^* \dot{w}' B^*}\}. \end{aligned}$$

Note that X (resp. \bar{X}) is naturally an open dense subset of \mathbf{X} (resp. $\bar{\mathbf{X}}$). Define $\sigma' : \mathbf{X} \rightarrow C$ by $(h_1 U^*, h_2 U^*, h_3 U^*) \mapsto (h_1 U^*, h_3 U^*)$. Define $\bar{\sigma}' : \bar{\mathbf{X}} \rightarrow C$ by $(h_1 U^*, h_2 B^*, h_3 U^*) \mapsto (h_1 U^*, h_3 U^*)$. Let $\mathcal{F} = \tau^*(\mathcal{L} \boxtimes \mathcal{L}')$, a local system on X . Then $\mathcal{F}^\# := IC(\mathbf{X}, \mathcal{F}) \in \mathcal{D}(\mathbf{X})$ is defined and we have $\mathbf{L} = \sigma'_! \mathcal{F}^\#$. We have a cartesian diagram

$$\begin{array}{ccccc} \mathbf{X} & \xrightarrow{\tilde{f}} & \bar{\mathbf{X}} & \xrightarrow{\bar{\sigma}'} & C \\ \uparrow & & \uparrow & & \\ X & \xrightarrow{f} & \bar{X} & & \\ \tau \downarrow & & \bar{\tau} \downarrow & & \\ T \times T & \xrightarrow{f'} & T & & \end{array}$$

where $X \rightarrow \mathbf{X}, \bar{X} \rightarrow \bar{\mathbf{X}}$ are the obvious imbeddings, $f, f', \bar{\tau}$ are as in 40.6 and \tilde{f} is the obvious map.

Assume first that 40.6(i) holds. Let $m' : T \times \mathbf{X} \rightarrow \mathbf{X}$ be the free T -action $t_1 : (h_1 U^*, h_2 U^*, h_3 U^*) \mapsto (h_1 U^*, h_2 t_1^{-1} U^*, h_3 U^*)$. This restricts to a free T -action $m : T \times X \rightarrow X$. Define a free T action $m_0 : T \times (T \times T) \rightarrow T \times T$ by $t_1 : (t, t') \mapsto (t_1^{-1} t, \text{Ad}(\dot{w}')^{-1}(t_1) t')$. Then m, m_0 are compatible with τ . By our assumption we have $m_0^*(\mathcal{L} \boxtimes \mathcal{L}') = \mathcal{L}_0 \boxtimes \mathcal{L} \boxtimes \mathcal{L}'$ where $\mathcal{L}_0 \in \mathfrak{s}(T)$, $\mathcal{L}_0 \not\cong \mathbf{Q}_l$. It follows that $m^*(\mathcal{F}) \cong \mathcal{L}_0 \boxtimes \mathcal{F}$. From the properties of intersection cohomology we then have $m'^*(\mathcal{F}^\#) \cong \mathcal{L}_0 \boxtimes \mathcal{F}^\#$. Let $r : T \times \mathbf{X} \rightarrow \mathbf{X}$ be the second projection. Since $\mathcal{L}_0 \in \mathfrak{s}(T)$, $\mathcal{L}_0 \not\cong \mathbf{Q}_l$, we have $r_!(\mathcal{L}_0 \boxtimes \mathcal{F}^\#) = 0$. Hence $r_! m'^*(\mathcal{F}^\#) = 0$. Since m', f', r, f' form a cartesian diagram we must have $f'^* f'(\mathcal{F}^\#) = 0$. Since f' is a principal T -bundle we deduce that $f'_!(\mathcal{F}^\#) = 0$. We have $\mathbf{L} = \bar{\sigma}'_! f'_!(\mathcal{F}^\#)$ hence $\mathbf{L} = 0$. In this case (a) is clear.

Assume next that 40.6(ii) holds. Then $\mathcal{L} \boxtimes \mathcal{L}' = f'^* \mathcal{L}'$ and $\mathcal{F} = \tau^* f'^* \mathcal{L}' = f^* \bar{\tau}^* \mathcal{L}'$. Since f' is a principal T -bundle and $X = f'^{-1}(\bar{X})$ it follows that $\mathcal{F}^\# =$

$f'^*IC(\bar{\mathbf{X}}, \bar{\tau}^* \mathcal{L}')$. Note that $f'_i \bar{\mathbf{Q}}_i \simeq \{\mathcal{H}^e(f'_i \bar{\mathbf{Q}}_i)[-e], \mathbf{r} \leq e \leq 2\mathbf{r}\}$,

$$\mathcal{H}^e(f'_i \bar{\mathbf{Q}}_i) \simeq \{\bar{\mathbf{Q}}_i(\mathbf{r} - e), \dots, \bar{\mathbf{Q}}_i(\mathbf{r} - e), \left(\binom{\mathbf{r}}{2\mathbf{r} - e} \text{ copies}\right)\}.$$

Hence setting $\bar{\mathbf{L}} = \bar{\sigma}'_i(IC(\bar{\mathbf{X}}, \bar{\tau}^* \mathcal{L}'))$ we have

$$\begin{aligned} \mathbf{L} &= \sigma'_i f'^* IC(\bar{\mathbf{X}}, \bar{\tau}^* \mathcal{L}') = \bar{\sigma}'_i f'_i f'^* IC(\bar{\mathbf{X}}, \bar{\tau}^* \mathcal{L}') = \bar{\sigma}'_i(IC(\bar{\mathbf{X}}, \bar{\tau}^* \mathcal{L}') \otimes f'_i \bar{\mathbf{Q}}_i), \\ \mathbf{L} &\simeq \{\bar{\mathbf{L}}(\mathbf{r} - e)[-e], \dots, \bar{\mathbf{L}}(\mathbf{r} - e)[-e], \left(\binom{\mathbf{r}}{2\mathbf{r} - e} \text{ copies}\right), \mathbf{r} \leq e \leq 2\mathbf{r}\}. \end{aligned}$$

Since $A \dashv \mathbf{L}$, this shows that $A \dashv \bar{\mathbf{L}}$. We regard \mathcal{L}' as a pure local system of weight 0. Then $\bar{\mathbf{L}} = \bar{\sigma}'_i(IC(\bar{\mathbf{X}}, \bar{\tau}^* \mathcal{L}'))$ is again pure of weight 0, since $\bar{\sigma}'$ is proper (see [BBD]). Hence the coefficient with which A appears in the expansion of $gr(\bar{\mathbf{L}})$ is a polynomial in $-v$ with coefficients given by the multiplicities of A in the various ${}^p H^j(\bar{\mathbf{L}})$; in particular, A appears with coefficient $\neq 0$ in $gr(\bar{\mathbf{L}})$. On the other hand, the arguments above show that $[w, \lambda] * [w', \lambda'] = (v^2 - 1)^r gr(\bar{\mathbf{L}})$. It follows that A appears with coefficient $\neq 0$ in $[w, \lambda] * [w', \lambda']$. This proves (a).

41. CHARACTER SHEAVES AND TWO-SIDED CELLS

41.1. In this section we preserve the notation of 40.3. We fix a connected component D of G and we pick $\delta \in N_D B^* \cap N_D T$. We write ϵ instead of $\epsilon_D : \mathbf{W} \rightarrow \mathbf{W}$. For $w \in \mathbf{W}$ we set

$$Z_{\emptyset, D}^w = \{(B, B', xU_B) \in Z_{\emptyset, D}; \text{pos}(B, B') = w\}.$$

(This is the same as ${}^{w^{-1}}Z_{\emptyset, D}$ in 36.2.) Define $\xi_D : C \rightarrow Z_{\emptyset, D}$ by $(hU^*, h'U^*) \mapsto (hB^*h^{-1}, h'B^*h'^{-1}, h'\delta h^{-1}U_{hB^*h^{-1}})$, a principal T -bundle for the free T -action on C given by $t : (hU^*, h'U^*) \rightarrow (htU^*, h'(\delta t \delta^{-1})U^*)$.

Since $\xi_D^{-1}(Z_{\emptyset, D}^w) = C_w$, ξ_D restricts to a principal T -bundle $\xi_{D, w} : C_w \rightarrow Z_{\emptyset, D}^w$. We have a commutative diagram

$$\begin{array}{ccccc} T & \xleftarrow{\psi} & C_w & \xrightarrow{=} & C_w \\ \zeta \downarrow & & j' \uparrow & & \xi_{D, w} \downarrow \\ \mathbf{d} & \xleftarrow{pr_2} & G^0/(U^* \cap \dot{w}U^* \dot{w}^{-1}) \times \mathbf{d} & \xrightarrow{j} & Z_{\emptyset, D}^w \end{array}$$

where ψ is as in 40.3,

$$\begin{aligned} \mathbf{d} &= \dot{w}\delta T, \\ j(f(U^* \cap \dot{w}U^* \dot{w}^{-1}), s) &= (fB^*f^{-1}, f\dot{w}B^*\dot{w}^{-1}f^{-1}, fsf^{-1}U_{fB^*f^{-1}}), \\ j'(f(U^* \cap \dot{w}U^* \dot{w}^{-1}), s) &= (fU^*, fs\delta^{-1}U^*), \\ \zeta(t) &= \dot{w}\delta(\delta^{-1}t\delta). \end{aligned}$$

Note that the lower row in the diagram is as in 36.2(a).

Define $\iota : \mathbf{d} \rightarrow T$ by $\iota(\dot{w}\delta t) = t$ where $t \in T$. If $\mathcal{L} \in \mathfrak{s}$ is such that $\text{Ad}((\dot{w}d)^{-1})^* \mathcal{L} \cong \mathcal{L}$, then $pr_2^* \iota^*(\mathcal{L})$ is a local system on $G^0/(U^* \cap \dot{w}U^* \dot{w}^{-1}) \times \mathbf{d}$, equivariant for the T -action $t_0 : (f(U^* \cap \dot{w}U^* \dot{w}^{-1}), s) = (ft_0^{-1}(U^* \cap \dot{w}U^* \dot{w}^{-1}), t_0 s t_0^{-1})$ on $G^0/(U^* \cap \dot{w}U^* \dot{w}^{-1}) \times \mathbf{d}$, which makes j a principal T -bundle. It follows that there is a well-defined local system $\dot{\mathcal{L}}_w$ (of rank 1) on $Z_{\emptyset, D}^w$ such that $j^* \dot{\mathcal{L}}_w = pr_2^* \iota^*(\mathcal{L})$. We show that

$$(a) \quad \xi_{D, w}^*(\dot{\mathcal{L}}_w) = (\text{Ad}(\delta^{-1})^* \mathcal{L})_w.$$

Since j' is an isomorphism, it is enough to show that $j'^*\xi_{D,w}^*(\dot{\mathcal{L}}_w) = j'^*(\text{Ad}(\delta^{-1})^*\mathcal{L})_w$ or that $j^*\dot{\mathcal{L}}_w = j'^*(\text{Ad}(\delta^{-1})^*\mathcal{L})_w$ or that $pr_2^*\iota^*\mathcal{L} = j'^*\psi^*(\text{Ad}(\delta^{-1})^*\mathcal{L})$ or that $j'^*\psi^*\zeta^*\iota^*\mathcal{L} = j'^*\psi^*(\text{Ad}(\delta^{-1})^*\mathcal{L})$. It is enough to show that $\zeta^*\iota^*\mathcal{L} = \text{Ad}(\delta^{-1})^*\mathcal{L}$. This follows from $\text{Ad}(\delta^{-1}) = \iota\zeta : T \rightarrow T$.

Let $h_w : Z_{\emptyset,D}^w \rightarrow Z_{\emptyset,D}$, $\bar{h}_w : \bar{Z}_{\emptyset,D}^w \rightarrow Z_{\emptyset,D}$ be the inclusions ($\bar{Z}_{\emptyset,D}^w = \bigcup_{w'; w' \leq w} Z_{\emptyset,D}^{w'}$ is the closure of $Z_{\emptyset,D}^w$ in $Z_{\emptyset,D}$). Let $\dot{\mathcal{L}}_w = h_{w!}\dot{\mathcal{L}}_w$, $\dot{\mathcal{L}}_w^\sharp = \bar{h}_{w!}\dot{\mathcal{L}}_w^\sharp$. Using (a) and the fact that ξ_D is a principal T -bundle we deduce

- (b) $\xi_D^*(\dot{\mathcal{L}}_w) = \underline{(\text{Ad}(\delta^{-1})^*\mathcal{L})}_w$,
- (c) $\xi_D^*(\dot{\mathcal{L}}_w^\sharp) = \underline{(\text{Ad}(\delta^{-1})^*\mathcal{L})}_w^\sharp$.

Now let D' be another connected component of G . We pick $\delta' \in N_{D'}B^* \cap N_{D'}T$. We have a commutative diagram with a cartesian right square

$$\begin{array}{ccccc} C \times C & \xleftarrow{r} & (G^0/U^*)^3 & \xrightarrow{s} & C \\ \xi_D \times \xi_{D'} \downarrow & & \xi_0 \downarrow & & \xi_{D'D} \downarrow \\ Z_{\emptyset,D} \times Z_{\emptyset,D'} & \xleftarrow{b_1} & Z_0 & \xrightarrow{b_2} & Z_{\emptyset,D'D} \end{array}$$

where r, s are as in 40.4, Z_0, b_1, b_2 are as in 32.5 (with $J = \emptyset$) and

$$\begin{aligned} &\xi_0(h_1U^*, h_2U^*, h_3U^*) \\ &= (h_1B^*h_1^{-1}, h_2B^*h_2^{-1}, h_3B^*h_3^{-1}, h_2\delta h_1^{-1}U_{h_1B^*h_1^{-1}}, h_3\delta' h_2^{-1}U_{h_2B^*h_2^{-1}}). \end{aligned}$$

Hence, if $A \in \mathcal{D}(Z_{\emptyset,D})$, $A' \in \mathcal{D}(Z_{\emptyset,D'})$, then $\xi_{D'D}^*b_2!b_1^*(A \boxtimes A') = s_!r^*(\xi_D^*A \boxtimes \xi_{D'}^*A')$, or equivalently

$$(d) \xi_{D'D}^*(A * A') = (\xi_D^*A) * (\xi_{D'}^*A').$$

41.2. Let $u \in \mathbf{W}$. Let

$$\begin{aligned} \Upsilon_u &= \{(B, B', g(U_B \cap U_{B'})); \\ & B \in \mathcal{B}, B' \in \mathcal{B}, g(U_B \cap U_{B'}) \in D/(U_B \cap U_{B'}), \text{pos}(B, B') = u\} \end{aligned}$$

and let $\Phi_u : \mathcal{D}(Z_{\emptyset,D}) \rightarrow \mathcal{D}(Z_{\emptyset,D})$ be the composition $\mathfrak{h}!j^*$ where $j : \Upsilon_u \rightarrow Z_{\emptyset,D}$ is $(B, B', g(U_B \cap U_{B'})) \mapsto (B, gBg^{-1}, gU_B)$ and $\mathfrak{h} : \Upsilon_u \rightarrow Z_{\emptyset,D}$ is

$$(B, B', g(U_B \cap U_{B'})) \mapsto (B', gB'g^{-1}, gU_{B'}).$$

(A special case of definitions in 37.1.) Let

$$\begin{aligned} \Upsilon' &= \{(B', B, \tilde{B}, \tilde{B}', gU_{B'}); B' \in \mathcal{B}, B \in \mathcal{B}, \tilde{B} \in \mathcal{B}, \tilde{B}' \in \mathcal{B}, \\ & gU_{B'} \in D/U_{B'}, \text{pos}(B', B) = u^{-1}, \text{pos}(\tilde{B}, \tilde{B}') = \epsilon(u), gB'g^{-1} = \tilde{B}'\}, \\ s : \Upsilon_u &\rightarrow \Upsilon', (B, B', g(U_B \cap U_{B'})) \mapsto (B', B, gBg^{-1}, gB'g^{-1}, gU_{B'}). \end{aligned}$$

Note that s is an isomorphism. (We show this only at the level of sets. Define $s' : \Upsilon' \rightarrow \Upsilon_u$ by $(B', B, \tilde{B}, \tilde{B}', gU_{B'}) \mapsto (B, B', x(U_B \cap U_{B'}))$ where $x \in D$ is such that $xBx^{-1} = \tilde{B}$, $xU_{B'} = gU_{B'}$. This is well defined and clearly an inverse of s .) It follows that $\mathfrak{h}!j^* = \mathfrak{h}'!j'^*$ where

$$\begin{aligned} \mathfrak{h}' &= \mathfrak{h}s' : \Upsilon' \rightarrow Z_{\emptyset,D} \text{ is } (B', B, \tilde{B}, \tilde{B}', gU_{B'}) \mapsto (B', \tilde{B}', gU_{B'}), \\ j' &= js' : \Upsilon' \rightarrow Z_{\emptyset,D} \text{ is } (B', B, \tilde{B}, \tilde{B}', gU_{B'}) \mapsto (B, \tilde{B}, xU_B) \end{aligned}$$

and $x \in D$ is such that $xBx^{-1} = \tilde{B}$, $xU_{B'} = gU_{B'}$ (then $x(U_B \cap U_{B'})$ is well defined). We have a commutative diagram with a cartesian right square

$$\begin{CD} C @<\tilde{j}<< \tilde{C} @>\tilde{h}>> C \\ @V\xi_DVV @VV\xi'V @VV\xi_DV \\ Z_{\emptyset,D} @<j'<< \Upsilon'_u @>\mathfrak{h}'>> Z_{\emptyset,D} \end{CD}$$

where ξ_D is as in 41.1,

$$\begin{aligned} \tilde{C} = & \{(h_1U^*, h_2B^*, h_3B^*, h_4U^*) \in (G^0/U^*)^4; \\ & h_1^{-1}h_2 \in B^*\dot{u}^{-1}B^*, h_3^{-1}h_4 \in B^*\delta\dot{u}\delta^{-1}B^*\}, \end{aligned}$$

\tilde{h} is $(h_1U^*, h_2B^*, h_3B^*, h_4U^*) \mapsto (h_1U^*, h_4U^*)$, ξ' is

$$\begin{aligned} & (h_1U^*, h_2B^*, h_3B^*, h_4U^*) \\ & \mapsto (h_1B^*h_1^{-1}, h_2B^*h_2^{-1}, h_3B^*h_3^{-1}, h_4B^*h_4^{-1}, h_4\delta h_1^{-1}U_{h_1B^*h_1^{-1}}), \end{aligned}$$

\tilde{j} is $(h_1U^*, h_2B^*, h_3B^*, h_4U^*) \mapsto (h_2t^{-1}U^*, h_3\tilde{t}U^*)$ where $t, \tilde{t} \in T$ are given by $h_1^{-1}h_2 \in U^*\dot{u}^{-1}tU^*$, $h_3^{-1}h_4 \in U^*\tilde{t}\delta\dot{u}\delta^{-1}U^*$.

We see that for $A \in \mathcal{D}(Z_{\emptyset,D})$ we have

$$\xi_D^* \Phi_u(A) = \xi_D \mathfrak{h}_i j^* A = \xi_D^* \mathfrak{h}'_i j'^* A = \tilde{h}_i \xi'^* j'^* A = \tilde{h}_i \tilde{j}^* \xi_D^* A.$$

Taking here $A = \dot{\underline{\mathcal{L}}}_w^\sharp$ (with $w \in \mathbf{W}$, $\lambda \in \mathfrak{s}$, $\mathcal{L} \in \lambda$ with $wD\lambda = \lambda$) and using 41.1(c) we obtain $\xi_D^* \Phi_u(\dot{\underline{\mathcal{L}}}_w^\sharp) = \tilde{h}_i \tilde{j}^* ((\text{Ad}(\delta^{-1})^* \mathcal{L})_w^\sharp)$ or equivalently $\xi_D^* \Phi_u(\dot{\underline{\mathcal{L}}}_w^\sharp) = \bar{\sigma}_i \tilde{j}'^* ((\text{Ad}(\delta^{-1})^* \mathcal{L})_w^\sharp)$ where

$$\bar{X} = \{(h_1U^*, h_2B^*, h_3B^*, h_4U^*) \in \tilde{C}; h_2^{-1}h_3 \in \overline{B^*\dot{w}B^*}\}$$

and $\tilde{j}' : \bar{X} \rightarrow \bar{C}_w$, $\bar{\sigma} : \bar{X} \rightarrow C$ are the restrictions of \tilde{j}, \tilde{h} . Let

$$\bar{X}_0 = \{(h_1U^*, h_2B^*, h_3B^*, h_4U^*) \in \tilde{C}; h_2^{-1}h_3 \in B^*\dot{w}B^*\}$$

and let $\tilde{j}'_0 : \bar{X}_0 \rightarrow C_w$ be the restriction of \tilde{j} . Let $\mathcal{F}_0 = \tilde{j}'_0^* (\text{Ad}(\delta^{-1})^* \mathcal{L})$, a local system on \bar{X}_0 . Since \tilde{j}' is a fibration with smooth connected fibres, we have $\tilde{j}'^* ((\text{Ad}(\delta^{-1})^* \mathcal{L})_w^\sharp) = IC(\bar{X}, \mathcal{F}_0)$. Thus, $\xi_D^* \Phi_u(\dot{\underline{\mathcal{L}}}_w^\sharp) = \bar{\sigma}_i (IC(\bar{X}, \mathcal{F}_0))$. From the definitions we see that $\mathcal{F}_0 = \bar{\tau}^* \mathcal{L}''$, hence $\bar{\sigma}_i (IC(\bar{X}, \mathcal{F}_0)) = \bar{K}$ and

$$(a) \quad \xi_D^* \Phi_u(\dot{\underline{\mathcal{L}}}_w^\sharp) = \bar{K}$$

where $\bar{\tau}^* \mathcal{L}'$, \bar{K} are given as in 40.10 in terms of

$$(u^{-1}, \mathcal{L}), (w, \text{Ad}(\delta^{-1})^* \mathcal{L}), (\epsilon(u), \text{Ad}(\delta\dot{u}\delta^{-1})^* \text{Ad}(\delta^{-1})^* \mathcal{L})$$

instead of $(w, \mathcal{L}), (w', \mathcal{L}'), (w'', \mathcal{L}'')$.

41.3. For $J \subset \mathbf{I}$ let $\mathcal{D}_J^{\text{cs}}(C)$ be the subcategory of $\mathcal{D}^{\text{cs}}(C)$ whose objects are those $K \in \mathcal{D}(C)$ such that for any j , any simple subquotient of ${}^p H^j K$ is isomorphic to $\dot{\underline{\mathcal{L}}}_w^\sharp$ for some $\mathcal{L} \in \mathfrak{s}$ and some $w \in \mathbf{W}_J$.

Let $J, J' \subset \mathbf{I}$. Let $K \in \mathcal{D}_J^{\text{cs}}(C), K' \in \mathcal{D}_{J'}^{\text{cs}}(C)$, and let $w', w'' \in \mathbf{W}$, $\lambda', \lambda'' \in \mathfrak{s}$, $\mathcal{L}' \in \lambda', \mathcal{L}'' \in \lambda''$. Let $A = \dot{\underline{\mathcal{L}}}_{w''}^{\sharp} [d_{w''}]$. We show that

(a) if (i) $A \dashv K * \dot{\underline{\mathcal{L}}}_{w'}^{\sharp} [d_{w'}]$ or (ii) $A \dashv \dot{\underline{\mathcal{L}}}_{w'}^{\sharp} [d_{w'}] * K'$ or (iii) $A \dashv K * \dot{\underline{\mathcal{L}}}_{w'}^{\sharp} [d_{w'}] * K'$, then $(w'', \lambda'') \preceq_{J, J'} (w', \lambda')$.

For the proof we may assume that \mathbf{k} is an algebraic closure of a finite field. Then the results in 40.7 are applicable. We first consider the case (i). In this case we can find $\mathcal{L} \in \mathfrak{s}, w \in \mathbf{W}_J$ such that $A \dashv \underline{\mathcal{L}}_w^\sharp[d_w] * \underline{\mathcal{L}}_{w'}^\sharp[d_{w'}]$. By 40.11(a), $[w'', \lambda'']$ appears with nonzero coefficient in the expansion of the product $[w, \lambda] * [w', \lambda']$ in terms of the basis $([y, \nu])$ of $\mathfrak{K}(C)$. Applying ω (see 40.7(b)) we see that $c_{w'', \lambda''}$ appears with nonzero coefficient in the expansion of the product $c_{w, \lambda} c_{w', \lambda'}$ in terms of the basis $(c_{y, \nu})$ of H and the desired result follows. Case (ii) is treated in an entirely similar way. We now consider case (iii). In this case we must have $A \dashv A' * K'$ for some simple perverse sheaf A' such that $A' \dashv K * \underline{\mathcal{L}}_{w'}^\sharp[d_{w'}]$. We have $A' = \underline{\mathcal{M}}_y^\sharp[d_y]$ where $y \in \mathbf{W}, \mathcal{M} \in \mathfrak{s}$. Let ν be the isomorphism class of \mathcal{M} . From case (ii) applied to $A \dashv A' * K'$ we see that $(w'', \lambda'') \preceq_{J, J'} (y, \nu)$. From case (i) applied to $A' \dashv K * \underline{\mathcal{L}}_{w'}^\sharp[d_{w'}]$ we see that $(y, \nu) \preceq_{J, J'} (w', \lambda')$. Combining these two inequalities we obtain $(w'', \lambda'') \preceq_{J, J'} (w', \lambda')$, as desired.

41.4. Let $J \subset \mathbf{I}$. In the remainder of this section we write $\mathfrak{f}, \mathfrak{e}$ instead of $\mathfrak{f}_{\emptyset, J} : \mathcal{D}(Z_{\emptyset, D} \rightarrow \mathcal{D}(Z_{J, D}), \mathfrak{e}_{\emptyset, J} : \mathcal{D}(Z_{J, D} \rightarrow \mathcal{D}(Z_{\emptyset, D}))$. We note that

(a) if $A \in \mathcal{D}(Z_{J, D})$, then $\mathfrak{f}\mathfrak{e}(A) \cong A[m] \oplus A'$ for some $m \in \mathbf{Z}$ and some $A' \in \mathcal{D}(Z_{J, D})$.

See [G], [MV] for the special case $D = G^0, J = \mathbf{I}$ and [L10, 6.6] for the general case. Now:

(b) Let A be a simple perverse sheaf on $Z_{J, D}$. Then $A \dashv \mathfrak{f}({}^p H^j(\mathfrak{e}(A)))$ for some $j \in \mathbf{Z}$.

Assume that this is not true. As in [BBD, p. 142], for any $n \in \mathbf{Z}$ we have a distinguished triangle $({}^p \tau_{\leq n-1} \mathfrak{e}A, {}^p \tau_{\leq n} \mathfrak{e}A, {}^p H^n(\mathfrak{e}A)[-n])$, hence a distinguished triangle

$$(\mathfrak{f}({}^p \tau_{\leq n-1} \mathfrak{e}A), \mathfrak{f}({}^p \tau_{\leq n} \mathfrak{e}A), \mathfrak{f}({}^p H^n(\mathfrak{e}A))[-n]).$$

Using our assumption, we see that $A \dashv \mathfrak{f}({}^p \tau_{\leq n-1} \mathfrak{e}A)$ if and only if $A \dashv \mathfrak{f}({}^p \tau_{\leq n} \mathfrak{e}A)$. Thus we have $A \dashv \mathfrak{f}({}^p \tau_{\leq n} \mathfrak{e}A)$ for some n if and only if $A \dashv \mathfrak{f}({}^p \tau_{\leq n} \mathfrak{e}A)$ for any n . Since ${}^p \tau_{\leq n} \mathfrak{e}A = 0$ for some n , we see that $A \dashv \mathfrak{f}({}^p \tau_{\leq n} \mathfrak{e}A)$ for any n . Since ${}^p \tau_{\leq n} \mathfrak{e}A = \mathfrak{e}A$ for some n , we deduce that $A \dashv \mathfrak{f}\mathfrak{e}A$. This contradicts (a); (b) is proved.

We show that

(c) if A is a simple perverse sheaf on $Z_{J, D}$, then there exists a simple perverse sheaf A' on $Z_{\emptyset, D}$ such that $A \dashv \mathfrak{f}(A'), A' \dashv \mathfrak{e}(A)$.

By (b) we can find $i, j \in \mathbf{Z}$ such that $A \dashv {}^p H^i(\mathfrak{f}(P))$ where $P = {}^p H^j(\mathfrak{e}(A))$.

Assume that $A \dashv {}^p H^i(\mathfrak{f}(A'))$ for any simple subquotient A' of P . We claim that $A \dashv {}^p H^i(\mathfrak{f}(P'))$ for any subobject P' of P . We argue by induction on the length of P' . If P' has length 1, the claim holds by assumption. If P' has length ≥ 2 , we can find a simple subobject P'' of P' . We have a distinguished triangle $(\mathfrak{f}(P''), \mathfrak{f}(P'), \mathfrak{f}(P'/P''))$. Hence we have an exact sequence ${}^p H^i(\mathfrak{f}(P'')) \rightarrow {}^p H^i(\mathfrak{f}(P')) \rightarrow {}^p H^i(\mathfrak{f}(P'/P''))$. By the induction hypothesis, we have $A \dashv {}^p H^i(\mathfrak{f}(P''))$, $A \dashv {}^p H^i(\mathfrak{f}(P'/P''))$. Hence $A \dashv {}^p H^i(\mathfrak{f}(P'))$. This proves the claim. In particular, $A \dashv {}^p H^i(\mathfrak{f}(P))$, contradicting the definition of i, P .

We see that there exists a simple subquotient A' of P such that $A \dashv {}^p H^i(\mathfrak{f}(A'))$. Then A' is as required by (c).

Let $\bar{d}_w = \dim Z_{\emptyset, D}^w$. Let

(d) $A' = \underline{\mathcal{L}}_w^\sharp[\bar{d}_w], A'' = \underline{\mathcal{M}}_y^\sharp[\bar{d}_y] \in \hat{Z}_{\emptyset, D}, \mathcal{L} \in \lambda, \mathcal{M} \in \nu$.

Here $w\underline{D}\lambda = \lambda, y\underline{D}\nu = \nu$. Now:

(e) *Let A be a character sheaf on $Z_{J,D}$ such that $A \dashv \mathfrak{f}(A'), A'' \dashv \mathfrak{e}(A)$. Then $(y, \underline{D}\nu) \preceq_{J,J'} (w, \underline{D}\lambda)$.*

Since \mathfrak{f} is proper, $\mathfrak{f}(A')$ is a semisimple complex (see [BBD]). Hence $\mathfrak{f}(A') \cong A[m] \oplus A_1$ for some $m \in \mathbf{Z}, A' \in \mathcal{D}(Z_{J,D})$ and $\mathfrak{e}\mathfrak{f}(A') \cong \mathfrak{e}(A)[m] \oplus \mathfrak{e}(A_1)$. Hence from $A'' \dashv \mathfrak{e}(A)$ we can deduce $A'' \dashv \mathfrak{e}\mathfrak{f}(A')$. By 37.2 we have $\mathfrak{e}\mathfrak{f}(A') \simeq \{\Phi_u(A')[[-m_u]]; u \in \mathbf{W}_J\}$ where m_u are certain integers. Hence for some $u \in \mathbf{W}_J$ we have $A'' \dashv \Phi_u(A')[[-m_u]]$, that is, $A'' \dashv \Phi_u(A')$ and $\xi_D^* A''[\mathbf{r}] \dashv \xi_D^* \Phi_u(A')[\mathbf{r}]$. Hence using 41.2(a) we have $\xi_D^* A''[\mathbf{r}] \dashv \bar{K}$ where \bar{K} is as in the end of 41.2. Thus, $\underline{\mathcal{M}}_y^\sharp[d_y] \dashv \bar{K}$. Using 40.10(b) we deduce that

$$\underline{\mathcal{M}}_y^\sharp[d_y] \dashv \underline{\text{Ad}(w)^{-1}}^* \underline{\text{Ad}(\delta^{-1})}^* \mathcal{L}_{u^{-1}} * (\underline{\text{Ad}(\delta^{-1})}^* \mathcal{L}_w^\sharp * \underline{\text{Ad}(\delta u \delta^{-1})}^* \underline{\text{Ad}(\delta^{-1})}^* \mathcal{L}_{\epsilon(u)}).$$

Using this and 41.3(a) we see that (e) holds.

Now:

(f) *Let A be a character sheaf on $Z_{J,D}$. In the setup of (d) assume that $A \dashv \mathfrak{f}(A'), A' \dashv \mathfrak{e}(A), A \dashv \mathfrak{f}(A''), A'' \dashv \mathfrak{e}(A)$. Then $(y, \underline{D}\nu) \sim_{J,J'} (w, \underline{D}\lambda)$.*

Applying (e) to A', A'' we see that $(y, \underline{D}\nu) \preceq_{J,J'} (w, \underline{D}\lambda)$. Applying (e) to A'', A' (instead of A', A'') we see that $(w, \underline{D}\lambda) \preceq_{J,J'} (y, \underline{D}\nu)$. Hence (f) holds.

From (c),(f) we see that there is a well-defined map $A \mapsto \mathbf{c}_A$ from the set of character sheaves on $Z_{J,D}$ (up to isomorphism) to the set of (J, J') -two-sided cells in $\mathbf{W} \times \underline{\mathfrak{F}}$ where \mathbf{c}_A is the unique (J, J') -two-sided cell that contains

$$\{(w, \underline{D}\lambda) \in \mathbf{W} \times \underline{\mathfrak{g}}; w\underline{D}\lambda = \lambda, A \dashv \mathfrak{f}(\underline{\mathcal{L}}_w^\sharp[\bar{d}_w]), \underline{\mathcal{L}}_w^\sharp[\bar{d}_w] \dashv A\}$$

(a nonempty set); here $\mathcal{L} \in \lambda$.

41.5. In the setup of 41.4, let A be a character sheaf on $Z_{J,D}$. We show that:

(a) *There exists $(w, \underline{D}\lambda) \in \mathbf{c}_A$ such that $w\underline{D}\lambda = \lambda, A \dashv \mathfrak{f}(\underline{\mathcal{L}}_w^\sharp[\bar{d}_w])$. If $(w', \underline{D}\lambda') \in \mathbf{W} \times \underline{\mathfrak{g}}$ is such that $w'\underline{D}\lambda' = \lambda', A \dashv \mathfrak{f}(\underline{\mathcal{L}}_{w'}^\sharp[\bar{d}_{w'}])$, then $(w, \underline{D}\lambda) \preceq_{J,J'} (w', \underline{D}\lambda')$. Here $\mathcal{L} \in \lambda, \mathcal{L}' \in \lambda'$.*

(b) *There exists $(w, \underline{D}\lambda) \in \mathbf{c}_A$ such that $w\underline{D}\lambda = \lambda, \underline{\mathcal{L}}_w^\sharp[\bar{d}_w] \dashv \mathfrak{e}(A)$. If $(w', \underline{D}\lambda') \in \mathbf{W} \times \underline{\mathfrak{g}}$ is such that $w'\underline{D}\lambda' = \lambda', \underline{\mathcal{L}}_{w'}^\sharp[\bar{d}_{w'}] \dashv \mathfrak{e}(A)$, then $(w', \underline{D}\lambda') \preceq_{J,J'} (w, \underline{D}\lambda)$. Here $\mathcal{L} \in \lambda, \mathcal{L}' \in \lambda'$.*

Note that (a) follows immediately from 41.4(c),(e) and the definition of \mathbf{c}_A . Similarly, (b) follows from 41.4(c),(e) and the definition of \mathbf{c}_A .

41.6. In this subsection we assume that $J = \mathbf{I}$. The \mathcal{A} linear map $H \rightarrow H$ given by

$$(a) \tilde{T}_w 1_\lambda \mapsto \tilde{T}_{\epsilon(w)} 1_{\underline{D}\lambda} \text{ for } w \in \mathbf{W}, \lambda \in \underline{\mathfrak{g}}$$

is an \mathcal{A} -algebra isomorphism. It carries $c_{w,\lambda}$ to $c_{\epsilon(w),\underline{D}\lambda}$ for any $w \in \mathbf{W}, \lambda \in \underline{\mathfrak{g}}$. It induces a bijection $\mathbf{c} \mapsto \mathbf{c}'$ from the set of two-sided cells in $\mathbf{W} \times \underline{\mathfrak{g}}$ onto itself. We show that

$$(b) \text{ if } A \text{ is a character sheaf on } D, \text{ then } (\mathbf{c}_A)' = \mathbf{c}_A.$$

Consider the automorphism $\text{Ad}(\delta) : D \rightarrow D$. From the definitions we see that for $(w, \lambda) \in \mathbf{W} \times \underline{\mathfrak{g}}$ such that $w\underline{D}\lambda = \lambda$ we have $A \dashv \mathfrak{f}(\underline{\mathcal{L}}_w^\sharp[\bar{d}_w])$ if and only if

$\text{Ad}(\delta^{-1})^*A \dashv \text{f}(\underline{\text{Ad}}(\underline{D}^{-1})^*\mathcal{L}_{\epsilon(w)}^\sharp[\bar{d}_w])$. Using this and 41.5(a) we see that

$$\mathbf{c}_{\text{Ad}(\delta^{-1})^*A} = (\mathbf{c}_A)'$$

It is then enough to show that $\text{Ad}(\delta^{-1})^*A \cong A$. By the G^0 -equivariance of A we have $m^*A \cong q^*A$ where $m : G^0 \times D \rightarrow D$ is $(x, g) \mapsto xgx^{-1}$ and $q : G^0 \times D \rightarrow D$ is $(x, g) \mapsto g$. Define $r : D \rightarrow G^0 \times D$ by $r(g) = (\delta g^{-1}, g)$. Then $r^*m^*A \cong r^*q^*A$ that is, $(mr)^*A \cong (qr)^*A$. We have $mr = \text{Ad}(\delta)$, $qr = 1$, hence $\text{Ad}(\delta)^*A \cong A$ and $\text{Ad}(\delta^{-1})^*A \cong A$, as required.

Note also that for (w, λ) as above we have

$$(c) \text{f}(\underline{\text{Ad}}(\underline{D}^{-1})^*\mathcal{L}_{\epsilon(w)}^\sharp[\bar{d}_w]) \cong \text{f}(\underline{\mathcal{L}}_w^\sharp[\bar{d}_w]).$$

Indeed, let $K = \text{f}(\underline{\mathcal{L}}_w^\sharp[\bar{d}_w])$. Clearly, we have $m^*K \cong q^*K$ with m, q as above. Then as in the proof of (b) we see that $\text{Ad}(\delta)^*K \cong K$. From the definitions we see that $\text{f}(\underline{\text{Ad}}(\underline{D}^{-1})^*\mathcal{L}_{\epsilon(w)}^\sharp[\bar{d}_w]) = \text{Ad}(\delta^{-1})^*K$. Since $\text{Ad}(\delta^{-1})^*K \cong K$, (c) follows.

41.7. In this and the next subsection we assume that \mathbf{k} is an algebraic closure of a finite field. From 41.1(c) we see that $\xi_D^* : \mathcal{D}(Z_{\emptyset, D}) \rightarrow \mathcal{D}(C)$ restricts to a functor $\mathcal{D}^{cs}(Z_{\emptyset, D}) \rightarrow \mathcal{D}^{cs}(C)$, hence, as in 36.8, the \mathcal{A} -linear map $gr(\xi_D^*) : \mathfrak{K}(Z_{\emptyset, D}) \rightarrow \mathfrak{K}(C)$ is well defined; from 41.1(c) we see also that

$$(a) \text{gr}(\xi_D^*)(\underline{\mathcal{L}}_w^\sharp[\bar{d}_w]) = (-v)^r[w; \underline{D}\lambda]$$

for $w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}$ such that $w\underline{D}\lambda = \lambda$ and $\mathcal{L} \in \lambda$. From (a) we see that $gr(\xi_D^*)$ is injective with image equal to $\mathfrak{K}(C)^D$, the \mathcal{A} -submodule of $\mathfrak{K}(C)$ spanned by $\{[w; \underline{D}\lambda]; w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}, w\underline{D}\lambda = \lambda\}$ or equivalently by $\{[w; \underline{D}\lambda]'; w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}, w\underline{D}\lambda = \lambda\}$. Thus, $gr(\xi_D^*)$ defines an isomorphism $\eta' : \mathfrak{K}(Z_{\emptyset, D}) \xrightarrow{\sim} \mathfrak{K}(C)^D$. Let $\eta = \eta'^{-1}$.

Let $n \in \mathbf{N}_{\mathbf{k}}^*$. Let $\mathfrak{K}(C)_n^D$ be the \mathcal{A} -submodule of $\mathfrak{K}(C)$ spanned by $\{[w; \underline{D}\lambda]; w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}_n, w\underline{D}\lambda = \lambda\}$ or equivalently by $\{[w; \underline{D}\lambda]'; w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}_n, w\underline{D}\lambda = \lambda\}$.

Let $u, w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}_n$ be such that $w\underline{D}\lambda = \lambda$ and let $\mathcal{L} \in \lambda$. From 37.3(c) we see that the \mathcal{A} -linear map $gr(\Phi_u) : \mathfrak{K}(Z_{\emptyset, D}) \rightarrow \mathfrak{K}(Z_{\emptyset, D})$ is well defined; we denote it again by Φ_u . From 40.10(a), 41.2(a) we have

$$[u^{-1}; \lambda]' * [w; \underline{D}\lambda]'^\sharp * [\epsilon_D(u); \underline{D}(u^{-1}\lambda)]' = (v^2 - 1)^{2r} \eta' \Phi_u \eta([w; \underline{D}\lambda]'^\sharp),$$

equality in $\mathfrak{K}(C)$. If $\lambda' \in \underline{\mathfrak{s}}_n, \lambda' \neq \lambda$, we have (from 40.7) that $[u^{-1}; \lambda']' * [w; \underline{D}\lambda]'^\sharp * [\epsilon_D(u); \underline{D}(u^{-1}\lambda)']' = 0$. It follows that

$$(v^2 - 1)^{2r} \eta' \Phi_u \eta([w; \underline{D}\lambda]'^\sharp) = \sum_{\lambda' \in \underline{\mathfrak{s}}_n} [u^{-1}; \lambda']' * [w; \underline{D}\lambda]'^\sharp * [\epsilon_D(u); \underline{D}(u^{-1}\lambda)']'.$$

Using this and the definition of $\mathfrak{K}(C)_n^D$ we see that

$$(v^2 - 1)^{2r} \eta' \Phi_u \eta(x) = \sum_{\lambda' \in \underline{\mathfrak{s}}_n} [u^{-1}; \lambda']' * x * [\epsilon_D(u); \underline{D}(u^{-1}\lambda)']'$$

for any $x \in \mathfrak{K}(C)_n^D$. Applying η to both sides we obtain

$$(b) \quad (v^2 - 1)^{2r} \Phi_u \eta'(x) = \sum_{\lambda' \in \underline{\mathfrak{s}}_n} \eta([u^{-1}; \lambda']' * x * [\epsilon_D(u); \underline{D}(u^{-1}\lambda)']')$$

for any $x \in \mathfrak{K}(C)_n^D$.

41.8. In the setup of 41.4, let A be a character sheaf on $Z_{J,D}$. From 36.9(b) we see that the condition that, if $(w', \underline{D}\lambda') \in \mathbf{W} \times \underline{\mathfrak{g}}$ is such that $w' \underline{D}\lambda' = \lambda'$, then we have $A \dashv \mathfrak{f}(\dot{\mathcal{L}}_{w'}^\#[\bar{d}_{w'}])$ if and only if A appears with coefficient $\neq 0$ in the expansion of $\mathfrak{f}(\dot{\mathcal{L}}_{w'}^\#[\bar{d}_{w'}]) \in \mathfrak{K}(Z_{J,D})$ as a linear combination of the canonical basis of $\mathfrak{K}(Z_{J,D})$. Hence from 41.5(a) we deduce:

(a) *There exists $(w, \underline{D}\lambda) \in \mathbf{c}_A$ such that $w \underline{D}\lambda = \lambda$ and A appears with nonzero coefficient in $\mathfrak{f}(\dot{\mathcal{L}}_w^\#[\bar{d}_w]) \in \mathfrak{K}(Z_{J,D})$. If $(w', \underline{D}\lambda') \in \mathbf{W} \times \underline{\mathfrak{g}}$ is such that $w' \underline{D}\lambda' = \lambda'$ and A appears with nonzero coefficient in $\mathfrak{f}(\dot{\mathcal{L}}_{w'}^\#[\bar{d}_{w'}]) \in \mathfrak{K}(Z_{J,D})$, then $(w, \underline{D}\lambda) \preceq_{J,J'} (w', \underline{D}\lambda')$. Here $\mathcal{L} \in \lambda, \mathcal{L}' \in \lambda'$.*

Clearly, property (a) characterizes \mathbf{c}_A .

41.9. Let $J \subset J' \subset \mathbf{I}$ and let D' be another connected component of G . Let $A_0 \in \mathcal{D}(Z_{J,D})$, $A' \in \mathcal{D}(Z_{\epsilon_D(J'), D'})$. We show that

$$(a) \mathfrak{f}_{J,J'}(A_0) * A' \cong \mathfrak{f}_{J,J'}(A_0 * \mathfrak{e}_{\epsilon_D(J), \epsilon_D(J')} A') \text{ in } \mathcal{D}(Z_{J', D'D}).$$

Indeed, from the definitions we see that both sides of (a) can be identified with $b_! c^*(A_0 \boxtimes A')$ where b, c are as in the diagram

$$Z_{J,D} \times Z_{\epsilon_D(J'), D'} \xleftarrow{c} Y \xrightarrow{b} Z_{J', D'D}$$

where

$$Y = \{(P, R, R', gU_R, g'U_{R'}); P \in \mathcal{P}_J, R \in \mathcal{P}_{J'}, R' \in \mathcal{P}_{\epsilon_D(J')}, \\ gU_R \in D/U_R, g'U_{R'} \in D'/U_{R'}, gRg^{-1} = R', P \subset R\},$$

c is $(P, R, R', gU_R, g'U_{R'}) \mapsto ((P, gU_P), (R', g'U_{R'}))$, b is $(P, R, R', gU_R, g'U_{R'}) \mapsto (R, g'gU_R)$.

An entirely similar proof shows that, if $A \in \mathcal{D}(Z_{J', D})$, $A'_0 \in \mathcal{D}(Z_{\epsilon_D(J), D'})$, then

$$(b) A * \mathfrak{f}_{\epsilon_D(J), \epsilon_D(J')} (A'_0) \cong \mathfrak{f}_{J,J'} (\mathfrak{e}_{J,J'} A * A'_0) \text{ in } \mathcal{D}(Z_{J', D'D}).$$

41.10. Let \mathbf{c} be a two-sided cell in $\mathbf{W} \times \underline{\mathfrak{g}}$. Let $\bar{\mathbf{c}}$ be the set of all $(w, \lambda) \in \mathbf{W} \times \underline{\mathfrak{g}}$ such that $(w, \lambda) \preceq_{\mathbf{I}, \mathbf{I}} (y, \nu)$ for some/any $(y, \nu) \in \mathbf{c}$.

If $K \in \mathcal{D}(Z_{\emptyset, D})$, we say that $K \in \mathcal{D}_{\bar{\mathbf{c}}}^{cs}(Z_{\emptyset, D})$ if for any $j \in \mathbf{Z}$ and simple subquotient A of ${}^p H^j(K)$ satisfies $\mathbf{c}_A \subset \bar{\mathbf{c}}$.

Let D' be another connected component of G . We show that

$$(a) \text{ if } K \in \mathcal{D}_{\bar{\mathbf{c}}}^{cs}(Z_{\emptyset, D}), K' \in \mathcal{D}^{cs}(Z_{\epsilon_D(J'), D'}), \text{ then } K * K' \in \mathcal{D}_{\bar{\mathbf{c}}}^{cs}(Z_{\emptyset, D'D}).$$

We may assume that \mathbf{k} is an algebraic closure of a finite field. We may assume that $K \in \hat{Z}_{\emptyset, D}$ and $\mathbf{c}_K \subset \bar{\mathbf{c}}$. Then there exists $(w, \underline{D}\lambda) \in \mathbf{c}_K$ such that $w \underline{D}\lambda = \lambda$, $K \dashv \mathfrak{f}(\dot{\mathcal{L}}_w^\#[\bar{d}_w])$, $\mathcal{L} \in \lambda$. It is enough to show that, if $\tilde{A} \in \hat{Z}_{\emptyset, D'D}$ is such that $\tilde{A} \dashv K * K'$, then $\mathbf{c}_{\tilde{A}} \subset \bar{\mathbf{c}}$. Since $\mathfrak{f}(\dot{\mathcal{L}}_w^\#[\bar{d}_w])$ is a semisimple complex (see the line after 41.4(e)) we have $\mathfrak{f}(\dot{\mathcal{L}}_w^\#[\bar{d}_w]) \cong K[m] \oplus \tilde{K}$ for some $m \in \mathbf{Z}$, $\tilde{K} \in \mathcal{D}(Z_{\emptyset, D'D})$. It follows that $\mathfrak{f}(\dot{\mathcal{L}}_w^\#[\bar{d}_w]) * K' \cong K * K'[m] \oplus \tilde{K} * K'$ hence $\tilde{A} \dashv \mathfrak{f}(\dot{\mathcal{L}}_w^\#[\bar{d}_w]) * K'$. By 41.9(a) we have $\mathfrak{f}(\dot{\mathcal{L}}_w^\#[\bar{d}_w]) * K' \cong \mathfrak{f}(\dot{\mathcal{L}}_w^\#[\bar{d}_w] * \mathfrak{e}(K'))$ hence $\tilde{A} \dashv \mathfrak{f}(\dot{\mathcal{L}}_w^\#[\bar{d}_w] * \mathfrak{e}(K'))$. We deduce that there exists $K'_0 \in \hat{Z}_{\emptyset, D'}$ such that $\tilde{A} \dashv \mathfrak{f}(\dot{\mathcal{L}}_w^\#[\bar{d}_w] * K'_0)$ and $K''_0 \in \hat{Z}_{\emptyset, D'D}$ such that $K''_0 \dashv \dot{\mathcal{L}}_w^\#[\bar{d}_w] * K'_0$, $\tilde{A} \dashv \mathfrak{f}(K''_0)$. We then have $\xi_{D'D}^* K''_0[\mathbf{r}] \dashv \xi_{D'D}^*(\dot{\mathcal{L}}_w^\#[\bar{d}_w] * K'_0)[\mathbf{r}]$, hence, using 41.1(d), $\xi_{D'D}^* K''_0[\mathbf{r}] \dashv (\xi_D^* \dot{\mathcal{L}}_w^\#[\bar{d}_w]) * \xi_{D'} K'_0$. Setting $gr(\xi_{D'D}^* K''_0[\mathbf{r}]) = [w_1, \underline{D}' \underline{D}\lambda_1] \in \mathfrak{K}(C)$ with $(w_1, \lambda_1) \in \mathbf{W} \times \underline{\mathfrak{g}}$ we see, using 41.3(a) that $(w_1, \underline{D}' \underline{D}\lambda_1) \preceq_{\mathbf{I}, \mathbf{I}} (w, \underline{D}\lambda)$. From $\tilde{A} \dashv \mathfrak{f}(K''_0)$ we see using 41.5(a) that

$\mathbf{c}_{\bar{A}} \preceq_{\mathbf{I}, \mathbf{I}} (w_1, \underline{D}' \underline{D} \lambda_1)$ (that is, some/any element of $\mathbf{c}_{\bar{A}}$ is $\preceq_{\mathbf{I}, \mathbf{I}} (w_1, \underline{D}' \underline{D} \lambda_1)$). Using the transitivity of $\preceq_{\mathbf{I}, \mathbf{I}}$ we see that $\mathbf{c}_{\bar{A}} \preceq_{\mathbf{I}, \mathbf{I}} (w, \underline{D} \lambda)$. This proves (a).

An entirely similar argument shows that

(b) if $K \in \mathcal{D}^{cs}(Z_{\emptyset, D})$, $K' \in \mathcal{D}^{cs}(Z_{\epsilon_D(J'), D'})$, then $K * K' \in \mathcal{D}^{cs}(Z_{\emptyset, D'D})$.

42. DUALITY AND THE FUNCTOR $\mathfrak{f}_{\emptyset, \mathbf{I}}$

42.1. In this section we fix a connected component D of G . We write ϵ instead of $\epsilon_D : \mathbf{W} \rightarrow \mathbf{W}$. We write \mathfrak{f} instead of $\mathfrak{f}_{\emptyset, \mathbf{I}} : \mathcal{D}(Z_{\emptyset, D} \rightarrow \mathcal{D}(Z_{\mathbf{I}, D}))$. We assume that \mathbf{k} is an algebraic closure of a finite field.

Let $J \subset \mathbf{I}$ be such that $\epsilon(J) = J$. Recall from 30.3 that $V_{J, D} = \{(P, gU_P); P \in \mathcal{P}_J, gU_P \in N_D P/U_P\}$. As in 30.4 (with $J' = \mathbf{I}$) we consider the diagram $V_{J, D} \xleftarrow{c} V_{J, \mathbf{I}, D} \xrightarrow{d} D$ where $V_{J, \mathbf{I}, D} = \{(P, g); P \in \mathcal{P}_J, g \in N_D P\}$, c is $(P, g) \mapsto (P, gU_P)$ and d is $(P, g) \mapsto g$. Define $\tilde{f}_J : \mathcal{D}(V_{J, D}) \rightarrow \mathcal{D}(D)$, $\tilde{e}_J : \mathcal{D}(D) \rightarrow \mathcal{D}(V_{J, D})$ by $\tilde{f}_J A = d_! c^* A$, $\tilde{e}_J A' = c_! d^* A'$. (In the notation of 30.4 we have $\tilde{f}_J = \tilde{f}_{J, \mathbf{I}}$, $\tilde{e}_J = \tilde{e}_{J, \mathbf{I}}$.) Define $f_J : \mathcal{D}(V_{J, D}) \rightarrow \mathcal{D}(D)$, $e_J : \mathcal{D}(D) \rightarrow \mathcal{D}(V_{J, D})$ by $f_J A = \tilde{f}_J A[[\alpha_J/2]]$, $e_J A = \tilde{e}_J A[[\alpha_J/2]]$ where $\alpha_J = \dim \mathcal{P}_J$. (In the notation of 30.4 we have $f_J A = f_{J, \mathbf{I}} A(\alpha_J/2)$, $e_J A = e_{J, \mathbf{I}} A(-\alpha_J/2)$. Thus, f_J, e_J are the same, up to a twist, as $f_{J, \mathbf{I}}, e_{J, \mathbf{I}}$.)

From 30.5 (with $J' = \mathbf{I}$) we see that for $A \in \mathcal{D}(V_{J, D})$, $A' \in \mathcal{D}(D)$ we have canonically

(a) $\text{Hom}_{\mathcal{D}(V_{J, D})}(e_J A', A) = \text{Hom}_{\mathcal{D}(D)}(A', f_J A)$.

Let $CS(V_{J, D}), CS(D)$ be as in 38.1. From 38.2, 38.3 we see that

(b) f_J, e_J restrict to functors $CS(V_{J, D}) \rightarrow CS(D)$, $CS(D) \rightarrow CS(V_{J, D})$ denoted again by f_J, e_J .

We show that

(c) if $A \in CS(V_{J, D})$ comes from a pure complex of weight 0 with respect to a rational structure over a finite subfield of \mathbf{k} , then $f_J A$ (naturally regarded as a mixed complex) is pure of weight 0.

Indeed, the functor c^* preserves pure complexes of weight 0 since c is smooth with connected fibres; the functor $d_!$ preserves pure complexes of weight 0 since d is proper (see [De, 6.2.6]) and $[[\alpha_J/2]]$ also preserves pure complexes of weight 0.

We show that

(d) if $A' \in CS(D)$ comes from a pure complex of weight 0 with respect to a rational structure over a finite subfield of \mathbf{k} , then $e_J A'$ (naturally regarded as a mixed complex) is pure of weight 0.

Using (b), it is enough to show that for any simple A as in (c), the natural action of Frobenius on the vector space $\text{Hom}_{\mathcal{D}(V_{J, D})}(e_J A', A)$ has weight 0. Using (a) we see that it is enough to show that the natural action of Frobenius on the vector space $\text{Hom}_{\mathcal{D}(D)}(A', f_J A)$ has weight 0. This follows from (c) using (b).

Define an imbedding $s : V_{J, D} \rightarrow Z_{J, D}$ by $(P, gU_P) \mapsto (P, P, gU_P)$. From the definitions we see that

(e) $\tilde{f}_J : \mathcal{D}(V_{J, D}) \rightarrow \mathcal{D}(D)$ is the composition $\mathcal{D}(V_{J, D}) \xrightarrow{s_!} \mathcal{D}(Z_{J, D}) \xrightarrow{\tilde{f}_{J, \mathbf{I}}} \mathcal{D}(D)$,

(f) $\tilde{e}_J : \mathcal{D}(D) \rightarrow \mathcal{D}(V_{J, D})$ is the composition $\mathcal{D}(D) \xrightarrow{\epsilon_{J, \mathbf{I}}} \mathcal{D}(Z_{J, D}) \xrightarrow{s^*} \mathcal{D}(V_{J, D})$.

Let $Y = \{(B, B', gU_B) \in Z_{\emptyset, D}; \text{pos}(B, B') \in \mathbf{W}_J\}$ and let $r : Y \rightarrow Z_{\emptyset, D}$ be the inclusion. From the definitions we have

(g) $s_! s^* \mathfrak{f}_{\emptyset, J} = \mathfrak{f}_{\emptyset, J} r_! r^* : \mathcal{D}(Z_{\emptyset, D}) \rightarrow \mathcal{D}(Z_{J, D})$.

Note that $V_{J,D} = {}^1Z_{J,D}$ (see 36.2); hence the “character sheaves” on $V_{J,D} = {}^1Z_{J,D}$ are defined as in 36.8 and $\mathcal{D}^{cs}(V_{J,D} = \mathcal{D}^{cs}({}^1Z_{J,D}))$ is defined as 36.8. In particular, $\mathfrak{K}(V_{J,D}) = \mathfrak{K}({}^1Z_{J,D})$ is defined. Let $\mathfrak{K}_0(V_{J,D}) = \bigoplus_A \mathbf{Z}A \subset \mathfrak{K}(V_{J,D})$ where A runs through the character sheaves on $V_{J,D}$ (up to isomorphism).

From (b) we see that \tilde{f}_J, \tilde{e}_J restrict to functors $\mathcal{D}^{cs}(V_{J,D}) \rightarrow \mathcal{D}^{cs}(D), \mathcal{D}^{cs}(D) \rightarrow \mathcal{D}^{cs}(V_{J,D})$, hence the \mathcal{A} -linear maps $gr(\tilde{f}_J) : \mathfrak{K}(V_{J,D}) \rightarrow \mathfrak{K}(D), gr(\tilde{e}_J) : \mathfrak{K}(D) \rightarrow \mathfrak{K}(V_{J,D})$ are well defined; we denote them by \tilde{f}_J, \tilde{e}_J . Define $f_J : \mathfrak{K}(V_{J,D}) \rightarrow \mathfrak{K}(D)$ by $f_J = (-v)^{-\alpha_J} \tilde{f}_J$ and $e_J : \mathfrak{K}(D) \rightarrow \mathfrak{K}(V_{J,D})$ by $e_J = (-v)^{-\alpha_J} \tilde{e}_J$. We show that

(h) $f_J : \mathfrak{K}(V_{J,D}) \rightarrow \mathfrak{K}(D), e_J : \mathfrak{K}(D) \rightarrow \mathfrak{K}(V_{J,D})$ restrict to group homomorphisms $\mathfrak{K}_0(V_{J,D}) \rightarrow \mathfrak{K}_0(D), \mathfrak{K}_0(D) \rightarrow \mathfrak{K}_0(V_{J,D})$ denoted again by f_J, e_J .

It is enough to prove the following statement. If x is a canonical basis element of $\mathfrak{K}(V_{J,D})$ (resp. $\mathfrak{K}(D)$), then $f_J(x)$ (resp. $e_J(x)$) is an \mathbf{N} -linear combination of canonical basis elements of $\mathfrak{K}(D)$ (resp. $\mathfrak{K}(V_{J,D})$). This is immediate from (c), (d).

Now, one checks easily that $r_1 r^* : \mathcal{D}(Z_{\emptyset,D}) \rightarrow \mathcal{D}(Z_{\emptyset,D})$ restricts to a functor $\mathcal{D}^{cs}(Z_{\emptyset,D}) \rightarrow \mathcal{D}^{cs}(Z_{\emptyset,D})$. (Note that, if $w \in \mathbf{W}, \lambda \in \underline{\mathfrak{g}}, \mathcal{L} \in \lambda$ and $w \underline{D} \lambda = \lambda$, then $r_1 r^*(\dot{\mathcal{L}}_w) = \dot{\mathcal{L}}_w$ for $w \in \mathbf{W}_J$ and $r_1 r^*(\dot{\mathcal{L}}_w) = 0$ for $w \in \mathbf{W} - \mathbf{W}_J$.) It follows that the \mathcal{A} -linear map $gr(r_1 r^*) : \mathfrak{K}(Z_{\emptyset,D}) \rightarrow \mathfrak{K}(Z_{\emptyset,D})$ (denoted by ρ_J) is well defined.

Let $\mathfrak{K}(C)^D, \eta$ be as in 41.7. Define an \mathcal{A} -linear map $\tilde{\rho}_J : \mathfrak{K}(C)^D \rightarrow \mathfrak{K}(C)^D$ by $[w; \underline{D} \lambda]' \mapsto [w; \underline{D} \lambda]'$ if $w \in \mathbf{W}_J, \lambda \in \underline{\mathfrak{g}}, w \underline{D} \lambda = \lambda$ and $[w; \underline{D} \lambda]' \mapsto 0$ if $w \in \mathbf{W} - \mathbf{W}_J, \lambda \in \underline{\mathfrak{g}}, w \underline{D} \lambda = \lambda$. From the definitions we see that

(i) $\rho_J \eta(x) = \eta \tilde{\rho}_J(x)$ for all $x \in \mathfrak{K}(C)^D$.

42.2. We define an \mathcal{A} -linear map $\mathbf{d} : \mathfrak{K}(D) \rightarrow \mathfrak{K}(D)$ by

$$\mathbf{d}(x) = \sum_{J; J \subset \mathbf{I}; \epsilon(J)=J} (-1)^{|J_\epsilon|} f_J e_J(x)$$

where f_J, e_J are as in 42.1(h) and J_ϵ is as in 38.1. Now:

(a) Let A be a character sheaf on D . Then $\mathbf{d}(A) = \pm A'$ where A' is a character sheaf on D . Moreover, \pm and A' are the same as in 38.11(a).

For any $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ let $\mathcal{K}(V_{J,D})$ be as in 38.9. We shall identify $\mathfrak{K}(V_{J,D})/(v-1)\mathfrak{K}(V_{J,D}) = \mathcal{K}(V_{J,D})$ as abelian groups in such a way that the image of A_1 (a character sheaf on $V_{J,D}$) in $\mathfrak{K}(V_{J,D})/(v-1)\mathfrak{K}(V_{J,D})$ is identified with the basis element A_1 of $\mathcal{K}(V_{J,D})$. From the definitions we see that the homomorphisms

$$\mathfrak{K}(D)/(v-1)\mathfrak{K}(D) \rightarrow \mathfrak{K}(V_{J,D})/(v-1)\mathfrak{K}(V_{J,D}) \rightarrow \mathfrak{K}(D)/(v-1)\mathfrak{K}(D)$$

induced by e_J, f_J in 42.1(h) are then identified with the homomorphisms

$$e_{J,\mathbf{I}} : \mathcal{K}(D) \rightarrow \mathcal{K}(V_{J,D}), f_{J,\mathbf{I}} : \mathcal{K}(V_{J,D}) \rightarrow \mathcal{K}(D)$$

in 38.2, 38.3. It follows that the endomorphism of $\mathfrak{K}(D)/(v-1)\mathfrak{K}(D)$ induced by $\mathbf{d} : \mathfrak{K}(D) \rightarrow \mathfrak{K}(D)$ is identified with the homomorphism $\mathcal{K}(D) \rightarrow \mathcal{K}(D)$ denoted in 38.10(a), 38.11 again by \mathbf{d} . Hence we have $\mathbf{d}(A) = \pm A' + (v-1)x$ (in $\mathfrak{K}(D)$) where \pm, A' are as in 38.11(a) and $x \in \mathfrak{K}(D)$. From 42.1(h) we see that $\mathbf{d}(A) \in \mathfrak{K}_0(D)$. Since $\pm A' \in \mathfrak{K}_0(D)$, we see that $(v-1)x \in \mathfrak{K}_0(D)$. Since $\mathfrak{K}_0(D) \cap (v-1)\mathfrak{K}(D) = 0$, we have $(v-1)x = 0$ and $x = 0$. This proves (a).

42.3. We have $H = H_D \oplus H'_D$ where H_D (resp. H'_D) is the \mathcal{A} -submodule of H_n spanned by $\{\tilde{T}_w 1_{\underline{D}\lambda}; w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}, w\underline{D}\lambda = \lambda\}$ (resp. by $\{\tilde{T}_w 1_{\underline{D}\lambda}; w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}, w\underline{D}\lambda \neq \lambda\}$). Equivalently,

$$H_D = \sum_{\lambda \in \underline{\mathfrak{s}}} 1_\lambda H 1_{\underline{D}\lambda} \subset H, \quad H'_D = \sum_{\lambda, \lambda' \in \underline{\mathfrak{s}}; \lambda \neq \lambda'} 1_{\lambda'} H 1_{\underline{D}\lambda} \subset H.$$

Recall that $\omega : \mathfrak{K}(C) \xrightarrow{\sim} H$ is defined in 40.7(b). Define an \mathcal{A} -linear map $\tilde{\omega} : H \rightarrow \mathfrak{K}(C)^D$ by

$$\begin{aligned} \tilde{\omega}(y) &= \omega^{-1}(y) \quad \text{if } y \in H_D, \\ \tilde{\omega}(y) &= 0 \quad \text{if } y \in H'_D. \end{aligned}$$

Then $\eta\tilde{\omega}(y) \in \mathfrak{K}(Z_{\emptyset, D})$ is well defined for any $y \in H$. Here η is as in 41.7.

Let $n \in \mathbf{N}_{\mathbf{k}}^*$. Let $H_{n, D} = H_D \cap H_n$. Note that $H_{n, D}$ is the \mathcal{A} -submodule of H_n spanned by $\{\tilde{T}_w 1_{\underline{D}\lambda}; w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}_n, w\underline{D}\lambda = \lambda\}$.

For $J \subset \mathbf{I}$ such that $\epsilon(J) = J$ we define an \mathcal{A} -linear map $\rho_{J, n} : H_{n, D} \rightarrow H_{n, D}$ by

$$\begin{aligned} \tilde{T}_w 1_{\underline{D}\lambda} &\mapsto \tilde{T}_w 1_{\underline{D}\lambda} \quad \text{if } w \in \mathbf{W}_J, \lambda \in \underline{\mathfrak{s}}_n, w\underline{D}\lambda = \lambda, \\ \tilde{T}_w 1_{\underline{D}\lambda} &\mapsto 0 \quad \text{if } w \in \mathbf{W} - \mathbf{W}_J, \lambda \in \underline{\mathfrak{s}}_n, w\underline{D}\lambda = \lambda. \end{aligned}$$

We have the following result.

Lemma 42.4. *For any $y \in H_{n, D}$ we have $\mathbf{d}(\mathfrak{f}\eta\tilde{\omega}(y)) = \mathfrak{f}\eta\tilde{\omega}(\delta(y))$ where*

$$\delta = \sum_{J \subset \mathbf{I}; \epsilon(J) = J} (-1)^{|J\epsilon|} \delta_J$$

with $\delta_J : H_{n, D} \rightarrow H_{n, D}$ given by

$$\delta_J(y) = \rho_{J, n} \left(\sum_{u \in \mathbf{W}^J} \tilde{T}_{u^{-1}} y \tilde{T}_{\epsilon_D(u)} \right)$$

(the sum in the right-hand side is computed in H_n but it belongs to $H_{n, D}$).

Applying 37.2 with K, K', J replaced by \emptyset, J, \mathbf{I} and with $A' \in \mathcal{D}^{cs}(Z_{\emptyset, D})$ we obtain

$$\mathbf{e}_{J, \mathbf{I}} \mathfrak{f} A' \simeq \{\mathfrak{f}_{\emptyset, J} \Phi_u A' [[-m_u]]; u \in \mathbf{W}^J\}$$

(in $\mathcal{D}(Z_{J, D})$), with $\Phi_u : \mathcal{D}(Z_{\emptyset, D}) \rightarrow \mathcal{D}(Z_{\emptyset, D})$ as in 37.1 and $m_u = \alpha_J - \lambda(u)$ where $\alpha_J = \dim \mathcal{P}_J$. Applying here s^* we obtain

$$s^* \mathbf{e}_{J, \mathbf{I}} \mathfrak{f} A' \simeq \{s^* \mathfrak{f}_{\emptyset, J} \Phi_u A' [[-m_u]]; u \in \mathbf{W}^J\}.$$

We replace $s^* \mathbf{e}_{J, \mathbf{I}}$ by \tilde{e}_J (see 42.1(f)) and we apply $\tilde{f}_J = \mathfrak{f}_{J, \mathbf{I}} s!$ (see 42.1(e)); we obtain

$$\tilde{f}_J \tilde{e}_J \mathfrak{f} A' \simeq \{\mathfrak{f}_{J, \mathbf{I}} s! s^* \mathfrak{f}_{\emptyset, J} \Phi_u A' [[-m_u]]; u \in \mathbf{W}^J\}.$$

Using now 42.1(g) we obtain

$$\tilde{f}_J \tilde{e}_J \mathfrak{f} A' \simeq \{\mathfrak{f}_{J, \mathbf{I}} \mathfrak{f}_{\emptyset, J} r! r^* \Phi_u A' [[-m_u]]; u \in \mathbf{W}^J\}.$$

Here we replace $\mathfrak{f}_{J, \mathbf{I}} \mathfrak{f}_{\emptyset, J}$ by \mathfrak{f} (see 36.4(b)). This (or rather its mixed analogue) gives rise to the following equality in $\mathfrak{K}(D)$:

$$\tilde{f}_J \tilde{e}_J \mathfrak{f}(x') = \sum_{u \in \mathbf{W}^J} v^{2m_u} \mathfrak{f}_{\rho_J} \Phi_u(x')$$

for any $x' \in \mathfrak{K}(Z_{\emptyset, D})$, or equivalently

$$f_J e_J \mathfrak{f}(x') = \sum_{u \in \mathbf{W}^J} v^{2m_u - 2\alpha_J} \mathfrak{f} \rho_J \Phi_u(x').$$

Taking $x' = \eta(x)$ where $x \in \mathfrak{K}(C)_n^D$ (see 41.7) and using 41.7(b) we obtain

$$\begin{aligned} & (v^2 - 1)^{2r} f_J e_J \mathfrak{f} \eta(x) \\ &= \sum_{u \in \mathbf{W}^J} \sum_{\lambda \in \underline{\mathfrak{s}}_n} v^{-2l(u)} \mathfrak{f} \rho_J \eta([u^{-1}; \lambda]' * x * [\epsilon_D(u); \underline{D}(u^{-1}\lambda)]') \end{aligned}$$

and using 42.1(i),

$$\begin{aligned} & (v^2 - 1)^{2r} f_J e_J \mathfrak{f} \eta(x) \\ &= \sum_{u \in \mathbf{W}^J} \sum_{\lambda \in \underline{\mathfrak{s}}_n} v^{-2l(u)} \mathfrak{f} \eta \tilde{\rho}_J([u^{-1}; \lambda]' * x * [\epsilon_D(u); \underline{D}(u^{-1}\lambda)]') \end{aligned}$$

for any $x \in \mathfrak{K}(C)_n^D$. Here we replace x by $\tilde{\omega}(y)$ where $y \in H_{n, D}$ and $\tilde{\rho}_J|_{\mathfrak{K}(C)_n^D}$ by $\tilde{\omega}|_{H_{n, D}} \rho_{J, n} \omega_{\mathfrak{K}(C)_n^D}$; using 40.7(b) we obtain:

$$\begin{aligned} f_J e_J \mathfrak{f} \eta \tilde{\omega}(y) &= \sum_{u \in \mathbf{W}^J} \sum_{\lambda \in \underline{\mathfrak{s}}_n} \mathfrak{f} \eta \tilde{\omega} \rho_{J, n}(\tilde{T}_{u^{-1}} 1_{\lambda} y \tilde{T}_{\epsilon_D(u)} 1_{\underline{D}(u^{-1}\lambda)}) \\ &= \mathfrak{f} \eta \tilde{\omega} \rho_{J, n} \left(\sum_{u \in \mathbf{W}^J} \tilde{T}_{u^{-1}} y \tilde{T}_{\epsilon_D(u)} \right). \end{aligned}$$

The lemma is proved.

42.5. As in 34.12 let \mathfrak{U} be the subfield of $\bar{\mathbf{Q}}_l$ generated by the roots of 1. Let $\Phi : H_n^D \rightarrow \mathcal{A} \otimes_{\mathbf{Z}} H_n^{D, \infty}$ be as in 34.12 (a special case of a definition in 34.1) and let $\Phi^1 : H_n^{D, 1} \rightarrow \mathfrak{U} \otimes_{\mathbf{Z}} H_n^{D, \infty}$ be the specialization of Φ for $v = 1$ (see 34.12(b)). Let $\tilde{\mathcal{A}} = \mathfrak{U}[v, v^{-1}]$, let $H_n^{D, \tilde{\mathcal{A}}} = \tilde{\mathcal{A}} \otimes_{\mathcal{A}} H_n^D$ and let $\Phi^{\tilde{\mathcal{A}}} : H_n^{D, \tilde{\mathcal{A}}} \rightarrow \tilde{\mathcal{A}} \otimes_{\mathbf{Z}} H_n^{D, \infty}$ be the homomorphism obtained from Φ by extending the scalars from \mathcal{A} to $\tilde{\mathcal{A}}$.

Let E be an $H_n^{D, 1}$ -module of finite dimension over \mathfrak{U} . Since Φ^1 is an isomorphism of \mathfrak{U} -algebras (see 34.12(b)) we may regard E as an $\mathfrak{U} \otimes_{\mathbf{Z}} H_n^{D, \infty}$ -module E^∞ via $(\Phi^1)^{-1}$. By extension of scalars, $\tilde{\mathcal{A}} \otimes_{\mathfrak{U}} E^\infty$ is naturally a module over

$$\tilde{\mathcal{A}} \otimes_{\mathfrak{U}} (\mathfrak{U} \otimes_{\mathbf{Z}} H_n^{D, iy}) = \tilde{\mathcal{A}} \otimes_{\mathbf{Z}} H_n^{D, iy}$$

and this can be regarded as an $H_n^{D, \tilde{\mathcal{A}}}$ -module $E^{\tilde{\mathcal{A}}}$ via $\Phi^{\tilde{\mathcal{A}}}$.

Let $J \subset \mathbf{I}$ be such that $\epsilon(J) = J$. Let $H_{J, n}^D$ be the \mathcal{A} -algebra of H_n^D generated by $1_\lambda, \lambda \in \underline{\mathfrak{s}}_n$ by $\tilde{T}_w, w \in \mathbf{W}_J$ and by $\tilde{T}_{\underline{D}}$. Note that $\{\tilde{T}_{w \underline{D}'} 1_\lambda; w \in \mathbf{W}_J, \underline{D}' = \text{power of } \underline{D}\}$ is an \mathcal{A} -basis of $H_{J, n}^D$. Let $H_{J, n}^{D, 1} = \mathfrak{U} \otimes_{\mathcal{A}} H_{J, n}^D$ where \mathfrak{U} is regarded as an \mathcal{A} -algebra via $v \mapsto 1$. Let $H_{J, n}^{D, \tilde{\mathcal{A}}} = \tilde{\mathcal{A}} \otimes_{\mathcal{A}} H_{J, n}^D$. Note that $H_{J, n}^{D, \tilde{\mathcal{A}}}$ is naturally a subalgebra of $H_n^{D, \tilde{\mathcal{A}}}$. Hence $E^{\tilde{\mathcal{A}}}$ may be regarded as an $H_{J, n}^{D, \tilde{\mathcal{A}}}$ -module $(E^{\tilde{\mathcal{A}}})_J$. This $H_{J, n}^{D, \tilde{\mathcal{A}}}$ -module may be induced to an $H_n^{D, \tilde{\mathcal{A}}}$ -module

$$\text{IND}((E^{\tilde{\mathcal{A}}})_J) := H_n^{D, \tilde{\mathcal{A}}} \otimes_{H_{J, n}^{D, \tilde{\mathcal{A}}}} E_J^{\tilde{\mathcal{A}}}.$$

Next, $H_{J, n}^{D, 1}$ is naturally a subalgebra of $H_n^{D, 1}$. Hence E may be regarded as an $H_{J, n}^{D, 1}$ -module E_J . This $H_{J, n}^{D, 1}$ -module may be induced to an $H_n^{D, 1}$ -module $\text{ind}(E_J) := H_n^{D, 1} \otimes_{H_{J, n}^{D, 1}} E_J$. Define an $H_{J, n}^{D, \tilde{\mathcal{A}}}$ -module $(\text{ind}(E_J))^{\tilde{\mathcal{A}}}$ in terms of $\text{ind}(E_J)$

in the same way as $E^{\tilde{\mathcal{A}}}$ was defined in terms of E . By extension of scalars from $\tilde{\mathcal{A}}$ to $\mathfrak{U}(v)$ (the quotient field of $\tilde{\mathcal{A}}$), $\text{IND}((E^{\tilde{\mathcal{A}}})_J)$, $(\text{ind}(E_J))^{\tilde{\mathcal{A}}}$ give rise to $\mathfrak{U}(v) \otimes_{\mathcal{A}} H_n^D$ -modules $\mathfrak{U}(v) \otimes_{\tilde{\mathcal{A}}} \text{IND}((E^{\tilde{\mathcal{A}}})_J)$, $\mathfrak{U}(v) \otimes_{\tilde{\mathcal{A}}} (\text{ind}(E_J))^{\tilde{\mathcal{A}}}$. We show that

(a) *these two $\mathfrak{U}(v) \otimes_{\mathcal{A}} H_n^D$ -modules are isomorphic.*

Since $\mathfrak{U}(v) \otimes_{\mathcal{A}} H_n^D$, $H_n^{D,1}$ are (finite dimensional) semisimple algebras (see 34.12) it follows by standard arguments that it is enough to show that $\text{IND}((E^{\tilde{\mathcal{A}}})_J)$, $(\text{ind}(E_J))^{\tilde{\mathcal{A}}}$ become isomorphic $H_n^{D,1}$ -modules under the specialization $v = 1$. First we note that under the specialization $v = 1$, $E^{\tilde{\mathcal{A}}}$ becomes the $H_n^{D,1}$ -module E . (This is because the specialization of $\Phi^{\tilde{\mathcal{A}}}$ at $v = 1$ cancels $(\Phi_1)^{-1}$.) In particular, the specialization of $(\text{ind}(E_J))^{\tilde{\mathcal{A}}}$ for $v = 1$ is $\text{ind}(E_J)$. Moreover, from the definition of induction, the specialization of $\text{IND}((E^{\tilde{\mathcal{A}}})_J)$ for $v = 1$ is the same as $\text{ind}(E'_J)$ where E' is the specialization of $E^{\tilde{\mathcal{A}}}$ for $v = 1$, that is, $E' = E$. This proves (a).

Lemma 42.6. *We preserve the setup of 42.5. Let $\mathfrak{U}(v) \otimes_{\tilde{\mathcal{A}}} E^{\tilde{\mathcal{A}}}$, $\mathfrak{U}(v) \otimes_{\tilde{\mathcal{A}}} (\text{ind}(E_J))^{\tilde{\mathcal{A}}}$ be the $\mathfrak{U}(v) \otimes_{\mathcal{A}} H_n^D$ -module obtained from $E^{\tilde{\mathcal{A}}}$, $(\text{ind}(E_J))^{\tilde{\mathcal{A}}}$ by extension of scalars from $\tilde{\mathcal{A}}$ to $\mathfrak{U}(v)$. Let $y \in H_{n,D}$. We have*

$$\text{tr}(\delta_J(y)\tilde{T}_D, \mathfrak{U}(v) \otimes_{\tilde{\mathcal{A}}} E^{\tilde{\mathcal{A}}}) = \text{tr}(y\tilde{T}_D, \mathfrak{U}(v) \otimes_{\tilde{\mathcal{A}}} (\text{ind}(E_J))^{\tilde{\mathcal{A}}}).$$

Let $H_{J,n}$ be the \mathcal{A} -subalgebra of H_n defined in 31.8. Define an \mathcal{A} -linear map $p_J : H_n \rightarrow H_{J,n}$ by $p_J(\tilde{T}_z 1_\lambda) = \tilde{T}_z 1_\lambda$ if $z \in \mathbf{W}_J, \lambda \in \underline{\mathfrak{s}}_n$, $p_J(\tilde{T}_z 1_\lambda) = 0$ if $z \in \mathbf{W} - \mathbf{W}_J, \lambda \in \underline{\mathfrak{s}}_n$. We show that

(a) $p_J(\tilde{T}_u h') = \delta_{u,1} h'$ if $u \in \mathbf{W}^J, h' \in H_{J,n}$.

We may assume that $h' = \tilde{T}_b 1_\lambda, b \in \mathbf{W}_J, \lambda \in \underline{\mathfrak{s}}_n$. Then $p_J(\tilde{T}_u \tilde{T}_b 1_\lambda) = p_J(\tilde{T}_{ub} 1_\lambda) = \delta_{u,1} \tilde{T}_{ub} 1_\lambda = \delta_{u,1} \tilde{T}_b 1_\lambda$, as required.

We show that

(b) $p_J(hh') = p_J(h)h'$ for any $h \in H_n, h' \in H_{J,n}$.

We may assume $h = \tilde{T}_u \tilde{T}_b 1_\nu, h' = \tilde{T}_a 1_\lambda, u \in \mathbf{W}^J, a, b \in \mathbf{W}_J, \lambda, \nu \in \underline{\mathfrak{s}}_n$. We must show that $p_J(\tilde{T}_u \tilde{T}_b 1_\nu \tilde{T}_a 1_\lambda) = p_J(\tilde{T}_u \tilde{T}_b 1_\nu) \tilde{T}_a 1_\lambda$. If $u \neq 1$, both sides are zero by (a). If $u = 1$, both sides are $\tilde{T}_b 1_\nu \tilde{T}_a 1_\lambda$. This proves (b).

By 34.13(a) we have

(c) $p_\emptyset(\tilde{T}_x \tilde{T}_{x'} 1_\lambda) = \delta_{xx',1}$ for $x, x' \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}_n$.

For $u, u' \in \mathbf{W}^J, \lambda \in \underline{\mathfrak{s}}_n$ we write $\tilde{T}_{u^{-1}} \tilde{T}_{u'} 1_\lambda = \sum_{a \in \mathbf{W}} f_a \tilde{T}_a 1_\lambda$ where $f_a \in \mathcal{A}$. For $a' \in \mathbf{W}_J$ we have

$$\tilde{T}_{a'^{-1}u^{-1}} \tilde{T}_{u'} 1_\lambda = \tilde{T}_{a'^{-1}} \tilde{T}_{u^{-1}} \tilde{T}_{u'} 1_\lambda = \sum_{a \in \mathbf{W}} f_a \tilde{T}_{a'^{-1}} \tilde{T}_a 1_\lambda.$$

Applying p_\emptyset to this and using (c) gives $f_{a'} = \delta_{u',ua'} = \delta_{a',1} \delta_{u,u'}$ so that

$$p_J(\tilde{T}_{u^{-1}} \tilde{T}_{u'} 1_\lambda) = \sum_{a \in \mathbf{W}_J} f_a \tilde{T}_a 1_\lambda = \delta_{u,u'} \tilde{T}_1 1_\lambda.$$

Since this holds for any $\lambda \in \underline{\mathfrak{s}}_n$ we have

(d) $p_J(\tilde{T}_{u^{-1}} \tilde{T}_{u'}) = \delta_{u,u'} \tilde{T}_1$.

Let $w \in \mathbf{W}, \lambda \in \underline{\mathfrak{s}}_n, u \in \mathbf{W}^J$. We have

$$\tilde{T}_w 1_\lambda \tilde{T}_u = \sum_{u' \in \mathbf{W}^J, a \in \mathbf{W}_J} c_{w,u,u',a,\lambda} \tilde{T}_{u'} \tilde{T}_a 1_{u^{-1}\lambda}$$

where $c_{w,u,u',a,\lambda} \in \mathcal{A}$ are uniquely determined. It follows that

$$\tilde{T}_{u^{-1}}\tilde{T}_w1_\lambda\tilde{T}_{\epsilon(u)} = \sum_{u' \in \mathbf{W}^J, a \in \mathbf{W}_J} c_{w,\epsilon(u),u',a,\lambda}\tilde{T}_{u^{-1}}\tilde{T}_{u'}\tilde{T}_a1_{\epsilon(u)^{-1}\lambda}.$$

Applying p_J and using (b),(d) we obtain

$$\begin{aligned} p_J(\tilde{T}_{u^{-1}}\tilde{T}_w1_\lambda\tilde{T}_{\epsilon(u)}) &= \sum_{u' \in \mathbf{W}^J, a \in \mathbf{W}_J} c_{w,\epsilon(u),u',a,\lambda}p_J(\tilde{T}_{u^{-1}}\tilde{T}_{u'})\tilde{T}_a1_{\epsilon(u)^{-1}\lambda} \\ \text{(e)} \quad &= \sum_{u' \in \mathbf{W}^J, a \in \mathbf{W}_J} c_{w,\epsilon(u),u',a,\lambda}\delta_{u,u'}\tilde{T}_a1_{\epsilon(u)^{-1}\lambda} = \sum_{a \in \mathbf{W}_J} c_{w,\epsilon(u),u,a,\lambda}\tilde{T}_a1_{\epsilon(u)^{-1}\lambda}. \end{aligned}$$

Let $(e_i)_{i \in X}$ be a basis of the free $\tilde{\mathcal{A}}$ -module $E^{\tilde{\mathcal{A}}}$. For $a \in \mathbf{W}_J, \lambda \in \underline{\mathfrak{g}}_n$ we have $\tilde{T}_a1_\lambda\tilde{T}_{\underline{D}}e_i = \sum_{i' \in X} \tilde{c}_{a,\lambda,i,i'}e_{i'}$ where $\tilde{c}_{a,\lambda,i,i'} \in \tilde{\mathcal{A}}$.

Since $H_n^{D,\tilde{\mathcal{A}}}$ is a free right $H_{J,n}^{D,\tilde{\mathcal{A}}}$ -module with basis $\{\tilde{T}_u; u \in \mathbf{W}^J\}$, we see that $\{\tilde{T}_u \otimes e_i; u \in \mathbf{W}^J, i \in X\}$ is a basis of the free $\tilde{\mathcal{A}}$ -module $\text{ind}((E^{\tilde{\mathcal{A}}})_J)$.

Let $w \in \mathbf{W}, \lambda \in \underline{\mathfrak{g}}_n, u \in \mathbf{W}^J$ be such that $w\underline{D}\lambda = \lambda$. In $\text{IND}((E^{\tilde{\mathcal{A}}})_J)$ we have

$$\begin{aligned} \tilde{T}_w1_\lambda\tilde{T}_{\underline{D}}(\tilde{T}_u \otimes e_i) &= (\tilde{T}_w1_\lambda\tilde{T}_{\epsilon(u)}\tilde{T}_{\underline{D}}) \otimes e_i \\ &= \sum_{u' \in \mathbf{W}^J, a \in \mathbf{W}_J} c_{w,\epsilon(u),u',a,\lambda}(\tilde{T}_{u'}\tilde{T}_a1_{\epsilon(u)^{-1}\lambda}\tilde{T}_{\underline{D}}) \otimes e_i \\ &= \sum_{u' \in \mathbf{W}^J, a \in \mathbf{W}_J} c_{w,\epsilon(u),u',a,\lambda}\tilde{T}_{u'} \otimes (\tilde{T}_a1_{\epsilon(u)^{-1}\lambda}\tilde{T}_{\underline{D}}e_i) \\ &= \sum_{u' \in \mathbf{W}^J, a \in \mathbf{W}_J, i' \in X} c_{w,\epsilon(u),u',a,\lambda}\tilde{c}_{a,\epsilon(u)^{-1}\lambda,i,i'}\tilde{T}_{u'} \otimes e_{i'}. \end{aligned}$$

Hence, using (e),

$$\begin{aligned} \text{tr}(\tilde{T}_w1_\lambda\tilde{T}_{\underline{D}}, \text{IND}((E^{\tilde{\mathcal{A}}})_J)) &= \sum_{u \in \mathbf{W}^J, a \in \mathbf{W}_J, i \in X} c_{w,\epsilon(u),u,a,\lambda}\tilde{c}_{a,\epsilon(u)^{-1}\lambda,i,i} \\ &= \sum_{u \in \mathbf{W}^J, a \in \mathbf{W}_J} c_{w,\epsilon(u),u,a,\lambda}\text{tr}(\tilde{T}_a1_{\epsilon(u)^{-1}\lambda}\tilde{T}_{\underline{D}}, E^{\tilde{\mathcal{A}}}) \\ &= \sum_{u \in \mathbf{W}^J} \text{tr}(\sum_{a \in \mathbf{W}_J} c_{w,\epsilon(u),u,a,\lambda}\tilde{T}_a1_{\epsilon(u)^{-1}\lambda}\tilde{T}_{\underline{D}}, E^{\tilde{\mathcal{A}}}) \\ &= \text{tr}(p_J(\sum_{u \in \mathbf{W}^J} \tilde{T}_{u^{-1}}\tilde{T}_w1_\lambda\tilde{T}_{\epsilon(u)})\tilde{T}_{\underline{D}}, E^{\tilde{\mathcal{A}}}) \\ &= \text{tr}(\rho_{J,n}(\sum_{u \in \mathbf{W}^J} \tilde{T}_{u^{-1}}\tilde{T}_w1_\lambda\tilde{T}_{\epsilon(u)})\tilde{T}_{\underline{D}}, E^{\tilde{\mathcal{A}}}) = \text{tr}(\delta_J(\tilde{T}_w1_\lambda)\tilde{T}_{\underline{D}}, E^{\tilde{\mathcal{A}}}). \end{aligned}$$

Thus we have

$$\text{tr}(\delta_J(\tilde{T}_w1_\lambda)\tilde{T}_{\underline{D}}, E^{\tilde{\mathcal{A}}}) = \text{tr}(\tilde{T}_w1_\lambda\tilde{T}_{\underline{D}}, \text{IND}((E^{\tilde{\mathcal{A}}})_J)) = \text{tr}(\tilde{T}_w1_\lambda\tilde{T}_{\underline{D}}, (\text{ind}(E_J))^{\tilde{\mathcal{A}}})$$

where the second equality follows from 42.5(a). Since the elements \tilde{T}_w1_λ as above generate the \mathcal{A} -module $H_{n,D}$, the lemma follows.

42.7. Let \mathcal{V} be the \mathbf{Q} -vector subspace of $\mathbf{Q} \otimes \text{Hom}(\mathbf{k}^*, \mathbf{T})$ spanned by the coroots. Let $\mathcal{V}_{\mathbf{R}} = \mathbf{R} \otimes_{\mathbf{Q}} \mathcal{V}$. The kernels of the roots $\mathcal{V}_{\mathbf{R}} \rightarrow \mathbf{R}$ a hyperplane arrangement which defines a partition of $\mathcal{V}_{\mathbf{R}}$ into facets in a standard way. Let \mathcal{F} be the set of facets. Now the orbits of \mathbf{W} on \mathcal{F} are naturally indexed by the various subsets J

of \mathbf{I} . This gives a partition $\mathcal{F} = \bigsqcup_{J \subset \mathbf{I}} \mathcal{F}_J$. For example, \mathcal{F}_\emptyset consists of all Weyl chambers. If $F \in \mathcal{F}_J$, then F is homeomorphic to a real affine space of dimension $|\mathbf{I} - J|$ hence we have $H_c^i(F) = 0$ if $i \neq |\mathbf{I} - J|$ and $H_c^{|\mathbf{I} - J|}(F) = \Lambda^{|\mathbf{I} - J|}[F]$; here we write $H_c^i(?)$ instead of $H_c^i(?, \mathbf{R})$, $[F]$ denotes the vector subspace of $\mathcal{V}_{\mathbf{R}}$ in which F is open dense and $\Lambda^{|\mathbf{I} - J|}[F]$ is the top exterior power of $[F]$. Note that $[F] = \mathbf{R} \otimes_{\mathbf{Q}} ([F]_{\mathbf{Q}})$ for a well-defined \mathbf{Q} -subspace $[F]_{\mathbf{Q}}$ of \mathcal{V} . For any \underline{D} -orbit η on the set of subsets of \mathbf{I} let $\mathcal{V}_{\mathbf{R}}^\eta = \bigcup_{J \in \eta} \bigcup_{F \in \mathcal{F}_J} F \subset \mathcal{V}_{\mathbf{R}}$ and let $r_\eta = |\mathbf{I} - J|$ for some/any $J \in \eta$. We have $H_c^i(\mathcal{V}_{\mathbf{R}}^\eta) = 0$ if $i \neq r_\eta$, $H_c^{r_\eta}(\mathcal{V}_{\mathbf{R}}^\eta) = \bigoplus_{J \in \eta} \bigoplus_{F \in \mathcal{F}_J} \Lambda^{r_\eta}[F]$. Note also that $H_c^i(\mathcal{V}_{\mathbf{R}}) = 0$ if $i \neq |\mathbf{I}|$ and $H_c^{|\mathbf{I}|}(\mathcal{V}_{\mathbf{R}}) = \Lambda^{|\mathbf{I}|}\mathcal{V}_{\mathbf{R}}$. The \mathbf{W}^D -action on \mathbf{T} induces a linear action of \mathbf{W}^D on $\mathcal{V}_{\mathbf{R}}$. This action restricts for any η to a \mathbf{W}^D -action on $\mathcal{V}_{\mathbf{R}}^\eta$ and this induces a \mathbf{W}^D -action on $H_c^{r_\eta}(\mathcal{V}_{\mathbf{R}}^\eta)$. It also induces a natural \mathbf{W}^D -action on $H_c^{|\mathbf{I}|}(\mathcal{V}_{\mathbf{R}}) = \Lambda^{|\mathbf{I}|}\mathcal{V}_{\mathbf{R}}$. The long cohomology exact sequences attached to the partition $\mathcal{V}_{\mathbf{R}} = \bigcup_{\eta} \mathcal{V}_{\mathbf{R}}^\eta$ show that $(-1)^{|\mathbf{I}|} H_c^{|\mathbf{I}|}(\mathcal{V}_{\mathbf{R}}) = \sum_{\eta} (-1)^{r_\eta} H_c^{r_\eta}(\mathcal{V}_{\mathbf{R}}^\eta)$ in the Grothendieck group of representations of \mathbf{W}^D over \mathbf{R} , that is,

$$\begin{aligned} & \Lambda^{|\mathbf{I}|}\mathcal{V}_{\mathbf{R}} \oplus \bigoplus_{\eta; r_\eta = |\mathbf{I}| + 1} \bigoplus_{\text{mod } 2} \left(\bigoplus_{J \in \eta} \bigoplus_{F \in \mathcal{F}_J} \Lambda^{r_\eta}[F] \right) \\ & \cong \bigoplus_{\eta; r_\eta = |\mathbf{I}|} \bigoplus_{\text{mod } 2} \left(\bigoplus_{J \in \eta} \bigoplus_{F \in \mathcal{F}_J} \Lambda^{r_\eta}[F] \right) \end{aligned}$$

as representations of \mathbf{W}^D over \mathbf{R} . All real representations in this formula come naturally from representations of \mathbf{W}^D over \mathbf{Q} . Hence the previous formula remains valid (as representations of \mathbf{W}^D over \mathbf{Q}) if $\mathcal{V}_{\mathbf{R}}, [F]$ are replaced by $\mathcal{V}, [F]_{\mathbf{Q}}$ and the exterior powers are taken over \mathbf{Q} . Tensoring both sides (over \mathbf{Q}) by \mathfrak{U} (as in 42.5) we obtain

$$\begin{aligned} & \Lambda^{|\mathbf{I}|}\mathcal{V}_{\mathfrak{U}} \oplus \bigoplus_{\eta; r_\eta = |\mathbf{I}| + 1} \bigoplus_{\text{mod } 2} \left(\bigoplus_{J \in \eta} \bigoplus_{F \in \mathcal{F}_J} \Lambda^{r_\eta}[F]_{\mathfrak{U}} \right) \\ \text{(a)} \quad & \cong \bigoplus_{\eta; r_\eta = |\mathbf{I}|} \bigoplus_{\text{mod } 2} \left(\bigoplus_{J \in \eta} \bigoplus_{F \in \mathcal{F}_J} \Lambda^{r_\eta}[F]_{\mathfrak{U}} \right) \end{aligned}$$

as representations of \mathbf{W}^D over \mathfrak{U} ; here $\mathcal{V}_{\mathfrak{U}} = \mathfrak{U} \otimes_{\mathbf{Q}} \mathcal{V}$, $[F]_{\mathfrak{U}} = \mathfrak{U} \otimes_{\mathbf{Q}} [F]_{\mathbf{Q}}$ and the exterior powers are taken over \mathfrak{U} . We may view (a) as an isomorphism of $H_n^{D,1}$ -modules: the \mathbf{W}^D -modules in (a) may be viewed as $H_n^{D,1}$ -modules via the algebra homomorphism $H_n^{D,1} \rightarrow \mathfrak{U}[\mathbf{W}^D]$ given by $\tilde{T}_w \mapsto w$ for $w \in \mathbf{W}^D$, $1_\lambda \mapsto 0$ for $\lambda \neq \lambda_0$, $1_{\lambda_0} \mapsto 1$ (here λ_0 is the neutral element of the abelian group $\underline{\mathfrak{s}}_n$; see 28.1).

We define an \mathfrak{U} -linear map $\Delta : H_n^{D,1} \rightarrow H_n^{D,1} \otimes H_n^{D,1}$ by $\Delta(\tilde{T}_w) = \tilde{T}_w \otimes \tilde{T}_w$ for $w \in \mathbf{W}^D$ and $\Delta(1_\lambda) = \sum_{\lambda_1, \lambda_2 \in \underline{\mathfrak{s}}_n; \lambda_1 \lambda_2 = \lambda} 1_{\lambda_1} \otimes 1_{\lambda_2}$ for $\lambda \in \underline{\mathfrak{s}}_n$. (Here we use the abelian group structure on $\underline{\mathfrak{s}}_n$, a subgroup of $\underline{\mathfrak{s}}$; see 28.1.) This makes $H_n^{D,1}$ into a Hopf algebra. (Note that the analogous formulas do not make H_n^D into a Hopf algebra.) It follows that for any two $H_n^{D,1}$ -modules E_1, E_2 , the \mathfrak{U} -vector space $E_1 \otimes E_2$ is naturally an $H_n^{D,1}$ -module.

Now let E be an $H_n^{D,1}$ -module of finite dimension over \mathfrak{U} . Then we can take the tensor product of each $H_n^{D,1}$ -module in (a) with E and we obtain an isomorphism of $H_n^{D,1}$ -modules

$$E \otimes \Lambda^{|\mathbf{I}|}\mathcal{V}_{\mathfrak{U}} \oplus \bigoplus_{\eta; r_\eta = |\mathbf{I}| + 1} \bigoplus_{\text{mod } 2} X_\eta \cong \bigoplus_{\eta; r_\eta = |\mathbf{I}|} \bigoplus_{\text{mod } 2} X_\eta$$

where $X_\eta = E \otimes \bigoplus_{J \in \eta} \bigoplus_{F \in \mathcal{F}_J} \Lambda^{r_\eta}[F]_{\mathfrak{U}}$. Applying to this the functor $E \mapsto E^{\tilde{\mathcal{A}}}$ (see 42.5), we deduce an isomorphism of $H_n^{D, \tilde{\mathcal{A}}}$ -modules

$$(E \otimes \Lambda^{|\mathbf{I}|} \mathcal{V}_{\mathfrak{U}})^{\tilde{\mathcal{A}}} \oplus \bigoplus_{\eta; r_\eta = |\mathbf{I}|+1 \pmod 2} X_\eta^{\tilde{\mathcal{A}}} \cong \bigoplus_{\eta; r_\eta = |\mathbf{I}| \pmod 2} X_\eta^{\tilde{\mathcal{A}}}.$$

We deduce that for $y \in H_{n,D}$ we have

$$(b) \quad \text{tr}(y\tilde{T}_{\underline{D}}, (E \otimes \Lambda^{|\mathbf{I}|} \mathcal{V}_{\mathfrak{U}})^{\tilde{\mathcal{A}}}) = \sum_{\eta} (-1)^{r_\eta + |\mathbf{I}|} \text{tr}(y\tilde{T}_{\underline{D}}, X_\eta^{\tilde{\mathcal{A}}}).$$

We have $X_\eta = \bigoplus_{J \in \eta} X^J$ where $X^J = E \otimes (\bigoplus_{F \in \mathcal{F}_J} \Lambda^{r_\eta}[F]_{\mathfrak{U}})$.

Assume first that η consists of at least two subsets of \mathbf{I} . Then each X_J is stable under $H_n^{D,1}$ and is mapped by $\tilde{T}_{\underline{D}}$ into $X_{J'}$ with $J \neq J'$. From the definitions we have $X_\eta^{\tilde{\mathcal{A}}} = \bigoplus_{J \in \eta} \tilde{\mathcal{A}} \otimes_{\mathfrak{U}} X_J$ as an $\tilde{\mathcal{A}}$ -module and each summand $\tilde{\mathcal{A}} \otimes_{\mathfrak{U}} X_J$ is stable under H_n and is mapped by $\tilde{T}_{\underline{D}}$ into a summand $\tilde{\mathcal{A}} \otimes_{\mathfrak{U}} X_{J'}$ with $J \neq J'$. It follows that for our η we have

$$(c) \quad \text{tr}(y\tilde{T}_{\underline{D}}, X_\eta^{\tilde{\mathcal{A}}}) = 0.$$

Next assume that η consists of a single subset J of \mathbf{I} . We have $\underline{D}(J) = J$. Let F_J be the unique facet in \mathcal{F}_J such that F_J is contained in the closure of the dominant Weyl chamber. Then F_J is stable under the the subgroup \mathbf{W}_J^D of \mathbf{W}^D generated by \mathbf{W}_J and \underline{D} and X_η may be identified with $E \otimes (H_n^{D,1} \otimes_{H_{J,n}^{D,1}} (\Lambda^{|\mathbf{I}-J|}[F_J]_{\mathfrak{U}}))$. Here $\Lambda^{|\mathbf{I}-J|}[F_J]_{\mathfrak{U}}$ is regarded as a WW_J^D -module and then is viewed as a $H_{J,n}^{D,1}$ -module via the canonical algebra homomorphism $H_{J,n}^{D,1} \rightarrow \mathfrak{U}[\mathbf{W}_J^D]$; thus 1_λ acts on it as 1 if $\lambda = \lambda_0$ and as 0 if $\lambda \neq \lambda_0$. Note that in the \mathbf{W}_J^D -module $\Lambda^{|\mathbf{I}-J|}[F_J]_{\mathfrak{U}}$, \mathbf{W}_J acts trivially (since \mathbf{W}_J acts trivially on $[F_J]_{\mathfrak{U}}$) and \underline{D} acts as multiplication by $(-1)^{|\mathbf{I}-J| - |(\mathbf{I}-J)^\epsilon|}$. Let $X'_\eta = E \otimes (H_n^{D,1} \otimes_{H_{J,n}^{D,1}} \mathfrak{U})$ where \mathfrak{U} is regarded as a $H_{J,n}^{D,1}$ -module coming from the trivial representation of \mathbf{W}_J^D . We see that we may identify X_η, X'_η in a way compatible with the H_n^1 -module structures and so that the action of $\tilde{T}_{\underline{D}}$ on X_η corresponds to $(-1)^{|\mathbf{I}-J| - |(\mathbf{I}-J)^\epsilon|}$ times the action of $\tilde{T}_{\underline{D}}$ on X'_η . Using the definitions we see that we may identify $X_\eta^{\tilde{\mathcal{A}}}, X'^{\tilde{\mathcal{A}}}$ in a way compatible with the H_n -module structures and so that the action of $\tilde{T}_{\underline{D}}$ on $X_\eta^{\tilde{\mathcal{A}}}$ corresponds to $(-1)^{|\mathbf{I}-J| - |(\mathbf{I}-J)^\epsilon|}$ times the action of $\tilde{T}_{\underline{D}}$ on $X'^{\tilde{\mathcal{A}}}$. From the definitions we have $X'_\eta = \text{ind}(E_J)$ (notation of 42.5). We see that for our η we have

$$(d) \quad \text{tr}(y\tilde{T}_{\underline{D}}, X_\eta^{\tilde{\mathcal{A}}}) = (-1)^{|\mathbf{I}-J| - |(\mathbf{I}-J)^\epsilon|} \text{tr}(y\tilde{T}_{\underline{D}}, (\text{ind}(E_J))^{\tilde{\mathcal{A}}}).$$

From the definitions (34.4) we see that there is a unique $\tilde{\mathcal{A}}$ -algebra homomorphism $\vartheta : H_n^{D, \tilde{\mathcal{A}}} \rightarrow H_n^{D, \tilde{\mathcal{A}}}$ such that

$$\begin{aligned} \vartheta(1_\lambda) &= 1_\lambda \text{ for any } \lambda \in \underline{\mathfrak{s}}_n, \\ \vartheta(\tilde{T}_w) &= (-1)^{l(w)} \tilde{T}_{w^{-1}}^{-1} \text{ for any } w \in \mathbf{W}, \\ \vartheta(\tilde{T}_{\underline{D}}) &= (-1)^{|\mathbf{I}| - |\mathbf{I}^\epsilon|} \tilde{T}_{\underline{D}}. \end{aligned}$$

We have $\vartheta^2 = 1$.

Using ϑ and $E^{\tilde{\mathcal{A}}}$ we can define a new $H_n^{D, \tilde{\mathcal{A}}}$ -module $E^{\tilde{\mathcal{A}}, \vartheta}$ with the same underlying $\tilde{\mathcal{A}}$ -module as $E^{\tilde{\mathcal{A}}}$ but with $x \in H_n^{D, \tilde{\mathcal{A}}}$ acting on $E^{\tilde{\mathcal{A}}, \vartheta}$ in the same way that $\vartheta(x)$ acts on $E^{\tilde{\mathcal{A}}}$. We show that

(e) under extension of scalars from $\tilde{\mathcal{A}}$ to $\mathfrak{U}(v)$, the $H_n^{D,\tilde{\mathcal{A}}}$ -modules $E^{\tilde{\mathcal{A}},\vartheta}$ and $(E \otimes \Lambda^{|\mathbf{I}|}\mathcal{V}_{\mathfrak{U}})^{\tilde{\mathcal{A}}}$ become isomorphic $\mathfrak{U}(v) \otimes_{\mathcal{A}} H_n^D$ -modules.

As in the proof of 42.5(a) it is enough to show that these $H_n^{D,\tilde{\mathcal{A}}}$ -modules become isomorphic $H_n^{D,1}$ -modules under the specialization $v = 1$. Thus it is enough to show that $E^{\tilde{\mathcal{A}},\vartheta}|_{v=1} \cong E \otimes \Lambda^{|\mathbf{I}|}\mathcal{V}_{\mathfrak{U}}$ as $H_n^{D,1}$ -modules. Now the underlying \mathfrak{U} -vector space of $E^{\tilde{\mathcal{A}},\vartheta}|_{v=1}$ is E but the action of $x \in H_n^{D,1}$ on $E^{\tilde{\mathcal{A}},\vartheta}|_{v=1}$ is the same as the action of $\vartheta_1(x)$ on E . Here $\vartheta_1 : H_n^{D,1} \rightarrow H_n^{D,1}$ is the specialization of ϑ_1 for $v = 1$. Note that $\vartheta_1(1_\lambda) = 1_\lambda$ for any $\lambda \in \underline{\mathfrak{g}}_n$ and $\vartheta_1(\tilde{T}_w) = \gamma_w \tilde{T}_w$ for any $w \in \mathbf{W}^D$, where $\gamma_w = \pm 1$ is the scalar by which w acts in the \mathbf{W}^D -module $\Lambda^{|\mathbf{I}|}\mathcal{V}_{\mathfrak{U}}$. The desired result follows.

Combining (b),(c),(d),(e) we see that for any $y \in H_{n,D}$ we have

$$(-1)^{|\mathbf{I}|+|\mathbf{I}_\epsilon|} \text{tr}(\vartheta(y\tilde{T}_{\underline{D}}), E^{\tilde{\mathcal{A}}}) = \sum_{J \subset \mathbf{I}; \epsilon(J)=J} (-1)^{|J_\epsilon|} \text{tr}(y\tilde{T}_{\underline{D}}, (\text{ind}(E_J))^{\tilde{\mathcal{A}}}).$$

Replacing here $(-1)^{|\mathbf{I}|+|\mathbf{I}_\epsilon|} \vartheta(y\tilde{T}_{\underline{D}})$ by $\vartheta(y)\tilde{T}_{\underline{D}}$ and using Lemma 42.6 we may rewrite this as

$$\text{tr}(\vartheta(y)\tilde{T}_{\underline{D}}, E^{\tilde{\mathcal{A}}}) = \sum_{J \subset \mathbf{I}; \epsilon(J)=J} (-1)^{|J_\epsilon|} \text{tr}(\delta_J(y)\tilde{T}_{\underline{D}}, E^{\tilde{\mathcal{A}}})$$

or equivalently (see 42.4) $\text{tr}(\vartheta(y)\tilde{T}_{\underline{D}}, E^{\tilde{\mathcal{A}}}) = \text{tr}(\delta(y)\tilde{T}_{\underline{D}}, E^{\tilde{\mathcal{A}}})$. Since any simple $\mathfrak{U}(v) \otimes_{\mathcal{A}} H_n^D$ -module can be obtained by extension of scalars (from $\tilde{\mathcal{A}}$ to $\mathfrak{U}(v)$) from some $E^{\tilde{\mathcal{A}}}$ as above, we deduce that

$$\text{tr}((\delta(y) - \vartheta(y))\tilde{T}_{\underline{D}}, \mathbf{E}) = 0$$

for any simple $\mathfrak{U}(v) \otimes_{\mathcal{A}} H_n^D$ -module \mathbf{E} . Since $\mathfrak{U}(v) \otimes_{\mathcal{A}} H_n^D$ is a semisimple algebra, it follows that $(\delta(y) - \vartheta(y))\tilde{T}_{\underline{D}}$ belongs to the $\mathfrak{U}(v)$ -subspace of $\mathfrak{U}(v) \otimes_{\mathcal{A}} H_n^D$ spanned by commutators $xx' - x'x$ with $x, x' \in \mathfrak{U}(v) \otimes_{\mathcal{A}} H_n^D$. Hence we have

$$g(\delta(y) - \vartheta(y))\tilde{T}_{\underline{D}} = \sum_{i=1}^m g_i(x_i \tilde{T}_{\underline{D}}^{s_i} x'_i \tilde{T}_{\underline{D}}^{1-s_i} - x'_i \tilde{T}_{\underline{D}}^{1-s_i} x_i \tilde{T}_{\underline{D}}^{s_i})$$

with $g \in \mathcal{A} - \{0\}$, $g_i \in \mathcal{A}$, $x_i \in H_n$, $x'_i \in H_n$, $s_i \in \mathbf{Z}$, that is,

$$(f) \quad g(\delta(y) - vt(y)) = \sum_{i=1}^m g_i(x_i \tilde{T}_{\underline{D}}^{s_i} x'_i \tilde{T}_{\underline{D}}^{-s_i} - x'_i \tilde{T}_{\underline{D}}^{1-s_i} x_i \tilde{T}_{\underline{D}}^{s_i-1}).$$

42.8. We show that for any $y, y' \in H_n$ we have

$$(a) \quad \mathfrak{f}\eta\tilde{\omega}(yy' - y'\tilde{T}_{\underline{D}}y\tilde{T}_{\underline{D}}^{-1}) = 0.$$

Let $w \in \mathbf{W}$, $\lambda \in \underline{\mathfrak{g}}_n$. Let $\mathcal{L} \in \lambda$. If $w\underline{D}\lambda = \lambda$, using notation and results in 31.6, 31.7, we have

$$\begin{aligned} \mathfrak{f}\eta\tilde{\omega}(v^{l(w)}\tilde{T}_w 1_{\underline{D}\lambda}) &= gr(K_{\mathbf{I},\underline{D}}^{w,\mathcal{L}}) \\ &= \sum_A \chi_v^A(K_{\mathbf{I},\underline{D}}^{w,\mathcal{L}}) = \sum_A \tilde{\zeta}^A(v^{l(w)}\tilde{T}_w 1_{\underline{D}\lambda}\tilde{T}_{\underline{D}}) = \sum_A \zeta^A(v^{l(w)}\tilde{T}_w 1_{\underline{D}\lambda}\tilde{T}_{\underline{D}}) \end{aligned}$$

(the last equation comes from 31.7(e); A runs over the objects in \hat{D} up to isomorphism such that $\zeta^A \neq 0$.) The equation

$$\mathfrak{f}\eta\tilde{\omega}(v^{l(w)}\tilde{T}_w 1_{\underline{D}\lambda}) = \sum_A \zeta^A(v^{l(w)}\tilde{T}_w 1_{\underline{D}\lambda}\tilde{T}_{\underline{D}})$$

holds also if $w\underline{D}\lambda \neq \lambda$ (in this case both sides are 0). It follows that

$$\mathfrak{f}\eta\tilde{\omega}(x) = \sum_A \zeta^A(x\tilde{T}_D) \text{ for any } x \in H_n.$$

We deduce

$$\mathfrak{f}\eta\tilde{\omega}(yy' - y'\tilde{T}_D y\tilde{T}_D^{-1}) = \sum_A (\zeta^A(yy'\tilde{T}_D) - \zeta^A(y\epsilon(y)\tilde{T}_D)) = 0$$

where the last equality follows from 31.8. This proves (a).

Proposition 42.9. *Let $y \in H$. We have $\mathbf{d}(\mathfrak{f}\eta\tilde{\omega}(y)) = \mathfrak{f}\eta\tilde{\omega}(\vartheta(y)) \in \mathfrak{K}(D)$ with $\mathbf{d} : \mathfrak{K}(D) \rightarrow \mathfrak{K}(\Delta)$ as in 42.2.*

If $y \in H'_D$ (see 42.3), both sides of the desired equality are 0. (Note that ϑ maps H_D into itself and H'_D into itself.) Hence we may assume that $y \in H_D$. We can assume that $y \in H_n$ where $n \in \mathbf{N}_k^*$. Then $y \in H_{n,D}$. By 42.4 it is enough to show that $\mathfrak{f}\eta\tilde{\omega}(\delta(y) - \vartheta(y)) = 0$. Let g, g_i, x_i, x'_i, s_i be as in 42.7(f). Since $g \neq 0$, it is enough to show that $g\mathfrak{f}\eta\tilde{\omega}(\delta(y) - \vartheta(y)) = 0$ or that $\mathfrak{f}\eta\tilde{\omega}(g(\delta(y) - \vartheta(y))) = 0$. Using 42.7 it is enough to show that

$$\mathfrak{f}\eta\tilde{\omega}\left(\sum_{i=1}^m g_i(x_i\tilde{T}_D^{s_i}x'_i\tilde{T}_D^{-s_i} - x'_i\tilde{T}_D^{1-s_i}x_i\tilde{T}_D^{s_i-1})\right) = 0.$$

Hence it is enough to show that

$$\mathfrak{f}\eta\tilde{\omega}(x\tilde{T}_D^s x'\tilde{T}_D^{-s} - x'\tilde{T}_D^{1-s} x\tilde{T}_D^{s-1}) = 0$$

for any $x, x' \in H_n$ and any $s \in \mathbf{Z}$. We have

$$x\tilde{T}_D^s x'\tilde{T}_D^{-s} - x'\tilde{T}_D^{1-s} x\tilde{T}_D^{s-1} = (z - \tilde{T}_D^{-s} z\tilde{T}_D^s) + (z'x' - x'\tilde{T}_D z'\tilde{T}_D^{-1})$$

where $z = x\tilde{T}_D^s x'\tilde{T}_D^{-s} \in H_n$ and $z' = \tilde{T}_D^{-s} x\tilde{T}_D^s \in H_n$. Hence it is enough to show that $\mathfrak{f}\eta\tilde{\omega}(z'x' - x'\tilde{T}_D z'\tilde{T}_D^{-1}) = 0$ (see 42.8(a)) and

$$(a) \mathfrak{f}\eta\tilde{\omega}(z - \tilde{T}_D^{-s} z\tilde{T}_D^s) = 0$$

for any $z \in H_n$. This follows from 41.6(c).

REFERENCES

- [BBD] A. Beilinson, J. Bernstein, P. Deligne, *Faisceaux pervers*, Astérisque **100** (1982). MR0751966 (86g:32015)
- [De] P. Deligne, *La conjecture de Weil*, II, Publ.Math. I.H.E.S. **52** (1980), 137-252. MR0601520 (83c:14017)
- [G] V. Ginzburg, *Admissible modules on a symmetric space*, Astérisque **173-174** (1989), 199-255. MR1021512 (91c:22030)
- [Gr] I. Grojnowski, *Character sheaves on symmetric spaces*, Ph.D. thesis, MIT, 1992.
- [L3] G. Lusztig, *Character sheaves*, I, Adv. Math. **56** (1985), 193-237; II, vol. 57, 1985, pp. 226-265; III, vol. 57, 1985, pp. 266-315; IV, vol. 59, 1986, pp. 1-63; V, vol. 61, 1986, pp. 103-155. MR0792706 (87b:20055); MR0806210 (87m:20118a); MR0825086 (87m:20118b); MR0866163 (87m:20118d)
- [L9] G. Lusztig, *Character sheaves on disconnected groups*, I, Represent. Theory. (electronic) **7** (2003), 374-403; II, vol. 8, 2004, pp. 72-124; III, vol. 8, 2004, pp. 125-144; IV, vol. 8, 2004, pp. 145-178; Errata, vol. 8, 2004, pp. 179-179; V, vol. 8, 2004, pp. 346-376; VI, vol. 8, 2004, pp. 377-413; VII, vol. 9, 2005, pp. 209-266; VIII, vol. 10, 2006. MR2017063 (2006d:20090a); MR2048588 (2006d:20090b); MR2048589 (2006d:20090c); MR2048590 (2006d:20090d); MR2077486 (2005h:20111); MR2084488 (2005h:20112); MR2133758 (2006e:20089)
- [L14] G. Lusztig, *Characters of reductive groups over a finite field*, Ann. Math. Studies, vol. 107, Princeton Univ. Press, 1984. MR0742472 (86j:20038)

- [L10] G. Lusztig, *Parabolic character sheaves*, I, *Moscow Math. J.* **4** (2004), 153-179. MR2074987 (2006d:20091a)
- [MV] I. Mirković, K. Vilonen, *Characteristic varieties of character sheaves*, *Invent. Math.* **93** (1988), 405-418. MR0948107 (89i:20066)

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE,
MASSACHUSETTS 02139