

## REGULARITY OF A FREE BOUNDARY IN PARABOLIC POTENTIAL THEORY

LUIS CAFFARELLI, ARSHAK PETROSYAN, AND HENRIK SHAHGHOLIAN

### CONTENTS

1. Introduction	828
1.1. Background	828
1.2. Problem	829
1.3. Notation	829
1.4. Local solutions	829
1.5. Global solutions	830
1.6. Scaling	830
1.7. Blow-up	830
2. Main results	830
2.1. Examples	830
2.2. Main theorems	832
3. Monotonicity formulas	833
4. Uniform $C_x^{1,1} \cap C_t^{0,1}$ regularity of solutions	836
5. Nondegeneracy	839
5.1. Nondegeneracy	839
5.2. Stability under the limit	841
5.3. Lebesgue measure of $\partial\Omega$	841
6. Homogeneous global solutions	842
7. Balanced energy	844
7.1. Zero energy points	845
7.2. High energy points	845
7.3. Low energy points	846
8. Positive global solutions	849
9. Classification of global solutions	851
10. Proof of Theorem I	861
11. Lipschitz regularity: global solutions	861
12. Balanced energy: local solutions	864
13. Lipschitz regularity: local solutions	864
14. $C^{1,\alpha}$ regularity	866
15. Higher regularity	867
16. Proof of Theorem II	868
References	868

---

Received by the editors December 20, 2002.

2000 *Mathematics Subject Classification*. Primary 35R35.

*Key words and phrases*. Free boundary problems, Stefan problem, regularity, global solutions, monotonicity formulas.

The first author was supported in part by the NSF.

The second author thanks the Göran Gustafsson Foundation and the Department of Mathematics, Royal Institute of Technology, for the visiting appointment.

The third author was supported in part by the Swedish Research Council.

## 1. INTRODUCTION

**1.1. Background.** In the last few years the free boundary regularity of both variational and nonvariational types has gained renewed attention. Due to developments of the so-called monotonicity formulas for elliptic and parabolic PDEs on one hand and developments of new techniques in free boundary regularity on the other hand several longstanding questions have been answered.

One of these questions, treated in this paper and with roots in parabolic potential theory, concerns the nature of those boundaries that allow caloric continuation of the heat potential from the free space into the space occupied by the density function. To clarify this, let  $U^f$  be the heat potential of a density function  $f$ :

$$U^f(x, t) = \int_{\mathbb{R}^n \times \mathbb{R}} f(y, s) G(x - y, s - t) dy ds,$$

where  $G(x, t)$  is the heat kernel. Then it is known that

$$HU^f = c_n f,$$

where  $H = \Delta - \partial_t$  is the heat operator and  $c_n < 0$  is some constant. Now suppose

$$f(x, t) = \frac{1}{c_n} \chi_\Omega$$

for some domain  $\Omega$  and denote the corresponding potential by  $U^\Omega$ . Then

$$HU^\Omega = \chi_\Omega.$$

Suppose now that there exist  $v$  such that

$$\begin{cases} Hv = 0 & \text{in } Q_r(x_0, t_0), \\ v = U^\Omega & \text{in } Q_r(x_0, t_0) \setminus \Omega \end{cases}$$

for some  $(x_0, t_0) \in \partial\Omega$  and  $r > 0$ , where  $Q_r(x_0, t_0) = B_r(x_0) \times (t_0 - r^2, t_0 + r^2)$ . Then we call  $v$  a caloric continuation of  $U^\Omega$ . Moreover, the function

$$u = U^\Omega - v$$

satisfies

$$(1.1) \quad \begin{cases} Hu = \chi_\Omega & \text{in } Q_r, \\ u = |\nabla u| = 0 & \text{in } Q_r \setminus \Omega. \end{cases}$$

So our question is, When does the boundary of a domain allow a caloric continuation of the potential?

It is well known, through the Cauchy-Kowalevskaya theorem, that analytic boundaries do allow such a continuation locally. Hence we ask the reverse of the Cauchy-Kowalevskaya theorem in the sense that the existence of the caloric continuation implies the regularity of the boundary.

In a particular case when  $u \geq 0$  and  $\partial_t u \geq 0$  problem (1.1) is the well-known *Stefan problem* (see e.g. [Fri88]), describing the melting of ice, and is treated extensively in the literature. However, even the variational inequality case  $u \geq 0$  (and not necessarily  $\partial_t u \geq 0$ ) has not been considered earlier.

In this paper we treat (1.1) in its full generality without any sign assumptions on either  $u$  or  $\partial_t u$ . The stationary case, i.e. when  $u$  is independent of  $t$ , was studied in [CKS00]. The results of this paper generalize those of [CKS00] to the time dependent case.

**1.2. Problem.** For a function  $u(x, t)$ , continuous with its spatial derivatives in a domain  $D$  of  $\mathbb{R}^n \times \mathbb{R}$ , define the *coincidence set* as

$$\Lambda := \{u = |\nabla u| = 0\}$$

and suppose that

$$(1.2) \quad Hu = \chi_\Omega \quad \text{in } D, \quad \Omega := D \setminus \Lambda.$$

Here  $H = \Delta - \partial_t$  is the heat operator and we assume that the equation is satisfied in the weak (distributional) sense, i.e.

$$\int_D u(\Delta\eta + \partial_t\eta) \, dxdt = \int_{D \cap \Omega} \eta \, dxdt$$

for all  $C^\infty$  test functions  $\eta$  with compact support in  $D$ . Then we are interested in the regularity of the so-called *free boundary*  $\Gamma$ , which consists of all  $(x, t) \in \partial\Omega \cap D$ , that are not parabolically interior for  $\Lambda$ , i.e. such that

$$Q_\varepsilon^-(x, t) \cap \Omega \neq \emptyset$$

for any small  $\varepsilon > 0$ , where  $Q_\varepsilon^-(x, t) = B_\varepsilon(x) \times (t - \varepsilon^2, t]$  is the lower parabolic cylinder.

**1.3. Notation.** Points in  $\mathbb{R}^n \times \mathbb{R}$  are denoted by  $(x, t)$ , where  $x \in \mathbb{R}^n$  and  $t \in \mathbb{R}$ .

Generic constants are denoted by  $C, C_0, C_n, \dots$ ;

$\mathbb{R}_a^- = (-\infty, a]$ ;  $\mathbb{R}^- = \mathbb{R}_0^-$ ;

$a_\pm = \max(\pm a, 0)$  for any  $a \in \mathbb{R}$ ;

$B_r(x)$  is the open ball in  $\mathbb{R}^n$  with center  $x$  and radius  $r$ ;  $B_r = B_r(0)$ ;

$Q_r(x, t) = B(x, r) \times (t - r^2, t + r^2)$  (parabolic cylinder);  $Q_r = Q_r(0, 0)$ ;

$Q_r^+(x, t) = B_r(x) \times [t, t + r^2)$  (the upper half-cylinder);  $Q_r^+ = Q_r^+(0, 0)$ ;

$Q_r^-(x, t) = B_r(x) \times (t - r^2, t]$  (the lower half-cylinder);  $Q_r^- = Q_r^-(0, 0)$ ;

$\partial_p Q_r(x, t)$  is the parabolic boundary, i.e. the topological boundary minus the top of the cylinder;

$\nabla$  denotes the spatial gradient,  $\nabla = (\partial_1, \dots, \partial_n)$ ;

$\Delta = \sum_{i=1}^n \partial_{ii}$  (the spatial Laplacian);

$H = \Delta - \partial_t$  (the heat operator);

$\chi_\Omega$  is the characteristic function of the set  $\Omega$ ;

$E(t) = \{x : (x, t) \in E\}$  is the  $t$ -section of the set  $E$  in  $\mathbb{R}^n \times \mathbb{R}$ .

Below we define classes of local and global solutions of (1.2) that we study in this paper.

#### 1.4. Local solutions.

**Definition 1.1.** For given  $r, M > 0$  and  $(x_0, t_0) \in \mathbb{R}^n \times \mathbb{R}$  let  $\mathcal{P}_r^-(x_0, t_0; M)$  be the class of functions  $u$  in  $Q^- = Q_r^-(x_0, t_0)$  such that

- (i)  $u$  satisfies (1.2) in  $D = Q^-$ ;
- (ii)  $|u| \leq M$  in  $Q^-$ ;
- (iii)  $(x_0, t_0) \in \Lambda$ .

In the case  $(x_0, t_0) = (0, 0)$  we will denote the corresponding class  $\mathcal{P}_r^-(0, 0; M)$  also by  $\mathcal{P}_r^-(M)$ .

Similarly, define the class  $\mathcal{P}_r(x, t; M)$  by replacing  $Q^- = Q_r^-(x_0, t_0)$  with  $Q = Q_r(x_0, t_0)$  in (i)–(iii) above.

### 1.5. Global solutions.

**Definition 1.2.** For a given  $M > 0$  let  $\mathcal{P}_\infty^-(M)$  be the class of functions  $u$  in  $\mathbb{R}^n \times \mathbb{R}^-$  such that

- (i)  $u$  satisfies (1.2) in  $D = \mathbb{R}^n \times \mathbb{R}^-$ ;
- (ii)  $|u(x, t)| \leq M(1 + |x|^2 + |t|)$ ;
- (iii)  $(0, 0) \in \Lambda$ .

Similarly, define the class  $\mathcal{P}_\infty(M)$  by replacing  $\mathbb{R}^n \times \mathbb{R}^-$  with  $\mathbb{R}^n \times \mathbb{R}$ .

The elements of  $\mathcal{P}_\infty^-(M)$  and  $\mathcal{P}_\infty(M)$  will be called *global solutions*.

It is also noteworthy that elements in  $\mathcal{P}_\infty^-(M)$  can be extended, in a natural way, to  $\mathbb{R}^n \times \mathbb{R}^+$  by solving the Cauchy problem for the equation  $Hu = 1$ . In particular, we may consider each element in  $\mathcal{P}_\infty^-(M)$  as an element of  $\mathcal{P}_\infty(M)$ , and vice versa.

The following operations will be extensively used throughout the paper.

**1.6. Scaling.** For a function  $u(x, t)$  set

$$u_r(x, t) = \frac{1}{r^2} u(rx, r^2t),$$

the *parabolic scaling* of  $u$  around  $(0, 0)$ . This scaling preserves equation (1.2) with

$$\Omega(u_r) = \Omega_r := \{(x, t) : (rx, r^2t) \in \Omega\}.$$

Also,  $u \in \mathcal{P}_r(M)$  implies  $u_r \in \mathcal{P}_1(M/r^2)$ .

Similarly, one can scale  $u$  around any point  $(x_0, t_0)$  by

$$\frac{1}{r^2} u(rx + x_0, r^2t + t_0).$$

**1.7. Blow-up.** As we show in Theorem 4.1, solutions  $u \in \mathcal{P}_1^-(M)$  are locally  $C_x^{1,1} \cap C_t^{0,1}$  regular in  $Q_1^-$ . Then the scaled functions  $u_r$  are defined and uniformly bounded in  $Q_R^-$  for any  $R < 1/r$ . Since  $Hu_r = \chi_{\Omega_r}$ , by standard compactness methods in parabolic theory (see e.g. [Fri64]), we may let  $r \rightarrow 0$  and obtain (for a subsequence) a global solution (see the stability discussion below). This process is referred to as *blowing-up*, and the global solution thus obtained is called a *blow-up* of  $u$ .

Similarly, we can define the blow-up of a local solution  $u$  at any free boundary point  $(x_0, t_0)$  by considering the parabolic scalings of  $u$  around  $(x_0, t_0)$ .

Also, if  $u$  is a global solution, we can define the blow-up at infinity, by considering the scaled functions  $u_r$  and letting  $r \rightarrow \infty$ . The blow-up at infinity will be called *shrink-down*.

## 2. MAIN RESULTS

Before stating our main results, we would like to illustrate the problem with the following examples.

### 2.1. Examples.

**1. Stationary** (i.e.  $t$ -independent) *solutions*. These include *halfspace solutions*

$$u(x, t) = \frac{1}{2}(x \cdot e)_+^2,$$

where  $e$  is a spatial unit vector, as well as other global stationary solutions of the obstacle problem that have ellipsoids and paraboloids as coincidence sets.

**2. Space-independent (i.e.  $x$ -independent) solutions**

$$u(x, t) = -t, \quad u(x, t) = -t_+, \quad u(x, t) = t_-.$$

In fact, it is easy to see that the solutions depending only on  $t$  have the form

$$u(x, t) = \begin{cases} -(t - T_2), & t > T_2, \\ 0, & T_1 \leq t \leq T_2, \\ -(t - T_1), & t < T_1 \end{cases}$$

for some constants  $-\infty \leq T_1 \leq T_2 \leq \infty$ . This is a particular case of our Theorem I below.

**3. Polynomial solutions of the type**

$$u(x, t) = P(x) + m t,$$

where  $P(x)$  is a quadratic polynomial satisfying  $\Delta P = m + 1$ . In particular, for a given constant  $c$ , the function

$$u(x, t) = c|x|^2 + (2nc - 1)t$$

is a solution of (1.2) in  $\mathbb{R}^n \times \mathbb{R}$ . The only free boundary point of this solution is the origin  $(0, 0)$ , unless  $c = 0$  or  $c = 1/2n$ . In the former case the free boundary is  $\mathbb{R}^n \times \{0\}$  and in the latter case it is  $\{0\} \times \mathbb{R}$ .

**4.** For the next example we modify the solution above for  $t \geq 0$  by solving a one phase free boundary problem: find a function  $f(\xi)$  on  $[0, \infty)$  such that

$$(2.1) \quad u(x, t) = t f\left(\frac{|x|}{\sqrt{t}}\right)$$

satisfies (1.2) for  $t > 0$ . This will be so if  $f$  vanishes on  $[0, a]$  for some  $a > 0$  and satisfies an ordinary differential equation

$$f''(\xi) + \left(\frac{n-1}{\xi} + \frac{\xi}{2}\right) f'(\xi) - f(\xi) - 1 = 0$$

on  $(a, \infty)$  with boundary conditions

$$f(a) = f'(a) = 0.$$

The solution can be given explicitly as

$$f(\xi) = (2n + \xi^2) \left( \frac{1}{2n + a^2} - 2a^n e^{a^2/4} \int_a^\xi \frac{e^{-s^2/4}}{s^{n-1}(2n + s^2)^2} ds \right).$$

It is easy to see that the limit

$$(2.2) \quad c = \lim_{\xi \rightarrow \infty} \frac{f(\xi)}{\xi^2}$$

exists and satisfies

$$0 < c < \frac{1}{2n}.$$

Moreover, changing  $a$  between 0 and  $\infty$ , we can get all values from  $(0, 1/2n)$ . Now (2.2) implies that

$$u(x, 0+) = c|x|^2.$$

Hence, if we define

$$u(x, t) = \begin{cases} t f\left(\frac{|x|}{\sqrt{t}}\right) & \text{for } t > 0, \\ c|x|^2 + (2nc - 1)t & \text{for } t \leq 0, \end{cases}$$

we again obtain a solution of (1.2) in  $\mathbb{R}^n \times \mathbb{R}$ . The free boundary in this case is the paraboloid  $\{|x|^2 = a^2t\}$ . The solution  $u$  is identically 0 inside and positive outside.

5. Finally, we point out that at any time  $t = T$  we have the freedom to choose not to have a free boundary. Namely, fix  $T > 0$  and let  $u$  be, for instance, as in the previous example. Now solve the Cauchy problem

$$Hv = 1 \quad \text{in } \mathbb{R}^n \times (T, \infty); \quad v(\cdot, T) = u(\cdot, T)$$

and let

$$w(x, t) = \begin{cases} v(x, t), & t > T, \\ u(x, t), & t \leq T. \end{cases}$$

Then  $w$  is a solution of (1.2) in  $\mathbb{R}^n \times \mathbb{R}$ . Its free boundary is the truncation of the paraboloid  $\{|x|^2 = a^2t\}$  for  $t \leq T$ . We remark that the disk  $B_{a^2T} \times \{T\}$  is not a part of the free boundary, even though it is a part of  $\partial\Omega$ .

As we will see later (Section 7), the points on  $B_{a^2T} \times \{T\}$  have zero (balanced) energy, the tip  $(0, 0)$  has high energy, and the rest of the points on the truncated paraboloid have low energy. We show in this paper that, in a sense, the regular free boundary points are the ones with low energy.

**2.2. Main theorems.** The solution that we constructed in the example above has the property that it is polynomial for  $t < 0$ , nonnegative and convex in space for  $0 \leq t \leq T$  and solves  $Hw = 1$  for  $t > T$ . Our first main theorem states that something similar is true for every global solution.

**Theorem I** (Classification of global solutions). *Let  $u$  be a solution of (1.2) in  $D = \mathbb{R}^n \times (-\infty, a]$  with at most quadratic growth at infinity:*

$$|u(x, t)| \leq M (|x|^2 + |t| + 1)$$

for some constant  $M > 0$ . Then there exist  $-\infty \leq T_1 \leq T_2 \leq a$  with the following properties:

(i) if  $-\infty < T_1$ , then

$$u(x, t) = P(x) + mt \quad \text{for } t < T_1$$

for a quadratic polynomial  $P(x)$  and a constant  $m$ ;

(ii) if  $T_1 < T_2$ , then

$$u \geq 0, \quad \partial_{ee}u \geq 0, \quad \partial_t u \leq 0 \quad \text{for } t < T_2,$$

where  $e$  is any spatial unit vector;

(iii) if  $T_2 < a$ , then  $u$  satisfies

$$Hu = 1 \quad \text{for } T_2 < t < a.$$

As in the case of the obstacle problem (see [Caf98]) the classification of global solutions implies the regularity of the free boundary for local solutions at points satisfying a certain density condition. Such a condition can be given in terms of the minimal diameter.

**Definition 2.1** (Minimal diameter). The *minimal diameter* of a set  $E$  in  $\mathbb{R}^n$ , denoted  $\text{md}(E)$ , is the infimum of distances between two parallel planes such that  $E$  is contained in the strip between these planes. The *lower density function* for the solution of  $u$  of (1.2) at  $(0, 0)$  is defined by

$$\delta_r^-(u) = \frac{\text{md}(\Lambda(-r^2) \cap B_r)}{r}.$$

**Theorem II** (Regularity of local solutions). *Let  $u \in \mathcal{P}_1^-(M)$  be a local solution, such that  $(0, 0) \in \Gamma$ . Then there is a universal modulus of continuity  $\sigma(r)$  and a constant  $c > 0$  such that if for one value of  $r$ , say  $r_0$ , we have*

$$\delta_{r_0}^-(u) > \sigma(r_0),$$

*then  $\Gamma \cap Q_{cr_0}^-$  is a  $C^\infty$  surface (in space and time).*

*Remark 2.2.* If we replace  $\delta_r^-(u)$  by a weaker density function

$$\delta_r^*(u) = \sup_{-r^2 \leq t \leq -r^2/2} \frac{\text{md}(\Lambda(t) \cap B_{2r})}{r},$$

then the conclusion of the theorem still remains true (perhaps with different constants.)

### 3. MONOTONICITY FORMULAS

So-called *monotonicity formulas* will play an important role in this paper and will appear in almost every section.

We will use two different kinds of monotonicity formulas, the first due to Caffarelli [Caf93] and the second due to Weiss [Wei99], both in global and local forms.

Let

$$G(x, t) := \frac{1}{(4\pi t)^{n/2}} e^{-|x|^2/4t} \quad \text{for } (x, t) \in \mathbb{R}^n \times (0, \infty)$$

be the heat kernel. Then for a function  $v$  and any  $t > 0$  define

$$I(t; v) = \int_{-t}^0 \int_{\mathbb{R}^n} |\nabla v(x, s)|^2 G(x, -s) \, dx ds.$$

**Theorem 3.1** (Caffarelli [Caf93]). *Let  $h_1$  and  $h_2$  be nonnegative subcaloric functions in the strip  $\mathbb{R}^n \times [-1, 0]$  with a polynomial growth at infinity such that*

$$h_1(0, 0) = h_2(0, 0) = 0 \quad \text{and} \quad h_1 \cdot h_2 = 0.$$

*Then the functional*

$$\Phi(t) = \Phi(t; h_1, h_2) := \frac{1}{t^2} I(t; h_1) I(t; h_2)$$

*is monotone nondecreasing in  $t$  for  $0 < t < 1$ .* □

For the proof see Theorem 1 in [Caf93]. This theorem is a generalization of the Alt-Caffarelli-Friedman monotonicity formula from [ACF84].

*Remark 3.2.* As follows from the proof, if  $\Phi(t) > 0$  and the supports of  $h_1(\cdot, t)$  and  $h_2(\cdot, t)$  are not complementary halfspaces, then  $\Phi'(t) > 0$ .

We will also use the following local counterpart of the monotonicity theorem above. It takes the form of an estimate.

**Theorem 3.3** (Caffarelli [Caf93]). *Let  $h_1$  and  $h_2$  be nonnegative subcaloric functions in  $Q_1^-$  such that*

$$h_1(0, 0) = h_2(0, 0) = 0 \quad \text{and} \quad h_1 \cdot h_2 = 0.$$

*Also let  $\psi(x) \geq 0$  be a  $C^\infty$  cut-off function with  $\text{supp } \psi \subset B_{3/4}$  and  $\psi|_{B_{1/2}} = 1$  and set  $w_i = h_i \psi$ . Then there exists a constant  $C = C(n, \psi) > 0$  such that*

$$\Phi(t; w_1, w_2) \leq C \|h_1\|_{L^2(Q_1^-)}^2 \|h_2\|_{L^2(Q_1^-)}^2$$

*for any  $0 < t < 1/2$ .*

For the proof see Theorem 2 in [Caf93] and the remark after it. See also Theorem 2.1.3 in [CK98] for the generalization of this estimate for parabolic equations with variable coefficients.

To formulate the second monotonicity formula, we define Weiss’s functional for a function  $u$  by

$$W(r; u) = \frac{1}{r^4} \int_{-4r^2}^{-r^2} \int_{\mathbb{R}^n} \left( |\nabla u(x, t)|^2 + 2u(x, t) + \frac{u(x, t)^2}{t} \right) G(x, -t) \, dxdt.$$

**Theorem 3.4** (Weiss [Wei99]). *Let  $u$  be a solution of (1.2) in  $\mathbb{R}^n \times (-4, 0]$  with a polynomial growth at infinity. Then  $W(r; u)$  is monotone nondecreasing in  $r$  for  $0 < r < 1$ .*

The proof can be found in [Wei99]. An easy proof can be given by using the following scaling property of  $W$ :

$$W(r; u_r) = W(1, u),$$

where  $u_r(x, t) = (1/r^2)u(rx, r^2t)$  is the parabolic scaling of  $u$ . It can be shown that

$$W'(r; u) = \frac{1}{r^5} \int_{-4r^2}^{-r^2} \int_{\mathbb{R}^n} (\mathcal{L}u)^2 \frac{G(x, -t)}{-t} \, dxdt \geq 0$$

for every  $0 < r < 1$ , where

$$\mathcal{L}u(x, t) := x \cdot \nabla u(x, t) + 2t \partial_t u(x, t) - 2u(x, t) = \left. \frac{d}{dr} u_r(x, t) \right|_{r=1}.$$

*Remark 3.5.* In Weiss’s monotonicity theorem  $W'(r; u) = 0$  iff  $\mathcal{L}u = 0$  a.e. in  $\mathbb{R}^n \times [-4r^2, -r^2]$ . In particular,  $W(r; u) \equiv \text{const} =: W(u)$  iff  $u$  is homogeneous, i.e.  $u(x, t) = u_r(x, t) = (1/r^2)u(rx, r^2t)$  for  $0 < r \leq 1$ .

Before we state a local form of Weiss’s monotonicity theorem, we remark that it will not be used in most of the paper and will appear only in the last sections.

**Theorem 3.6.** *Let  $u \in \mathcal{P}_1^-(M)$  and  $\psi(x) \geq 0$  be a  $C^\infty$  cut-off function in  $\mathbb{R}^n$  with  $\text{supp } \psi \subset B_{3/4}$  and  $\psi|_{B_{1/2}} = 1$ . Then there exists  $C = C(n, \psi, M) > 0$  such that for  $w = u\psi$  the function*

$$W(r; w) + CF_n(r)$$

*is monotone nondecreasing in  $r$  for  $0 < r < 1/2$ , where*

$$F_n(r) = \int_0^r s^{-n-3} e^{-1/(16s^2)} \, ds.$$

The proof is based on the following lemma.

**Lemma 3.7.** *Let  $w$  be of the Sobolev class  $W_x^{2,p} \cap W_t^{1,p}(Q_R^-)$  for some  $p \geq 2$  and  $\text{supp } w(\cdot, t) \subset\subset B_R$  for every  $-R^2 \leq t \leq 0$ . Then*

$$W'(r; w) = \frac{1}{r^5} \int_{-4r^2}^{-r^2} \int_{\mathbb{R}^n} \mathcal{L}w(x, t) \left( \frac{\mathcal{L}w(x, t)}{-t} - 2(Hw(x, t) - 1) \right) G(x, -t) \, dxdt$$

*for  $0 < r < R/2$ .*

*Proof.* The computations below are formal but well justified, since  $w$  is a  $W_x^{2,p} \cap W_t^{1,p}$  function. Using the scaling property  $W(r; w) = W(1; w_r)$ , we obtain for  $r = 1$

$$\begin{aligned} W'(1; w) &= \frac{d}{dr} W(1; w_r) \\ &= \int_{-4}^{-1} \int_{\mathbb{R}^n} \left( \mathcal{L}(|\nabla w|^2) + 2\mathcal{L}w + 2\frac{w}{t}\mathcal{L}w \right) G(x, -t) \, dxdt \\ &= \int_{-4}^{-1} \int_{\mathbb{R}^n} \left( 2\nabla w \cdot \nabla(\mathcal{L}w) + 2\mathcal{L}w + 2\frac{w}{t}\mathcal{L}w \right) G(x, -t) \, dxdt, \end{aligned}$$

where we have used the (easily verified) identity

$$\mathcal{L}(|\nabla w|^2) = 2\nabla w \cdot \nabla(\mathcal{L}w).$$

Now integrating the term

$$2\nabla w \cdot \nabla(\mathcal{L}w) G(x, -t)$$

by parts and using that

$$\nabla G(x, -t) = -\frac{1}{2t} x \cdot G(x, -t),$$

we obtain

$$\begin{aligned} W'(1; w) &= 2 \int_{-4}^{-1} \int_{\mathbb{R}^n} \mathcal{L}w \left( -\Delta w - \frac{1}{2t} x \cdot \nabla w + 1 + \frac{w}{t} \right) G(x, -t) \, dxdt \\ &= \int_{-4}^{-1} \int_{\mathbb{R}^n} \mathcal{L}w \left( \frac{\mathcal{L}w(x, t)}{-t} - 2(Hw(x, t) - 1) \right) G(x, -t) \, dxdt, \end{aligned}$$

which proves the lemma for  $r = 1$  and by the rescaling argument, for all  $r$ .  $\square$

*Proof of Theorem 3.6.* By standard parabolic estimates (see e.g. [Lie96], Chapter VII) we have that  $u$  is of class  $W_x^{2,p} \cap W_t^{1,p}$  locally in  $Q_1^-$  for any  $1 < p < \infty$ , since  $\chi_\Omega \in L^\infty$ . As an immediate corollary from Lemma 3.7 we obtain that

$$(3.1) \quad W'(r; w) \geq -\frac{2}{r^5} \int_{-4r^2}^{-r^2} \int_{\mathbb{R}^n} \mathcal{L}w (Hw(x, t) - 1) G(x, -t) \, dxdt.$$

Next, from the representation  $w(x, t) = u(x, t) \psi(x)$  in  $Q_1^-$ , we have the following identities

$$\begin{aligned} \mathcal{L}w &= u(x \cdot \nabla \psi) + \psi \mathcal{L}u, \\ Hw &= u \Delta \psi + \psi Hu + 2\nabla \psi \cdot \nabla u. \end{aligned}$$

Since  $u$  satisfies (1.2) and  $\text{supp } \psi \subset B_{3/4}$ , it is easy to see that the integrand in (3.1) vanishes a.e. in  $B_{1/2} \times [-1, 0]$  and  $B_{3/4}^c \times [-1, 0]$ . Hence we obtain

$$W'(r; w) \geq -\frac{1}{r^5} \int_{-4r^2}^{-r^2} \int_{B_{3/4} \setminus B_{1/2}} f(x, t) G(x, -t) \, dxdt$$

with  $\|f\|_{L^1(Q_{3/4}^-)} \leq C = C(n, \psi, M) < \infty$  and consequently

$$W'(r; w) \geq -\frac{C}{r^{n+3}} e^{-1/(16r^2)}.$$

Therefore the function

$$W(r; w) + CF_n(r)$$

is nondecreasing, where

$$F_n(r) = \int_0^r s^{-n-3} e^{-1/(16s^2)} ds.$$

The proof is complete. □

#### 4. UNIFORM $C_x^{1,1} \cap C_t^{0,1}$ REGULARITY OF SOLUTIONS

In this section we establish uniform local  $C_x^{1,1} \cap C_t^{0,1}$  regularity of bounded solutions of (1.2).

**Theorem 4.1.** *Let  $u \in \mathcal{P}_1^-(x_0, t_0; M)$ . Then  $u \in C_x^{1,1} \cap C_t^{0,1}(Q_{1/4}^-(x_0, t_0))$ , uniformly. More precisely, there exists a universal constant  $C_0 = C_0(n)$  such that if  $u \in \mathcal{P}_1^-(x_0, t_0; M)$ , then*

$$\sup_{\Omega \cap Q_{1/4}^-(x_0, t_0)} (|\partial_{ij}u(x, t)| + |\partial_t u(x, t)|) \leq C_0 M.$$

In the general theory of the Stefan problem (where the additional assumptions  $u \geq 0$  and  $\partial_t u \geq 0$  are imposed by the problem) it can be show that  $\partial_t u$  is continuous with logarithmic modulus of continuity. In fact, if we knew more about the regularity of  $\partial_t u$  ( $C^\alpha$  is enough), we could threat the problem as an elliptic problem writing

$$\Delta u = \chi_\Omega f(x, t),$$

where  $f(x, t) = (1 + \partial_t u)$ .

Here we choose to approach the problem in its parabolic setting. The core of the proof of Theorem 4.1 is the following lemma, establishing the quadratic growth of solutions near the free boundary.

**Lemma 4.2.** *Let  $u \in \mathcal{P}_1^-(M)$ . Then there exists a constant  $C = C(n)$  such that*

$$(4.1) \quad \sup_{Q_r^-} |u| \leq CMr^2$$

for any  $0 \leq r \leq 1$ .

*Proof.* We use a method adopted from [CKS00]. Set

$$(4.2) \quad S_j(u) = \sup_{Q_{2^{-j}}^-} |u|$$

and define  $N(u)$  to be the set of all nonnegative integers satisfying the doubling condition

$$(4.3) \quad 2^2 S_{j+1}(u) \geq S_j(u).$$

Suppose now for some universal constant  $C_0 \geq 1$

$$(4.4) \quad S_{j+1}(u) \leq C_0 M 2^{-2j} \quad \text{for all } j \in N(u).$$

Then we claim

$$(4.5) \quad S_j(u) \leq C_0 M 2^{-2j+2} \quad \text{for all } j \in \mathbb{N}.$$

Obviously (4.5) holds for  $j = 1$ . Next, let (4.5) hold for some  $j$ . Then it holds also for  $j + 1$ . Indeed, if  $j \in N(u)$ , it follows from (4.4). If  $j \notin N(u)$ , (4.3) fails and we obtain

$$S_{j+1}(u) \leq 2^{-2} S_j(u) \leq C_0 M 2^{-2j}.$$

Therefore (4.5) holds for all  $j \in \mathbb{N}$ . This implies

$$\sup_{Q_r^-} |u| \leq 8C_0Mr^2$$

for any  $r \leq 1$ , and the lemma follows with  $C = 8C_0$ .

Now to complete the proof, we need to show (4.4). Suppose it fails. Then there exist sequences  $u_j \in \mathcal{P}_1^-(M)$ , and  $k_j \in N(u_j)$ ,  $j = 1, 2, \dots$ , such that

$$(4.6) \quad S_{k_j+1}(u_j) \geq jM2^{-2k_j}.$$

Define  $\tilde{u}_j$  as

$$\tilde{u}_j(x, t) = \frac{u_j(2^{-k_j}x, 2^{-2k_j}t)}{S_{k_j+1}(u_j)} \quad \text{in } Q_1^-.$$

Then

$$(4.7) \quad \sup_{Q_1^-} |H(\tilde{u}_j)| \leq \frac{2^{-2k_j}}{S_{k_j+1}(u_j)} \leq \frac{1}{jM} \rightarrow 0,$$

$$(4.8) \quad \sup_{Q_{1/2}^-} |\tilde{u}_j| = 1, \quad (\text{by (4.2)}),$$

$$(4.9) \quad \sup_{Q_1^-} |\tilde{u}_j| \leq \frac{S_{k_j}(u_j)}{S_{k_j+1}(u_j)} \leq 4 \quad (\text{by (4.3)}),$$

$$(4.10) \quad \tilde{u}_j(0, 0) = |\nabla \tilde{u}_j(0, 0)| = 0.$$

Now by (4.7)–(4.10) we will have a subsequence of  $\tilde{u}_j$  converging in  $C_x^{1,\alpha} \cap C_t^{0,\alpha}(Q_1^-)$  to a nonzero caloric function  $u_0$  in  $Q_1^-$ , satisfying  $u_0(0, 0) = |\nabla u_0(0, 0)| = 0$ . Moreover, from (4.8), we will have

$$(4.11) \quad \sup_{Q_{1/2}^-} |u_0| = 1.$$

For any spatial unit vector  $e$  define

$$v = \partial_e u_0, \quad v_j = \partial_e u_j, \quad \tilde{v}_j = \partial_e \tilde{u}_j.$$

Then, over a subsequence,  $\tilde{v}_j$  converges in  $C_x^{0,\alpha} \cap C_t^{0,\alpha}(Q_1^-)$  to  $v$ . Moreover  $H(v) = 0$ . Now, for a fixed cut-off function  $\psi(x)$  with  $\psi|_{B_{1/2}} = 1$  and  $\text{supp } \psi \subset B_{3/4}$  and  $u \in \mathcal{P}_1(M)$  consider

$$\Phi(t; (\partial_e u)\psi) = \frac{1}{t^2} I(t; (\partial_e u)^+ \psi) I(t; (\partial_e u)^- \psi).$$

Then to apply the monotonicity formula of [Caf93] (see Theorem 3.3 above), we need to verify that the functions  $(\partial_e u)^\pm$  are subcaloric; we leave this to the reader. Then, for all  $0 < t < t_0$ , we obtain

$$(4.12) \quad \Phi(t; (\partial_e u)\psi) \leq C \|\nabla u\|_{L^2(Q_1^-)}^4 \leq C_0,$$

for a universal constant  $C_0$ , which, by classical estimates, depends on the class only.

Now choose  $\psi$  as above and set  $\psi_j(x) = \psi(2^{-k_j}x)$ . Then estimate (4.12) applied to  $\tilde{v}_j\psi_j$  gives

$$(4.13) \quad \Phi(1; \tilde{v}_j\psi_j) \leq \left(\frac{2^{-2k_j}}{S_{k_j+1}}\right)^4 \Phi(2^{-2k_j}; v_j\psi) \leq C_0 \left(\frac{2^{-2k_j}}{S_{k_j+1}}\right)^4$$

for  $k_j$  large enough. Since  $\psi_j = 1$  in  $B_{2^{k_j-1}}$ , we will have

$$|\nabla(\tilde{v}_j\psi_j)|^2 \geq |\nabla \tilde{v}_j|^2 \chi_{B_1}.$$

Hence for  $\varepsilon > 0$  (small and fixed) we have

$$C_{n,\varepsilon} \int_{-1}^{-\varepsilon} \int_{B_1} |\nabla \tilde{v}_j^\pm|^2 \, dxdt \leq \int_{-1}^0 \int_{B_1} |\nabla \tilde{v}_j^\pm \psi_j|^2 G(x, -t) \, dxdt = I(1, \tilde{v}_j^\pm \psi_j).$$

This estimate, in combination with Poincaré’s inequality, gives

$$\int_{-1}^{-\varepsilon} \int_{B_1} |\tilde{v}_j^\pm - M^\pm(t)|^2 \, dxdt \leq C_n \int_{-1}^{-\varepsilon} \int_{B_1} |\nabla \tilde{v}_j^\pm|^2 \, dxdt \leq C(n, \varepsilon) I(1, \tilde{v}_j^\pm \psi_j),$$

where  $M_j^\pm(t)$  denotes the corresponding mean value of  $\tilde{v}_j^\pm$  on the  $t$ -section.

Using this and (4.13), we will have

$$\begin{aligned} & \left( \int_{-1}^{-\varepsilon} \int_{B_1} |\tilde{v}_j^+ - M_j^+(t)|^2 \, dxdt \right) \left( \int_{-1}^{-\varepsilon} \int_{B_1} |\tilde{v}_j^- - M_j^-(t)|^2 \, dxdt \right) \\ & \leq C(n, \varepsilon) \Phi(1, v_j \psi) \leq C(n, \varepsilon) \left( \frac{2^{-2k_j}}{S_{k_j+1}} \right)^4. \end{aligned}$$

Using (4.6) and letting  $j \rightarrow \infty$  (and then  $\varepsilon \rightarrow 0$ ), we obtain

$$(4.14) \quad \int_{-1}^0 \int_{B_1} |v^+ - M^+(t)|^2 \int_{-1}^0 \int_{B_1} |v^- - M^-(t)|^2 = 0,$$

where  $M^\pm(t)$  denotes the corresponding mean value of  $v^\pm$  on  $t$ -sections over  $B_1$ . Obviously, (4.14) implies that either of  $v^\pm$  is equivalent to  $M^\pm(t)$  in  $Q_1^-$  and is thus independent of the spatial variables. Let us assume  $v^- = M^-(t)$ . Then  $-\partial_t v^- = H(v^-) = 0$ , i.e.  $M^-$  is constant in  $Q_1^-$ . Since  $v(0, 0) = 0$ , we must have  $M^- = 0$ , i.e.  $v \geq 0$  in  $Q_1^-$ . Hence by the minimum principle  $v \equiv 0$  in  $Q_1^-$ . Since  $v = \partial_e u_0$ , and  $e$  is an arbitrary direction, we conclude that  $u_0$  is constant in  $Q_1^-$ . Also  $u_0(0, 0) = 0$  implies that the constant must be zero, i.e.  $u_0 \equiv 0$  in  $Q_1^-$ . This contradicts (4.11) and the lemma is proved.  $\square$

In fact, for the proof of Theorem 4.1 we will also need an extension of Lemma 4.2 to the “upper half” as well.

**Lemma 4.3.** *Let  $u \in \mathcal{P}_1(M)$ . Then there exists a constant  $C = C(n)$  such that*

$$(4.15) \quad \sup_{Q_r} |u| \leq CMr^2$$

for any  $0 \leq r \leq 1$ .

*Proof.* Define  $w_1 = CM(|x|^2 + 2nt)$ , where  $C$  is as in Lemma 4.2. Then  $H(w_1) = 0 \leq H(u)$  in  $Q_1^+$ . Also, by (4.1),  $w_1 \geq u$  on the parabolic boundary  $\partial_p Q_1^+$ . Hence by the comparison principle we will have  $w_1 \geq u$  in  $Q_1^+$ .

Similarly we define  $w_2 = -CM(|x|^2 + 2nt) - t$ , which satisfies  $H(w_2) = 1 \geq H(u)$  in  $Q_1^+$ . Also, by (4.1), on the parabolic boundary  $\partial_p Q_1^+$  we have  $w_2 \leq u$ . Hence by the comparison principle  $w_2 \leq u$  in  $Q_1^+$ . This completes the proof of the lemma.  $\square$

Now, we are ready to prove Theorem 4.1.

*Proof of Theorem 4.1.* For  $(x_0, t_0) \in \Omega \in Q_{1/4}^-$  let

$$d = d^-(x_0, t_0) = \sup\{r : Q_r^-(x_0, t_0) \subset \Omega \cap Q_1^-\},$$

the parabolic distance to the free boundary. Then Lemma 4.3 implies that

$$|u(x, t)| \leq CMd^2 \quad \text{in } Q_d^-(x_0, t_0).$$

Now consider the function

$$v(x, t) = \frac{1}{d^2} u(dx + x_0, d^2t + t_0) \quad \text{in } Q_1^-.$$

Then  $v$  satisfies  $Hv = 1$ , and  $|v|$  is uniformly bounded in  $Q_1^-$ . Hence by standard parabolic estimates (see for instance [Fri64])  $\partial_{ij}v(0, 0) = \partial_{ij}u(x_0, t_0)$ , and  $\partial_tv(0, 0) = \partial_tu(x_0, t_0)$  are uniformly bounded (independent of  $x_0$  and  $t_0$ ), which is the desired result. The theorem is proved.  $\square$

The above theorem has the following obvious implication.

**Corollary 4.4.** *Let  $u \in \mathcal{P}_\infty^-(M)$ . Then*

$$|\partial_{ij}u(x, t)| + |\partial_tu(x, t)| \leq C_0M \quad \text{in } \Omega.$$

*Proof.* Let  $u_r$  be a scaling of  $u$  at the origin, i.e.

$$u_r(x, t) = \frac{1}{r^2} u(rx, r^2t) \quad \text{in } Q_1^-.$$

Then  $u \in \mathcal{P}_\infty^-(M)$  implies  $u_r \in \mathcal{P}_1^-(3M)$  for  $r \geq 1$ . Hence by Theorem 4.1 we have

$$\sup_{\Omega_r \cap Q_{1/4}^-} (|\partial_{ij}u_r(x, t)| + |\partial_tu_r(x, t)|) \leq C_0M,$$

i.e.

$$\sup_{\Omega \cap Q_{r/4}^-} (|\partial_{ij}u(x, t)| + |\partial_tu(x, t)|) \leq C_0M.$$

Letting  $r \rightarrow \infty$ , we will obtain the statement of the corollary.  $\square$

## 5. NONDEGENERACY

**5.1. Nondegeneracy.** The reader may have wondered what happens if the functions  $u_r$  under the blow-up process converge identically to zero. This happens if the function  $u$  decays to zero faster than quadratically. This, however, does not happen if we blow-up at a free boundary point.

**Lemma 5.1.** *Let  $u$  be a solution of (1.2) and let  $(x_0, t_0) \in \Gamma$ . Then there exists a universal constant  $C_n > 0$  such that*

$$(5.1) \quad \sup_{Q_r^-(x_0, t_0)} u \geq C_n r^2$$

for any  $r > 0$  such that  $Q_r^-(x_0, t_0) \subset D$ . More generally, for any  $(x_0, t_0) \in \Lambda$  we have that either (5.1) holds or  $u \equiv 0$  in  $Q_{r/2}^-(x_0, t_0)$  for any  $r > 0$  as above.

*Proof.* Consider first  $(x_1, t_1) \in \{u > 0\}$  and set

$$w(x, t) = u(x, t) - u(x_1, t_1) - \frac{1}{2n+1} (|x - x_1|^2 - (t - t_1)).$$

Then  $w$  is caloric in  $\Omega \cap Q_r^-(x_1, t_1)$  and strictly negative on  $\partial\Omega \cap Q_r^-(x_1, t_1)$ . Since  $w(x_1, t_1) = 0$ , the maximum of  $w$  on the parabolic boundary of the cylinder  $Q_r^-(x_1, t_1)$  is nonnegative. In particular, we obtain

$$\sup_{\partial_p Q_r^-(x_1, t_1)} \left( u(x, t) - u(x_1, t_1) - \frac{r^2}{2n+1} \right) \geq 0.$$

Hence

$$(5.2) \quad \sup_{Q_r^-(x_1, t_1)} u \geq u(x_1, t_1) + \frac{r^2}{2n + 1}.$$

Then a limiting argument shows that (5.1) holds if  $(x_0, t_0)$  is in the closure of  $\{u > 0\}$  with  $C_n = 1/(2n + 1)$ . Moreover, if  $Q_{r/2}^-(x_0, t_0)$  contains a point  $(x_1, t_1)$  in  $\{u > 0\}$ , we still have

$$\sup_{Q_r^-(x_0, t_0)} u(x, t) \geq \sup_{Q_{r/2}^-(x_1, t_1)} u(x, t) \geq u(x_1, t_1) + \frac{(r/2)^2}{2n + 1} \geq C_n r^2.$$

Finally, in the case when  $u \leq 0$  in  $Q_{r/2}^-(x_0, t_0)$ , the maximum principle implies that  $u \equiv 0$  in  $Q_{r/2}^-(x_0, t_0)$ , since  $u(x_0, t_0) = 0$ . Thus  $(x_0, t_0)$  is not a free boundary point.  $\square$

The next lemma shows that we also have a certain nondegeneracy at the points of  $\partial\Omega \cap D$  even if they are not in  $\Gamma$ .

**Lemma 5.2.** *Let  $u$  be a solution of (1.2) and let  $(x_0, t_0) \in \partial\Omega \cap D$ . Then there exists a constant  $C_n > 0$  such that*

$$(5.3) \quad \sup_{Q_r(x_0, t_0)} |u| \geq C_n r^2$$

for any  $r > 0$  with  $Q_r(x_0, t_0) \subset D$ .

*Proof.* Consider two cases: (i)  $Q_{r/2}(x_0, t_0)$  contains a point  $(x_1, t_1)$  in  $\{u > 0\}$  and (ii)  $u \leq 0$  in  $Q_{r/2}$ . As in the proof of the previous lemma, we obtain that in the first case

$$\sup_{Q_r(x_0, t_0)} u \geq C_n r^2$$

(and we are done) and in the second case that  $u \equiv 0$  in  $Q_{r/2}^-(x_0, t_0)$ . Moreover, in the second case we claim that

$$\inf_{Q_{r/2}^+(x_0, t_0)} u \leq -C_n r^2.$$

Indeed, first observe that  $u < 0$  in  $Q_{r/2}(x_0, t_0) \cap \{t > t_0\}$ ; otherwise we would have  $u \equiv 0$  in  $B_{r/2}(x_0) \times (t_0 - r^2/4, t_1)$  for some  $t_1 > t_0$ , which contradicts the assumption that  $(x_0, t_0) \in \partial\Omega$ . The parabolic scaling

$$v(x, t) = \frac{1}{r^2} u(rx + x_0, r^2t + t_0)$$

satisfies

$$Hv = 1, \quad v < 0 \quad \text{in } Q_{1/2}^+.$$

But then

$$\inf_{Q_{1/2}^+} v \leq -C_n;$$

otherwise we would have a sequence of functions  $-1/k \leq v_k \leq 0$  in  $Q_{1/2}^+$  satisfying  $Hv_k = 1$ . This is impossible, since the limit function  $v_0$ , which is identically 0, should also satisfy  $Hv_0 = 1$ .

Scaling back, we complete the proof of the lemma.  $\square$

**5.2. Stability under the limit.** Let  $u_j$  be any converging sequence in the class  $\mathcal{P}_1^-(M)$  and let  $u_0 = \lim_{j \rightarrow \infty} u_j$ . Then we claim  $u_0 \in \mathcal{P}_1^-(M)$ .

To prove this statement, we may assume that the convergence is in  $C_x^{1,\alpha} \cap C_t^{0,\alpha}$ . Hence we have

$$(5.4) \quad \limsup_{j \rightarrow \infty} \Lambda(u_j) \subset \Lambda(u_0),$$

where  $\limsup_{j \rightarrow \infty} E_j$  for the sequence of sets  $E_j$  is defined as the set of all limit points of sequences  $(x_{j_k}, t_{j_k}) \in E_{j_k}$ ,  $j_k \rightarrow \infty$ . Then for any  $(x, t) \in \Omega(u_0)$  there exists  $\varepsilon > 0$  such that  $Q_\varepsilon^-(x, t) \subset \Omega(u_j)$ , thus implying that

$$Hu_0 = 1 \quad \text{in } \Omega(u_0).$$

Also since  $u_0$  is  $C_x^{1,1} \cap C_t^{0,1}$  regular, it follows that  $u_0$  is a solution of (1.2).

Next, we claim that

$$(5.5) \quad \limsup_{j \rightarrow \infty} \Gamma(u_j) \subset \Gamma(u_0).$$

In particular, if  $(0, 0) \in \Gamma(u_j)$ , then  $(0, 0) \in \Gamma(u_0)$ . This follows immediately from the nondegeneracy Lemma 5.1.

In fact, we also have a similar inclusion for  $\partial\Omega$ :

$$(5.6) \quad \limsup_{j \rightarrow \infty} \partial\Omega(u_j) \subset \partial\Omega(u_0).$$

Indeed, (5.6) will follow once we show that if  $u_0 = 0$  in  $Q_r(x_0, t_0)$ , then  $u_j = 0$  in  $Q_{r/2}(x_0, t_0)$  for sufficiently large  $j$ . Assume the contrary. Then we will have either  $Q_{r/2}(x_0, t_0) \subset \Omega(u_j)$  or  $Q_{r/2}(x_0, t_0) \cap \Omega(u_j) \neq \emptyset$  over infinitely many  $j = j_k$ . In the first case we will obtain that  $u_0$  satisfies  $H(u_0) = 1$  in  $Q_{r/2}(x_0, t_0)$  and in the second that  $\sup_{Q_r(x_0, t_0)} |u_0| \geq C_n r^2$ , both of which are impossible for  $u_0 = 0$  in  $Q_r(x_0, t_0)$ .

The same argument as above also shows that

$$(5.7) \quad \limsup_{j \rightarrow \infty} \Omega(u_j) \subset \overline{\Omega(u_0)}.$$

Generally, we cannot prove inclusions similar to (5.4)–(5.7) for  $t$ -levels of the sets  $\Omega$  and  $\Lambda$ . The reason is the lack of the “elliptic version” of (5.3) on  $t$ -sections. However, for the Stefan problem when one assumes that  $\partial_t u \geq 0$ , one has  $\Delta u = \chi_\Omega + \partial_t u \geq \chi_\Omega$ , which allows us to prove the elliptic version of (5.3).

**5.3. Lebesgue measure of  $\partial\Omega$ .** From nondegeneracy and  $C_x^{1,1} \cap C_t^{0,1}$  regularity one can deduce that  $\partial\Omega$  (hence also the free boundary  $\Gamma$ ) has  $(n + 1)$ -dimensional Lebesgue measure zero. It is enough to show that  $\partial\Omega$  has Lebesgue density less than 1 at its every point.

Indeed, take any  $(x_0, t_0) \in \partial\Omega$ . Using (5.3), we can find  $(x_1, t_1) \in Q_{r/4}(x_0, t_0)$  such that  $|u(x_1, t_1)| \geq Cr^2$ . On the other hand, by Theorem 4.1, we have  $|u(x_1, t_1)| \leq C_1 d^2(x_1, t_1)$  (where  $d$  is the parabolic distance to  $\Lambda$ ). Hence  $d(x_1, t_1) \geq Cr$ . In particular the set  $\Omega \cap Q_r(x_0, t_0)$  contains a cylinder of a size proportional to  $Q_r(x_0, t_0)$ .

In fact, we claim that for any parabolic cylinder  $Q_r(x, t)$ , not necessarily centered at a point on  $\partial\Omega$ ,  $Q_r(x, t) \setminus \partial\Omega$  contains a cylinder proportional to  $Q_r(x, t)$ . To prove it, consider the following two alternatives: either  $Q_{r/2}(x, t)$  contains a point

on  $\partial\Omega$  or it does not. In the first case the claim follows from the arguments above and in the second case  $Q_{r/2}(x, t)$  itself is contained in  $Q_r(x, t) \setminus \partial\Omega$ .

Furthermore, that  $\partial\Omega$  has Lebesgue density less than 1 at  $(x_0, t_0)$  will follow if we show that for every hyperbolic cylinder

$$C_r(x_0, t_0) := B_r(x_0) \times (t_0 - r, t_0 + r)$$

$C_r(x_0, t_0) \setminus \partial\Omega$  contains a set  $E$  of Lebesgue measure proportional to that of  $C_r(x_0, t_0)$ . Note that it is enough to show this statement for  $r = 1/k$ ,  $k = 1, 2, \dots$ . Subdivide  $C_{1/k}(x_0, t_0)$  into  $k$  parabolic cylinders

$$Q_i = Q_{1/k}(x_0, t_i), \quad t_i = t_0 + 1 - (2i - 1)/k, \quad i = 1, 2, \dots, k.$$

Then  $Q_i \setminus \partial\Omega$  contains a cylinder  $\tilde{Q}_i$  proportional to  $Q_i$  and one can take

$$E = \bigcup_{i=1}^k \tilde{Q}_i.$$

Thus,  $\partial\Omega$  has  $(n + 1)$ -dimensional Lebesgue measure 0.

## 6. HOMOGENEOUS GLOBAL SOLUTIONS

**Definition 6.1.** We say that the solution  $u(x, t)$  is *homogeneous* (with respect to the origin) if

$$\frac{1}{r^2} u(rx, r^2t) = u(x, t)$$

for every  $r > 0$ .

Simple examples of homogeneous solutions are the polynomial solutions of the type

$$u(x, t) = mt + P(x),$$

where  $m$  is a constant and  $P$  is a homogeneous quadratic polynomial satisfying  $\Delta P = m + 1$ , and the halfspace solutions

$$u(x, t) = \frac{1}{2} (x \cdot e)_+^2$$

for spatial unit vectors  $e$ . As we will see below these are the only nonzero homogeneous solutions in  $\mathbb{R}^n \times \mathbb{R}^-$ .

As was already mentioned in Remark 3.5, solutions  $u$ , homogeneous in the past, have the property that their Weiss functional  $W(r; u)$  is a constant (and vice versa). We denote this constant by  $W(u)$ .

**Lemma 6.2.**

(i) For every spatial direction  $e$

$$W\left(\frac{1}{2} (x \cdot e)_+^2\right) = W\left(\frac{1}{2} (x_1)_+^2\right) =: A.$$

(ii) For every constant  $m$  and homogeneous quadratic polynomial  $P(x)$  satisfying  $\Delta P = m + 1$

$$W(mt + P(x)) = W\left(\frac{1}{2} (x_1)^2\right) = 2A.$$

*Proof.* Part (i) is obvious because of the rotational symmetry of the functional  $W$ . Part (ii) follows from direct computations. Indeed, for given  $t < 0$ ,

$$\begin{aligned} & \int_{\mathbb{R}^n} \left( |\nabla u(x, t)|^2 + 2u(x, t) + \frac{u(x, t)^2}{t} \right) G(x, -t) \, dx \\ &= \int_{\mathbb{R}^n} u \left( -\Delta u - \frac{1}{2t} x \cdot \nabla u + \frac{u}{t} + 2 \right) G(x, -t) \, dx \\ &= \int_{\mathbb{R}^n} u(x, t) G(x, -t) \, dx = -t, \end{aligned}$$

where we first integrated the term  $|\nabla u|^2 G = \nabla u \cdot (G \nabla u)$  by parts and then used that

$$\Delta u + \frac{1}{2t} x \cdot \nabla u - \frac{u}{t} - 1 = 0$$

for homogeneous solutions. This equation can be obtained, for instance, from  $\Delta u - \partial_t u = 1$  and the homogeneity property  $x \cdot \nabla u + 2t \partial_t u = 2u$  by eliminating  $\partial_t u$ .

Hence

$$W(u, r) = \frac{1}{r^4} \int_{-4r^2}^{-r^2} (-t) dt = \frac{15}{2}$$

and the lemma follows with  $A = 15/4$ . □

The importance of this lemma is emphasized by the following fact.

**Lemma 6.3.** *The only nonzero homogeneous solutions of (1.2) in  $D = \mathbb{R}^n \times \mathbb{R}^-$  are of the type*

- (i)  $u(x, t) = \frac{1}{2}(x \cdot e)_+^2$  for a certain spatial unit vector  $e$ ;
- (ii)  $u(x, t) = mt + P(x)$ , where  $m$  is a constant and  $P(x)$  is a homogeneous quadratic polynomial satisfying  $\Delta P = m + 1$ .

*Proof.* From the homogeneity of  $u$  it follows that the time sections  $\Omega(t) = \{(x, t) \in \Omega\}$  satisfy

$$\Omega(r^2 t) = r \Omega(t)$$

for any  $r > 0$ . We consider two different cases.

*Case 1.*  $\Omega^c$  has an empty interior. This will happen if and only if  $\Omega(t)^c$  has an empty interior for one and thus for all  $t$ . Since both  $u$  and  $|\nabla u|$  vanish on  $\Omega^c$ , it follows that  $u$  satisfies  $\Delta u - \partial_t u = 1$  in all of  $\mathbb{R}^n \times \mathbb{R}^-$ . But then  $\partial_t u$  is a bounded caloric function in  $\mathbb{R}^n \times \mathbb{R}^-$ , thus a constant by the Liouville theorem. Similarly,  $\partial_{ij} u$  are constants. Therefore we obtain the representation

$$u(x, t) = mt + P(x)$$

where  $P$  is a homogeneous quadratic polynomial such that  $\Delta P = m + 1$ .

*Case 2.*  $\Omega^c$  has nonempty interior. By homogeneity, for every unit spatial direction  $e$ ,

$$\Phi(t; \partial_e u) \equiv \text{const}$$

where  $\Phi$  is as in Theorem 3.1. However, this is possible only if the spatial supports of  $(\partial_e u)_+$  and  $(\partial_e u)_-$  are complementary halfspaces at almost all  $t$ , or if  $\Phi(t; \partial_e u) \equiv 0$ , see Remark 3.2. The former case cannot occur since  $\Omega(t)^c$  is nonempty for all  $t < 0$ , and the latter case implies  $\partial_e u \geq 0$  or  $\partial_e u \leq 0$  in all of  $\mathbb{R}^n \times \mathbb{R}^-$ . Since this is true for all spatial directions  $e$ , it follows that  $u(x, t)$  is one-dimensional, i.e. in suitable spatial coordinates  $u(x, t) = u(x_1, t)$ . We may assume therefore that in the

rest of the proof the spatial dimension  $n = 1$ . From homogeneity we also have the representation

$$u(x, t) = -t f\left(\frac{x}{\sqrt{-t}}\right),$$

where  $f = u(\cdot, -1)$ . The function  $f$  satisfies

$$\begin{aligned} f''(\xi) - \frac{\xi}{2} f'(\xi) + f(\xi) - 1 &= 0 \quad \text{in } \Omega(-1), \\ f(\xi) = f'(\xi) &= 0 \quad \text{on } \Omega(-1)^c. \end{aligned}$$

The general solution to the ordinary differential equation above is given by

$$f(\xi) = 1 + C_1 (\xi^2 - 2) + C_2 \left( -2\xi e^{\xi^2/4} + (\xi^2 - 2) \int_0^\xi e^{s^2/2} ds \right)$$

and if  $a$  is a finite endpoint of a connected component of  $\Omega(-1)$ , we have

$$C_2 = \frac{1}{4} a e^{-a^2/4}, \quad C_1 = \frac{1}{2} - \frac{1}{4} a e^{-a^2/4} \int_0^a e^{-s^2/4} ds.$$

In particular, we see that there could not be two different values of  $a$ ; hence each connected component of  $\Omega(-1)$  is unbounded.

Next, on the unbounded interval we must have  $C_2 = 0$ , since  $f$  has at most quadratic growth at infinity. This implies  $C_1 = 1/2$  and the only possible value of  $a$  is 0. Thus,  $\Omega(-1)$  is either  $(0, \infty)$  or  $(-\infty, 0)$  and  $f(\xi) = (1/2) \xi_+^2$  or  $f(\xi) = (1/2) \xi_-^2$ , respectively.  $\square$

*Remark 6.4.* As the example before Theorem I shows, Lemma 6.3 is valid only for solutions in lower-half space  $\mathbb{R}^n \times \mathbb{R}^-$  but not for the solutions in all of  $\mathbb{R}^n \times \mathbb{R}$ . In fact, if we take any homogeneous solution in  $\mathbb{R}^n \times \mathbb{R}^-$  and continue it to  $\mathbb{R}^n \times \mathbb{R}$  by solving the Cauchy problem for  $Hu = 1$  in  $\mathbb{R}^n \times \mathbb{R}^+$ , then the resulting function will still be homogeneous, but will not have one of the forms in Lemma 6.3 in  $\mathbb{R}^n \times \mathbb{R}^+$ .

### 7. BALANCED ENERGY

Let  $u \in \mathcal{P}_\infty^-(M)$  be a global solution. Then we define the *balanced energy*

$$(7.1) \quad \omega = \lim_{r \rightarrow 0} W(r; u),$$

which exists due to Weiss's monotonicity formula. Recall that the functional  $W$  has the scaling property

$$(7.2) \quad W(rs; u) = W(s; u_r),$$

where

$$u_r(x, t) = \frac{1}{r^2} u(rx, r^2t).$$

Since the functions  $u_r$  are locally uniformly in class  $C_x^{1,1} \cap C_t^{0,1}$  in  $\mathbb{R}^n \times \mathbb{R}^-$ , we can extract a converging subsequence  $u_{r_k}$  to a global solution  $u_0$  in  $\mathbb{R}^n \times \mathbb{R}^-$ . Then passing to the limit in (7.2), we will obtain that

$$\omega = W(s, u_0)$$

for any  $s > 0$ . This implies that the blow-up  $u_0$  is a homogeneous global solution. Moreover, from Lemmas 6.2 and 6.3 it follows that  $\omega$  can take only three values: 0,  $A$ , or  $2A$ .

Similarly we define the balanced energy at any point  $(x, t) \in \Lambda$  for a global solution  $u \in \mathcal{P}_\infty^-(M)$  by

$$\omega(x, t) = \lim_{r \rightarrow 0} W(r, x, t; u).$$

**Definition 7.1.** We say that a point  $(x, t) \in \partial\Omega$  is a *zero, low, or high energy point* of the global solution  $u \in \mathcal{P}_\infty^-(M)$  if

$$\omega(x, t) = 0, \quad A, \quad 2A,$$

respectively. Here  $A$  is as in Lemma 6.2.

*Remark 7.2.* The balanced energy function  $\omega$  is upper semicontinuous, since

$$W(r, \cdot, \cdot; u) =: \omega_r \searrow \omega \quad \text{as } r \searrow 0$$

and functions  $\omega_r$  are continuous on  $\partial\Omega$ . Hence, if

$$\mathcal{E}_j = \{\omega = jA\}, \quad \text{for } j = 0, 1, 2,$$

then  $\mathcal{E}_0, \mathcal{E}_0 \cup \mathcal{E}_1$  are relatively open and  $\mathcal{E}_2$  is closed.

**7.1. Zero energy points.** By definition,  $(x_0, t_0) \in \partial\Omega$  is a zero energy point for  $u$  if there exists a blow-up  $u_0$  of  $u$  at  $(x_0, t_0)$ , such that  $u_0 \equiv 0$  in  $\mathbb{R}^n \times \mathbb{R}^-$ . From Lemma 5.1 we have that either

$$\sup_{Q_r^-(x_0, t_0)} u \geq C_n r^2,$$

for all  $r > 0$ , or  $u \equiv 0$  in  $Q_r^-(x_0, t_0)$  and  $u < 0$  in  $Q_r^+(x_0, t_0)$  for some  $r > 0$ . In the first case the point  $(x_0, t_0)$  is either of low or high energy, since no blow-up  $u_0$  at  $(x_0, t_0)$  can vanish identically in  $\mathbb{R}^n \times \mathbb{R}^-$ . Also, only in the second case is the point  $(x_0, t_0)$  of zero energy. Thus, zero energy points are parabolically interior points of  $\Lambda$ .

Also, we obtain that the free boundary  $\Gamma$  cannot contain zero energy points and in fact

$$\Gamma = \partial\Omega \setminus \mathcal{E}_0 = \mathcal{E}_1 \cup \mathcal{E}_2.$$

In other words, the free boundary points consist of low and high energy points of  $\partial\Omega$ .

Finally, we remark that if  $u \geq 0$ ,  $\partial\Omega$  coincides with the free boundary  $\Gamma$ , since there could be no zero energy points.

**7.2. High energy points.**

**Lemma 7.3.** *Let  $(x_0, t_0) \in \partial\Omega$  be a high energy point for a global solution  $u$ . Then*

$$(7.3) \quad u(x, t) = m(t - t_0) + P(x - x_0)$$

*in  $\mathbb{R}^n \times \mathbb{R}_{t_0}^-$ , where  $m$  is a constant and  $P$  is a homogeneous quadratic polynomial.*

*Proof.* Without loss of generality we may assume that  $(x_0, t_0) = (0, 0)$ . Consider then the functional  $W(r; u)$ . It is nondecreasing in  $r$  and therefore

$$(7.4) \quad \omega = W(0+; u) \leq W(\infty; u) = W(u_\infty),$$

where  $u_\infty$  is a shrink-down limit over a subsequence of scaled functions  $u_r$  as  $r \rightarrow \infty$ . Since  $(0, 0)$  is a high energy point,  $\omega = 2A$  and we obtain  $W(u_\infty) \geq 2A$ . The shrink-down function  $u_\infty$  is homogeneous and therefore from Lemmas 6.2 and 6.3 we also have  $W(u_\infty) \leq 2A$ . Hence  $W(u_\infty) = 2A$ , which is possible if and only if

$$W(r; u) = 2A \quad \text{for all } r > 0.$$

This implies that  $u(x, t)$  is homogeneous with respect to  $(0, 0)$  in  $\mathbb{R}^n \times \mathbb{R}^-$ . Applying Lemma 6.3, we finish the proof.  $\square$

*Remark 7.4.* A simple corollary from the lemma above is that all high energy points, if they exist, are on the same time level  $t = t_0$ , if  $m \neq 0$  in the representation (7.3). Moreover  $x_0$  must be on the hyperplane  $\{\nabla u(\cdot, t_0) = 0\} = \{\nabla P(\cdot - x_0)\}$ . Except for the case when  $P \equiv 0$  (equivalently  $u(x, t) = -(t - t_0)$ ),  $\{\nabla u(\cdot, t_0) = 0\}$  is a lower-dimensional hyperplane in  $\mathbb{R}^n$ .

If  $m = 0$  in (7.3), then there exists a maximal  $t_* \geq t_0$  (possibly infinite) such that  $u(x, t) = P(x - x_0)$  for all  $x$  and  $t \leq t_*$ . If  $t_*$  is finite, then  $u$  has no high energy points  $(x, t)$  with  $t > t_*$ .

**7.3. Low energy points.**

**Theorem 7.5.** *Let  $u \in \mathcal{P}_\infty^-(M)$  be a global solution and  $(x_0, t_0) \in \partial\Omega$  be a low energy point. Then there exists  $r = r(x_0, t_0) > 0$  such that  $u \geq 0$  in  $Q_r^-(x_0, t_0)$ . Moreover, we can choose  $r > 0$  such that  $\partial\Omega \cap Q_r^-(x_0, t_0)$  is a Lipschitz (in space and time) surface.*

The proof is based on the following two useful lemmas.

**Lemma 7.6.** *Let  $u$  be a bounded solution of (1.2) in  $Q_1^-$  and let  $h$  be caloric in  $Q_1^- \cap \Omega$ . Suppose moreover that*

- (i)  $h \geq 0$  on  $\partial\Omega \cap Q_1^-$  and
- (ii)  $h - u \geq -\varepsilon_0$  in  $Q_1^-$ , for some  $\varepsilon_0 > 0$ .

*Then  $h - u \geq 0$  in  $Q_{1/2}^-$ , provided  $\varepsilon_0 = \varepsilon_0(n)$  is small enough.*

*Proof.* Suppose the conclusion of the lemma fails. Then there are  $u$  and  $h$  satisfying the conditions of the lemma such that

$$(7.5) \quad h(x_0, t_0) - u(x_0, t_0) < 0$$

for some  $(x_0, t_0) \in Q_{1/2}^-$ . Let

$$w(x, t) = h(x, t) - u(x, t) + \frac{1}{2n + 1}(|x - x_0|^2 - (t - t_0)).$$

Then  $w$  is caloric in  $\Omega \cap Q_{1/2}^-(x_0, t_0)$ ,  $w(x_0, t_0) < 0$  by (7.5) and  $w \geq 0$  on  $\partial\Omega \cap Q_{1/2}^-(x_0, t_0)$ . Hence by the minimum principle the negative infimum of  $w$  is attained on  $\partial_p Q_{1/2}^-(x_0, t_0)$ . We thus obtain

$$-\varepsilon_0 \leq \inf_{\partial Q_{1/2}^-(x_0, t_0) \cap \Omega} (h - u) \leq -\frac{1}{4(2n + 1)},$$

which is a contradiction as soon as  $\varepsilon_0 < 1/4(2n + 1)$ . This proves the lemma.  $\square$

**Lemma 7.7.** *Let  $u \in \mathcal{P}_\infty^-(M)$  be a global solution and let  $(x_0, t_0) \in \partial\Omega$  be such that  $\overline{Q}_\varepsilon(x_0, t_0) \cap \partial\Omega$  consists only of low energy points for some  $\varepsilon > 0$ . Then the time derivative  $\partial_t u$  vanishes continuously at  $(x_0, t_0)$ :*

$$\lim_{(x, t) \rightarrow (x_0, t_0)} \partial_t u(x, t) = 0.$$

*Proof.* Consider the family of continuous functions  $\omega_r$  defined on  $\overline{Q}_\varepsilon(x_0, t_0) \cap \partial\Omega$  for every  $r > 0$  by

$$\omega_r(x, t) = W(r, x, t; u).$$

Functions  $\omega_r$  are continuous and converge pointwise to the balanced energy function  $\omega$  as  $r \rightarrow 0$ . Since  $\overline{Q}_\varepsilon(x_0, t_0) \cap \partial\Omega$  consists only of low energy points,  $\omega = A$  there. Hence

$$\omega_r \searrow A \quad \text{as } r \searrow 0 \text{ on } \overline{Q}_\varepsilon \cap \partial\Omega,$$

as follows from Weiss's monotonicity formula. From Dini's monotone convergence theorem it follows that the convergence  $\omega_r \rightarrow A$  is uniform. In particular, for any sequences  $(y_j, s_j) \rightarrow (x_0, t_0)$  and  $r_j \rightarrow 0$  we have

$$W(r_j, y_j, s_j; u) \rightarrow A.$$

Let now  $(x_j, t_j) \rightarrow (x_0, t_0)$  be the maximizing sequence for  $\partial_t u$  in the sense that

$$\partial_t u(x_j, t_j) \rightarrow m := \limsup_{(x,t) \rightarrow (x_0, t_0)} \partial_t u(x, t).$$

Let  $d_j = d^-(x_j, t_j) = \sup\{r : Q_r^-(x_j, t_j) \subset \Omega\}$  and  $(y_j, s_j) \in \partial_p Q_{d_j}^-(x_j, t_j) \cap \partial\Omega$ . Without loss of generality we may assume that

$$\frac{1}{d_j^2} u(d_j x + x_j, d_j^2 t + t_j) =: u_j(x, t) \rightarrow u_0(x, t)$$

in  $\mathbb{R}^n \times \mathbb{R}^-$  and

$$((y_j - x_j)/d_j, (s_j - t_j)/d_j^2) =: (\tilde{y}_j, \tilde{s}_j) \rightarrow (\tilde{y}_0, \tilde{s}_0) \in \partial Q_1^-.$$

Observe that since  $Q_1^- \subset \Omega(u_j)$ , we will have  $Q_1^- \subset \Omega(u_0)$  and we may assume that the convergence in  $Q_1^-$  is locally uniform in  $C_x^2 \cap C_t^1$ -norm. Thus

$$\partial_t u_0(0, 0) = \lim_{j \rightarrow \infty} \partial_t u_j(0, 0) = \lim_{j \rightarrow \infty} \partial_t u(x_j, t_j) = m$$

and

$$\partial_t u_0(x, t) = \lim_{j \rightarrow \infty} \partial_t u_j(x, t) = \lim_{j \rightarrow \infty} \partial_t u(d_j x + x_j, d_j^2 t + t_j) \leq m$$

for any  $(x, t) \in Q_1^-$ . Also since  $\partial_t u_0 = 0$  in  $Q_1^-$ , the maximum principle implies

$$(7.6) \quad \partial_t u_0 = m \quad \text{in } Q_1^-.$$

On the other hand

$$(7.7) \quad W(r, \tilde{y}_0, \tilde{s}_0; u_0) = \lim_{j \rightarrow \infty} W(r, \tilde{y}_j, \tilde{s}_j; u_j) = \lim_{j \rightarrow \infty} W(d_j r, y_j, s_j; u) = A$$

for every  $r > 0$ . In particular,  $u_0$  is homogeneous with respect to  $(\tilde{y}_0, \tilde{s}_0) \in \partial\Omega(u_0)$  in  $\mathbb{R}^n \times \mathbb{R}_{\tilde{s}_0}^-$ , and from Lemmas 6.2 and 6.3 we obtain that

$$(7.8) \quad u_0(x, t) = \frac{1}{2} ((x - \tilde{y}_0) \cdot e)_+^2 \quad \text{in } \mathbb{R}^n \times \mathbb{R}_{\tilde{s}_0}^-,$$

for a spatial unit vector  $e$ .

We want to show now that (7.6) and (7.8) contradict each other, unless  $m = 0$ . Indeed, if  $\tilde{s}_0 > -1$ ,  $Q_1^- \cap (\mathbb{R}^n \times \mathbb{R}_{\tilde{s}_0}^-)$  is nonempty and the contradiction is immediate. Next, if  $\tilde{s}_0 = -1$  and  $B_1 \cap \{(x - \tilde{y}_0) \cdot e > 0\} =: E$  is nonempty, we obtain that  $\partial_t u_0$  is discontinuous on  $E \times \{-1\} \subset \Omega(u_0)$ , which is not possible since  $u_0$  is caloric in  $\Omega(u_0)$ . Hence, the remaining case is when  $\tilde{s}_0 = -1$  and  $u_0$  vanishes on  $B_1 \times \{-1\}$ , which implies that  $u_0(x, t) = m(t + 1)$  in  $Q_1^-$ . Now, using that  $u_0(x, t)$  is analytic in variable  $x$  in  $\Omega(u_0)$ , we obtain that the whole strip  $\mathbb{R}^n \times (0, 1)$  is contained in

$\Omega(u_0)$  and that  $u_0(x, t) = m(t + 1)$  there. This is a contradiction, since  $u_0$  must be continuous across  $\{(x - \tilde{y}_0) \cdot e > 0\} \times \{-1\}$ .

This shows that  $\limsup \partial_t u(x, t) = 0$  as  $(x, t) \rightarrow (x_0, t_0)$ . Similarly one can prove that  $\liminf \partial_t v(x, t) = 0$ , which will conclude the proof of the lemma.  $\square$

*Proof of Theorem 7.5.*

*Step 1.* Without loss of generality we may assume  $(x_0, t_0) = (0, 0)$ . Consider then the rescaled functions  $u_r$ , which converge to a homogeneous global solution  $u_0 \in \mathcal{P}_\infty^-(M)$  over a subsequence  $r = r_j \rightarrow 0$ . Since  $(0, 0)$  is a low energy point,  $u_0(x, t) = \frac{1}{2}(x \cdot e_0)_+^2$  for some spatial direction  $e_0$ . Choose now a spatial direction  $e$  such that  $e \cdot e_0 > 1/2$ . Then

$$\partial_e u_0 - u_0 \geq (e \cdot e_0)(x \cdot e_0) - \frac{1}{2}(x \cdot e_0)^2 \geq 0 \quad \text{in } Q_1^-.$$

Since the functions  $u_r$  converge uniformly in  $C_x^{1,\alpha} \cap C_t^{0,\alpha}$ -norm on  $Q_1^-$ , for  $r = r_j$  sufficiently small, we will have

$$\partial_e u_r - u_r \geq -\varepsilon_0 \quad \text{in } Q_1^-,$$

where  $\varepsilon_0$  is the same as in Lemma 7.6. Applying Lemma 7.6 with  $h = \partial_e u_r$ , we obtain

$$(7.9) \quad \partial_e u_r - u_r \geq 0 \quad \text{in } Q_{1/2}^-.$$

*Step 2.* Next, we claim that for any  $\varepsilon > 0$  and small  $r < r(\varepsilon)$

$$(7.10) \quad u_r = 0 \quad \text{on } \{x \cdot e_0 \leq -\varepsilon\} \cap Q_{1/2}^-.$$

Indeed, this follows easily from the nondegeneracy Lemma 5.1.

Now, observe that (7.9) can be written as

$$(7.11) \quad \partial_e \left( e^{-(x \cdot e)} u_r \right) \geq 0.$$

So, integrating this along the direction  $e$  and using (7.10), we obtain that  $u_r \geq 0$  in  $Q_{1/2}^-$ , which after scaling back translates into  $u \geq 0$  in  $Q_{r/2}^-$ .

Moreover, (7.11) implies that for any point  $x_0 \in \partial\Omega_r(t) \cap B_{1/4}$  and  $-1/4 \leq t \leq 0$ , the cone  $x_0 + \mathcal{C}$ , where  $\mathcal{C} = \{-se : 0 \leq s \leq 1/4, e \cdot e_0 > 1/2\}$  is contained in  $\Omega_r^c(t)$ . Hence the time sections  $\partial\Omega(t)$  are Lipschitz regular in  $B_{r/4}$  for  $-r^2/4 \leq t \leq 0$ .

*Step 3.* We have proved now that  $u \geq 0$  in  $Q_{r/2}^-$ . A simple consequence of this is that for some  $r_1 < r$ , the intersection  $\partial\Omega \cap Q_{r_1}^-$  consists of low energy points. Indeed, first observe that there could be no zero energy points in  $Q_{r_1}^-$ , which follows from nonnegativity of  $u$ ; see Subsection 7.1. Next, if there are high energy points, they all should be below some  $t$ -level, with  $t = t_* < 0$ . Hence, if we take  $r_1 < \min(r/2, \sqrt{-t_*})$ , the intersection  $\partial\Omega \cap Q_{r_1}^-$  may consist only of low energy points.

Scaling parabolically with  $r < r_1$ , we see that  $\partial\Omega_r \cap Q_1^-$  consists of low energy points of  $u_r$ . Applying Lemma 7.7, we obtain an important fact that the time derivative  $\partial_t u_r$  vanishes continuously on  $\partial\Omega_r \cap Q_1^-$ . Consider then a caloric function

$$h = \partial_e u_r + \eta \partial_t u_r$$

in  $Q_1^- \cap \Omega_r$ , where  $|\eta| < \eta_0$  is a small constant. Observe that  $h = 0$  continuously on  $\partial\Omega_r \cap Q_1^-$ . Since  $\partial_t u_r$  are uniformly bounded, arguing as in Step 1 above, we

obtain that

$$(\partial_e u_r + \eta \partial_t u_r) - u_r \geq 0 \quad \text{in } Q_{1/2}^-$$

for  $r$  sufficiently small. Then, as in Step 2, we obtain the existence of space-time cones at every point on  $\partial\Omega$ , which implies the joint space-time Lipschitz regularity of  $\partial\Omega \cap Q_{r/2}$ .

The theorem is proved. □

### 8. POSITIVE GLOBAL SOLUTIONS

**Theorem 8.1.** *Let  $u \in \mathcal{P}_\infty^-(M)$  be a global solution and assume that  $u \geq 0$  in  $\mathbb{R}^n \times \mathbb{R}^-$ . Then  $\partial_t u \leq 0$  and  $\partial_{ee} u \geq 0$  in  $\Omega \cap (\mathbb{R}^n \times \mathbb{R}^-)$  for any spatial direction  $e$ . In particular, the time sections  $\Lambda(t)$  are convex for any  $t \leq 0$ .*

*Proof. Part 1.*  $\partial_t u \leq 0$ .

Indeed, assume the contrary, and let

$$m := \sup_{\Omega \cap (\mathbb{R}^n \times \mathbb{R}^-)} \partial_t u > 0.$$

Choose a maximizing sequence  $(x_j, t_j) \in \Omega \cap (\mathbb{R}^n \times \mathbb{R}^-)$  for the value  $m$ , i.e.

$$\lim_{j \rightarrow \infty} \partial_t u(x_j, t_j) = m.$$

Let  $d_j = d^-(x_j, t_j) = \sup\{r : Q_r^-(x_j, t_j) \subset \Omega\}$  and consider the scaled functions

$$(8.1) \quad u_j(x, t) = \frac{1}{d_j^2} u(d_j x + x_j, d_j^2 t + t_j).$$

Then functions  $u_j$  are uniformly  $C_x^{1,1} \cap C_t^{0,1}$  regular in  $\mathbb{R}^n \times \mathbb{R}^-$  and we can extract a converging subsequence to a global solution  $u_0$ . Since we assume  $u \geq 0$ , we have  $u_0 \geq 0$ . Therefore  $\Omega(u_0) = \{u_0 > 0\}$ . Next, observe that since  $Q_1^- \subset \Omega(u_j)$  by definition, we will have  $Q_1^- \subset \Omega(u_0)$ . In particular,  $H(u_0) = 1$  in  $Q_1^-$  and the convergence of  $u_j$  to  $u_0$  will be at least in  $C_x^2 \cap C_t^1$ -norm on  $Q_{1/2}^-$  and more generally on compact subsets of  $\Omega(u_0)$ . Hence

$$\partial_t u_0(0, 0) = \lim_{j \rightarrow \infty} \partial_t u_j(0, 0) = \lim_{j \rightarrow \infty} \partial_t u(x_j, t_j) = m.$$

On the other hand for every  $(x, t) \in \Omega(u_0)$

$$\partial_t u_0(x, t) = \lim_{j \rightarrow \infty} \partial_t u_j(x, t) = \lim_{j \rightarrow \infty} \partial_t u(d_j x + x_j, d_j^2 t + t_j) \leq m.$$

Since  $\partial_t u_0$  is caloric in  $Q_1^-$ , from the maximum principle we immediately obtain that  $\partial_t u_0 = m$  everywhere in  $Q_1^-$  and therefore

$$(8.2) \quad u_0(x, t) = mt + f(x)$$

in  $Q_1^-$ . Moreover, (8.2) is valid in the parabolically connected component of  $\Omega(u_0)$  that contains the origin. It is easy to see that this implies the representation (8.2) for every  $t \in (-f(x)/m, 0)$  with  $x \in B_1$ . Indeed, starting at  $(x, 0)$  and moving down along the vertical line  $\{x\} \times \mathbb{R}^-$  as long as  $u(x, t) > 0$ , we can extend the equality  $\partial_t u_0 = m$  (and thus (8.2)) from the point  $(x, t)$  to its small neighborhood by applying the maximum principle.

Thus, the free boundary becomes the graph of a function  $t = t(x) := -f(x)/m$ . Since  $(x, t(x)) \in \Lambda(u_0)$ , we must have  $\nabla u_0(x, t) = 0$  at  $t = t(x)$ . But  $\nabla u_0(x, t) = \nabla f(x)$  for  $0 > t > -t(x)$  and since  $\nabla u_0$  is continuous, we obtain

$$\nabla f(x) = 0$$

for every  $x \in B_1$ . Hence  $f(x) = c_0$  is constant in  $B_1$  and  $u(x, t) = mt + c_0$  in  $Q_1^-$ . Then  $H(u_0) = -m$  in  $Q_1^-$ , which means  $m = -1$ . This contradicts the assumption that  $m > 0$  and the first statement of Theorem 8.1 is proved.

*Part 2.*  $\partial_{ee}u \geq 0$  for any spatial unit vector  $e$ .

The reasoning is very similar to Part 1 above, so we will skip some details. Without loss of generality assume  $e = e_n = (0, \dots, 0, 1)$ . Let

$$-m = \inf_{\Omega \cap (\mathbb{R}^n \times \mathbb{R}^-)} \partial_{nn}u < 0$$

and let  $(x_j, t_j) \in \Omega \cap (\mathbb{R}^n \times \mathbb{R}^-)$  be the minimizing sequence for  $-m$ , i.e.

$$\lim_{j \rightarrow \infty} \partial_{nn}u(x_j, t_j) = -m.$$

Considering the rescaled functions  $u_j$  as in (8.1), extract a converging subsequence to a function  $u_0 \geq 0$ . As in Part 1, we have  $Q_1^- \subset \Omega(u_0)$ . From the locally  $C_x^2 \cap C_t^1$  convergence of  $u_j$  to  $u_0$  in  $\Omega(u_0)$ , we obtain that

$$\partial_{nn}u_0(0, 0) = -m$$

and

$$\partial_{nn}u_0(x, t) \geq -m$$

in  $\Omega(u_0)$ . Since  $\partial_{nn}u_0$  is caloric in  $\Omega(u_0)$ , the minimum principle implies that  $\partial_{nn}u_0 = -m$  in the parabolically connected component of  $\Omega(u_0)$ , in particular in  $Q_1^-$ . From there we obtain the representations

$$(8.3) \quad \partial_n u_0(x, t) = f(x', t) - m x_n$$

and

$$(8.4) \quad u_0(x, t) = g(x', t) + f(x', t) x_n - \frac{m}{2} x_n^2$$

in  $Q_1^-$  where  $x' = (x_1, \dots, x_{n-1})$ . Now let us choose a point  $(x', 0, t) \in Q_1^-$  and start moving in the direction  $e_n$ , as long as we stay in  $\Omega(u_0)$ . By applying the minimum principle for  $\partial_{nn}u_0$  while we move, we can prove  $\partial_{nn}u_0 = -m$  and both of the representations (8.3) and (8.4) as long as we stay in  $\Omega(u_0)$ . Observe however, sooner or later we will hit  $\Lambda(u_0)$ ; otherwise, if  $x_n$  becomes very large, (8.4) will imply  $u_0 < 0$ , which is impossible. Therefore let  $x_n = \xi(x', t)$  be the first value of  $x_n$  for which we hit  $\Lambda(u_0)$ . Then  $\partial_n u_0(x', x_n, t) = 0$  for  $x_n = \xi(x', t)$ ; hence  $\xi(x', t) = f(x', t)/m$ . Since we also have the condition  $u_0(x', x_n, t) = 0$  for  $x_n = \xi(x', t)$ , the representation (8.4) takes the form

$$u_0(x, t) = -\frac{m}{2} (x_n - \xi(x', t))^2,$$

which is not possible since  $u_0 \geq 0$ . This concludes the proof of the theorem.  $\square$

9. CLASSIFICATION OF GLOBAL SOLUTIONS

In this section we classify global solutions in  $\mathbb{R}^n \times \mathbb{R}^-$ . First we make some observations that follow from the previous sections.

For a given  $t_0 \leq 0$  consider the set  $\Lambda(t_0)$ . We claim that for  $x_0 \in \partial\Lambda(t_0)$ , the corresponding point  $(x_0, t_0) \in \partial\Lambda$  cannot be a zero energy point. Indeed, for zero energy point  $(x_0, t_0)$  there would exist  $r$  such that  $u = 0$  in  $Q_r^-(x_0, t_0)$  and in particular  $B_r(x_0) \subset \Lambda(t_0)$ ; see Subsection 7.1.

Next, if  $(x_0, t_0)$  is a high energy point, necessarily  $u(x, t) = m(t - t_0) + P(x - x_0)$  for  $t \leq t_0$ ,  $\Lambda(t_0)$  is a  $k$ -dimensional plane,  $k = 0, \dots, n$ , and all points on  $\Lambda(t_0)$  are high energy points; see Lemma 7.3.

Hence, if there is a low energy point  $(x_0, t_0)$ , then all points on  $\partial\Lambda(t_0) \times \{t_0\}$  are low energy. Also, according to Theorem 7.5, in that case the boundary  $\partial\Lambda(t_0)$  is a locally Lipschitz surface. In particular, the set  $\Lambda(t_0)$  is a regular closed set, i.e. the closure of its interior. We can say even more. Theorem 7.5 implies that for given  $R > 0$  there exists  $\delta = \delta(u, R, t_0) > 0$  such that  $u \geq 0$  in  $U_{\delta, R} := \mathcal{N}_\delta(\Lambda(t_0) \cap B_R) \times (t_0 - \delta^2, t_0)$ , where  $\mathcal{N}_\delta(E)$  denotes the  $\delta$ -neighborhood of the set  $E$ , and  $\partial\Lambda \cap U_{\delta, R}$  is a Lipschitz in space and time surface. Another important fact is that  $\partial_t u$  will be continuous up to  $\partial\Lambda$  in  $B_R \times (t_0 - \delta^2, t_0)$ . The latter implies that  $(\partial_t u)^\pm$  are subcaloric functions in  $B_R \times (t_0 - \delta^2, t_0)$ .

The main theorem of this section is as follows.

**Theorem 9.1.** *Let  $u \in \mathcal{P}_\infty^-(M)$  be a global solution and suppose that  $(0, 0)$  is a low energy point. Then  $u \geq 0$  in  $\mathbb{R}^n \times \mathbb{R}^-$ .*

Before proving the theorem, it is convenient to introduce the *balanced energy at  $\infty$*  of a global solution  $u \in \mathcal{P}_\infty^-(M)$ . We define it as

$$\omega_\infty = \lim_{r \rightarrow \infty} W(r; u).$$

In analogy with the construction from Section 7, consider the rescaled functions  $u_r$  and let  $r \rightarrow \infty$ . Then, over a sequence  $r = r_k \rightarrow \infty$ ,  $u_r$  will converge to a function  $u_\infty$  in  $\mathbb{R}^n \times \mathbb{R}^-$ , which will be a solution of (1.2). Moreover,  $u_\infty$  will be a homogeneous solution and

$$\omega_\infty = W(u_\infty).$$

Thus  $\omega_\infty$  can take only values 0,  $A$ , and  $2A$  by Lemmas 6.3 and 6.2. We say that  $\infty$  is a zero, low, and high energy point, respectively.

**Lemma 9.2.** *For  $u \in \mathcal{P}_\infty(M)$*

- (i)  $\omega_\infty = 0$  implies that  $u = 0$  in  $\mathbb{R}^n \times \mathbb{R}^-$ ;
- (ii) if both  $\omega = \omega_\infty = A$ , then  $u$  is a stationary halfspace solution of the form  $\frac{1}{2}(x \cdot e)_+^2$ .

*Proof.* In both cases we obtain that  $W(r; u)$  is constant, since  $\omega \leq W(r; u) \leq \omega_\infty$ . Hence  $u$  is a homogeneous solution with energy 0 or  $A$  and the lemma follows.  $\square$

When  $u \in \mathcal{P}_\infty(M)$  and  $\omega_\infty = 2A$ , then the shrink-down  $u_\infty$  is a polynomial solution

$$u_\infty(x, t) = mt + P_\infty(x),$$

where  $P_\infty(x)$  is a homogeneous quadratic polynomial. The next lemma shows what information we can extract, if  $P_\infty(x)$  is degenerate in some direction.

**Lemma 9.3.** *Suppose  $P_\infty(x)$  is degenerate in the direction  $e$ , i.e.  $\partial_e P_\infty = 0$ . Then  $\partial_e u$  has a sign in  $\mathbb{R}^n \times \mathbb{R}^-$ , i.e.  $\partial_e u \geq 0$  or  $\partial_e u \leq 0$  everywhere.*

*Proof.* Let  $\psi$  be a cut-off function with  $\psi = 1$  on  $B_{1/2}$ ,  $\text{supp } \psi \subset B_{3/4}$  and let  $\psi_r(x) = \psi(x/r)$ . Consider the scaled functions  $u_r$  in  $Q_1^-$ . Then  $(\partial_e u_r)^\pm$  are subcaloric and vanish at  $(0, 0)$ . Hence we can apply the [Caf93] monotonicity formula (Theorem 3.3) to obtain the estimate

$$\Phi(t; (\partial_e u_r)\psi) \leq C_0 \|(\partial_e u_r)\|_{L^2(Q_1^-)}^4,$$

for any  $0 < t < \tau_0$ , where  $\tau_0, C_0$  do not depend on  $r$ . Observe now that over a subsequence of  $r = r_k \rightarrow \infty$  for which  $u_r \rightarrow u_\infty$ ,  $\partial_e u_r$  converges uniformly to  $\partial_e u_\infty = 0$  in  $Q_1^-$ . Hence

$$\Phi(t; (\partial_e u_r)\psi) \rightarrow 0 \quad \text{as } r = r_k \rightarrow \infty$$

uniformly for  $0 < t < \tau_0$ . Scaling back to the function  $u$ , using that  $\partial_e u_r(x, t) = (1/r)\partial_e u(rx, r^2t)$ , we obtain

$$\Phi(t; (\partial_e u)\psi_r) = \Phi(t/r^2; (\partial_e u_r)\psi) \rightarrow 0 \quad \text{as } r = r_k \rightarrow \infty$$

uniformly for  $0 < t < \tau_0 r^2$ . Therefore for any fixed  $t > 0$ ,

$$\left( \frac{1}{t} \int_{-t}^0 \int_{B_{r/2}} |\nabla(\partial_e u)^+|^2 G(x, -s) dx ds \right) \left( \frac{1}{t} \int_{-t}^0 \int_{B_{r/2}} |\nabla(\partial_e u)^-|^2 G(x, -s) dx ds \right) \rightarrow 0$$

as  $r = r_k \rightarrow \infty$ , since  $\psi_r = 1$  on  $B_{r/2}$ . Passing to the limit, we obtain

$$\Phi(t; \partial_e u) = 0$$

for any  $t > 0$ . This is possible if and only if one of the functions  $(\partial_e u)^\pm$  vanishes in  $\mathbb{R}^n \times \mathbb{R}^-$ .

The lemma is proved. □

We will also need the following modification of Lemma 7.7.

**Lemma 9.4.** *Let  $u \in \mathcal{P}_\infty^-(M)$  be a global solution. Suppose that  $(x_0, t_0) \in \partial\Omega$  and  $\overline{Q}_\varepsilon(x_0, t_0) \cap \partial\Omega$  contains no high energy points for some  $\varepsilon > 0$ . Then*

$$m := \limsup_{(x,t) \rightarrow (x_0,t_0)} \partial_t u \leq 0.$$

*Proof.* The proof will follow the lines of the proof of Lemma 7.7.

As there, consider the continuous functions

$$\omega_r(x, t) = W(r, x, t; u)$$

for small  $r > 0$  and  $(x, t) \in \overline{Q}_\varepsilon(x_0, t_0) \cap \partial\Omega$ . Since there are no high energy points in a small neighborhood of  $(x_0, t_0)$ , we have

$$\lim_{r \searrow 0} \omega_r(x, t) \leq A.$$

Therefore, setting

$$\tilde{\omega} = \max(\omega_r, A),$$

we obtain

$$\tilde{\omega}_r(x, t) \searrow A \quad \text{as } r \searrow A$$

on  $\overline{Q_\varepsilon}(x_0, t_0) \cap \partial\Omega$ . Then, by Dini's theorem, the convergence  $\tilde{\omega}_r(x, t) \searrow A$  is uniform on  $\overline{Q_\varepsilon}(x_0, t_0) \cap \partial\Omega$ . Therefore, if  $(y_j, s_j) \rightarrow (x_0, t_0)$  and  $r_j \rightarrow 0$  are any sequences, we have

$$\lim_{j \rightarrow \infty} \tilde{\omega}_{r_j}(y_j, s_j) = A.$$

This implies that

$$\limsup_{j \rightarrow \infty} W(r_j, y_j, s_j; u) \leq A.$$

Now, having this, take a maximizing sequence  $(x_j, t_j) \rightarrow (x_0, t_0)$  such that

$$\lim_{j \rightarrow \infty} \partial_t u(x_j, t_j) = m.$$

Assume  $m > 0$ . Then scaling  $u$  around  $(x_j, t_j)$  by the parabolic distance  $d_j$ , precisely as in the proof of Lemma 7.7, we can extract a converging subsequence to a global solution  $u_0$ . Then, again, we can prove the identity (7.6). However, instead of equality (7.7) we will have the inequality

$$W(r, \tilde{y}_0, \tilde{s}_0; u_0) = \lim_{j \rightarrow \infty} W(r, \tilde{y}_j, \tilde{s}_j; u_j) = \lim_{j \rightarrow \infty} W(r_j, y_j, s_j; u) \leq A$$

for any  $r > 0$ . Applying Lemma 9.2, we see that either

$$u_0(x, t) = \frac{1}{2}((x - \tilde{y}_0) \cdot e)_+^2 \quad \text{for } t \leq \tilde{s}_0$$

or

$$u_0(x, t) = 0 \quad \text{for } t \leq \tilde{s}_0.$$

In the first case, we finish the proof as in Lemma 7.7. In the second case, we have necessarily  $\tilde{s}_0 = -1$ , since  $H(u_0) = 1$  in  $Q_1^-$  and we obtain the representation

$$u_0(x, t) = m(t + 1) \quad \text{in } Q_1^-.$$

Hence  $H(u_0) = -m < 0$ , while we must have  $H(u_0) = 1$ .

This concludes the proof of the lemma. □

*Remark 9.5.* Under the conditions of Lemma 9.4, the limit

$$\liminf_{(x,t) \rightarrow (x_0,t_0)} \partial_t u$$

can have only two possible values: 0 or  $-1$ . The same proof as above applies.

*Proof of Theorem 9.1.* First, we note that  $u$  cannot have zero balanced energy at  $\infty$ , and if it has low energy at  $\infty$ , the theorem readily follows from Lemma 9.2. So we need to consider only the case when  $\omega_\infty = 2A$ . Then let  $u_\infty(x, t) = mt + P_\infty(x)$  be as above, a shrink-down limit of rescaled functions  $u_r$ , as  $r = r_k \rightarrow \infty$ .

*Step 1: Dimension reduction.* Suppose that there exists a shrink-down  $u_\infty(x, t)$  for which the homogeneous quadratic polynomial  $P_\infty(x)$  is degenerate in the direction  $e$ . Then by Lemma 9.3, we may assume  $\partial_e u \geq 0$  (otherwise we will have  $\partial_{-e} u \geq 0$  and will just change the direction of  $e$ ). Also, without loss of generality, let  $e = e_n = (0, \dots, 0, 1)$ . Since we assume that  $(0, 0)$  is a low energy point, from Theorem 7.5 it follows that  $\partial\Lambda(0)$  is a Lipschitz surface in  $\mathbb{R}^n$  and hence the interior of  $\Lambda(0)$  is nonempty. Let  $B_\delta(x_0) \subset \Lambda(0)$ ,  $x_0 = (x'_0, a)$ . We claim now that

$$(9.1) \quad u(x, 0) = 0 \quad \text{for } x = (x', x_n) \in B'_\delta(x'_0) \times (-\infty, a).$$

Indeed, for  $x' \in B'_\delta(x'_0)$  define

$$b(x') = \inf\{b : u(x', s, 0) = 0 \text{ for all } s \in [b, a]\}.$$

Then obviously  $b(x') \leq a$  and  $\xi := (x', b(x')) \in \partial\Lambda(0)$ , if  $b(x')$  is finite. Moreover,  $(\xi, 0)$  can be only a low energy point; see the discussion preceding Theorem 9.1. Then, by Theorem 7.5, there is  $r > 0$  such that  $u \geq 0$  in  $Q_r^-(\xi, 0)$ , in particular  $u(x, 0) \geq 0$  in  $B_r(\xi)$ . On the other hand, since  $u(\xi, 0) = 0$  and  $\partial_n u \geq 0$ ,  $u(x', s, 0) \leq 0$  for  $s \in (b(x') - r, b(x'))$ . Hence  $u(x', s, 0) = 0$  for  $s \in (b(x') - r, b(x'))$  and we arrive at a contradiction, if  $b(x')$  is finite. Thus, (9.1) follows.

Now, for every  $\tau \geq 0$  define the shifts of  $u$  in the direction  $e_n$

$$v_\tau(x', x_n, t) = u(x', x_n - \tau, t).$$

Since  $\partial_n u \geq 0$ , the functions  $v_\tau$  decrease as  $\tau \rightarrow \infty$  and therefore there exist the limit

$$v = \lim_{\tau \rightarrow \infty} v_\tau.$$

Moreover, as follows from (9.1),  $v_\tau(x_0, 0) = 0$  for all  $\tau \geq 0$  and we have the uniform estimate

$$|v_\tau(x, t)| \leq C(M)(|x - x_0|^2 + |t|)$$

in  $\mathbb{R}^n \times \mathbb{R}^-$ . Hence  $v$  is finite everywhere in  $\mathbb{R}^n \times \mathbb{R}^-$  and thus is a solution of (1.2). Moreover, clearly,  $v$  is independent of the direction  $e_n$ . So we may think of  $v = v(x', t)$  as a solution of (1.2) in  $\mathbb{R}^{n-1} \times \mathbb{R}^-$ . Observe also that

$$u(x', x_n, t) \geq v(x', t),$$

so if we prove that  $v \geq 0$  in  $\mathbb{R}^{n-1} \times \mathbb{R}^-$ , we will be done.

Now, consider several cases. Suppose that for every  $\varepsilon > 0$ ,  $v$  has a low energy point  $(x', t)$  with  $-\varepsilon^2 \leq t \leq 0$ . Taking such a point as the origin, we arrive at the conditions of the theorem, but with the reduced dimension. So if the theorem is true for the dimension  $n - 1$ , we conclude that  $v \geq 0$  for  $t \leq -\varepsilon^2$  and letting  $\varepsilon \rightarrow 0$  we complete the proof.

The next case would be that there are no low energy points of  $v$  in  $\mathbb{R}^{n-1} \times (-\varepsilon^2, 0]$  for some  $\varepsilon > 0$ . Observe that (9.1) implies  $B'_\delta(x'_0) \subset \Lambda_v(0)$ . Suppose for a moment that  $(0, 0) \in \Gamma(v)$ . Then it is a high energy point and  $v$  is a polynomial solution. Since  $\Lambda_v(0)$  has nonempty interior, it will be possible only if  $v(x, t) = -t > 0$  and we will be done. Thus, we may assume that for some small  $0 < \eta < \varepsilon$ ,  $Q'_\eta \subset \Lambda_v$ . Since there are no low energy points for  $-\eta^2 < t \leq 0$ ,  $\partial\Lambda_v(t)$  must be empty, thus implying that  $\Lambda_v(t) = \mathbb{R}^{n-1}$  for  $-\eta^2 < t \leq 0$ . The latter means that  $v = 0$  in  $\mathbb{R}^{n-1} \times (-\eta^2, 0]$ .

Next, let  $\eta = \eta_*$  be maximal (possibly infinite) with the property that  $v = 0$  on  $\mathbb{R}^{n-1} \times (-\eta^2, 0]$ . If  $\eta_* = \infty$ , we will have that  $v = 0$  identically. If  $\eta_*$  is finite, then the arguments above show that  $v(x', t) = (\eta_*^2 - t)_+$  and we are done.

Thus, in all possible cases  $v \geq 0$ , which implies  $u \geq 0$ .

To complete the dimension reduction, we note that for  $n = 0$  the statement of the theorem is trivial. Indeed,  $\mathbb{R}^0 = \{0\}$ ,  $u(0, 0) = 0$  and  $H(u) = -\partial_t u \geq 0$  imply that  $u(0, t) \geq 0$  for  $t \leq 0$ .

*Step 2: Nondegenerate  $P_\infty$ .* The reasoning above allows us to reduce the problem to the case when  $P_\infty(x)$  is nondegenerate for every shrink-down  $u_\infty$  over every sequence  $r = r_k \rightarrow \infty$ .

**Lemma 9.6.** *Suppose  $u$  has no high energy points in  $\mathbb{R}^n \times \mathbb{R}^-$  and for every shrink-down  $u_\infty(x, t) = mt + P_\infty(x)$  the polynomial  $P_\infty(x)$  is nondegenerate. Then there exist a shrink-down with  $m = 0$ .*

*Proof.* First, suppose that  $\Lambda \cap (\mathbb{R}^n \times \mathbb{R}^-)$  is bounded. Let  $(x_0, t_0)$  be a point with minimal  $t$ -coordinate. Then, obviously,  $(x_0, t_0)$  is a high energy point, contradicting the assumption. Therefore, there exists a sequence  $(x_k, t_k) \in \Lambda$  such that

$$r_k := \max(|x_k|, \sqrt{-t_k}) \rightarrow \infty.$$

Consider then the scale functions  $u_{r_k}$ . Then

$$(x_k/r_k, t_k/r^2) \in \partial_p Q_1^- \cap \Lambda(u_{r_k}).$$

Hence if  $u_\infty$  is a shrink-down over a subsequence of  $r = r_k \rightarrow \infty$ , we will have

$$\partial_p Q_1^- \cap \Lambda(u_\infty) \neq \emptyset.$$

Since  $P_\infty(x)$  is nondegenerate, this may happen only if  $m = 0$ . □

Lemma 9.6 has the consequence that only the following three cases are possible if for every shrink-down  $u_\infty(x, t) = mt + P_\infty(x)$  the polynomial  $P_\infty$  is nondegenerate:

1. there exists a high energy point  $(x_0, t_0) \in \mathbb{R}^n \times \mathbb{R}^-$  and  $m > 0$ ;
2. there exists a high energy point  $(x_0, t_0) \in \mathbb{R}^n \times \mathbb{R}^-$  and  $m \leq 0$ ;
3. there are no high energy points in  $\mathbb{R}^n \times \mathbb{R}^-$  and  $m = 0$  for a shrink-down over a sequence  $r = r_k \rightarrow \infty$ .

We will treat these three cases separately.

*Case 1.* There exists a high energy point  $(x_0, t_0) \in \mathbb{R}^n \times \mathbb{R}^-$  and  $m > 0$ .

We show below that this configuration is not possible. Since  $(x_0, t_0)$  is a high energy point, we have that  $u(x, t) = c(t - t_0) + P(x - x_0)$  for  $t \leq t_0$  and  $t_0 < 0$ . Moreover, considering the shrink-down, we realize that  $c = m$  and  $P = P_\infty$ . Hence

$$u(x, t) = m(t - t_0) + P_\infty(x - x_0) \quad \text{for } t \leq t_0.$$

Consider next a blow-up of  $u$  at  $(x_0, t_0)$

$$w(x, t) = \lim_{r=r_k \rightarrow 0} \frac{1}{r^2} u(rx + x_0, r^2 t + t_0),$$

which is defined in  $\mathbb{R}^n \times \mathbb{R}$ . By a version of Weiss's monotonicity formula for the upper halfspace, see Theorem 3.1 in [Wei99], analogous to our Theorem 3.4, and by arguments as in the beginning of Section 7, we obtain that  $w$  is a homogeneous solution not only in  $\mathbb{R}^n \times \mathbb{R}^-$  but also in  $\mathbb{R}^n \times \mathbb{R}$ . We actually claim that

$$w(x, t) = mt + P_\infty(x)$$

in  $\mathbb{R}^n \times \mathbb{R}$ . Observe that the equality is obvious for  $t \leq 0$ . Moreover, by the comparison principle,  $w(x, t) \geq mt + P_\infty(x)$  for  $t \geq 0$ , since  $H(w) \leq 1$ . The equality for  $t \geq 0$  will follow (by uniqueness) once we show that the coincidence set  $\Lambda_w$  consists only of  $(0, 0)$ . If there is a point  $\Lambda_w$  with  $t > 0$ , it is necessarily a low energy point. It cannot be a high energy point, since  $P_\infty$  is nondegenerate and  $m > 0$ . It cannot be a zero energy point either, since  $w$  is homogeneous for  $t > 0$ . Thus, by Lemma 7.7,

$$\partial_t w = 0 \quad \text{on } \partial \Lambda_w \cap (\mathbb{R}^n \times (0, \infty)).$$

Besides,

$$\partial_t w = m > 0 \quad \text{on } (\mathbb{R}^n \setminus \{0\}) \times \{0\}, \quad H(\partial_t w) = 0 \quad \text{in } (\mathbb{R}^n \times (0, \infty)) \setminus \Lambda_w.$$

Also since  $\partial_t w$  is uniformly bounded, we obtain that

$$\partial_t w > 0 \quad \text{in } (\mathbb{R}^n \times \mathbb{R}^+) \setminus \Lambda_w.$$

Now, suppose that  $(x_1, t_1) \in \Lambda_w$  with  $t_1 > 0$ . Since  $w(x, t) \geq mt + P_\infty(x)$ , it follows that  $x_1 \neq 0$  and we can assume that  $t_1 > 0$  is minimal for fixed  $x_1$ . Then  $w(x_1, t_1) = 0$  and  $w(x_1, t) < 0$  for  $0 \leq t < t_1$ . But this is impossible by Theorem 7.5, since we know that  $(x_1, t_1)$  has a low energy.

Thus, we obtain that  $w(x, t) = mt + P_\infty(x)$  everywhere in  $\mathbb{R}^n \times \mathbb{R}$ . Going back to the function  $u$ , we use this fact to show that  $(x_0, t_0)$  is an isolated free boundary point of  $u$ . If there is a sequence  $(x_k, t_k) \in \Lambda_u \setminus \{(x_0, t_0)\}$  converging to  $(x_0, t_0)$ , let

$$r_k = \max\{|x_k - x_0|, \sqrt{|t_k - t_0|}\} \rightarrow 0.$$

Then for the rescalings  $w_{r_k}(x, t) = (1/r_k^2)u(r_k x + x_0, r_k^2 t + t_0)$  we have

$$((x_k - x_0)/r_k, (t_k - t_0)/r_k^2) \in \Lambda(w_{r_k}).$$

Thus, in the limit we obtain that  $\Lambda_w \cap (\partial Q_1)$  is nonempty, which is a contradiction.

By a similar argument, we can establish that  $\Lambda_u$  is bounded. Indeed, if there is a sequence  $(x_k, t_k)$  with  $r_k = |x_k| \rightarrow \infty$  and  $t_0 \leq t_k \leq 0$ , then

$$(|x_k|/r_k, t_k/r_k^2) \in \Lambda(u_{r_k})$$

for the rescalings  $u_{r_k}$  around the origin. Thus,  $\Lambda(u_\infty) \cap (\partial B_1 \times \{0\})$  is nonempty for a shrink-down  $u_\infty$ , which is not possible. Thus, the set  $\Lambda_u$  is compact. Since  $(x_0, t_0) \in \Lambda_u$  is isolated, there exists a point  $(x_*, t_*) \in \Lambda_u \setminus \{(x_0, t_0)\}$  with a minimal  $t$ -coordinate. Note that  $t_* > t_0$ . The point  $(x_*, t_*)$  can neither be a zero energy point nor a low energy point (by Theorem 7.5). It cannot be a high energy point either, since  $m > 0$  and  $P_\infty$  is nondegenerate. Thus, we arrive at a contradiction, which shows that this case is not possible.

Before we proceed to the remaining two cases, we prove the following lemma.

**Lemma 9.7.** *Suppose in the representation  $u_\infty(x, t) = mt + P_\infty(x)$  the polynomial  $P_\infty(x)$  is nondegenerate and  $m \leq 0$ . Then  $\partial_t u \leq 0$  in  $\mathbb{R}^n \times \mathbb{R}^-$ .*

*Proof.* The basic idea is that  $\partial_t u$  is 0 on the “regular” part of the free boundary and nonpositive at infinity. We subdivide the proof into two cases.

(i) There are no high energy points of  $u$  in  $\mathbb{R}^n \times \mathbb{R}^-$ .

Then Lemma 9.4 implies that  $(\partial_t u)_+$  is continuous and therefore subcaloric in  $\mathbb{R}^n \times \mathbb{R}^-$ . Consider then the scaled functions  $u_r \rightarrow u_\infty$  in  $Q_2^-, r = r_k \rightarrow \infty$ . Since  $\Lambda(u_\infty) = \{0\} \times \mathbb{R}^-$ , the convergence will be at least  $C_x^2 \cap C_t^1$  in  $Q_1^- \setminus (B_\varepsilon \times [-1, 0])$  for any  $\varepsilon > 0$ . In particular, for  $r = r_k$  very large,

$$\begin{aligned} (\partial_t u_r)_+ &\leq \varepsilon && \text{on } \partial_p Q_1 \setminus (B_\varepsilon \times \{-1\}), \\ (\partial_t u_r)_+ &\leq C(M) && \text{on } B_\varepsilon \times \{-1\}. \end{aligned}$$

Hence if  $v_\varepsilon$  is the solution of the Dirichlet problem for the heat equation with boundary data

$$\begin{aligned} v &= \varepsilon && \text{on } \partial_p Q_1 \setminus (B_\varepsilon \times \{-1\}), \\ v &= C(M) && \text{on } B_\varepsilon \times \{-1\}, \end{aligned}$$

we will have

$$(\partial_t u_r)_+ \leq v_\varepsilon \quad \text{in } Q_1^-.$$

It is not hard to see that  $v_\varepsilon \leq c(\varepsilon) \rightarrow 0$  as  $\varepsilon \rightarrow 0$  uniformly in  $Q_{1/2}^-$ , and hence

$$(\partial_t u_r)_+ \leq c(\varepsilon) \quad \text{in } Q_{1/2}^-.$$

Scaling back to  $u$ , we obtain

$$(\partial_t u)_+ \leq c(\varepsilon) \quad \text{in } Q_{r/2}^-.$$

Letting  $r = r_k \rightarrow \infty$  and then  $\varepsilon \rightarrow 0$ , we obtain the claim of the lemma.

(ii) There is a high energy point  $(x_0, t_0) \in \mathbb{R}^n \times \mathbb{R}^-$ . Observe that  $x_0$  is unique, since  $P_\infty$  is nondegenerate. Also  $t_0$  is unique, unless  $m = 0$ . If the latter is the case, we will assume  $t_0$  is the maximal value of  $t$  for which  $(x_0, t)$  is a high energy point.

If  $t_0 = 0$ , we are done. If  $t_0 < 0$ , Lemma 9.4 implies that  $(\partial_t u)_+$  is continuous and thus subcaloric in  $\mathbb{R}^n \times (t_0, 0)$ . We want to show that it in fact vanishes there.

Considering as above the scaled functions  $u_r$  and their convergence to  $u_\infty$ , as well as the fact that  $\partial_t u_\infty = m \leq 0$ , scaling back to  $u$ , we obtain that

$$(\partial_t u)_+ \leq \varepsilon \quad \text{on } \partial B_r \times (t_0, 0).$$

Moreover, since every point  $(x, t_0)$  is in  $\Omega$  except for  $(x_0, t_0)$ , we will also have

$$\begin{aligned} (\partial_t u)_+ &= 0 && \text{on } (B_r \setminus B_\varepsilon(x_0)) \times \{t_0\}, \\ (\partial_t u)_+ &\leq C(M) && \text{on } (B_\varepsilon(x_0)) \times \{t_0\}. \end{aligned}$$

Hence if  $w_{\varepsilon,r}$  is a solution to the Dirichlet problem for the heat equation in  $B_r \times (t_0, 0)$  with the boundary values

$$\begin{aligned} w &= \varepsilon && \text{on } \partial_p B_r \times (t_0, 0) \setminus (B_\varepsilon(x_0) \times \{t_0\}), \\ w &= C(M) && \text{on } B_\varepsilon(x_0) \times \{t_0\}, \end{aligned}$$

we will have

$$(\partial_t u)_+ \leq w_{\varepsilon,r} \quad \text{in } B_r \times (t_0, 0).$$

It is easy to see that as  $r \rightarrow \infty$  and  $\varepsilon \rightarrow 0$ ,  $w_{\varepsilon,r} \rightarrow 0$  uniformly on compact subsets of  $\mathbb{R}^n \times (t_0, 0)$ . Hence  $\partial_t u \leq 0$  in  $\mathbb{R}^n \times (t_0, 0)$  as well and the proof is complete.  $\square$

We now resume the proof of Theorem 9.1.

*Case 2.* There exists a high energy point  $(x_0, t_0) \in \mathbb{R}^n \times \mathbb{R}^-$  and  $m \leq 0$ .

Unless  $m = 0$ ,  $(x_0, t_0)$  is unique. If  $m = 0$ , assume that  $t_0$  is the maximal  $t$  such that there exists a high energy point at time  $t = t_0$ .

Arguing as in Case 1 above, we obtain that

$$u(x, t) = m(t - t_0) + P_\infty(x - x_0) \quad \text{for } t \leq t_0$$

and that  $\Lambda \subset \mathbb{R}^n \times [t_0, 0]$  is bounded. Since  $m \leq 0$ , Lemma 9.7 implies that  $\partial_t u \leq 0$  in  $\mathbb{R}^n \times \mathbb{R}^-$ . Next we consider the cross-sections  $\Lambda(t)$ . Since there are no high energy points for  $t_0 < t < 0$ ,  $\partial\Lambda(t)$  are Lipschitz surfaces and the interiors of  $\Lambda(t)$  are nonempty, provided  $\Lambda(t)$  are nonempty. Now let  $U(0)$  be a connected component of the interior of  $\Lambda(0)$ . Then Theorem 7.5 implies that there exist an open set  $W$  such that  $U(0) \subset\subset W$  and  $u \geq 0$  in  $W \times (-\delta, 0)$  for a small  $\delta > 0$ . Then  $\partial_t u \leq 0$  implies  $u \geq 0$  in  $W \times \mathbb{R}^-$ . Moreover  $\partial_t u \leq 0$  implies also that

$$U(t) := W \cap \text{Int}(\Lambda(t)) \searrow \text{ as } t \searrow.$$

Since  $U(t_0) = \emptyset$ , there exist  $t_* \in [t_0, 0)$  such that  $U(t) = \emptyset$  for  $t < t_*$  and  $U(t) \neq \emptyset$  for  $t_* < t < 0$ . Then the intersection

$$K_* = \bigcap_{t_* < t < 0} \overline{U(t)}$$

is nonempty. Choose now any  $x_* \in K_*$ . Then obviously  $(x_*, t_*) \in \partial\Lambda$ , and we claim that  $(x_*, t_*)$  is a high energy point. Clearly, it is not a zero energy point. Also, it is not a low energy point by Theorem 7.5. Hence  $(x_*, t_*)$  is a high energy

point. Since  $P_\infty$  is nondegenerate, necessarily  $x_* = x_0$ . (We also have  $t_* = t_0$ .) In particular,

$$x_0 \in \overline{U(0)}.$$

This implies immediately that  $U(0)$  is the only connected component of the interior of  $\Lambda(0)$ . Starting at any time  $t \in (t_0, 0)$ , we can prove a similar statement for  $\Lambda(t)$ . Thus

$$\Lambda(t) = \overline{U(t)}.$$

Moreover, we have that  $u \geq 0$  in  $W \times \mathbb{R}^- \supset \Lambda$ . In particular,  $P_\infty(x - x_0) = u(x, t_0) \geq 0$  in a neighborhood  $W$  of  $x_0$ , thus implying that  $P_\infty$  is positive definite.

Now let  $c > 0$  be small enough such that

$$P_\infty(x) \geq c|x|^2.$$

Also let

$$\partial_t u \geq -C = -C(M).$$

Then

$$u(x, t) \geq c|x - x_0|^2 - C(t - t_0)$$

in  $\mathbb{R}^n \times (t_0, 0)$ . Consequently,

$$(9.2) \quad u(x, t) > 0 \quad \text{for } x \in \mathbb{R}^n \setminus B_{\kappa(t)}(x_0) \text{ and } t_0 < t < 0,$$

where  $\kappa(t) = \sqrt{C(t - t_0)/c}$ . Consider now the set  $\Omega_- = \{u < 0\} \subset \mathbb{R}^n \times (t_0, 0)$  and suppose it is nonempty. Then

$$\Omega_-(t) \subset B_{\kappa(t)}$$

for  $t_0 < t < 0$ , by (9.2). In particular,  $\Omega_-$  is bounded. Hence there exists a point  $(x_1, t_1) \in \overline{\Omega_-}$  with minimal  $t$ -coordinate. Then  $t_1 \geq t_0$  and  $u(x, t_1) \geq 0$  for all  $x$ . Also since  $u(x_1, t_1) = 0$ , we obtain that  $\nabla u(x_1, t_1) = 0$ . Hence

$$(x_1, t_1) \in \Lambda \cap \overline{\Omega_-}.$$

This, however, contradicts the fact that  $u \geq 0$  in a neighborhood of  $\Lambda$ . Thus,  $\Omega_- = \emptyset$  and  $u \geq 0$  in  $\mathbb{R}^n \times \mathbb{R}^-$ .

*Case 3.* There are no high energy points in  $\mathbb{R}^n \times \mathbb{R}^-$  and  $m = 0$ .

Then Lemma 9.7 implies that  $\partial_t u \leq 0$  in  $\mathbb{R}^n \times \mathbb{R}^-$ .

We start with the claim that  $\Lambda(t)$  are bounded sets for  $-t$  sufficiently large. More specifically, we claim

$$(9.3) \quad \Lambda(t) \subset B_{\sqrt{-t}}$$

for  $t \leq t_0$ . Indeed, assume the contrary. Then there is a sequence  $t = t_k \rightarrow -\infty$  such that (9.3) does not hold, and therefore we can find  $x_k \in \Lambda(t_k)$  with

$$|x_k| \geq \sqrt{-t_k}.$$

Now let  $r = r_k = |x_k|$  and consider the scaled functions  $u_r$ . Then

$$(\tilde{x}_k, \tilde{t}_k) := (x_k/r_k, t_k/r_k^2) \in \Lambda(u_r) \cap (\partial B_1 \times [-1, 0]).$$

Hence passing to the limit over a subsequence of  $r = r_k \rightarrow \infty$ , we obtain that  $\Lambda(u_\infty) \cap (\partial B_1 \times [-1, 0])$  is nonempty. However, this is impossible if  $P_\infty$  is nondegenerate. Hence (9.3) should hold for  $t \leq t_0$ .

Next, suppose  $\Lambda(t)$  is empty for all  $t \leq t_0$ . Then  $u$  will satisfy  $H(u) = 1$  everywhere in  $\mathbb{R}^n \times \mathbb{R}_{t_0}^-$  and thus will have the form  $u(x, t) = ct + P(x)$  for  $t \leq t_0$ . Considering the shrink-down (recall we assume  $m = 0$ ), we find that

$$u(x, t) = P_\infty(x - x_1)$$

for  $t \leq t_0$ , where  $x_1 \in \mathbb{R}^n$  is some point. But then  $(x_1, t_0)$  is a high energy point, and we assume there are none.

Hence, without loss of generality we may assume that  $\Lambda(t_0)$  is nonempty. Since there are no high energy points for  $t \leq 0$ , the sets  $\Lambda(t)$  have nonempty interiors and  $\partial\Lambda(t)$  are Lipschitz surfaces, provided  $\Lambda(t)$  themselves are nonempty.

Let  $U(t_0)$  be a connected component of the interior of  $\Lambda(t_0)$ . Then we make a construction similar to the one in Case 2 above. There exists an open set  $W \subset \mathbb{R}^n$  such that  $U(t_0) \subset\subset W$  and  $u \geq 0$  in  $W \times (t_0 - \delta, t_0)$  for a small  $\delta > 0$ . Then from  $\partial_t u \leq 0$  we obtain that in fact  $u \geq 0$  in  $\mathbb{R}^n \times \mathbb{R}_{t_0}^-$ . Moreover  $u > 0$  in  $W \setminus \overline{U(t_0)} \times \mathbb{R}_{t_0}^-$ . Also,  $\partial_t u \leq 0$  implies that

$$U(t) := W \cap \text{Int}(\Lambda(t)) \searrow \text{ as } t \searrow.$$

Consider then the intersection

$$K = \bigcap_{t \leq t_0} \overline{U(t)}.$$

Since the sets  $\overline{U(t)}$  are compact,  $K$  is empty if and only if  $\overline{U(t)} = W \cap \Lambda(t)$  is empty for some  $t \leq t_0$ . If this is so, let  $t_*$  be such that  $W \cap \Lambda(t) = \emptyset$  for  $t < t_*$  and  $W \cap \Lambda(t_*)$  is nonempty. Take any  $x_* \in \Lambda(t_*)$ . Since  $W \cap \Lambda(t) = \emptyset$  for  $t < t_*$ ,  $u > 0$  in  $W \times (-\infty, t_*)$  and in particular  $(x_*, t_*)$  is a high energy point, and we assume there are none.

Thus  $K$  is nonempty and we can choose  $x_0 \in K$ . Then  $(x_0, t) \in \Lambda$  for all  $t \leq t_0$  and we obtain the estimate

$$u(x, t) \leq C(M)(|x - x_0|^2)$$

for  $x \in \mathbb{R}^n$  and  $t \leq t_0$ . Consider now the time shifts

$$v_\tau(x, t) = u(x, t - \tau)$$

defined in  $\mathbb{R}^n \times \mathbb{R}_{t_0}^-$ . Then from the estimate above and the monotonicity of  $u(x, t)$  in  $t$  the limit

$$v_\infty(x, t) = \lim_{\tau \rightarrow \infty} v_\tau(x, t)$$

exists and is finite everywhere in  $\mathbb{R}^n \times \mathbb{R}_{t_0}^-$ . Thus  $v_\infty$  is also a solution of (1.2). Moreover, it is easy to see that  $v_\infty$  is independent of  $t$ , so  $v_\infty = v_\infty(x)$  is a stationary global solution of (1.2).

As follows from [CKS00], the stationary global solutions are either polynomial or nonnegative. Observe now that  $x_0 \in \Lambda(v_\infty)$  and  $v \geq 0$  in the neighborhood  $W$  of  $x_0$ . Hence if  $v_\infty$  is a polynomial solution, the polynomial must be positive semidefinite. Therefore in any case we have  $v_\infty \geq 0$ .

Now, for the positive global solutions it is known that the set  $\Lambda(v_\infty)$  is convex, hence connected. Since  $x_0 \in \Lambda(v_\infty)$ ,  $x_0 \in \overline{U(t_0)} \subset W$  and  $v_\infty > 0$  in  $W \setminus \overline{U(t_0)}$ , the only possibility is that

$$\Lambda(v_\infty) \subset \overline{U(t_0)}.$$

A simple consequence of this is that the interior of  $\Lambda(t_0)$  consists only of one connected component. Now, if we made our construction starting at any  $t \leq t_0$ , we

would come to the conclusion that the interior of  $\Lambda(t)$  has at most one component. Hence

$$\Lambda(t) = \overline{U(t)}.$$

Also, we obtain that  $u \geq 0$  in the neighborhood  $W \times \mathbb{R}_{t_0}^-$  of  $\Lambda \cap (\mathbb{R}^n \times \mathbb{R}_{t_0}^-)$ . Then all the free boundary points in  $\mathbb{R}^n \times \mathbb{R}_{t_0}^-$  are low energy and Lemma 7.7 implies that  $\partial_t u = 0$  continuously on  $\partial\Lambda \cap (\mathbb{R}^n \times \mathbb{R}_{t_0}^-)$  and therefore  $\partial_t u$  is supercaloric in  $\mathbb{R}^n \times \mathbb{R}_{t_0}^-$ . In fact, we claim that

$$(9.4) \quad \partial_t u = 0 \quad \text{in } \mathbb{R}^n \times \mathbb{R}_{t_0}^-.$$

Consider the scaled functions  $u_r \rightarrow u_\infty$  in  $Q_2^-$ . Since  $\Lambda(u_\infty) = \{0\} \times \mathbb{R}^-$ , the convergence will be at least  $C_x^2 \cap C_t^1$  in  $Q_1^- \setminus (B_\varepsilon \times [-1, 0])$  for any  $\varepsilon > 0$ . In particular, for  $r = r_k$  very large,

$$\begin{aligned} -\varepsilon \leq \partial_t u_r \leq 0 & \quad \text{on } \partial_p Q_1 \setminus (B_\varepsilon \times \{-1\}), \\ -C(M) \leq \partial_t u_r \leq 0 & \quad \text{on } B_\varepsilon \times \{-1\}. \end{aligned}$$

Moreover,  $\partial_t u_r$  is supercaloric in  $B_1^- \times (-1, t_0/r^2)$ , so if  $v_\varepsilon$  is the solution of the Dirichlet problem for the heat equation with boundary data

$$\begin{aligned} v &= \varepsilon & \text{on } \partial_p Q_1 \setminus (B_\varepsilon \times \{-1\}), \\ v &= C(M) & \text{on } B_\varepsilon \times \{-1\}, \end{aligned}$$

we will have

$$-v_\varepsilon \leq \partial_t u_r \leq 0 \quad \text{in } B_1 \times (-1, t_0/r^2).$$

It is not hard to see that  $v_\varepsilon \leq c(\varepsilon) \rightarrow 0$  as  $\varepsilon \rightarrow 0$  uniformly in  $Q_{1/2}^-$ ; hence

$$-c(\varepsilon) \leq \partial_t u_r \leq 0 \quad \text{in } B_{1/2} \times (-1/4, t_0/r^2).$$

Scaling back to  $u$ , we obtain

$$-c(\varepsilon) \leq \partial_t u \leq 0 \quad \text{in } B_{r/2} \times (r/4, t_0).$$

Letting  $r = r_k \rightarrow \infty$  and then  $\varepsilon \rightarrow 0$ , we obtain (9.4). Thus,

$$(9.5) \quad u(x, t) = v_\infty(x) \quad \text{for any } x \in \mathbb{R}^n \text{ and } t \leq t_0.$$

It remains to prove that  $u \geq 0$  in  $\mathbb{R}^n \times (t_0, 0)$ , since we know that  $v_\infty \geq 0$ . In fact, we claim

$$(9.6) \quad v_\infty(x) \geq c|x|^2$$

for some fixed  $c > 0$  small and  $|x| > R$  large. If this fails, we could easily construct a shrink-down of  $v_\infty$  (which is always a polynomial) that vanishes at a point on  $\partial B_1$ . Hence this polynomial is degenerate. But from (9.5) we see that any shrink-down of  $v_\infty$  corresponds to the one of  $u$ , for which we assume that  $P_\infty$  is nondegenerate, a contradiction. Hence the estimate (9.6) holds. Consequently,  $u(x, t) > 0$  for  $|x| > R_1$  and  $t \in [t_0, 0]$ . Hence  $\Lambda(t)$  is bounded for all  $t \leq 0$ . Also  $\Lambda(0)$  is nonempty. So we could take  $t_0 = 0$  in all the arguments above. Thus,  $u$  is stationary and

$$u(x, t) = v_\infty(x) \geq 0 \quad \text{for any } x \in \mathbb{R}^n \text{ and } t \in \mathbb{R}^-.$$

The theorem is proved. □

10. PROOF OF THEOREM I

For a global solution  $u$  let us define

$$\begin{aligned} T_1 &= \sup\{t : (x, t) \in \partial\Omega \text{ is a high energy point}\}, \\ T_* &= \sup\{t : (x, t) \in \partial\Omega \text{ is a low or high energy point}\}, \\ T_2 &= \sup\{t : (x, t) \in \partial\Omega\}. \end{aligned}$$

Then

$$-\infty \leq T_1 \leq T_* \leq T_2 \leq a.$$

We claim that Theorem I holds with the values of  $T_1$  and  $T_2$  defined above. Parts (i) and (iii) of the theorem are easily verified, so we need to show only that (ii) holds.

Thus, if  $T_1 = T_2$ , we are done.

Suppose now that  $T_1 < T_2$ . If it happens that  $T_* = T_2$ , then we will find a sequence  $(x_k, t_k)$  of low energy points with  $t_k \nearrow T_2$  and applying Theorem 9.1, we will obtain that  $u \geq 0$  in  $\mathbb{R}^n \times (-\infty, T_2]$  and (ii) will follow from Theorem 8.1.

Suppose therefore that  $T_* < T_2$ . We claim now that

$$u = 0 \quad \text{in } \mathbb{R}^n \times (T_*, T_2).$$

By the very construction, the only points of  $\partial\Omega$  in  $\mathbb{R}^n \times (T_*, T_2)$  are of zero energy. A consequence is that for  $t \in (T_*, T_2)$ ,  $\partial\Lambda(t)$  is empty, implying that either  $\Lambda(t)$  is empty itself or it is the whole space  $\mathbb{R}^n$ . Moreover, if  $\Lambda(t_0) = \mathbb{R}^n$  for some  $t_0 \in (T_*, T_2]$ , then  $\Lambda(t) = \mathbb{R}^n$  for any  $t \in (t_0 - \varepsilon^2, t_0]$ , where  $\varepsilon$  is sufficiently small. Indeed, consider a point  $(x_0, t_0)$ . Then it is either an interior point of  $\Lambda$  or a zero energy point. In both cases there is  $\varepsilon > 0$  such that  $u = 0$  on  $Q_\varepsilon^-(x_0, t_0)$ . Then  $\Lambda(t)$  is nonempty for  $t \in (t_0 - \varepsilon^2, t_0]$ , thus implying  $\Lambda(t) = \mathbb{R}^n$ . In fact, by using a continuation argument, we obtain that  $\Lambda(t_0) = \mathbb{R}^n$  implies that  $u = 0$  on  $\mathbb{R}^n \times (T_*, t_0]$ .

Next, by the definition of  $T_2$  there is a sequence of points  $(x_k, t_k) \in \partial\Omega$  with  $t_k \nearrow T_2$ . Then  $\Lambda(t_k) = \mathbb{R}^n$  by the argument above and we obtain that  $u = 0$  in  $\mathbb{R}^n \times (T_*, T_2)$ .

To complete the proof, we consider the following two cases:  $T_1 < T_*$  and  $T_1 = T_*$ . In the first case we finish the proof by applying Theorems 9.1 and 8.1. In the second case, there are two possibilities. If  $T_1 = T_* = -\infty$ , we obtain that  $u = 0$  in  $\mathbb{R}^n \times (-\infty, T_2)$ , and if  $T_1 = T_* > -\infty$ , from the representation  $u(x, t) = P(x) + mt$  we obtain that  $u(x, t) = T_1 - t$ , since  $u$  vanishes for  $t = T_1$ . Thus, in all cases (ii) holds and the proof is complete.  $\square$

11. LIPSCHITZ REGULARITY: GLOBAL SOLUTIONS

**Theorem 11.1.** *Let  $u \in \mathcal{P}_\infty^-(M)$  be such that  $(0, 0) \in \Gamma$  and suppose that  $\Lambda$  contains a cylinder  $B \times [-1, 0]$ , where  $B = B_\rho(-se_n)$  for some  $0 \leq s \leq 1$ . Set  $K(\delta, s, h) = \{|x'| < \delta, -s \leq x_n \leq h\}$  for any  $\delta, h > 0$ . Then*

- (i)  $u \geq 0$  in  $\mathbb{R}^n \times \mathbb{R}^-$ .
- (ii) For any spatial unit vector  $e$  with  $|e - e_n| < \rho/8$  we have

$$\partial_e u \geq 0 \quad \text{in } K(\rho/8, s, 1) \times [-1/2, 0].$$

- (iii) Moreover, there exists  $C_0 = C_0(n, M, \rho) > 0$  such that

$$C_0 \partial_e u - u \geq 0 \quad \text{in } K(\rho/16, s, 1/2) \times [-1/2, 0].$$

(iv) *The free boundary  $\partial\Omega \cap (K(\rho/32, s, 1/4) \times [-1/4, 0])$  is a space-time Lipschitz graph*

$$x_n = f(x', t),$$

where  $f$  is concave in  $x'$  and

$$|\nabla_{x'} f| \leq \frac{C}{\rho}, \quad |\partial_t f| \leq C(n, M, \rho).$$

*Proof.* The origin is either a low or high energy point. The existence of the cylinder  $B \times [-1, 0]$  in  $\Lambda$  excludes the possibility of high energy. Moreover, for the same reason,  $u$  has no high energy points in  $\mathbb{R}^n \times [-1, 0]$ . Hence (i) follows from Theorem 9.1.

Next, applying Theorem 8.1, we obtain from (i) that  $\partial_{ee}u \geq 0$  and  $\partial_t u \leq 0$  in  $\mathbb{R}^n \times \mathbb{R}^-$ . Now suppose that  $|e - e_n| < \rho/8$ . Since  $\partial_e u = 0$  on  $B$  and every halfline in the direction  $-e$  originating at a point in  $K(\rho/8, s, 1)$  intersects  $B$ , the convexity of  $u$  implies (ii).

Furthermore, since  $0 \in \partial\Lambda(0)$ , we obtain that the cone  $\mathcal{C} = \{x_n > (8/\rho)|x'|\}$  is contained in  $\Omega(0)$ , and thus in every  $\Omega(t)$  for  $-1 \leq t \leq 0$ . Then, together with (ii) we find the representation  $x_n = f(x', t)$  in  $K(\rho/8, s, 1)$ , with the spatial Lipschitz estimate  $|\nabla_{x'} f| \leq C/\rho$ . This proves the first part of (iv). To prove the estimate on  $\partial_t f$  in (iv), as well as the estimate (iii), we need an additional lemma.

**Lemma 11.2.** *Let  $u$  be as in Theorem 11.1. Then there exist  $r_0 = r_0(n, \rho) > 0$  and  $\varepsilon_0 = \varepsilon_0(n, \rho) > 0$  such that*

$$W(r, x, t; u) \leq 2A - \varepsilon_0$$

for any  $(x, t) \in \partial\Omega \cap Q_{1/2}^-$  and  $0 < r \leq r_0$ .

*Proof.* Assume the contrary. Then there exists a sequence of functions  $u_k$  satisfying the assumptions of the lemma and  $(x_k, t_k) \in \partial\Omega(u_k) \cap Q_{1/2}^-$  such that

$$W(1/k, x_k, t_k; u_k) \geq 2A - 1/k.$$

From the uniform estimates on  $u_k$  we can extract a subsequence such that the functions

$$v_k(x, t) = u_k(x + x_k, t + t_k)$$

converge to a global solution  $v_0$  with  $(0, 0) \in \Lambda(v_0)$ . Then for every  $r > 0$  we have

$$W(r; v_0) = \lim_{k \rightarrow \infty} W(r, x_k, t_k; u_k) \geq \limsup_{k \rightarrow \infty} W(1/k, x_k, t_k; u_k) \geq 2A.$$

Also since  $W(r; v_0) \leq 2A$ , we obtain that  $W(r; v_0) = 2A$  for every  $r > 0$ . Then  $(0, 0)$  is a high energy point and therefore

$$v_0(x, t) = ct + P(x)$$

where  $c$  is a constant and  $P$  is a homogeneous quadratic polynomial. On the other hand, since  $u_k$  vanishes on the cylinders  $B_\rho(-s_k e_n) \times (-1, 0)$ ,  $v_0$  vanishes on a cylinder  $B \times (-1/2, 0)$ , where  $B$  is a certain ball of radius  $\rho$ . But this is impossible unless  $v_0$  is identically 0, a contradiction.

The lemma is proved. □

We continue the proof of Theorem 11.1. To show (iii) we assume the contrary. Then there exist a sequence of functions  $u_k$  satisfying the assumptions of the theorem and points  $(x_k, t_k) \in \Omega(u_k) \cap (K(\rho/16, s, 1/2) \times (-1/2, 0))$  such that

$$(11.1) \quad k \partial_e u_k(x_k, t_k) - u_k(x_k, t_k) \leq 0, \quad e = e(k).$$

Now let

$$\tilde{x}_k = (x'_k, f_k(x'_k)) \in \partial\Omega_k(t_k), \quad h_k = (x_k)_n - f_k(x'_k)$$

and consider

$$v_k(x, t) = \frac{1}{h_k^2} u(h_k x + \tilde{x}_k, h_k^2 t + t_k).$$

Then from (11.1) we have

$$(11.2) \quad \partial_e v_k(e_n, 0) \leq \frac{h_k}{k} v_k(e_n, 0).$$

Functions  $v_k$  are locally uniformly bounded in  $\mathbb{R}^n \times \mathbb{R}^-$ ; hence over a subsequence  $v_k$  converge to a global solution  $v$ . If we also assume that  $e(k) \rightarrow e$ , we will have

$$\partial_e v(e_n, 0) = 0.$$

Next, since  $h_k \leq 2$ , each of the sets  $\partial\Omega_{v_k}(0) \cap \{|x'| < \rho/32\}$  is a graph of a concave Lipschitz function, containing 0 and with the Lipschitz constant  $L \leq C/\rho$ . Also, since  $\Omega_{v_k}(t)$  expand as  $t$  decreases, we obtain that  $D_k \times (-1/4, 0) \subset \Omega_{v_k}$ , where  $D_k = \Omega_{v_k}(0) \cap K(\rho/32, s_k/h_k, 1/h_k)$ . In particular  $H(\partial_{e_k} v_k) = 0$  in  $D_k \times (-1/4, 0)$ . Moreover, (ii) also implies that  $\partial_{e_k} v_k \geq 0$  there. Passing to the limit, we can assume that  $D_k$  converge to a set  $D$ , having similar properties. Then

$$H(\partial_e v) = 0, \quad \partial_e v \geq 0 \quad \text{in } D \times (-1/4, 0).$$

Since  $e_n \in D$ , the maximum principle implies that  $\partial_e v = 0$  in  $D \times (-1/4, 0)$ . Then we also obtain that  $v(x, 0) = 0$  in  $D \times \{|x'| < \rho/64\}$  and as a consequence that  $v(x, 0)$  vanishes in a neighborhood of the origin.

The stability property implies that  $(0, 0) \in \Gamma(v)$ . Moreover, it cannot be low energy, since then  $v(x, 0)$  would not vanish in a neighborhood of the origin. So, the only possibility is that  $(0, 0)$  is a high energy point of  $v$ . The latter is possible only if

$$v(x, t) = -t \quad \text{in } \mathbb{R}^n \times \mathbb{R}^-.$$

To exclude this possibility, we apply Lemma 11.2. Indeed, we have

$$W(r, v) = \lim_{k \rightarrow \infty} W(r; v_k) = \lim_{k \rightarrow \infty} W(h_k r, \tilde{x}_k, t_k; u_k) \leq 2A - \varepsilon_0,$$

provided  $r < r_0/2$  (recall  $h_k \leq 2$ .) Hence  $(0, 0)$  cannot be a high energy point. This proves (iii).

Finally, to prove the estimate on  $\partial_t f$  in (iv), we apply the following generalization of Lemma 7.6.

**Lemma 11.3.** *Let  $u$  be a bounded solution of (1.2) in*

$$\mathcal{N}_\delta^-(E) = \bigcup \{Q_\delta^-(x, t) : (x, t) \in E\},$$

for a set  $E$  in  $\mathbb{R}^n \times \mathbb{R}^-$  and let  $h$  be caloric in  $\mathcal{N}_\delta^-(E) \cap \Omega$ . Suppose moreover that

- (i)  $h \geq 0$  on  $\mathcal{N}_\delta^-(E) \cap \partial\Omega$  and
- (ii)  $h - u \geq -\varepsilon_0$  in  $\mathcal{N}_\delta^-(E)$ , for some  $\varepsilon_0 > 0$ .

Then  $h - u \geq 0$  in  $\mathcal{N}_{\delta/2}^-(E)$ , provided  $\varepsilon_0 = \varepsilon_0(\delta, n)$  is small enough.

*Proof.* Consider  $h$  and  $u$  in every  $Q_{\delta}^-(x, t)$  with  $(x, t) \in E$ , parabolically scale to functions in  $Q_1^-$  and apply Lemma 7.6.  $\square$

Now, for small  $|\eta| < \eta_0(\rho, n, M)$  we obtain from (iii) in Theorem 11.1 that

$$(C_0 \partial_e u + \eta \partial_t u) - u \geq -\varepsilon_0$$

in  $K(\rho/16, s, 1/2) \times [-1/2, 0]$ . From Lemma 11.3 we have

$$(C_0 \partial_e u + \eta \partial_t u) - u \geq 0$$

in  $K(\rho/32, s, 1/4) \times [-1/4, 0]$ . Note that Lemma 11.3 is applicable with  $h = C_0 \partial_e u + \eta \partial_t u$ , since both  $\partial_e u$  and  $\partial_t u$  vanish on  $\partial\Omega$ . The latter follows from Lemma 7.7.

Then, as in the proof of Theorem 7.5, we obtain the existence of space-time cones with uniform openings at any point on  $\partial\Omega$  in  $K(\rho/32, s, 1/4) \times [-1/4, 0]$  and this proves the estimate on  $\partial_t f$  in (iv).

The proof of the theorem is complete.  $\square$

### 12. BALANCED ENERGY: LOCAL SOLUTIONS

In this short section we discuss how one can generalize the balanced energy that we defined for global solutions (see Section 7) for local solutions.

Let  $u$  be a solution of (1.2) in  $Q_1^-$  and let  $\psi(x) \geq 0$  be a  $C^\infty$  cut-off function with  $\text{supp } \psi \subset B_1$  and  $\psi|_{B_{3/4}} = 1$ . Then for  $w = u\psi$  and any  $(x_0, t_0) \in \partial\Omega \cap Q_{1/2}^-$  the functional

$$W(r, x_0, t_0; w) + F_n(r)$$

is nondecreasing by the local form of Weiss’s monotonicity theorem (Theorem 3.6). Hence there exists a limit

$$\omega(x_0, t_0) = \lim_{r \rightarrow 0} W(r, x_0, t_0; w),$$

since  $F_n(r) \rightarrow 0$  as  $r \rightarrow 0$ . Moreover, if  $u_0$  is a blow-up limit of parabolic scalings  $u_r(x, t) = (1/r^2)u(rx + x_0, r^2t + t_0)$ , then  $u_0$  is also a limit of corresponding scalings  $w_r$  of  $w$ , since  $\psi = 1$  on  $B_{1/4}(x_0)$  and we obtain that

$$W(s; u_0) = \lim_{r \rightarrow 0} W(s; w_r) = \lim_{r \rightarrow 0} W(sr, x_0, t_0; w) = \omega(x_0, t_0).$$

In particular,  $u_0$  is a homogeneous global solution and  $\omega(x_0, t_0)$  does not depend on the choice of the cut-off function  $\psi$ . The quantity  $\omega(x_0, t_0)$  will be called the *balanced energy* of  $u$  at  $(x_0, t_0)$ . Note that when  $u$  is a global solution, this definition coincides with the one from Section 7.

As in the global case, since  $\omega(x_0, t_0) = W(u_0)$  and  $u_0$  is homogeneous, we have only three possible values for the balanced energy: 0,  $A$ , and  $2A$ . We classify the point  $(x_0, t_0) \in \partial\Omega$  as of zero, low or high energy, respectively.

### 13. LIPSCHITZ REGULARITY: LOCAL SOLUTIONS

**Theorem 13.1.** *For every  $\sigma > 0$  there exists  $R_0 = R_0(\sigma, n, M)$  such that if  $u \in \mathcal{P}_R^-(MR^2)$  for  $R \geq R_0$ ,  $(0, 0) \in \Gamma$  and  $\delta_1^-(u) \geq \sigma$ , then  $\partial\Omega \cap Q_{1/2}^-$  is space-time Lipschitz regular with Lipschitz constant  $L \leq L(\sigma, n, M)$ .*

We use the following approximation lemma and then apply the results from Section 11.

**Lemma 13.2.** *Fix  $\sigma > 0$  and  $\varepsilon > 0$ . Then there exists  $R_0 = R_0(\varepsilon, \sigma, n, M)$  such that if  $u \in \mathcal{P}_R^-(MR^2)$  for  $R \geq R_0$ ,  $(0, 0) \in \Gamma$  and  $\delta_1^-(u) \geq \sigma$ , then we can find a global solution  $v \in \mathcal{P}_\infty^-(C_nM)$ ,  $(0, 0) \in \Gamma(v)$ , with the properties*

- (i)  $\|u - v\|_{C_x^1 \cap C_t^0(Q_1^-)} \leq \varepsilon$ ;
- (ii) *there exists a ball  $B = B_\rho(x) \subset B_1$  of radius  $\rho = \sigma/(4n)$  such that  $v$  vanishes on  $B \times [-1, 0]$ ;*
- (iii)  *$u$  vanishes on  $B_{\rho/2}(x) \times [-1/2, 0]$ .*

*Proof.* The proof is by compactness. Assume the contrary. Then for every  $k > 0$  we can find a solution  $u_k \in \mathcal{P}_k^-(Mk^2)$  with  $(0, 0) \in \Gamma(u_k)$  and  $\delta_1^-(u_k) \geq \sigma$  such that for any global solution  $v \in \mathcal{P}_\infty^-(C_nM)$  such that  $(0, 0) \in \Gamma(v)$  and conditions (ii) and (iii) are satisfied, we have

$$(13.1) \quad \|u_k - v\|_{C_x^1 \cap C_t^0(Q_1^-)} \geq \varepsilon.$$

Solutions  $u_k$  are locally uniformly bounded, so we can extract a subsequence converging to a global solution  $u_0$  in  $C_x^{1,\alpha} \cap C_t^{0,\alpha}$ -norm on compact subsets of  $\mathbb{R}^n \times \mathbb{R}^-$ . We claim now that  $\Lambda_{u_0}(-1) \cap B_1$  contains a ball of radius  $\rho = \sigma/(4n)$ . Indeed, first note that  $\delta_1^-(u_0) \geq \sigma/2$ ; otherwise we would have  $\delta_1^-(u_k) < \sigma$  for large  $k$ . Next, from the stability note that  $(0, 0)$  is not a zero energy point of  $u_0$ , since it is not for any of solutions  $u_k$ . Also, it is not a high energy point of  $u_0$  since  $\delta_1^-(u_0) \geq \sigma/2$ . Hence the only possibility is that  $(0, 0)$  is a low energy point of  $u_0$ . Then Theorem 9.1 implies that  $u_0 \geq 0$  in  $\mathbb{R}^n \times \mathbb{R}^-$  and hence the set  $\Lambda_{u_0}(-1) \cap B_1$  is convex by Theorem 8.1. Invoking F. John’s lemma, we obtain the existence of a ball  $B = B_\rho(x)$  of radius  $\rho = \sigma/(4n)$  in  $\Lambda_{u_0}(-1) \cap B_1$ . Moreover,  $u_0 \geq 0$  implies that  $\partial_t u_0 \leq 0$  and that the sets  $\Lambda_{u_0}(t)$  shrink as  $t$  decreases. Hence  $B \times [-1, 0]$  is contained in  $\Lambda(u_0)$ . Since  $u_k \rightarrow u_0$ , from the stability (see Subsection 5.2) we obtain that  $u_k$  vanishes on  $B_{\rho/2}(x) \times [-1/2, 0]$  for large  $k$ .

So, conditions (ii) and (iii) are satisfied for the global solution  $v = u_0$  and  $u = u_k$  for large  $k$ . But we also have  $\|u_k - u_0\|_{C_x^1 \cap C_t^0(Q_1^-)} \rightarrow 0$ , which contradicts (13.1). Hence the lemma follows.  $\square$

*Proof of Theorem 13.1.* Let  $\varepsilon = \varepsilon(\sigma, n, M) > 0$  be small (to be specified later) and let  $R_0 = R_0(\varepsilon, \sigma)$  be as in Lemma 13.2 and suppose that  $R \geq R_0$ . Also for  $u \in \mathcal{P}_R^-(MR^2)$  with  $\delta_1^-(u) \geq \sigma$  let the global solution  $v$  and the ball  $B \subset B_1$  be as in the conclusion of Lemma 13.2.

Rotating the spatial coordinate axes, we may assume that  $B = B_\rho(-se_n)$  for  $0 \leq s \leq 1$ ,  $\rho = \sigma/(4n)$ . Then by estimate (iii) in Theorem 11.1 applied to the global solution  $v$  we have

$$C_0 \partial_e v - v \geq 0 \quad \text{in } K(\rho/8, s, 1/2) \times [-1/2, 0]$$

for any spatial unit vector  $e$  with  $|e - e_n| \leq \rho/8$ . Since  $|u - v| \leq \varepsilon$ ,  $|\nabla u - \nabla v| \leq \varepsilon$  and  $|\partial_t u| \leq C_n M$  in  $Q_1^-$ , we obtain automatically that

$$(C_0 \partial_e u - \eta \partial_t u) - u \geq -C_0 \varepsilon - \varepsilon - C_n M \eta_0 \geq -\varepsilon_0$$

if  $\varepsilon = \varepsilon(\sigma, n, M)$  and  $|\eta| \leq \eta_0(n, M)$  are small, where  $-\varepsilon_0$  is as in Lemma 11.3. Next, we claim that

$$(13.2) \quad (C_0 \partial_e u - \eta \partial_t u) - u \geq 0 \quad \text{in } K(\rho/16, s, 1/4) \times [-1/4, 0].$$

This will follow from Lemma 11.3 with  $h = C_0 \partial_e u - \eta \partial_t u$  if we know that  $h \geq 0$  on  $\partial\Omega$ . We show next that this is indeed so.

**Lemma 13.3.** *Let  $u$  be as in Lemma 13.2 with  $R \geq R_0$ . Also let  $\psi(x) \geq 0$  be a  $C^\infty$  cut-off function with  $\text{supp } \psi \subset B_1$  and  $\psi|_{B_{3/4}} = 1$ . Then for  $w = u\psi$  and any  $(x_0, t_0) \in \partial\Omega \cap Q_{1/2}^-$  we have*

- (i)  $W(r, x, t; w) \leq 2A - \varepsilon_0$  for  $\varepsilon_0 = \varepsilon_0(\sigma, n, M) > 0$  and  $r \leq r_0(\sigma, n, M)$ ;
- (ii)  $\partial_t u$  vanishes continuously at  $(x_0, t_0)$ :  $\lim_{(x,t) \rightarrow (x_0,t_0)} \partial_t u(x, t) = 0$ .

*Proof.* (i) is a generalization of Lemma 11.2. The proof is basically the same, only instead of Weiss’s monotonicity theorem (Theorem 3.4) one have to use its local form (Theorem 3.6.)

(ii) is a generalization of Lemma 7.7. We note that  $u$  has no high energy points by (i) above. Then the proof is the same as that of Lemma 7.7, with application of the local form of Weiss’s theorem instead of the global form.  $\square$

The lemma above implies that we indeed have (13.2). In particular, we obtain that  $\partial\Omega \cap (K(\rho/16, s, 1/4) \times [-1/4, 0])$  is Lipschitz in space and in time with a Lipschitz constant  $L(\sigma, n, M)$ .

To finish the proof of the theorem, we observe that we will come to the same conclusion as above (perhaps with different constants) if instead of  $\delta_1^-(u) \geq \sigma$  we assume, say,

$$\delta_1^*(u) := \sup_{-1 \leq t \leq -1/2} \text{md}(\Lambda(t) \cap B_2) \geq \sigma.$$

This gives a little bit more flexibility. Now let  $(x_0, t_0) \in \partial\Omega \cap Q_{1/2}^-$  and consider the function  $u^*(x, t) = u(x + x_0, t + t_0)$ . We will have  $\delta_1^*(u^*) \geq \sigma$ ; thus after an appropriate choice of coordinate axes we will find that  $\partial\Omega$  is  $L(\sigma, n, M)$ -Lipschitz in a parabolic neighborhood of  $(x_0, t_0)$ . This finishes the proof of the theorem.  $\square$

14.  $C^{1,\alpha}$  REGULARITY

**Theorem 14.1.** *Under the conditions of Theorem 13.1,  $\partial\Omega \cap Q_{1/4}^-$  is space-time  $C^{1,\alpha}$  regular with the norm  $C \leq C(\sigma, n, M)$ .*

*Proof.* We are going to apply the result of [ACS96], Corollary 1, on mutual boundary regularity of positive caloric functions in Lipschitz domains.

We assume that  $R \geq R_0$  and that the ball  $B = B_\rho(-se_n)$  is as in Lemma 13.2, so that  $u$  vanishes on  $B_{\rho/2}(-se_n) \times [-1/2, 0]$ . As follows from the proof of Theorem 13.1, we have

$$(C_0 \partial_e u + \eta \partial_t u) - u \geq 0 \quad \text{in } K(\rho/16, s, 1/4) \times [-1/4, 0]$$

for any spatial unit vector  $e$  with  $|e - e_n| < \rho/8$  and  $|\eta|$  sufficiently small. In particular, we have that

$$\partial_e u + \varepsilon \partial_t u \geq 0,$$

where  $\varepsilon = \eta/C_0$ . Consider now two functions of the type above

$$\begin{aligned} u_1 &= \partial_n u, \\ u_2 &= \partial_e u + \varepsilon \partial_t u \end{aligned}$$

with  $e$  sufficiently close to  $e_n$  and  $\varepsilon$  small. Then [ACS96], Corollary 1, implies that the ratio

$$\frac{u_2}{u_1}$$

is  $C^\alpha$  regular (both in  $x$  and in  $t$ ) in  $\Omega \cap (K(\rho/32, s, 1/8) \times [-(\rho/32)^2, 0])$  up to  $\partial\Omega$ , with  $0 < \alpha < 1$  and  $C^\alpha$ -norm depending on  $\rho, n, M$ , the Lipschitz norm of  $\partial\Omega$ , as well as on the bound from below on

$$m_i = u_i(A^-), \quad A^- = ((3/16)e_n, -(\rho/16)^2).$$

We claim that

$$m_i \geq c_0(\rho, n, M) > 0.$$

It is enough to prove the bound only for  $m_1$ , since  $m_2$  can be made as close to  $m_1$  as we wish. Thus, we have to show that

$$\partial_n u \geq c_0 > 0$$

at  $A^-$ . Indeed, if it were not so, by compactness we would easily construct a function  $u$  as above with  $\partial_n u = 0$  at  $A^-$ . Then by the minimum principle  $\partial_n u$  and consequently  $u$  would vanish in  $K(\rho/32, s, 1/8) \times [-1/4, -(\rho/16)^2]$ , a contradiction.

Hence,

$$\frac{\partial_e u + \varepsilon \partial_t u}{\partial_n u}$$

is  $C^\alpha$  up to  $\partial\Omega$  in  $\Omega \cap Q_{\rho/32}^-$ . Then varying  $e$  and  $\varepsilon$ , we obtain that the ratios

$$\frac{\partial_i u}{\partial_n u}, \quad i = 1, \dots, n-1, \quad \frac{\partial_t u}{\partial_n u}$$

are  $C^\alpha$ . This implies that  $\partial\Omega \cap (K(\rho/32, s, 1/8) \times [-(\rho/32)^2, 0])$  is a graph  $x_n = f(x', t)$  with

$$\|f\|_{C^{1,\alpha}} \leq C(\rho, n, M)$$

since

$$\partial_i f = \frac{\partial_i u}{\partial_n u}, \quad i = 1, \dots, n-1, \quad \partial_t f = \frac{\partial_t u}{\partial_n u}.$$

Arguing as in the end of the proof of Theorem 13.1, we obtain that in fact  $\partial\Omega \cap Q_{1/4}^-$  is  $C^{1,\alpha}$  regular.

The proof is complete. □

### 15. HIGHER REGULARITY

**Theorem 15.1.** *Under the conditions of Theorem 13.1,  $\partial\Omega \cap Q_{1/8}^-$  is space-time  $C^\infty$  regular.*

*Proof.* First, we prove the higher regularity for  $u$ .

As follows from the proof of Theorems 13.1 and 14.1, we have that

$$\partial_n u + \varepsilon \partial_t u \geq 0 \quad \text{in } K(\rho/16, s, 1/4) \times [-1/4, 0]$$

for small  $|\varepsilon| \leq \varepsilon(\rho, n, M)$ . Then for large  $C = C(\rho, n, M) > 0$

$$-C \partial_n u(x, t) \leq \partial_t u(x, t) \leq C \partial_n u(x, t).$$

Thus,  $\partial_t u$  will grow linearly in  $\Omega \cap Q_{1/4}^-$  away from  $\partial\Omega$ , implying that  $|\nabla \partial_t u|$  will be uniformly bounded in  $\Omega \cap Q_{1/8}^-$ .

Next, we claim that  $u$  is  $C_x^{2,\alpha}$  in  $Q_{1/8}^-$ . In fact, something stronger is true: if  $w$  is any partial derivative (in space or in time) of  $u$ , then  $w$  is  $C_x^{1,\alpha} \cap C_t^{0,\alpha}$  regular in  $\Omega \cap Q_{1/8}^-$  up to  $\partial\Omega$ . Indeed  $w$  satisfies the heat equation in  $\Omega \cap Q_{1/4}^-$ , it is uniformly bounded there, and it vanishes continuously on the  $C^{1,\alpha}$ -graph  $\partial\Omega$ . Then the classical boundary regularity implies that  $w$  is  $C_x^{1,\alpha} \cap C_t^{0,\alpha}$  regular.

Now we have enough regularity to apply the Kinderlehrer-Nirenberg technique [KN77].

Without loss of generality assume that  $e_1$  is the (spatial) exterior normal to  $\partial\Omega(0)$  at 0. Since  $|\nabla u| = 0$  on  $\partial\Omega$ , all spatial second order derivatives vanish at  $(0, 0)$ , except for  $\partial_{11}u(0, 0)$ . Also since  $\partial_t u$  vanishes at  $(0, 0)$ , we obtain that

$$\partial_{11}u(0, 0) = 1.$$

Hence, in  $\Omega \cap Q_{2r}^-$  for  $r = r(\rho, n, M) > 0$  small we will have that

$$\partial_{11}u \geq \frac{1}{2}.$$

Then consider there the partial hodograph transform

$$(x, t) \mapsto (y, t) = (-\partial_1 u, x_2, \dots, x_n, t),$$

which is  $C^1$  and has a nonsingular Jacobian, and the associated Legendre transform

$$v = x_1 y_1 + u = -x_1 \partial_1 u + u.$$

Then  $\partial\Omega$  transforms to a portion of  $\{y_1 = 0\}$  and the equation for  $v$  takes the form

$$Lv := -\frac{1}{\partial_{11}v} - \frac{1}{\partial_{11}v} \sum_{i=2}^n (\partial_{i1}v)^2 + \sum_{i=2}^n \partial_{ii}v - \partial_t v = 1.$$

As can be shown,  $Lv$  is a uniformly parabolic equation on the image of  $\Omega \cap Q_{2r}^-$  under the hodograph transform. Moreover  $v$  vanishes on the image of  $\partial\Omega \cap Q_{2r}^-$ , which is a subset of  $\{y_1 = 0\}$ . Hence  $v$  is  $C^\infty$  regular on the image of  $\partial\Omega \cap Q_r^-$  and considering the inverse transformation

$$(y, t) \mapsto (x, t) = (\partial_1 v, y_2, \dots, y_n, t),$$

we find that  $\partial\Omega \cap Q_r^-$ , as well as  $u$ , is  $C^\infty$  regular. For details we refer to [KN77].

To finish the proof, we note that by similar reasoning one can show that  $\partial\Omega \cap Q_r^-(x, t)$  is  $C^\infty$  regular near every point  $(x, t) \in \partial\Omega \cap Q_{1/8}^-$ . Hence the theorem follows.  $\square$

*Remark 15.2.* In fact, one can show that  $\partial\Omega \cap Q_{1/8}^-$  is not only  $C^\infty$  but also analytic in the space variables and in the second Gevrey class with respect to the time variable; see [KN78].

## 16. PROOF OF THEOREM II

For the solution  $u \in \mathcal{P}_1^-(M)$  satisfying the assumptions of the theorem consider the parabolic scaling

$$u_{r_0}(x, t) = \frac{1}{r_0^2} u(r_0 x, r_0^2 t).$$

Then we arrive at the conditions of Theorem 13.1. Thus Theorems 14.1 and 15.1 are also applied. Scaling back, we conclude the proof of the theorem.  $\square$

## REFERENCES

- [ACF84] Hans Wilhelm Alt, Luis A. Caffarelli, and Avner Friedman, *Variational problems with two phases and their free boundaries*, Trans. Amer. Math. Soc. **282** (1984), no. 2, 431–461. MR0732100 (85h:49014)
- [ACS96] I. Athanasopoulos, L. Caffarelli, and S. Salsa, *Caloric functions in Lipschitz domains and the regularity of solutions to phase transition problems*, Ann. of Math. (2) **143** (1996), no. 3, 413–434. MR1394964 (97e:35074)

- [Caf93] Luis A. Caffarelli, *A monotonicity formula for heat functions in disjoint domains*, Boundary value problems for partial differential equations and applications, Masson, Paris, 1993, pp. 53–60. MR1260438 (95e:35096)
- [Caf98] L. A. Caffarelli, *The obstacle problem revisited*, J. Fourier Anal. Appl. **4** (1998), no. 4-5, 383–402. MR1658612 (2000b:49004)
- [CK98] Luis A. Caffarelli and Carlos E. Kenig, *Gradient estimates for variable coefficient parabolic equations and singular perturbation problems*, Amer. J. Math. **120** (1998), no. 2, 391–439. MR1613650 (99b:35081)
- [CKS00] Luis A. Caffarelli, Lavi Karp, and Henrik Shahgholian, *Regularity of a free boundary with application to the Pompeiu problem*, Ann. of Math. (2) **151** (2000), no. 1, 269–292. MR1745013 (2001a:35188)
- [Fri64] Avner Friedman, *Partial differential equations of parabolic type*, Prentice-Hall Inc., Englewood Cliffs, N.J., 1964. MR0181836 (31:6062)
- [Fri88] ———, *Variational principles and free-boundary problems*, second ed., Robert E. Krieger Publishing Co. Inc., Malabar, FL, 1988. MR1009785 (90k:35271)
- [KN77] D. Kinderlehrer and L. Nirenberg, *Regularity in free boundary problems*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **4** (1977), no. 2, 373–391. MR0440187 (55:13066)
- [KN78] David Kinderlehrer and Louis Nirenberg, *The smoothness of the free boundary in the one phase Stefan problem*, Comm. Pure Appl. Math. **31** (1978), no. 3, 257–282. MR0480348 (82b:35152)
- [Lie96] Gary M. Lieberman, *Second order parabolic differential equations*, World Scientific Publishing Co. Inc., River Edge, NJ, 1996. MR1465184 (98k:35003)
- [Wei99] G. S. Weiss, *Self-similar blow-up and Hausdorff dimension estimates for a class of parabolic free boundary problems*, SIAM J. Math. Anal. **30** (1999), no. 3, 623–644 (electronic). MR1677947 (2000d:35267)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT AUSTIN, AUSTIN, TEXAS 78712  
*E-mail address:* [caffarel@math.utexas.edu](mailto:caffarel@math.utexas.edu)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT AUSTIN, AUSTIN, TEXAS 78712  
*E-mail address:* [arshak@math.utexas.edu](mailto:arshak@math.utexas.edu)  
*Current address:* Department of Mathematics, Purdue University, West Lafayette, Indiana 47907  
*E-mail address:* [arshak@math.purdue.edu](mailto:arshak@math.purdue.edu)

DEPARTMENT OF MATHEMATICS, ROYAL INSTITUTE OF TECHNOLOGY, 100 44, STOCKHOLM, SWEDEN  
*E-mail address:* [henriksh@math.kth.se](mailto:henriksh@math.kth.se)