

VALUATIONS AND MULTIPLIER IDEALS

CHARLES FAVRE AND MATTIAS JONSSON

CONTENTS

Introduction
1. Background
2. Multiplier ideals of tree potentials
3. Multiplier ideals of ideals
4. Multiplier ideals of psh functions
5. Singularity exponents and Kiselman numbers
6. The ascending chain condition
7. Ideals as multiplier ideals
8. Equisingularity
Appendix A. Subadditivity and Skoda's Theorem
References

INTRODUCTION

This article is the third of a series of work on a new approach to the study of singularities of various objects in a local, two-dimensional setting. Our focus in the present paper is on multiplier ideals and singularity exponents.

In the discussion below, we fix an equicharacteristic zero, two-dimensional regular local ring (R, \mathfrak{m}) with algebraically closed residue field. An important example is the ring $R = \mathcal{O}_0$ of holomorphic germs at the origin in \mathbf{C}^2 .

In [FJ1], we introduced the space \mathcal{V} consisting of all $\mathbf{R}_+ \cup \{+\infty\}$ -valued valuations on R centered at \mathfrak{m} , and normalized by $\nu(\mathfrak{m}) = 1$. This space is naturally a tree: it is a union of (uncountably many) real segments patched together in such a way that \mathcal{V} remains homotopic to a point. It is also an \mathbf{R} -tree in the classical sense for a natural metric. We hence call \mathcal{V} the *valuative tree*. It encodes in a natural way all possible blowups of R centered at \mathfrak{m} and therefore gives a way of measuring quite precisely singularities of different kinds of objects. The points in \mathcal{V} that are not ends form the subtree \mathcal{V}_{qm} of *quasimonomial* valuations. These valuations, which can alternatively be characterized as Abhyankar valuations of rank 1 or as

Received by the editors January 16, 2004.

2000 *Mathematics Subject Classification*. Primary 14B05; Secondary 32U25, 13H05.

Key words and phrases. Valuations, multiplier ideals, singularity exponents, Arnold multiplicity, Lelong numbers, Kiselman numbers, trees, Laplace operator.

The second author was partially supported by NSF Grant No. DMS-0200614.

©2005 American Mathematical Society
Reverts to public domain 28 years from publication

push-forwards of monomial valuations under birational morphisms, play a central role in our approach.

We used the valuative tree to study singularities of ideals $I \subset R$ in [FJ1], and of plurisubharmonic (psh) functions u in [FJ2] (in which case $R = \mathcal{O}_0$). We showed that I and u both give rise to functions g_I and g_u on \mathcal{V}_{qm} , called the *tree transforms* of I and u . These are defined as follows. For an ideal I and $\nu \in \mathcal{V}_{\text{qm}}$, we set $g_I(\nu) := \nu(I) = \min\{\nu(\phi) : \phi \in I\}$. For a psh function, the definition of $g_u(\nu) = \nu(u)$ is more involved but can be interpreted either as a generalized Lelong number in the sense of Demailly or as the push-forward of a Kiselman number under a birational morphism.

It turns out that the tree transforms g_I and g_u enjoy strong concavity properties. We denote by \mathcal{P} the closure of the cone generated by the functions on \mathcal{V}_{qm} of the form $c g_I$ over all $c > 0$ and all ideals I (or equivalently by all g_u 's for psh functions u). An element in $g \in \mathcal{P}$ is called a *tree potential*. Tree potentials can also be identified with positive measures on \mathcal{V} . In the case of an ideal, the measure is atomic and its decomposition into atoms is equivalent to Zariski's factorization of integrally closed ideals. In the case of a psh function, the measure determines the most important features of the singularity at the origin—a somewhat vague assertion that the paper at hand serves to substantiate. A general tree potential may be loosely viewed as a formal analogue of a psh function.

Multiplier ideals have emerged in recent years as a fundamental tool in algebraic and analytic geometry. We refer to the book [La] for a detailed account on this subject, on its range of applications, and for precise references. See also [D] for a more analytic point of view. Here we content ourselves with recalling a few definitions adapted to our setting.

Fix a psh function u defined near the origin in \mathbf{C}^2 . The multiplier ideal $\mathcal{J}(u)$ is the set of holomorphic germs $\phi \in \mathcal{O}_0$ such that $|\phi|^2 \exp(-2u)$ is locally integrable. As u may take the value $-\infty$, $\mathcal{J}(u)$ is in general strictly included in \mathcal{O}_0 . To worse behavior of u at 0, i.e., for u decreasing faster to $-\infty$, correspond deeper ideals $\mathcal{J}(u)$. The multiplier ideal hence measures, in some sense, the degree of singularity of u .

To an ideal $I \subset R$ (or more generally a formal power I^c , $c > 0$) we can also associate a multiplier ideal $\mathcal{J}(I)$. One way to do this uses resolution of singularities, whereas Lipman [Li] gives a more intrinsic definition. It is also possible to associate an asymptotic multiplier ideal to a graded system of ideals, i.e., a sequence $(I_k)_{k=1}^{\infty}$ of ideals in R such that $I_k I_l \subset I_{k+l}$.

Our first objective is to show that all these multiplier ideals can be analyzed in a unified way using tree potentials. More precisely, to any tree potential g on \mathcal{V}_{qm} we associate a multiplier ideal $\mathcal{J}(g) \subset R$ as follows: an element $\psi \in R$ belongs to $\mathcal{J}(g)$ iff the function $\chi_{\psi,g}(\nu) = g(\nu)/(\nu(\psi) + A(\nu))$ on \mathcal{V}_{qm} is uniformly bounded by a constant < 1 . Here A is a function on \mathcal{V}_{qm} called *thinness*. It contains information on the relative canonical divisor of suitable birational models of $\text{Spec } R$, as in the classical algebraic definition of multiplier ideals (see Section 1.2.9 for details).

We then show that the multiplier ideal of an ideal, a psh function, or a graded system of ideals, coincides with that of the corresponding tree transform. Let us be more precise. If I is an ideal and $c > 0$, then we show that the multiplier ideal of I^c coincides with that of the tree potential $c g_I$. Our proof essentially consists of translating classical conditions for defining multiplier ideals into our language. In

the case of a graded system of ideals, the condition $I_k I_l \subset I_{k+l}$ implies a statement $g_{k+l} \leq g_k + g_l$ on the level of tree potentials (g_k is the tree potential of I_k). As we show, the sequence $k^{-1}g_k$ converges to a tree potential g whose multiplier ideal coincides with that of the graded system.

Similarly, we prove that if u is psh, then the multiplier ideals $\mathcal{J}(g_u)$ and $\mathcal{J}(u)$ coincide. The proof is now more involved. We first use Demailly's approximation technique to reduce to the case of a psh function with logarithmic singularities. The latter case is then reduced to the statement $\mathcal{J}(I^c) = \mathcal{J}(c g_I)$, proved earlier, for a suitable chosen ideal I and $c > 0$. Many of the arguments involved in the reductions draw from [DK], but our proof also depends heavily on the fact that the tree transform $u \mapsto g_u$ behaves well under Demailly approximation.

Apart from providing a unifying framework, tree potentials can be used as a powerful tool for studying many questions regarding multiplier ideals. As support to this claim, we give three applications of our approach.

The first two concern singularity exponents. To a general tree potential g is associated a number $c(g)$, called the *singularity exponent* or *log-canonical threshold* of g . It is defined by $c(g) = \sup\{c > 0 : \mathcal{J}(c g) = R\}$. If $g = g_u$ is the tree transform of a psh function u , then the singularity exponent is given by $c(u) = \sup\{c > 0 : \exp(-2cu) \in L_{\text{loc}}^1\}$ and is a number measuring the "strength" of the singularity of u at the origin. Various bounds were known for $c(u)$ in terms of the Lelong number [S] and Kiselman numbers [Ki2] of u . We sharpen these bounds, and show that $c(u)$ can in fact be computed explicitly in terms of all Kiselman numbers of u . We also prove that for $c = c(u)$, the function $\exp(-2cu)$ is *never* integrable. This provides an affirmative answer to the openness conjecture (in dimension 2) by Demailly and Kollár (see [DK, Remark 5.3]). In fact, we establish the more precise estimate $\text{Vol}\{u < \log r\} \gtrsim r^{2c(u)}$ as $r \rightarrow 0$, and we improve a recent result by Blé and Mimouni [BM, Mi] by proving that if u has Lelong number one, then $\exp(-2u)$ fails to be locally integrable at the origin iff $dd^c u$ is the sum of the current of integration on a smooth curve and a current with zero Lelong number.

We then consider the set of complex singularity exponents $\mathbf{c} = \{c(\log |\psi|)\}$ when ψ ranges over all holomorphic functions. Shokurov [S] used Mori's minimal program to show that \mathbf{c} satisfies the so-called *ascending chain condition*: any increasing sequence in \mathbf{c} is eventually stationary. Kuwata [Ku] subsequently improved this result by characterizing explicitly all real numbers lying in \mathbf{c} . In 2000, Phong-Sturm in [PS] gave a completely analytic proof of this result. We present an algebraic proof in Section 6, independent of both approaches above. It is conjectured (see [DK, Remark 3.5]) that the ascending chain condition applies to the set of complex singularity exponents in any dimension (see [MP] for the most recent result in this direction). We hope that our result might lead to further developments in higher dimensions.

As a third application, we prove that any integrally closed ideal in R is the multiplier ideal of some formal power of an ideal. This result was recently proved, independently, by Lipman and Watanabe [LW].

Finally we prove quite generally that a tree potential is completely characterized by the collection of multiplier ideals $\{\mathcal{J}(tg)\}_{t \geq 0}$. As a consequence, two psh functions u and v have identical multiplier ideals $\mathcal{J}(tu) = \mathcal{J}(tv)$ for all $t \geq 0$ iff they are *equisingular* by which we mean that they have the same transforms $g_u = g_v$. Equisingularity may be geometrically interpreted as follows: for any composition π

of blowups, the Lelong numbers of the pull-backs π^*u and π^*v are the same at any point on the exceptional divisor $\pi^{-1}(0)$; see [FJ2, Proposition 6.2].

Since the results on singularity exponents for psh functions are arguably the most striking ones in the paper, we wish to briefly explain our approach, not using the language of valuations. Fix a psh function u and assume, for simplicity, that $dd^c u$ does not charge any curve. To any irreducible (possibly singular) curve D at the origin and any real number $t \geq 1$ we associate a family of punctured, conical regions $\mathcal{A}(r)$ of diameter r . Here t determines the “thinness” of the region. See Figure 2. We have $\text{Vol } \mathcal{A}(r) \sim r^{2A}$ for some $A > 0$, and in $\mathcal{A}(r)$ we have $u \leq \nu(u) \log r + O(1)$, for some (maximal) real number $\nu(u) \geq 0$. If $\exp(-2cu)$ is locally integrable at the origin, it is integrable in $\mathcal{A}(r)$, which easily implies $c\nu(u) < A$. The crucial fact is the existence of an *optimal* region $\mathcal{A}(r)$ detecting integrability: for this region we have $c(u)\nu(u) = A$, which yields nonintegrability of $\exp(-2c(u)u)$, i.e., the openness conjecture. Moreover, the curve D associated to this region is *smooth*, and then $\nu(u)$ is a Kiselman number of u . It is to establish the existence and main properties of the optimal region \mathcal{A} that we bring valuations into the picture. In particular, the concavity properties of the tree transform of u play a key role.

Most results presented here are probably not particular to dimension 2. The main difficulty in extending our approach to higher dimensions lies in understanding the analogue of the valuative tree. We hope to tackle this problem in future research.

The present article relies in an essential way on the analysis and formalism in our previous work: we recall in Section 1 the main results of [FJ1], [FJ2] that will be used. The rest of the paper is then organized as follows: in Section 2 we give the definition and main properties of multiplier ideals of tree potentials. As we show in Sections 3 and 4, this notion naturally generalizes multiplier ideals of formal powers of ideals, graded systems of ideals, and psh functions. Sections 5, 6 and 7 contain the applications mentioned above. We study equisingularity in Section 8 and end the paper with an appendix containing two proofs.

1. BACKGROUND

In this section we give a brief review of the valuative tree and its applications to the study of ideals and plurisubharmonic (psh) functions. For details, we refer to [FJ1] for Sections 1.1 to 1.3, and to [FJ2] for Section 1.4.

1.1. Conventions. In general, (R, \mathfrak{m}) denotes an equicharacteristic zero, two-dimensional regular local ring with algebraically closed residue field k . We will refer to this as the *general case*.

Whenever we talk about psh functions, we will always be in the *analytic case*, meaning that $R = \mathcal{O}_0$ is the ring of holomorphic germs at the origin in \mathbf{C}^2 . Then $k = \mathbf{C}$ and \mathfrak{m} is the maximal ideal of germs vanishing at the origin.

In general we write $(\hat{R}, \hat{\mathfrak{m}})$ for the completion of R . It is the ring of formal power series in two variables with coefficients in k .

1.2. The valuative tree. Our starting point is the approach to valuations worked out in [FJ1].

1.2.1. Valuations ([FJ1, Section 1.2]). We consider the space \mathcal{V} of centered, normalized valuations on R , i.e., the set of functions $\nu : R \rightarrow [0, \infty]$ satisfying:

- (i) $\nu(\psi\psi') = \nu(\psi) + \nu(\psi')$ for all ψ, ψ' ;
- (ii) $\nu(\psi + \psi') \geq \min\{\nu(\psi), \nu(\psi')\}$ for all ψ, ψ' ;
- (iii) $\nu(0) = \infty, \nu|_{\mathcal{C}^*} = 0, \nu(\mathfrak{m}) := \min\{\nu(\psi) ; \psi \in \mathfrak{m}\} = 1$.

Then \mathcal{V} is equipped with a natural *partial ordering*: $\nu \leq \mu$ iff $\nu(\psi) \leq \mu(\psi)$ for all $\psi \in \mathfrak{m}$. The *multiplicity valuation* $\nu_{\mathfrak{m}}$ defined by $\nu_{\mathfrak{m}}(\psi) = m(\psi) = \max\{k ; \psi \in \mathfrak{m}^k\}$ is the unique minimal element of \mathcal{V} . (See [FJ1, Section 1.5.1].)

Any valuation on R extends uniquely to a valuation in its completion \hat{R} ; hence the valuation spaces attached to R and \hat{R} are isomorphic. (See [Sp, Theorem 3.1].)

1.2.2. *Curve valuations* ([FJ1, Section 1.5.5]). Some natural maximal elements are the *curve valuations* defined as follows. To each irreducible (possibly formal) curve C we associate $\nu_C \in \mathcal{V}$ defined by $\nu_C(\psi) = C \cdot \{\psi = 0\}/m(C)$, where “ \cdot ” denotes intersection multiplicity and m multiplicity. If C is defined by $\phi \in \hat{\mathfrak{m}}$, then we also write $\nu_C = \nu_{\phi}$. Note that $\nu_{\phi}(\phi) = \infty$.

The set \mathcal{C} of local irreducible curves carries a natural (ultra)metric in which \mathcal{C} has diameter 1. It is given by $d_{\mathcal{C}}(C, D) = m(C)m(D)/C \cdot D$. (See [FJ1, Lemma 3.56].)




FIGURE 1. The valuative tree. The segments consist of valuations of the form $\nu_{\phi,t}$, where $\phi = x, y^2 - x^3, \dots, y$ and the skewness parameter t ranges from 1 to ∞ . Skewness $t = 1$ gives the multiplicity valuation $\nu_{\mathfrak{m}}$ and skewness $t = \infty$ the curve valuation ν_{ϕ} . The integer label above a segment indicates multiplicity (Section 1.2.6).

1.2.3. *Quasimonomial valuations* ([FJ1, Section 1.5.4]). Arguably the most important valuations in \mathcal{V} are the *quasimonomial* ones.¹ They are of the form $\nu_{C,t}$, where

¹A quasimonomial valuation can be made monomial (i.e., completely determined by its values on a pair of local coordinates (x, y)) by a birational morphism. Quasimonomial valuations are also known as Abhyankar valuations of rank 1; see [ELS].

$C \in \mathcal{C}$ and $t \in [1, \infty)$, and satisfy $\nu_{C,t}(\psi) = \min\{\nu_D(\psi) : d_C(C, D) \leq t^{-1}\}$. We have $\nu_{C,s} = \nu_{D,t}$ iff $s = t \geq d_C(C, D)^{-1}$. (See [FJ1, Proposition 3.55].) Thus \mathcal{V}_{qm} , the set of all quasimonomial valuations, is naturally a quotient of $\mathcal{C} \times [1, \infty)$ and has a natural tree structure: if $\nu, \nu' \in \mathcal{V}_{\text{qm}}$ and $\nu < \nu'$, then the *segment* $[\nu, \nu'] = \{\mu \in \mathcal{V}_{\text{qm}} : \nu \leq \mu \leq \nu'\}$ is isomorphic to a compact real interval. See [FJ1, Theorem 3.57] and Figure 1. We set $\nu_{\phi,t} := \nu_{C,t}$ when $C = \{\phi = 0\}$.

Quasimonomial valuations are of two types: *divisorial* and *irrational*, depending on whether the parameter t is rational or irrational.² We write \mathcal{V}_{div} for the set of divisorial valuations.

The full space \mathcal{V} is the completion of \mathcal{V}_{qm} in the sense that every element in \mathcal{V} is the limit of an increasing sequence in \mathcal{V}_{qm} . It is hence also naturally a tree, called the *valuative tree*. The ends of \mathcal{V} are exactly the elements of $\mathcal{V} \setminus \mathcal{V}_{\text{qm}}$ and are either curve valuations or *infinitely singular* valuations.³

1.2.4. *Skewness and intersection multiplicity* ([FJ1, Section 3.3]). An important invariant of a valuation is its *skewness* α defined by $\alpha(\nu) = \sup\{\nu(\phi)/m(\phi) : \phi \in \mathfrak{m}\}$. Skewness naturally *parameterizes* the trees \mathcal{V}_{qm} and \mathcal{V} in the sense that $\alpha : \mathcal{V}_{\text{qm}} \rightarrow [1, \infty)$ is strictly increasing and restricts to a bijection onto its image on any segment; indeed $\alpha(\nu_{\phi,t}) = t$ for any $\nu_{\phi,t} \in \mathcal{V}_{\text{qm}}$. Thus divisorial (irrational) valuations have rational (irrational) skewness. Curve valuations have infinite skewness whereas the skewness of an infinitely singular valuation may or may not be finite.

The tree structure on \mathcal{V} implies that any collection $(\nu_i)_{i \in I}$ of valuations in \mathcal{V} admits an *infimum* $\wedge_i \nu_i$; see [FJ1, Corollary 3.15]. Together with skewness, this allows us to define an *intersection product* on \mathcal{V} : we set $\nu \cdot \mu := \alpha(\nu \wedge \mu) \in [1, \infty]$. This is a normalized extension of the intersection product on \mathcal{C} as $C \cdot D = (\nu_C \cdot \nu_D)m(C)m(D)$. If $\nu \in \mathcal{V}$ and $\phi \in \mathfrak{m}$ is irreducible, then $\nu(\phi) = m(\phi)(\nu \cdot \nu_\phi)$. Moreover, if $\nu(\phi)$ is irrational, then $\nu = \nu_{\phi,t}$ with $t = \nu(\phi)/m(\phi)$.

1.2.5. *Tangent space and weak topology* ([FJ1, Sections 3.1-3.2]). Let μ be a valuation in \mathcal{V} . Declare $\nu, \nu' \in \mathcal{V} \setminus \{\mu\}$ to be equivalent if the segments $]\mu, \nu]$ and $]\mu, \nu']$ intersect. An equivalence class is called a *tangent vector* at μ and the set of tangent vectors at μ , the *tangent space*, is denoted by $T\mu$. If \vec{v} is a tangent vector, we denote by $U(\vec{v})$ the set of points in \mathcal{V} defining the equivalence class \vec{v} . The points in $U(\vec{v})$ are said to *represent* \vec{v} .

A point μ in the tree \mathcal{V} is an *end*, a *regular point*, or a *branch point* when $T\mu$ contains one, two, or three or more points, respectively. In terms of valuations: the ends of \mathcal{V} are curve and infinitely singular valuations; the regular points are irrational valuations; and the branch points are divisorial valuations, at which the tangent space is in bijection with the complex projective line \mathbf{P}^1 and hence uncountable. See [FJ1, Theorem 3.20].

We endow \mathcal{V} with the *weak topology*, generated by the sets $U(\vec{v})$ over all tangent vectors \vec{v} ; this turns \mathcal{V} into a compact (Hausdorff) space. If $\nu_k \rightarrow \nu$, then $\nu_k \wedge \mu \rightarrow \nu \wedge \mu$ for all $\mu \in \mathcal{V}$. The weak topology on \mathcal{V} is characterized by $\nu_k \rightarrow \nu$ iff $\nu_k(\phi) \rightarrow \nu(\phi)$ for all $\phi \in R$; see [FJ1, Theorem 5.1].

²A quasimonomial valuation ν is irrational iff $\nu(R) \not\subset \mathbf{Q}$, hence the name.

³The latter are represented by infinite Puiseux series whose exponents are rational numbers with unbounded denominators.

1.2.6. *Multiplicities* ([FJ1, Section 3.4]). By setting $m(\nu) := \min\{m(C) : C \in \mathcal{C}, \nu_C \geq \nu\}$ we extend the notion of *multiplicity* from \mathcal{C} to \mathcal{V}_{qm} . Clearly $m : \mathcal{V}_{\text{qm}} \rightarrow \mathbf{N}$ is increasing and hence extends to all of \mathcal{V} . In fact $m(\nu)$ divides $m(\mu)$ whenever $\nu \leq \mu$. The infinitely singular valuations are characterized as having infinite multiplicity.

As m is increasing and integer-valued, it is piecewise constant on any segment $[\nu_m, \nu_\phi]$, where $\phi \in \mathcal{C}$. This implies that $m(\vec{v})$ is naturally defined for any tangent vector \vec{v} . If ν is nondivisorial, then $m(\vec{v}) = m(\nu)$ for any $\vec{v} \in T\nu$.

If ν is divisorial, then the situation is more complicated. Suffice it to say that there exists an integer $b(\nu)$, divisible by $m(\nu)$, such that $m(\vec{v}) = b(\nu)$ for all but at most two tangent vectors \vec{v} at ν . We call $b(\nu)$ the *generic multiplicity* of ν .

1.2.7. *Approximating sequences* ([FJ1, Section 3.5]). Consider a quasimonomial valuation $\nu \in \mathcal{V}_{\text{qm}}$. The multiplicity m is integer-valued and piecewise constant on the segment $[\nu_m, \nu]$, hence has a finite number g (possibly zero) of jumps. Thus there are divisorial valuations ν_i , $0 \leq i \leq g$ and integers m_i , such that

$$(1.1) \quad \nu_m = \nu_0 < \nu_1 < \cdots < \nu_g < \nu_{g+1} = \nu$$

and $m(\mu) = m_i$ for $\mu \in]\nu_i, \nu_{i+1}]$, $0 \leq i \leq g$. We call the sequence $(\nu_i)_{i=1}^g$ the *approximating sequence* associated to ν . It plays a prominent role in [Sp].

The concept of approximating sequences extends naturally to valuations that are not quasimonomial: for curve valuations the sequences are still finite, for infinitely singular valuations they are infinite.

1.2.8. *Thinness* ([FJ1, Section 3.6]). Skewness α is a parameterization of \mathcal{V} that does not “see” multiplicities. Another parameterization, of crucial importance, is *thinness* A , defined as follows. If $\nu \in \mathcal{V}_{\text{qm}}$, then

$$(1.2) \quad A(\nu) = 2 + \int_{\nu_m}^{\nu} m(\mu) d\alpha(\mu).$$

In terms of (1.1) we have $A(\nu) = 2 + \sum_0^g m_i(\alpha_{i+1} - \alpha_i)$ with $\alpha_i = \alpha(\nu_i)$. Note that $A(\nu) \leq 1 + m(\nu)\alpha(\nu)$. Just like skewness, we may define $A(\nu)$ also for $\nu \notin \mathcal{V}_{\text{qm}}$.

The inequality $A(\nu) \geq 1 + \alpha(\nu)$ always holds, with equality iff $m(\nu) = 1$. Moreover, $A(\nu) < m(\nu)\alpha(\nu)$ as soon as $m(\nu) > 1$, and $A(\nu) - m(\nu)\alpha(\nu) \rightarrow -\infty$ as ν increases towards an infinitely singular valuation.

1.2.9. *Geometric interpretation of divisorial valuations* ([FJ1, Chapter 6]). Every divisorial valuation ν arises as follows: there exists a finite composition of point blowups $\pi : X \rightarrow \text{Spec } R$ (i.e., $\pi : X \rightarrow (\mathbf{C}^2, 0)$ in the analytic case) and an *exceptional component* E (i.e., an irreducible component of $\pi^{-1}(0)$) such that $\nu = b^{-1}\pi_* \text{div}_E$, where $b = b(\nu)$ is the generic multiplicity at ν and div_E denotes the order of vanishing along E . The generic multiplicity of ν is then equal to the multiplicity of any *curvette* of ν , i.e., the image by π of any smooth curve intersecting E transversely at a smooth point. In fact, every generic tangent vector at ν is represented by the curve valuation associated to a curvette [FJ1, Section 6.6.1]. We also have $A(\nu) = a/b$, where $a - 1$ is equal to the order of vanishing along E of the Jacobian determinant of π [FJ1, Theorem 6.22].

1.2.10. *Borel measures and tree potentials* ([FJ1, Chapter 7] or [FJ2, Section 5]). Let \mathcal{M} be the space of (weak, regular) positive Borel measures on \mathcal{V} ,⁴ endowed with the topology of vague convergence. We will identify $\nu \in \mathcal{V}$ with its Dirac mass $\delta_\nu \in \mathcal{M}$, thus viewing \mathcal{V} as a (compact) subset of \mathcal{M} .

Any $\rho \in \mathcal{M}$ determines a real-valued function g_ρ on \mathcal{V}_{qm} defined by $g_\rho(\nu) = \int_{\mathcal{V}} \mu \cdot \nu d\rho(\mu)$. A function of the form g_ρ is called a *tree potential*. The space \mathcal{P} of tree potentials is a closed convex cone in $\mathcal{V}_{\text{qm}}^{\mathbf{R}}$ (under pointwise convergence) and the map $\rho \mapsto g_\rho$ is a homeomorphism of \mathcal{M} onto \mathcal{P} whose inverse naturally defines a *Laplace operator* $\Delta : \mathcal{P} \rightarrow \mathcal{M}$.⁵ The mass of Δg is $g(\nu_{\mathbf{m}})$. See [FJ1, Theorem 7.64].

If $g \in \mathcal{P}$, then g is increasing and concave on any segment $[\nu_{\mathbf{m}}, \nu_0]$ (parameterized by skewness) in \mathcal{V} . If $\nu \in]\nu_{\mathbf{m}}, \nu_0[$, then the left derivative of g at ν with respect to skewness is equal to $\rho\{\mu \geq \nu\}$, whereas the right derivative is given by $\rho(U(\vec{v}))$, where $\rho = \Delta g$ and \vec{v} denotes the tangent vector at ν represented by ν_0 . We always have $\rho\{\mu \geq \nu\}\alpha(\nu) \leq g(\nu) \leq g(\nu_{\mathbf{m}})\alpha(\nu)$ with equality (in either inequality) iff $\rho = \Delta g$ is supported on $\{\mu \geq \nu\}$.

A *subtree* of \mathcal{V} is a subset \mathcal{T} such that $\nu \in \mathcal{T}$ and $\mu \leq \nu$ implies $\mu \in \mathcal{T}$. A subtree is *finite* if it has finitely many ends. If $g \in \mathcal{P}$ and \mathcal{T} is a subtree of \mathcal{V} , then $g_{\mathcal{T}}$ denotes the infimum of all tree potentials coinciding with g on \mathcal{T} .

The *support* of $g \in \mathcal{P}$ is the smallest subtree \mathcal{T} for which $g = g_{\mathcal{T}}$. Alternatively, it is the smallest subtree containing $\text{supp } \Delta g$. The support of any tree potential is included in the closure of a countable union of finite trees.

1.3. **Ideals** ([FJ1, Section 8.1]). An ideal I in R is *primary* if $\mathfrak{m}^n \subset I$ for some $n > 0$. The *integral closure* of I is the ideal \bar{I} of $\phi \in R$ such that $\phi^n + a_1\phi^{n-1} + \dots + a_n = 0$ for some $n \geq 1$ and $a_i \in I^i$. If $I = \bar{I}$, then I is *integrally closed*.

Any ideal I in R has an associated *tree transform* g_I , defined by $g_I(\nu) = \nu(I) = \min\{\nu(\phi) : \phi \in R\}$. This function g_I belongs to \mathcal{P} , and thus defines a *tree measure* $\rho_I = \Delta g_I \in \mathcal{M}$. The latter measure has mass $m(I) := \nu_{\mathbf{m}}(I)$.

If $I = \phi R$ is principal, then we write $g_\phi = g_I$ and $\rho_\phi = \rho_I$: the latter measure is given by $\rho_\phi = \sum_i n_i m_i \nu_i$, where $\phi = \prod \phi_i^{n_i}$ is the factorization of ϕ into irreducible factors, $m_i = m(\phi_i)$ and ν_i is the curve valuation associated to ϕ_i .

If I is primary, then $\rho_I = \sum_i n_i b_i \nu_i$, where $n_i \in \mathbf{N}$ and ν_i are divisorial valuations with generic multiplicity b_i . The valuations ν_i are exactly the *Rees valuations* of I .

A general ideal I is the product of a principal ideal and a primary ideal, hence has a tree measure of the form $\rho_I = \sum_i n_i b_i \nu_i + \sum_i n_i m_i \nu_i$.

To any measure ρ is associated an ideal $I_\rho = \{\phi \in R : g_\phi \geq g_\rho\}$. When ρ is of the previous form, then $\rho_{I_\rho} = \rho$, whereas I_{ρ_I} is the integral closure of I . The decomposition of ρ_I above gives an interpretation of Zariski's factorization theorem: if I is integrally closed, then $I = \prod_i I_{\nu_i}^{n_i}$. If furthermore I is primary, i.e., all the ν_i are divisorial, then $I = \bigcap_i \{\phi \in R : g_\phi(\nu_i) \geq g_\rho(\nu_i)\}$.

In general, the mass of ρ_I on a curve valuation ν_ϕ is the product of $m(\phi)$ and $\text{div}_\phi(I) := \max\{k : \phi^k | \psi \text{ for all } \psi \in I\}$.

1.4. **Psh functions.** All plurisubharmonic (psh) functions are defined near the origin in \mathbf{C}^2 . A psh function u is said to have *logarithmic singularities* if there exist $c > 0$ and holomorphic functions $\phi_i \in R$ such that $u = \frac{c}{2} \log \sum_1^n |\phi_i|^2 + O(1)$.

⁴In [FJ1], \mathcal{M} is denoted by \mathcal{M}^+ .

⁵In [FJ1], \mathcal{P} is denoted by \mathcal{P}^+ , and tree potentials are called *positive tree potentials*.

1.4.1. *Lelong and Kiselman numbers.* For a fixed choice of coordinates (x, y) , and weights $a, b > 0$ the *Kiselman number* [Ki1, Ki2] of u is defined to be

$$\nu_{a,b}^{x,y}(u) = \lim_{r \rightarrow 0} \frac{ab}{\log r} \sup\{u : |x| < r^{1/a}, |y| < r^{1/b}\}.$$

We have $\nu_{\lambda a, \lambda b}^{x,y}(u) = \lambda \nu_{a,b}^{x,y}(u)$ for any $\lambda > 0$. When $a = b = 1$, the Kiselman number reduces to the *Lelong number* $\nu^L(u)$. The latter does not depend on the choice of (x, y) .




FIGURE 2. Characteristic regions, see Section 1.4.2. To the left is a characteristic region \mathcal{A} around a singular curve $\{\phi = 0\}$. If $\pi : X \rightarrow (\mathbb{C}^2, 0)$ desingularizes this curve, the preimage of \mathcal{A} is a characteristic region around a smooth curve.

1.4.2. *Evaluating valuations on psh functions* ([FJ2, Section 3]). One can evaluate a quasimonomial valuation on a psh function as the normalized push-forward of a Kiselman number under a birational morphism. Concretely, this goes as follows. If $\nu \in \mathcal{V}_{\text{qm}}$ is quasimonomial, write $\nu = \nu_{\phi,t}$ for $\phi \in \mathfrak{m}$ irreducible with $m(\phi) = m(\nu) =: m$. Pick a coordinate x transverse to ϕ , i.e., $x \in \mathfrak{m}$ is irreducible, $m(x) = 1$ and $\nu_x \wedge \nu_\phi = \nu_m$. Also pick constants $0 < C_1 < C_2 < \infty$ and consider the region

$$\mathcal{A}(r) = \mathcal{A}_{\phi,t,x,C}(r) := \{|x| < r : C_1|x|^{mt} < |\phi| < C_2|x|^{mt}\}$$

for small r . See Figure 2. We have $\text{Vol } \mathcal{A}(r) \simeq r^{2A}$, where $A = A(\nu)$ is the thinness of ν . The value of ν on a psh function u is then given by

$$\nu(u) = \lim_{r \rightarrow 0} \frac{1}{\log r} \sup_{\mathcal{A}(r)} u;$$

this does not depend on the choices of ϕ, x or C . We always have the upper bound $u(q) \leq \nu(u) \log \|q\| + O(1)$ in $\mathcal{A}(r)$. If $u = \frac{c}{2} \log \sum_1^n |\phi_i|^2 + O(1)$ has logarithmic singularities, then $\nu(u) = c \min_i \nu(\phi_i) = c \nu(I)$, where I is the ideal generated by the ϕ_i 's. In this case we also have the lower bound $u(q) \geq \nu(u) \log \|q\| + O(1)$ for suitable choices of C_1 and C_2 . See [FJ2, Proposition 4.3].

If $\nu \in \mathcal{V}_{\text{qm}}$ is monomial, i.e., of multiplicity one, then there exist local coordinates (x, y) such that $\nu = \nu_{y,t}$, where $t = \alpha(\nu) \geq 1$. In this case $\nu(u)$ equals the Kiselman number $\nu_{t,1}^{x,y}(u)$. In particular $\nu_m(u)$ is the Lelong number of u .

1.4.3. *Geometric interpretation.* The value of divisorial valuations on psh functions can be interpreted along the lines of Section 1.2.9. Using the notation of that section, $b(\nu)\nu(u)$ is the Lelong number of the pull-back π^*u at a generic point $p \in E$ [FJ2, Proposition 4.1]. Moreover, if $p \in E$ is a smooth point on $\pi^{-1}(0)$, then the Lelong number of the strict transform of u under π at p is bounded from above by $b(\nu)^{-1} \rho_u(U(\vec{v}))$ [FJ2, Lemma 7.6]. Here ρ_u is the tree measure of u (see below)

and $\vec{v} = \vec{v}_p$ is the tangent vector at ν associated to p ; that is, \vec{v} is represented by any curve valuation ν_C such that the strict transform of C is smooth and intersects E transversely at p .

1.4.4. *Demailly approximation.* Demailly devised a method, crucially important to our analysis, of approximating a general psh function u with a sequence u_n of psh functions with logarithmic singularities. This goes as follows: see Theorem 4.2 and its proof in [DK] for details. Suppose u is psh on a fixed ball B containing the origin. For $n > 0$ let \mathcal{H}_n be the Hilbert space of holomorphic functions ψ on B such that $\int_B |\psi|^2 e^{-2nu} < \infty$. Let $(g_{nl})_1^\infty$ be an orthonormal basis for \mathcal{H}_n and set $u_n = \frac{1}{2n} \log \sum_l |g_{nl}|^2$. If $0 \ni B' \Subset B$ is a smaller ball, then there exists $k = k(u, n, B') < \infty$ such that $u_n - \frac{1}{2n} \log \sum_1^k |g_{nl}|^2$ is bounded in B' ; in particular u_n has a logarithmic singularity at the origin.

Demailly approximation interacts very well with the evaluation of quasimonomial valuations on psh functions: we have $0 \leq \nu(u) - \nu(u_n) \leq A(\nu)/n$, where A denotes thinness [FJ2, Proposition 3.12].

1.4.5. *Tree transforms* ([FJ2, Section 6.1]). Any psh function u has an associated *tree transform* $g_u \in \mathcal{P}$, defined by $g_u(\nu) = \nu(u)$ for $\nu \in \mathcal{V}_{\text{qm}}$, as well as an associated *tree measure* $\rho_u = \Delta g_u$ on \mathcal{V} . That these are well defined follows from Demailly approximation. If u is psh, then ρ_u cannot put mass on a formal (i.e., non-analytic) curve valuation. When ν_C is a curve valuation associated to an analytic curve $C = \{\phi = 0\}$, the mass of ρ_u on ν_C is exactly $m(C)$ times the mass of the positive closed $(1, 1)$ current $dd^c u$ on C . By Siu's theorem, this is equivalent to saying that $u = u' + c \log |\phi|$ with $c = m(C)^{-1} \rho_u \{\nu_\phi\}$ and u' a psh function with $\rho_{u'} \{\nu_C\} = 0$.

2. MULTIPLIER IDEALS OF TREE POTENTIALS

In this section we define multiplier ideals of tree potentials and examine their main properties. As we will show in subsequent sections, this notion contains all previously known (to the authors) definitions of multiplier ideals, in a local, two-dimensional setting. We also introduce some related singularity exponents.

Definition 2.1. Let $h : \mathcal{V}_{\text{qm}} \rightarrow [0, \infty)$ be a tree potential on \mathcal{V} , i.e., $h \in \mathcal{P}$. We define the *multiplier ideal* $\mathcal{J}(h)$ of h to be the ideal of elements $\psi \in R$ such that

$$(2.1) \quad \sup_{\nu \in \mathcal{V}_{\text{qm}}} \frac{h(\nu)}{\nu(\psi) + A(\nu)} < 1.$$

Definition 2.2. Let $h \in \mathcal{P}$. The *singularity exponent*, or *log-canonical threshold*, of h is the number $c(h) := \sup\{c > 0 : \mathcal{J}(ch) = R\}$. The *Arnold multiplicity* is the number $\lambda(h) = c(h)^{-1}$. Thus we have

$$(2.2) \quad \lambda(h) = \sup_{\nu \in \mathcal{V}_{\text{qm}}} \frac{h(\nu)}{A(\nu)} \quad \text{and} \quad c(h) = \inf_{\nu \in \mathcal{V}_{\text{qm}}} \frac{A(\nu)}{h(\nu)}.$$

Remark 2.3. It follows immediately from the definition that $\mathcal{J}(c(h)h) \subsetneq R$.

It will be useful to have a less elegant but more concrete criterion for membership in the multiplier ideal.

Proposition 2.4. Fix a tree potential $h \in \mathcal{P}$ and $\psi \in R$. Let ρ_h and ρ_ψ be the associated measures on \mathcal{V} . Then $\psi \in \mathcal{J}(h)$ iff the following hold:

- (a) $\nu(\psi) + A(\nu) > h(\nu)$ for all quasimonomial valuations ν ;

(b) $\rho_\psi\{\nu\} + m(\nu) > \rho_h\{\nu\}$ for all curve valuations ν .

Moreover, we have

- (i) if $\psi \notin \mathfrak{m}$, then it suffices to check (a)-(b) for ν of multiplicity one;
- (ii) if $h = c g_I$ for some ideal I and $c > 0$, then it suffices to check (a)-(b) for ν in the approximating sequence of some valuation in the support of ρ_I .

The proof is given below. Note that (2.1) is not a trivial consequence of (a).

2.1. Properties. Let us now state some basic properties of multiplier ideals associated to tree potentials. The proofs are given below. However, two of them are relegated to the appendix as they do not follow directly from our approach, but rely on the corresponding statements for ideals already proved in the literature.

Proposition 2.5. *The multiplier ideal $\mathcal{J}(h)$ is integrally closed.*

It is clear from the definition that if $g \leq h$, then $\mathcal{J}(h) \subset \mathcal{J}(g)$. A deeper result is the following semicontinuity statement.

Proposition 2.6. *If $(h_n)_1^\infty$ is a decreasing sequence of tree potentials converging in \mathcal{P} (i.e., pointwise) to a tree potential h , then $\mathcal{J}(h_n) = \mathcal{J}(h)$ for $n \gg 1$.*

Notice that since R is Noetherian, the sequence $(\mathcal{J}(h_n))_1^\infty$ is stationary. This does not, however, immediately imply the proposition.

We can bound the Arnold multiplicity as follows.

Proposition 2.7. *Consider $h \in \mathcal{P}$ with associated measure $\rho = \Delta h$. Then:*

- (i) $\frac{1}{2}h(\nu_{\mathfrak{m}}) \leq \lambda(h) \leq h(\nu_{\mathfrak{m}})$;
- (ii) $\lambda(h) > \frac{1}{2}h(\nu_{\mathfrak{m}})$ iff there exists a tangent vector \vec{v} at $\nu_{\mathfrak{m}}$ with $\rho U(\vec{v}) > \frac{1}{2}$;
- (iii) $\lambda(h) = h(\nu_{\mathfrak{m}})$ iff ρ is a point mass at a curve valuation of multiplicity one.

Next we have the following subadditivity property.

Proposition 2.8. *If $h_1, h_2 \in \mathcal{P}$, then $\mathcal{J}(h_1 + h_2) \leq \mathcal{J}(h_1)\mathcal{J}(h_2)$.*

Our proof uses a reduction to the case when the h_i 's are proportional to tree transforms of ideals and is given in the appendix. The same holds for the following version of Skoda's Theorem:

Proposition 2.9. *Let ν_0 be a divisorial valuation with associated (simple complete) ideal $I_0 = I_{\nu_0}$ and tree transform $g_0 = g_{I_0}$. Then we have*

$$\mathcal{J}(h + 2g_0) = I_0\mathcal{J}(h + g_0)$$

for any tree potential $h \in \mathcal{P}$.

On the other hand, Proposition 2.10 is a version of Skoda's Theorem that we can prove directly.

Proposition 2.10. *Consider $\phi \in \mathfrak{m}$ with associated tree potential g_ϕ . Then*

$$\mathcal{J}(h + g_\phi) = \phi\mathcal{J}(h)$$

for any tree potential $h \in \mathcal{P}$.

2.2. Proofs. We now turn to the proofs. The fact that multiplier ideals are integrally closed is easy to establish.

Proof of Proposition 2.5. Let $(\psi_i)_1^k$ be generators of $I = \mathcal{J}(h)$. If $\psi \in \bar{I}$, then $\nu(\psi) \geq \min_i \nu(\psi_i)$ for every $\nu \in \mathcal{V}_{\text{qm}}$. This implies that

$$\sup_{\nu} \frac{h(\nu)}{A(\nu) + \nu(\psi)} \leq \max_i \sup_{\nu} \frac{h(\nu)}{\nu(\psi_i) + A(\nu)} < 1$$

so that $\psi \in I$. □

We now turn to the characterization of $\mathcal{J}(h)$ given in Proposition 2.4. The main point is that the supremum in the definition of $\mathcal{J}(h)$ is actually attained. In fact, the following lemma is the key to the openness conjecture for psh functions.

Lemma 2.11. *Fix a tree potential $h \in \mathcal{P}$ and let $\psi \in R$. The function*

$$(2.3) \quad \chi(\nu) = \chi_{h,\psi}(\nu) = \frac{h(\nu)}{\nu(\psi) + A(\nu)},$$

defined on \mathcal{V}_{qm} , extends to a function on \mathcal{V} continuous on segments. Its supremum is attained at a valuation ν_ . We may choose ν_* to be a quasimonomial or a curve valuation, and to belong to the support of h . Further, we may take ν_* such that:*

- (i) *if $\psi \notin \mathfrak{m}$, then ν_* is of multiplicity one;*
- (ii) *if $h = g_I$ for some ideal I in R , then ν_* is an element in the approximating sequence of some valuation in $\text{supp } \rho_I$; in particular, ν_* is either divisorial or a curve valuation at an element of R ;*
- (iii) *if $h = g_u$ for some psh function u , then ν_* is either quasimonomial or an analytic curve valuation.*

The main observation in the proof of Lemma 2.11 is that it suffices to consider h on a finite subtree of \mathcal{V} . In the same spirit we have:

Lemma 2.12. *For $h \in \mathcal{P}$ and $0 < \varepsilon < 1$, the set*

$$\mathcal{T} = \mathcal{T}_{h,\varepsilon} := \{\nu \in \mathcal{V} : \rho_h\{\mu \geq \nu\} \geq (1 - \varepsilon)m(\nu)\}$$

is a finite subtree of \mathcal{V} and $\mathcal{J}(h) = \mathcal{J}(h_{\mathcal{T}})$, where $h_{\mathcal{T}}$ is the smallest tree potential coinciding with h on \mathcal{T} .

Proof of Lemma 2.11. We may assume that $\rho := \Delta h$ has mass 1. Set

$$\mathcal{T} = \mathcal{T}_{h,\psi} := \{\nu : \rho\{\mu \geq \nu\} \geq \chi(\nu_{\mathfrak{m}})m(\nu)\}.$$

If $\nu \leq \nu'$, then $m(\nu) \leq m(\nu')$ and $\rho\{\mu \geq \nu\} \geq \rho\{\mu \geq \nu'\}$. Hence \mathcal{T} is a tree. The number of ends of \mathcal{T} is finite, bounded by $1/\chi(\nu_{\mathfrak{m}})$. Thus \mathcal{T} is a finite tree. Its ends have finite multiplicity, hence consist of quasimonomial or curve valuations.

We claim that $\sup_{\mathcal{V}} \chi = \sup_{\mathcal{T}} \chi$. To see this, pick any $\nu_1 \notin \mathcal{T}$. Let $\nu_0 = \max\{\nu \in \mathcal{T} : \nu \leq \nu_1\}$ and let $\vec{\nu}$ be the tangent vector at ν_0 represented by ν_1 . Write $m(\vec{\nu}) = m$, $\alpha(\nu_0) = \alpha_0$ and, for $\nu \in [\nu_0, \nu_1]$, $\alpha = \alpha(\nu)$. The right derivative of h at ν_0 with respect to the skewness equals $\rho U(\vec{\nu})$ (see Section 1.2.10). As h is concave on $[\nu_0, \nu_1]$ we have $h(\nu) \leq h(\nu_0) + \rho U(\vec{\nu})(\alpha - \alpha_0)$ for $\nu \in [\nu_0, \nu_1]$. On the other hand, $A(\nu) \geq A(\nu_0) + m(\alpha - \alpha_0)$ and $\nu(\psi) \geq \nu_0(\psi)$; hence

$$\chi(\nu) \leq \frac{h(\nu_0) + \rho U(\vec{\nu})(\alpha - \alpha_0)}{\nu_0(\psi) + A(\nu_0) + m(\alpha - \alpha_0)} := M(\alpha), \quad \text{for all } \nu \in [\nu_0, \nu_1].$$

Here $\alpha \mapsto M(\alpha)$ is Möbius in α ; hence

$$\sup_{[\nu_0, \nu_1]} \chi \leq \sup_{[\alpha_0, \infty]} M = \max\{\chi(\nu_0), m^{-1}\rho U(\vec{v})\} \leq \max\{\chi(\nu_0), \chi(\nu_m)\}.$$

To prove the last inequality, we note that $U(\vec{v}) \cap \mathcal{T} = \emptyset$. If $\rho U(\vec{v}) > m\chi(\nu_m)$, we could find $\nu' \in U(\vec{v})$ close enough to ν_0 , of multiplicity m and such that $\rho\{\mu \geq \nu'\} \geq m\chi(\nu_m)$. This would imply $\nu' \in \mathcal{T}$, a contradiction.

Thus $\sup_{\mathcal{V}} \chi = \sup_{\mathcal{T}} \chi$.

As each of the functions $h(\nu)$, $\nu(\psi)$ and $A(\nu)$ is continuous on segments in \mathcal{V}_{qm} , so is χ . We extend χ to \mathcal{V} by taking limits along segments. To see that these limits exist, pick $\nu_0 \notin \mathcal{V}_{\text{qm}}$ and let $\nu \rightarrow \nu_0$ along the segment $[\nu_m, \nu_0[$. If ν_0 is a curve valuation, then $h(\nu)/\alpha(\nu) \rightarrow \rho\{\nu_0\}$, $\nu(\psi)/\alpha(\nu) \rightarrow \rho_\psi\{\nu_0\}$ and $A(\nu)/\alpha(\nu) \rightarrow m(\nu_0)$. This gives $\chi(\nu) \rightarrow \rho\{\nu_0\}(\rho_\psi\{\nu_0\} + m(\nu_0))^{-1}$.

When ν_0 is infinitely singular, we claim that χ is decreasing when $\nu \in [\nu_m, \nu_0]$ is sufficiently close to ν_0 . This implies that $\chi(\nu)$ converges when $\nu \rightarrow \nu_0$. To prove the claim, note that $\nu \mapsto \nu(\psi)$ is constant equal to $\nu_0(\psi)$ near ν_0 and that the left derivative of h with respect to skewness at ν equals $\rho\{\mu \geq \nu\}$. The left derivative of χ at ν is hence equal to

$$(2.4) \quad \frac{d\chi}{d\alpha} = \frac{(A(\nu) + \nu_0(\psi))\rho\{\mu \geq \nu\} - m(\nu)h(\nu)}{(A(\nu) + \nu_0(\psi))^2}.$$

By Section 1.2.8, $A(\nu) - m(\nu)\alpha(\nu) \rightarrow -\infty$ as $\nu \rightarrow \nu_0$ and $h(\nu) \geq \rho\{\mu \geq \nu\}\alpha(\nu)$. This easily implies $d\chi/d\alpha < 0$ and completes the proof of the claim.

Thus the function χ extends to \mathcal{V} . Its restriction to the finite tree \mathcal{T} is continuous; hence $\sup_{\mathcal{V}} \chi$ is attained at some $\nu_* \in \mathcal{T}$.

Now suppose $\psi \notin \mathfrak{m}$, which amounts to $\nu(\psi) \equiv 0$. Consider a segment $]\nu_0, \nu_1[$ on which the multiplicity is constant equal to $m_0 \geq 2$. Then (2.4) and the two inequalities $A(\nu) < m(\nu)\alpha(\nu) = m_0\alpha(\nu)$ and $h(\nu) \geq \rho\{\mu \geq \nu\}\alpha(\nu)$ imply $\frac{d\chi}{d\alpha} < 0$ on $]\nu_0, \nu_1[$. This proves that χ is decreasing off the subtree $\{\nu : m(\nu) = 1\}$. Hence $m(\nu_*) = 1$, proving (i).

We saw above that if ν is a curve valuation, then $\chi(\nu) = 0$ unless ρ puts mass on ν . This proves (iii) as a measure represented by a psh function cannot charge formal curve valuations (see Section 1.4.5).

Finally, if $h = g_I$ for an ideal I , then ρ is atomic, supported on finitely many valuations that are either divisorial or curve valuations; see Section 1.3. On any open segment in \mathcal{V} not intersecting the support of ρ , and on which the multiplicity is constant, h is Möbius, hence attains its maximum at its boundary points. This implies (ii). □

Proof of Lemma 2.12. Suppose first that \mathcal{T} is empty. This happens exactly when $h(\nu_m) < 1 - \varepsilon$. We have $A(\nu) \geq 1 + \alpha(\nu)$ and $h(\nu) \leq h(\nu_m)\alpha(\nu)$ for all ν , whence $\sup_{\mathcal{V}} \chi_{h,1} \leq h(\nu_m) < 1$ so that $\mathcal{J}(h) = \mathcal{J}(h_{\mathcal{T}}) = R$.

Suppose now that \mathcal{T} is nonempty; i.e., $h(\nu_m) \geq 1 - \varepsilon$. We follow the proof of Lemma 2.11. Clearly \mathcal{T} is a subtree of \mathcal{V} with at most $h(\nu_m)/(1 - \varepsilon)$ ends. As $h \geq h_{\mathcal{T}}$ we have $\mathcal{J}(h) \subset \mathcal{J}(h_{\mathcal{T}})$. To prove the reverse inclusion, we proceed as follows. Suppose $\sup_{\mathcal{T}} \chi = t < 1$ where $\chi = \chi_{h,\psi}$. We must show that $\sup_{\mathcal{V}} \chi < 1$. Consider $\nu_0 \in \mathcal{T}$ and a segment $]\nu_0, \nu_1[$, disjoint from \mathcal{T} . Let $m = m(\vec{v})$, where \vec{v} is the tangent vector at ν_0 represented by ν_1 . For $\nu \in]\nu_0, \nu_1[$ we get

$$\chi(\nu) \leq \max\{\chi(\nu_0), \rho\{\mu \geq \nu\}/m\} \leq \max\{t, 1 - \varepsilon\}.$$

Hence $\sup_{\mathcal{V}} \chi \leq \max\{t, 1 - \varepsilon\} < 1$, which completes the proof. □

Proof of Proposition 2.4. This is an immediate consequence of Lemma 2.11 and the formula $\chi(\nu) = \rho_n\{\nu\}/(\rho_\psi\{\nu\} + m(\nu))$ for a curve valuation ν . □

Next we address semicontinuity. The main ingredient in the proof is the following lemma. It can be viewed as a general statement about parameterized trees, but we shall formulate it only for the valuative tree \mathcal{V} parameterized by skewness.

Lemma 2.13. *Under the assumptions of Proposition 2.6, the functions $\frac{h_n(\nu)}{\alpha(\nu)}$ decrease uniformly on \mathcal{V}_{qm} to the function $h(\nu)/\alpha(\nu)$.*

Proof of Proposition 2.6. Clearly $\mathcal{J}(h_n)$ forms an increasing sequence of ideals in R and $\mathcal{J}(h_n) \subset \mathcal{J}(h)$ for all n . Since R is Noetherian, it suffices to show that $\bigcup \mathcal{J}(h_n) \supset \mathcal{J}(h)$. Consider $\phi \in \mathcal{J}(h)$ and write $\chi(\nu) = h(\nu)/(\nu(\phi) + A(\nu))$ and $\chi_n(\nu) = h_n(\nu)/(\nu(\phi) + A(\nu))$. Then $\sup \chi < 1$. Since $A \geq \alpha$, Lemma 2.13 gives

$$0 \leq \chi_n(\nu) - \chi(\nu) \leq \frac{h_n(\nu)}{\alpha(\nu)} - \frac{h(\nu)}{\alpha(\nu)} \rightarrow 0$$

uniformly on \mathcal{V}_{qm} , as $n \rightarrow \infty$. This implies $\sup \chi_n < 1$, i.e., $\phi \in \mathcal{J}(h_n)$ for large n , completing the proof. □

Proof of Lemma 2.13. As mentioned above, a version of the lemma holds for arbitrary parameterized trees. For a totally ordered tree it translates into the following elementary statement, the proof of which is left to the reader.

Lemma 2.14. *Let g_n, g be positive concave functions on $[1, \infty)$ such that g_n decreases to g pointwise. Then $g_n(x)/x$ decreases to $g(x)/x$ uniformly.*

Of course, a tree such as \mathcal{V}_{qm} has a lot of branching. The following lemma will be used to control the behavior of tree potentials at branch points.

Lemma 2.15. *Consider tree potentials g, h and quasimonomial valuations μ, ν with $\mu < \nu$. Assume that*

$$\frac{g(\nu)}{\alpha(\nu)} \geq \frac{h(\nu)}{\alpha(\nu)} + \varepsilon \quad \text{and} \quad \frac{g(\mu)}{\alpha(\mu)} \leq \frac{h(\mu)}{\alpha(\mu)} + \varepsilon$$

for some $\varepsilon > 0$. Then $\rho U(\vec{v}) \geq \varepsilon$, where $\rho = \rho_g$ is the tree measure of g and \vec{v} is the tangent vector at μ represented by ν .

Proof. On the one hand, since h is increasing, the assumptions give

$$g(\nu) \geq h(\nu) + \varepsilon\alpha(\nu) \geq h(\mu) + \varepsilon\alpha(\nu) = h(\mu) + \varepsilon\alpha(\mu) + \varepsilon(\alpha(\nu) - \alpha(\mu)).$$

On the other hand, as g is a tree potential, we have

$$g(\nu) \leq g(\mu) + \rho U(\vec{v})(\alpha(\nu) - \alpha(\mu)) \leq h(\mu) + \varepsilon\alpha(\mu) + \rho U(\vec{v})(\alpha(\nu) - \alpha(\mu)).$$

Thus $\rho U(\vec{v}) \geq \varepsilon$. □

We continue the proof of Lemma 2.13 and argue by contradiction. If h_n/α does not converge uniformly to h/α , then, after passing to a subsequence if necessary, there exist $\nu_n \in \mathcal{V}_{\text{qm}}$ and $\varepsilon > 0$ such that $h_n(\nu_n)/\alpha(\nu_n) > h(\nu_n)/\alpha(\nu_n) + \varepsilon$ for all n . Using the compactness of \mathcal{V} we may assume that ν_n converges weakly to some valuation $\nu_* \in \mathcal{V}$. Since h_n converges to h pointwise on \mathcal{V}_{qm} we may assume that the ν_n are all distinct and different from ν_* .

By Lemma 2.14 we may assume $h_n/\alpha \leq h/\alpha + \varepsilon$ on $[\nu_m, \nu_\star[$. Hence $\nu_n \notin [\nu_m, \nu_\star[$ so that $\mu_n := \nu_\star \wedge \nu_n$ is quasimonomial and $\mu_n < \nu_n$. We apply Lemma 2.15 to $g = h_n$, $h = h$, $\mu = \mu_k$ and $\nu = \nu_k$, for $n \geq k$. Recall that $h_n \geq h_k$. We conclude that $\rho_n U_k \geq \varepsilon$, where $\rho_n = \rho_{h_n}$ and $U_k = U(\vec{v}_k)$, where \vec{v}_k is the tangent vector at μ_k represented by ν_k .

If $k \neq l$, then $U_k = U_l$ or $U_k \cap U_l = \emptyset$. As mass $\rho_n = h_n(\nu_m) \rightarrow h(\nu_m)$ is uniformly bounded, this implies that, after passing to a subsequence, all the U_k are equal. In particular all the μ_k are equal. As $\mu_k \rightarrow \nu_\star$, this gives $\mu_k = \nu_\star$ for all k . In particular, ν_\star is quasimonomial and $\nu_n > \nu_\star$.

For $k > 1$ we have $\nu_k \wedge \nu_1 \in]\nu_\star, \nu_1]$ as $U_k = U_1$. Again by Lemma 2.14, $\nu_k \wedge \nu_1 < \nu_k$ for all $k \geq 2$. As $\nu_1 \wedge \nu_k \rightarrow \nu_\star$ as $k \rightarrow \infty$ we may assume that the valuations $\nu_1 \wedge \nu_k$ are all distinct. We again apply Lemma 2.15, now with $\mu = \nu_1 \wedge \nu_k$, $\nu = \nu_k$ and still using $h = h$ and $g = h_n$ for $n \geq k$. We conclude that $\rho_n U'_k \geq \varepsilon$, where $U'_k = U(\vec{v}'_k)$ and \vec{v}'_k is the tangent vector at $\nu_1 \wedge \nu_k$ represented by ν_k . But the regions U'_k are mutually disjoint, so since the ρ_n has uniformly bounded mass, this gives a contradiction. \square

Next we prove the bounds for the singularity exponents.

Proof of Proposition 2.7. Recall that ρ has mass $h(\nu_m)$.

Write $\chi(\nu) = h(\nu)/A(\nu)$ so that $\lambda(h) = \sup \chi$. The lower bound $\lambda(h) \geq \chi(\nu_m) = \frac{1}{2}h(\nu_m)$ is immediate. Recall that $A(\nu) \geq 1 + \alpha(\nu)$. If $\nu \in \mathcal{V}_{\text{qm}}$, $\nu \neq \nu_m$, then $h(\nu) \leq h(\nu_m) + \rho U(\vec{v})(\alpha(\nu) - 1)$, where \vec{v} is the tangent vector at ν_m represented by ν . In general, this gives $h(\nu) \leq h(\nu_m)\alpha(\nu)$, so that $\chi(\nu) < h(\nu_m)$.

If $\rho U(\vec{v}) \leq \frac{1}{2}h(\nu_m)$, then $h(\nu) \leq \frac{1}{2}h(\nu_m)(1 + \alpha(\nu))$ so that $\chi(\nu) \leq \frac{1}{2}h(\nu_m)$. On the other hand, if $\rho U(\vec{v}) > \frac{1}{2}h(\nu_m)$, then we may move ν closer to ν_m , still keeping $\nu \in U(\vec{v})$, so that $m(\nu) = 1$, $A(\nu) = 1 + \alpha(\nu)$ and $h(\nu) > h(\nu_m) + \frac{1}{2}h(\nu_m)(\alpha(\nu) - 1)$. This gives $\lambda(h) \geq \chi(\nu) > \frac{1}{2}h(\nu_m)$.

Finally suppose $\lambda(h) = h(\nu_m)$. By Lemma 2.11 we have $\lambda(h) = \chi(\nu)$ for some valuation $\nu \in \mathcal{V}$ of multiplicity one. We have seen above that $\chi < h(\nu_m)$ on \mathcal{V}_{qm} . Hence ν is a curve valuation. But then $\chi(\nu) = \rho\{\nu\}/m(\nu) = \rho\{\nu\}$ so that $\rho\{\nu\} = h(\nu_m)$ and ρ is a point mass at ν . \square

Finally we prove the second version of Skoda's Theorem.

Proof of Proposition 2.10. By unique factorization we may assume that ϕ is irreducible. We first claim that $J(h + g_\phi) \subset \phi R$. Indeed, if $\psi \in \mathcal{J}(h + g_\phi)$, then by Proposition 2.4 we have $\rho_\psi\{\nu_\phi\} + m(\phi) > \rho_h\{\nu_\phi\} + \rho_\phi\{\nu_\phi\} \geq 0 + m(\phi)$. Thus $\rho_\psi\{\nu_\phi\} > 0$ so that ϕ divides ψ .

It hence suffices to show that if $\psi \in R$, then $\phi\psi \in \mathcal{J}(h + g_\phi)$ iff $\psi \in \mathcal{J}(h)$. But this follows easily from Proposition 2.4. Indeed, if ν is quasimonomial, then $\nu(\psi\phi) - (h + g_\phi)(\nu) = \nu(\psi) - h(\nu)$ and if ν is a curve valuation, then $\rho_{\psi\phi}\{\nu\} - (\rho_h + \rho_\phi)\{\nu\} = \rho_\psi\{\nu\} - \rho_h\{\nu\}$. \square

3. MULTIPLIER IDEALS OF IDEALS

We now show that if h is the tree transform of a formal power I^c , where I is an ideal and $c > 0$, then the multiplier ideal of h agrees with the multiplier ideal of I^c with the standard definitions given in the literature. First we consider the approach based on resolution of singularities as described in [La, Part III]. Then we explain how to recover Lipman's definition given in [Li]. Finally we show how

to understand asymptotic multiplier ideals of graded systems of ideals in terms of tree potentials.

3.1. Multiplier ideals through log-resolutions. The exposition follows [La, Part III]. See also [LW] and [TW].

A *log-resolution* of an ideal $I \subset R$ is a birational morphism $\pi : X \rightarrow \text{Spec } R$ such that X is non-singular and $I \cdot \mathcal{O}_X$ is locally principal: $I \cdot \mathcal{O}_X = \mathcal{O}_X(-F)$ for an effective divisor F , and $F + E$ has simple normal crossing singularities, where E is the exceptional divisor of π .

The existence of such a resolution in our setting is well known. In fact, π can be chosen as a composition of blowups at (closed) points. In the analytic case $R = \mathcal{O}_0$ the reader may think of $\text{Spec } R$ as a neighborhood of the origin in \mathbf{C}^2 .

Another ingredient in the definition is *the relative canonical divisor* of π . It is the unique divisor $K_{X/R}$, with support in E and whose class represents $J\pi$, the Jacobian determinant of π :

$$K_{X/R} = K_X - \pi^* K_{\text{Spec } R}.$$

Here K_X and $K_{\text{Spec } R}$ denote the sheaf of 2-differentials over X and $\text{Spec } R$, respectively.

We also denote by $\lfloor c \rfloor$ the round-down of a positive real number, i.e., the greatest integer less than or equal to c .

Definition 3.1. Let $I \subset R$ be an ideal, and c be a positive real number. Fix a log-resolution $\pi : X \rightarrow \text{Spec } R$ of I such that $I \cdot \mathcal{O}_X = \mathcal{O}_X(-F)$ with $F = \sum r_i E_i$, and write $K_{X/R} = \sum \beta_i E_i$. We define the *multiplier ideal* of I^c to be

$$(\star) \quad \mathcal{J}(I^c) := \{\psi \in R : \text{div}_{E_i}(\pi^* \psi) \geq \lfloor cr_i \rfloor - \beta_i \text{ for all } i\}.$$

Here div_{E_i} denotes the generic order of vanishing along E_i of a function on X .

This definition does not depend on the choice of log-resolution. In the analytic case $R = \mathcal{O}_0$, the definition says that a holomorphic germ ψ belongs to the multiplier ideal iff the pull-back of ψ vanishes to sufficiently high orders along the irreducible components of $\pi^{-1}(V)$, where $V = \bigcap_{\phi \in I} \phi^{-1}(0)$.

Proposition 3.2. For any ideal $I \subset R$ and any positive real number $c > 0$, the multiplier ideal $\mathcal{J}(I^c)$ as defined in Definition 3.1 coincides with the multiplier ideal of the tree transform $c g_I$ as defined in Definition 2.1.

Remark 3.3. Similarly, we may associate a multiplier ideal to an expression of the form $I_1^{c_1} \cdots I_n^{c_n}$ for ideals I_j and real numbers c_j . The proof of Proposition 3.2 is easily adapted to show that $J(I_1^{c_1} \cdots I_n^{c_n}) = J(\sum_j c_j g_{I_j})$.

Proof. We first need to translate the condition of vanishing of $\pi^* \psi$ along $E_i \subset X$ in terms of valuations. We shall denote this condition by $(\star)_i$. Note that div_{E_i} is a valuation with values in \mathbf{Z} . We have $r_i = \text{div}_{E_i}(\pi^* I)$ and $\beta_i = \text{div}_{E_i}(J\pi)$ where $J\pi$ is the Jacobian determinant of π .

First suppose E_i does not belong to the exceptional divisor of π . Its image by π is then an irreducible curve. We let $\nu_i \in \mathcal{V}$ be the associated curve valuation (see Section 1.2.2) and m_i its multiplicity. Denote by ρ_I and ρ_ψ the tree measures of I and ψ , respectively. It follows from Section 1.3 that $\text{div}_{E_i}(\pi^* I)$ is equal to $m_i^{-1} \rho_I\{\nu_i\}$. In the same way, $\text{div}_{E_i}(\pi^* \psi) = m_i^{-1} \rho_\psi\{\nu_i\}$. On the other hand, it is clear that $\beta_i = \text{div}_{E_i}(J\pi) = 0$. The condition $(\star)_i$ is hence equivalent to

$m_i^{-1}\rho_\psi\{\nu_i\} \geq \lfloor cm_i^{-1}\rho_I\{\nu_i\} \rfloor$. As the left-hand side is an integer, this amounts to $m_i^{-1}\rho_\psi\{\nu_i\} > cm_i^{-1}\rho_I\{\nu_i\} - 1$. We thus conclude that if $E_i \not\subset \pi^{-1}(0)$, then

$$(3.1) \quad (\star)_i \Leftrightarrow \rho_\psi\{\nu_i\} + m(\nu_i) > c\rho_I\{\nu_i\}.$$

Now suppose E_i is an irreducible component of $\pi^{-1}(0)$. Then $\pi_* \operatorname{div}_{E_i}(\psi) := \operatorname{div}_{E_i}(\pi^*\psi)$ defines a divisorial valuation with values in \mathbf{Z} . It is proportional to a unique normalized valuation $\nu_i \in \mathcal{V}$: $\pi_* \operatorname{div}_{E_i} = b_i\nu_i$ where $b_i = b(\nu_i)$ is the generic multiplicity of ν_i (see Section 1.2.9). The thinness $A(\nu_i)$ is given by a_i/b_i , where $a_i = a(\nu_i) = \operatorname{div}_{E_i}(J\pi) + 1$ (again by Section 1.2.9). We infer that if $E_i \subset \pi^{-1}(0)$, then

$$\begin{aligned} (\star)_i &\Leftrightarrow \operatorname{div}_{E_i}(\pi^*\psi) \geq \lfloor c \operatorname{div}_{E_i}(\pi^*I) \rfloor - \operatorname{div}_{E_i}(J\pi) \\ &\Leftrightarrow \operatorname{div}_{E_i}(\pi^*\psi) > c \operatorname{div}_{E_i}(\pi^*I) - \operatorname{div}_{E_i}(J\pi) - 1 \\ &\Leftrightarrow b_i\nu_i(\psi) > cb_i\nu_i(I) - a_i, \end{aligned}$$

i.e.,

$$(3.2) \quad (\star)_i \Leftrightarrow \nu_i(\psi) + A(\nu_i) > c g_I(\nu_i).$$

Now consider $\psi \in \mathcal{J}(c g_I)$. It follows immediately from Proposition 3.2 that condition $(\star)_i$ in (3.1), (3.2) holds for all i . Thus $\psi \in \mathcal{J}(I^c)$.

Conversely suppose $\psi \notin \mathcal{J}(c g_I)$. Again we use Proposition 2.4. This tells us that either (a) or (b) fails in that proposition. If (b) fails, then there exists a curve valuation ν such that $\rho_\psi\{\nu\} + m(\nu) \leq c\rho_I\{\nu\}$. This inequality implies that the element in R corresponding to ν is a factor of I . Hence ν is one of the curve valuations ν_i above. Thus $(\star)_i$ in (3.1) fails and $\psi \notin \mathcal{J}(I^c)$. On the other hand, if (a) fails, then there exists a divisorial valuation ν such that $\nu(\psi) + A(\nu) \leq c\nu(I)$. We may take a log-resolution π of I such that ν corresponds to an exceptional divisor of π . This can be achieved by further blowups of a given log-resolution of I . Thus $\nu = \nu_i$ for some i , so that $(\star)_i$ in (3.2) fails and $\psi \notin \mathcal{J}(I^c)$. \square

3.2. Lipman’s approach. The definition of Lipman⁶ in [Li] has the advantage of not depending on a log-resolution and hence makes sense in great generality.

We describe Lipman’s construction in our setting, namely for an equicharacteristic zero, two-dimensional, regular local ring R with algebraically closed residue field k . Write K for the fraction field of R .

We first need some definitions. For a (not necessarily centered) valuation μ on R , we let $R_\mu = \{\phi \in K : \mu(\phi) \geq 0\}$ be the valuation ring, $\mathfrak{m}_\mu = \{\phi \in K : \mu(\phi) > 0\}$ the (unique) maximal ideal in R_μ and $k_\mu := R_\mu/\mathfrak{m}_\mu$ the residue field. Note that k_μ is a field extension of k . A *prime divisor* is by definition a valuation on R whose residue field has transcendence degree $l - 1$ over k , where l is the height of $\mathfrak{m}_\mu \cap R$, i.e., the dimension of the center of μ in $\operatorname{Spec} R$. Let us describe explicitly in our situation what a prime divisor means. Two different cases may appear.

- (i) $l = 1$. Then the center of μ is a principal ideal generated by an irreducible element $\psi \in R$. In this case for any $\phi \in R$, we have $\mu(\phi) = \operatorname{div}_C(\phi)$ for some irreducible $C \subset \operatorname{Spec} R$.
- (ii) $l = 2$. Then μ is centered at the maximal ideal \mathfrak{m} of R . The transcendence degree of k_μ being 1 implies that μ is a divisorial valuation (see [FJ1, Proposition 1.12]).

⁶Lipman used the term *adjoint ideals* and defined them for ideals rather than formal powers of ideals.

Lipman’s construction uses the *Jacobian ideal* $J_{R_\mu/R}$, whose definition is purely algebraic. In our setting it is given as follows, following the two cases above:

- (i) When $l = 1$, the ideal $J_{R_\mu/R}$ is trivial.
- (ii) When $l = 2$, μ is a divisorial valuation. Fix a composition of blowups $\pi : X \rightarrow \text{Spec } R$ such that the center of μ in X is a divisor E . For a closed point $p \in E$, the ring $\mathcal{O}_{X,p}$ can be naturally viewed as a subring of R_μ by the isomorphism π_* between function fields of X and $\text{Spec } R$. Then the ideal $J_{R_\mu/R} \subset R_\mu$ is generated by the Jacobian determinant of π .

We can now set

Definition 3.4. Let $I \subset R$ be an ideal and let $c > 0$. The multiplier ideal of I^c is the ideal

$$\mathcal{J}(I^c) := \bigcap_{\mu} \{ \psi \in R : \mu(\psi) \geq \lfloor c\mu(I) \rfloor - \mu(J_{R_\mu/R}) \}$$

where the intersection is taken over all prime divisors μ of R .⁷

If I is an ideal and $c > 0$, then we may consider both the multiplier ideal of I^c above and the multiplier ideal of the tree potential $c g_I$. Both are defined in terms of valuations. Hence the following result is perhaps not so surprising.

Proposition 3.5. *For any ideal $I \subset R$ and any $c > 0$, the multiplier ideal $\mathcal{J}(I^c)$ in the sense of Definition 3.4 coincides with the multiplier ideal of the tree potential $c g_I$ as defined in Definition 2.1.*

Proof. Pick $\psi \in R$. First consider a curve valuation $\nu = \nu_C$ in \mathcal{V} . We denote the prime divisor associated to ν by div_C . Note that div_C is not centered at the maximal ideal $\mathfrak{m} \subset R$. We then have the sequence of equivalences

$$\begin{aligned} \text{div}_C(\psi) \geq \lfloor c \text{div}_C(I) \rfloor - \text{div}_C(J_{R_{\text{div}_C}/R}) &\Leftrightarrow \text{div}_C(\psi) > c \text{div}_C(I) - 1 \\ &\Leftrightarrow \rho_\psi\{\nu_C\} > c \rho_I\{\nu_C\} - m(C). \end{aligned}$$

Now consider a divisorial valuation $\nu \in \mathcal{V}$. Fix a composition of blowups π such that $b\nu = \pi_* \text{div}_E$ for some exceptional component E , b being the generic multiplicity of ν (see Section 1.2.9). Let $a - 1$ be the order of vanishing of the Jacobian determinant of π along E . By the discussion above,

$$\begin{aligned} \nu(\psi) \geq \lfloor c\nu(I) \rfloor - \nu(J_{R_\nu/R}) &\Leftrightarrow \text{div}_E(\pi^*\psi) \geq \lfloor c \text{div}_E(\pi^*I) \rfloor - \text{div}_E(J\pi) \\ &\Leftrightarrow \text{div}_E(\pi^*\psi) > c \text{div}_E(\pi^*I) - \text{div}_E(J\pi) - 1 \\ &\Leftrightarrow b\nu(\psi) > bc\nu(I) - a \\ &\Leftrightarrow \nu(\psi) > c g_I(\nu) - A(\nu). \end{aligned}$$

This concludes the proof in view of Proposition 2.4. □

3.3. Multiplier ideals of graded systems. Another type of multiplier ideals has been defined for graded systems of ideals. Here we show that these multiplier ideals can also be understood through tree potentials.

Let $I_\bullet = (I_k)_{k \geq 1}$ be a graded system of ideals. This means that I_k is an ideal in R for every k and that $I_k I_l \subset I_{k+l}$ for all k, l . Fix $c > 0$ rational. It is proved in [La] (see also below) that the sequence of ideals $(\mathcal{J}(I_k^{c/k}))_{k \geq 1}$ has a unique maximal element.

⁷Lipman’s original definition was for $c = 1$ in which case the “[\cdot]” can be omitted.

Definition 3.6. The *asymptotic multiplier ideal* $\mathcal{J}(I_\bullet^c)$ is the unique maximal element of the sequence $(\mathcal{J}(I_k^{c/k}))_{k \geq 1}$.

Definition 3.7. The *singularity exponent* (or *log-canonical threshold*) of I_\bullet is $c(I_\bullet) := \sup\{c > 0 : \mathcal{J}(I_\bullet^c) = R\}$. The *Arnold multiplicity* is $\lambda(I_\bullet) := c(I_\bullet)^{-1}$.

Let g_k be the tree transform of I_k . The condition $I_k I_l \subset I_{k+l}$ then implies $g_{k+l} \leq g_k + g_l$. It is then elementary that the sequence $k^{-1}g_k$ converges in \mathcal{P} (i.e., pointwise) to the tree potential $g := \inf k^{-1}g_k$. Moreover, the subsequence $2^{-j}g_{2^j}$ is decreasing, so by semicontinuity (Proposition 2.6) we have $\mathcal{J}(c2^{-j}g_{2^j}) = \mathcal{J}(ch)$ for $j \gg 0$. On the other hand, $\mathcal{J}(ck^{-1}g_k) = \mathcal{J}(I_k^{c/k})$ for all k by Proposition 3.2. This gives

Proposition 3.8. *Given a graded sequence I_\bullet of ideals there exists a tree potential $g = g_{I_\bullet}$ such that the asymptotic multiplier ideal $\mathcal{J}(I_\bullet^c)$ coincides with the multiplier ideal $\mathcal{J}(cg)$ for any $c > 0$. As a consequence, $c(I_\bullet) = c(g)$ and $\lambda(I_\bullet) = \lambda(g)$.*

Remark 3.9. It follows from Theorem 8.1 that $g = g_{I_\bullet}$ is uniquely determined by the properties stated in Proposition 3.8.

Remark 3.10. Not every tree potential g is associated to a graded system of ideals I_\bullet . An example is given by $g = \alpha(\mu \wedge \cdot)$, where μ is infinitely singular of infinite skewness. Indeed, $g(\mu) = \infty$, whereas $\mu(I) < \infty$ for every ideal I ; hence $g_{I_\bullet}(\mu) < \infty$ for any graded system I_\bullet . It would be interesting to characterize all tree potentials associated to graded systems of ideals. See [FJ2, Section 6.2] for related questions.

Example 3.11. If $I \subset R$ is a fixed ideal and $k_0 \in \mathbb{N}$, then $I_k := I^{k+k_0}$ defines a graded system of ideals. In this case $k^{-1}g_{I_k} = (1+k_0/k)g_I \rightarrow g_I$ so $\mathcal{J}(I_\bullet^c) = \mathcal{J}(I^c)$ independently of k_0 .

Example 3.12. If $\nu \in \mathcal{V}$, then $I_k := \{\phi \in R : \nu(\phi) \geq k\}$ defines a graded system of ideals. In this case the tree potential h in Proposition 3.8 satisfies

$$(3.3) \quad \Delta h = \alpha(\nu)^{-1}\nu.$$

Moreover, $c(I_\bullet) = \alpha(\nu)(1 + \alpha(\nu_1)^{-1})$, where ν_1 is the first element in the approximating sequence of ν . In particular, if $\alpha(\nu) = \infty$, then $g = 0$, and $c(I_\bullet) = \infty$.

Let us sketch a proof of (3.3). It is easy to see in general that $g(\mu) \geq h(\mu) := \alpha(\nu)^{-1}\alpha(\mu \wedge \nu)$. For the reverse inequality, suppose first that ν is a divisorial valuation of generic multiplicity b . Then the tree transform of $I_{b\alpha(\nu)}$ is precisely $\mu \rightarrow b\alpha(\mu \wedge \nu)$; see Section 1.3. Whence $g = h$ in this case. In general, take an increasing sequence of divisorial valuations ν_n tending to ν , and apply the preceding result.

4. MULTIPLIER IDEALS OF PSH FUNCTIONS

We now turn to multiplier ideals of psh functions. For this to make sense we work in the analytic case $R = \mathcal{O}_0$. Recall that if u is psh, then the *multiplier ideal* $\mathcal{J}(u)$ consists of the holomorphic germs $\psi \in R$ such that $|\psi|^2 e^{-2u}$ is locally integrable at the origin (see [N, DK]). The (complex) *singularity exponent* (or *log-canonical threshold*) of u at the origin is $c(u) = \sup\{c > 0 : e^{-2cu} \in L_{\text{loc}}^1\}$. The *Arnold multiplicity* is $\lambda(u) = c(u)^{-1}$. Our goal is to prove the following theorem.

Theorem 4.1. *If u is a psh function, then the multiplier ideal of u equals that of its tree transform. In other words, if ψ is a holomorphic germ, then*

$$(4.1) \quad \psi \in \mathcal{J}(u) \quad \text{iff} \quad \sup_{\nu \in \mathcal{V}_{\text{qm}}} \frac{\nu(u)}{\nu(\psi) + A(\nu)} < 1.$$

The proof makes essential use of the Demailly approximation technique in Section 1.4.4. In fact we have

Corollary 4.2. *If u_n are the Demailly approximants of u , then $\mathcal{J}(u_n) = \mathcal{J}(u)$ for $n \gg 1$.*

Proof. We have $0 \leq \nu(u) - \nu(u_n) \leq A(\nu)/n$ for all $\nu \in \mathcal{V}_{\text{qm}}$; hence

$$0 \leq \frac{\nu(u)}{\nu(\psi) + A(\nu)} - \frac{\nu(u_n)}{\nu(\psi) + A(\nu)} \leq \frac{1}{n},$$

which easily implies the corollary in view of (4.1). \square

We split the proof of the theorem into two parts: non-integrability and integrability. Set

$$\chi(\nu) = \chi_{u,\psi}(\nu) = \frac{\nu(u)}{\nu(\psi) + A(\nu)}.$$

By Lemma 2.11, χ attains its supremum on \mathcal{V} . We emphasize that this fact is really the key to the openness conjecture (Corollary 5.1). At any rate, Theorem 4.1 follows immediately from Proposition 4.3 and Proposition 4.4 below.

Proposition 4.3. *Let u and ψ be as in Theorem 4.1. Suppose there exists $\nu \in \mathcal{V}$ such that $\chi(\nu) \geq 1$. Then $|\psi|^2 e^{-2u}$ is not locally integrable at the origin. In fact, we then have $\text{Vol}\{|\psi|^2 e^{-2u} > R\} \gtrsim R^{-1}$ as $R \rightarrow \infty$.*

Proposition 4.4. *Let u and ψ be as in Theorem 4.1. Suppose that $\sup \chi < 1$. Then $|\psi|^2 e^{-2u}$ is locally integrable at the origin.*

Proof of Proposition 4.3. It suffices to show the estimate $\text{Vol}\{|\psi|^2 e^{-2u} > R\} \gtrsim R^{-1}$ as $R \rightarrow \infty$. By Lemma 2.11, we may assume ν is not infinitely singular.

First suppose ν is quasimonomial. Then $\nu(\psi) - \nu(u) \leq -A(\nu) < 0$. We use the analysis in Section 1.4.2. Write $\nu = \nu_{\phi,t}$ and pick a coordinate x transverse to ϕ . Let $\mathcal{A} = \mathcal{A}_{\phi,t,x,C}(r)$ be a characteristic region. Then $\text{Vol}(\mathcal{A}) \gtrsim r^{2A(\nu)}$. Moreover, $u(p) \leq \nu(u) \log \|p\| r + O(1)$ and $\log |\psi| \geq \nu(\psi) \log \|p\| + O(1)$, so

$$\log |\psi| - u \geq (\nu(\psi) - \nu(u)) \log \|p\| + O(1) \geq -A(\nu) \log r + O(1)$$

in \mathcal{A} . The desired estimate follows by choosing $R \sim r^{-2A(\nu)}$.

Now suppose $\nu = \nu_\phi$ is a curve valuation. Then ϕ is a holomorphic germ. By Section 1.4.5, we have $u = c \log |\phi| + u'$ with $c = \rho_u\{\nu_\phi\}/m(\phi)$ and u' psh. Similarly, $\psi = \phi^a \psi'$, where $a \geq 0$ and ϕ does not divide ψ' . Then $1 \leq \chi(\nu_\phi) = c/(a+1)$, so that $c \geq a+1$. Since u' is bounded from above we have

$$|\psi|^2 e^{-2u} \gtrsim |\phi|^{2(a-c)} |\psi'|^2 \gtrsim \frac{|\psi'|^2}{|\phi|^2}$$

near the origin. Pick a composition of blowups π , and a smooth point $p \in \pi^{-1}(0)$ such that the strict transform of $\phi^{-1}(0)$ is smooth and intersects the exceptional divisor transversely at p , and such that the strict transform of $(\psi')^{-1}(0)$ does not contain p . This can be done by successively blowing up the intersection point of the

strict transform of $\phi^{-1}(0)$ and the exceptional divisor. Fix coordinates (z, w) at p such that $\pi^{-1}(0) = \{z = 0\}$ and $\phi^{-1}(0) = \{zw = 0\}$. We may assume $\pi^*\phi = z^k w$, $J\pi = z^l$ and $\pi^*\psi' = z^m \tilde{\psi}$, where $k, l, m \in \mathbf{N}^*$ and $\tilde{\psi}(0) \neq 0$. Fix $r_0 > 0$ small, and consider $0 < r \ll r_0$. Set $\Omega(r) := \pi\{|z| < r_0, |w| < r\}$. By the change of variables formula, $\text{Vol}\Omega(r) \gtrsim r^2$. If $p = \pi(z, w) \in \Omega(r)$, then

$$|\psi(p)|^2 e^{-2u(p)} \gtrsim \frac{|\pi^*\psi'(z, w)|^2}{|\pi^*\phi(z, w)|^2} \gtrsim |w|^{-2} \geq r^{-2}.$$

The desired estimate follows by choosing $R \sim r^{-2}$. □

Proof of Proposition 4.4. First suppose u has logarithmic singularities, say $u = \frac{c}{2} \log \sum_{i=1}^k |\phi_i|^2$ for $\phi_i \in R$ and $c > 0$. The tree transform of u then coincides with the tree transform of I^c , where I is the ideal generated by the ϕ_i 's. Integrability can now be proved as in [DK, Proposition 1.7], using the results of Section 3.

This goes as follows. Let $\pi : X \rightarrow (\mathbf{C}^2, 0)$ be a log-resolution of the ideal I . Thus the total transform of the curve $V = \{\prod_l \phi_l = 0\}$ is a union of smooth components with normal crossing singularities. By the change of variables formula, the function $\Phi^2 = |\psi|^2 \exp(-2u)$ is locally integrable iff $(\pi^*\Phi \cdot |J\pi|)^2$ is locally integrable at any point $p \in \pi^{-1}(0)$.

Let $\{E_i\}$ be the set of irreducible components of $\pi^{-1}(V)$. For any i , we let β_i, r_i and γ_i be the order of vanishing along E_i of $J\pi, \pi^*I$ and $\pi^*\psi$, respectively. As in Section 3 the condition $\sup \chi < 1$ implies that $\delta_i := \gamma_i + \beta_i - \lfloor cr_i \rfloor \geq 0$; hence $\delta_i > -1$ for all i .

Now pick $p \in \pi^{-1}(0)$ and local coordinates (z, w) at p such that $\pi^{-1}V \subset \{zw = 0\}$. The calculations above give $\pi^*\Phi \cdot |J\pi| \gtrsim |z|^\delta |w|^{\delta'}$ where $\delta, \delta' > -1$. Thus $(\pi^*\Phi \cdot |J\pi|)^2$ is locally integrable at p , completing the proof when u has logarithmic singularities.

We now consider the case of an arbitrary psh function. As in the proof of Theorem 4.2 in [DK] we use Demailly approximation to reduce to the preceding case. By Section 1.4.4 there exists a small neighborhood B' of the origin, and, for each $n > 0$, finitely many holomorphic functions $\{g_{nl}\}_{l=1}^k$ with $\int_{B'} |g_{nl}|^2 \exp(-2nu) \leq 1$, such that the psh function $u_n = (2n)^{-1} \log \sum |g_{nl}|^2$ (which has logarithmic singularities) satisfies $|\nu(u) - \nu(u_n)| \leq n^{-1} A(\nu)$ for any $\nu \in \mathcal{V}_{\text{qm}}$. Thus Hölder's inequality with $p = n$ and $q = n/(n - 1)$ gives

$$\begin{aligned} \int_{B'} |\psi|^2 \exp(-2u) &= \int_{B'} \left(\sum_1^k |g_{nl}|^2 \exp(-2pu) \right)^{1/p} \left(|\psi|^{2q} \left(\sum_1^k |g_{nl}|^2 \right)^{-q/p} \right)^{1/q} \\ &\leq k^{1/p} \left(\int_{B'} |\psi|^{2q} \left(\sum_1^k |g_{nl}|^2 \right)^{-q/p} \right)^{1/q} = k^{1/p} \left(\int_{B'} |\psi|^{2q} \exp(-2qu_n) \right)^{1/q}. \end{aligned}$$

For any quasimonomial valuation ν , we have

$$|\chi_{u,\psi}(\nu) - \chi_{qu_n,\psi^q}(\nu)| = \left| \frac{\nu(u)}{\nu(\psi) + A(\nu)} - \frac{q\nu(u_n)}{q\nu(\psi) + A(\nu)} \right| \lesssim \frac{1}{n}.$$

By assumption $\sup \chi_{u,\psi} < 1$; hence $\sup \chi_{qu_n,\psi^q} < 1$ for large n . By the preceding argument, $|\psi|^{2q} \exp(-2qu_n)$ is locally integrable, thus so is $|\psi|^2 \exp(-2u)$. □

5. SINGULARITY EXPONENTS AND KISELMAN NUMBERS

We now apply our machinery to study the singularities of psh functions.

Theorem 4.1 immediately implies that the complex singularity exponent $c(u)$ and Arnold multiplicity $\lambda(u)$ of a psh function u equal those of its tree transform g_u as defined by (2.2). In view of Remark 2.3 and Lemma 2.11 we then have

Corollary 5.1. *For any psh function u we have*

$$(5.1) \quad \lambda(u) = \sup_{\nu \in \mathcal{V}_{\text{qm}}} \frac{\nu(u)}{A(\nu)} \quad \text{and} \quad c(u) = \inf_{\nu \in \mathcal{V}_{\text{qm}}} \frac{A(\nu)}{\nu(u)},$$

and the supremum and infimum are attained at a valuation of multiplicity one.

Further, if $c = c(u)$, then $\exp(-2cu)$ is not locally integrable at the origin. In fact, $\text{Vol}\{u < \log r\} \gtrsim r^{2c(u)}$ as $r \rightarrow 0$.

Remark 5.2. This corollary contains an affirmative answer to the openness conjecture for psh functions in dimension 2; see [DK, Remarks 5.3 and 4.4].

Example 5.3. If $\psi = y^n + x^m$ with $m < n$, m, n relatively prime, and $u = \log |\psi|$, then the supremum in (5.1) must be attained at a valuation of the form $\nu_{y,t}$, $1 \leq t \leq \infty$. Thus $\lambda(\log |\psi|) = \sup_t \max\{nt, m\}/(1+t) = nm/(n+m)$.

Let us rephrase Corollary 5.1 in terms of Kiselman numbers, using Section 1.4.1. If $m(\nu) = 1$, then $\nu = \nu_{y,t}$ where (x, y) are local coordinates and $t \geq 1$. Moreover, $\nu(u) = \nu_{t,1}^{x,y}(u)$ and $A(\nu) = 1+t$. Hence $\nu(u)/A(\nu) = \nu_{t/(1+t), 1/(1+t)}^{x,y}(u)$.

Corollary 5.4. *The Arnold multiplicity of u is the supremum of all Kiselman numbers $\nu_{a,b}^{x,y}(u)$ over all choices of local coordinates (x, y) and all choices of weights (a, b) with $a + b = 1$.*

By fixing the local coordinates (x, y) and letting the weights vary we obtain a lower bound for the Arnold multiplicity proved by Kiselman [Ki2] (in any dimension). We can also bound it in terms of the Lelong number $\nu^L(u)$:

Corollary 5.5. *The Arnold multiplicity $\lambda(u)$ of u satisfies*

- (i) $\frac{1}{2}\nu^L(u) \leq \lambda(u) \leq \nu^L(u)$;
- (ii) $\lambda(u) = \nu^L(u)$ iff $dd^c u$ is the current of integration on a smooth curve plus a current with zero Lelong number;
- (iii) if $\lambda(u) = \frac{1}{2}\nu^L(u)$, then the Lelong number of the strict transform of u under a single blowup π of the origin is at most $\frac{1}{2}\nu^L(u)$ at any point on the exceptional divisor $\pi^{-1}(0)$.

Remark 5.6. The bounds in (i) are due (in any dimension) to Skoda [S]. The characterization in (ii) sharpens a recent result by Blel and Mimouni [BM] (see also [Mi]), who proved that if u has Lelong number one, then $\exp(-2u)$ is locally integrable unless $dd^c u$ puts mass on an analytic curve. The implication in (iii) seems to be new.

Remark 5.7. The converse to (iii) is false, as is shown by the example

$$u = \log \max\{|y|, |x|^{1+\varepsilon}\},$$

where $0 < \varepsilon < 1$. Here $\nu^L(u) = 1$, $\lambda(u) = (1+\varepsilon)/(2+\varepsilon) > 1/2$, but the strict transform of u has Lelong number zero at all points of $\pi^{-1}(0)$, but one, at which the Lelong number is ε .

Proof of Corollary 5.5. All of this is a consequence of Theorem 4.1 and Proposition 2.7. Indeed (i) follows immediately, as does (iii) in view of Section 1.4.3. As for (ii) we get that the tree measure ρ_u is a point mass at a curve valuation, which must be associated to a smooth analytic curve $\{\phi = 0\}$. Siu’s Theorem now yields $u = \log |\phi| + u'$ with u' psh. Clearly u' has zero Lelong number. \square

6. THE ASCENDING CHAIN CONDITION

We now wish to give a new proof of a result describing the structure of the set of complex singularity exponents of holomorphic functions.

As our approach is algebraic, we work in the general case; that is, R is an equicharacteristic zero, two-dimensional, regular local ring with algebraically closed residue field. If $\psi \in R$, let $c(\psi) = c(g_\psi)$ and $\lambda(\psi) = c(\psi)^{-1}$ be the singularity exponent and Arnold multiplicity of ψ . Here g_ψ is the tree potential of ψ . Thus $\lambda(\psi) = \sup_{\nu_{\text{qm}}} \chi_\psi(\nu)$, where $\chi_\psi(\nu) = \nu(\psi)/A(\nu)$. In the analytic case $R = \mathcal{O}_0$ we have $c(\psi) := c(\log |\psi|)$ and $\lambda(\psi) = \lambda(\log |\psi|)$.

To motivate the result, first consider $\psi \in \mathfrak{m}$ irreducible and let $(\nu_i)_0^g$ be the approximating sequence of ν_ψ . It follows from Corollary 5.1 that the supremum of χ_ψ is attained at ν_1 . This gives $c(\psi) = \frac{1}{\nu_1(\psi)} + \frac{1}{m(\psi)}$. By Section 1.2.7, $\nu_1(\psi)$ is an integer, so that $c(\psi) \in \frac{1}{\mathbf{N}^*} + \frac{1}{\mathbf{N}^*}$, a result that seems to have first been proved by Igusa [I]; see also [Ku]. When ψ is no longer assumed to be irreducible, $c(\psi) \notin \frac{1}{\mathbf{N}^*} + \frac{1}{\mathbf{N}^*}$ in general (see [PS] for a counterexample). Nevertheless we have

Theorem 6.1 ([S, 1992], [Ku, 1999], [PS, 2000]). *The set $\mathbf{c} = \{c(\psi) : \psi \in \mathfrak{m}\}$ satisfies the ascending chain condition (ACC): any increasing sequence in \mathbf{c} is eventually stationary.*

Further, the limit points of \mathbf{c} are 0 and the rational numbers $1/a$ for $a \geq 1$.

Proof. We will show that $\boldsymbol{\lambda} := \{\lambda(\psi) ; \psi \in \mathfrak{m}\}$ satisfies the *decreasing* chain condition (DCC) and that its limit points are $\mathbf{N} \cup \{\infty\}$. Our proof mainly consists of a careful study of the functions χ_ψ on \mathcal{V}_{qm} , for $\psi \in \mathfrak{m}$.

Fix ψ and let $\psi = \prod \phi_k^{\alpha_k}$ be the decomposition into irreducible factors. For any k we define the valuation $\nu_k = \max\{\nu \in \mathcal{V} : m(\nu) = 1, \nu \leq \nu_{\phi_k}\}$. This is either ν_{ϕ_k} or a divisorial valuation. Write $\nu_{kl} = \nu_k \wedge \nu_l$ and $\alpha_{kl} = \alpha(\nu_{kl})$.

Fact 1. For all k, l , either $\alpha_{kl}m_k$ or $\alpha_{kl}m_l$ is an integer where $m_k = m(\phi_k)$.

We first note that $m_k\alpha_k$ is always an integer (if $\alpha_k < \infty$). For this we refer repeatedly to Section 1.2.7. As $\nu_{\phi_k} > \nu_k$, the generic multiplicity b_k of ν_k divides m_k . Thus $\alpha(\nu_k)b(\nu_k) = \alpha_k b_k \in \mathbf{N}$; hence $\alpha_k m_k \in \mathbf{N}$. This shows Fact 1 when $\nu_{kl} = \nu_k$ or ν_l . If this is not the case, then $\nu_{kl} < \nu_k, \nu_{kl} < \nu_l$. As $m(\nu_k) = m(\nu_l) = 1$, $b(\nu_{kl}) = 1$ and again $\alpha_{kl} \in \mathbf{N}$.

Fact 2. The supremum of χ_ψ is attained at a valuation ν_{kl} for some k, l .

After unwinding definitions, Lemma 2.11 (i) and (ii) imply that the supremum of χ_ψ is attained either at $\nu_{\mathfrak{m}}$ or at one of the ν_{kl} . If $\nu_{kl} = \nu_{\mathfrak{m}}$ for some k and l , the proof is complete, so assume that $\nu_{kl} > \nu_{\mathfrak{m}}$ for all k, l . Then there exists a unique tangent vector \vec{v} at $\nu_{\mathfrak{m}}$ such that $\nu_{kl} \in U(\vec{v})$ for all k, l . It is then straightforward to verify that $D_{\vec{v}}\chi_\psi > 0$. Hence the maximum is not attained at $\nu_{\mathfrak{m}}$ and we are done.

Fact 3. For any k, l , we have $\nu_{kl}(\psi) \in \frac{\mathbf{N}^*}{m_k} \cup \frac{\mathbf{N}^*}{m_l}$.

If $\nu_{\phi_i} \geq \nu_{kl}$, we have $\nu_{kl}(\phi_i) = \alpha_{kl}m_i$, which belongs to $\mathbf{N}^*/m_k \cup \mathbf{N}^*/m_l$ by Fact 1. If $\nu_{\phi_i} \not\geq \nu_{kl}$, then $\nu_{kl}(\phi_i) = \nu_{ik}(\phi_i)$. Then either $\nu_{ik} = \nu_i$, and $\nu_{kl}(\phi_i) \in \mathbf{N}^*$; or $\nu_{ik} < \nu_i$, $\nu_{ik} < \nu_{kl}$ and $\alpha_{ik} \in \mathbf{N}^*$ so that $\nu_{kl}(\phi_i) \in \mathbf{N}^*$.

Fact 4. Suppose $\lambda(\psi) = \chi_\psi(\nu_{kl})$. Set $I = \{i : \nu_{\phi_i} \geq \nu_{kl}\}$ and let J be its complement. Then

$$(6.1) \quad \sum_J a_j \nu_{kl}(\phi_j) \leq \sum_I a_i m_i.$$

This inequality follows from the assumption $\lambda(\psi) = \chi_\psi(\nu_{kl})$. For any t in an interval $T = (\alpha_{kl} - \eta, \alpha_{kl})$ for η sufficiently small, we have

$$\chi(\nu_{\psi_i, t}) = \frac{t(\sum_I a_i m_i) + \sum_J a_j \nu_{kl}(\phi_j)}{1 + t}.$$

As the supremum of χ is attained when $t = \alpha_{kl}$, we infer that $\chi' \geq 0$ on T . A direct computation then shows (6.1).

We are now able to prove the theorem. First note that it suffices to prove that $\lambda \cap (0, C)$ satisfies the DCC for any $C > 0$. So pick a sequence $\psi_n \in \mathfrak{m}$ such that $\lambda(\psi_n)$ is decreasing and bounded from above by C . Since $\lambda(\psi) = \sup \nu(\psi)/A(\nu) \geq m(\psi)/2$, we can bound the multiplicity $m(\psi_n) \leq 2C$. For any n , we introduce ϕ_k^n , α_{kl}^n , I^n , J^n as above. To simplify notation we drop the superscript n . We also denote by $\mathcal{N} = \{\frac{p}{q} : p, q \in \mathbf{N}^*, q \leq 2C\}$. This is a discrete semigroup. Fact 2 implies that we can write

$$\begin{aligned} \lambda(\psi_n) &= \frac{(\sum_I a_i m_i)\alpha_{kl} + \sum_J a_j \nu_{kl}(\phi_j)}{1 + \alpha_{kl}} \\ &= \left(\sum_I a_i m_i\right) - \frac{1}{1 + \alpha_{kl}} \left[\sum_I a_i m_i - \sum_J a_j \nu_{kl}(\phi_j)\right]. \end{aligned}$$

We have $\sum_I a_i m_i \leq m(\psi_n) \leq 2C$, so we may assume that $\sum_I a_i m_i$ is constant, say equal to D , for all n . Fact 3 implies that $\sum_J a_j \nu_{kl}(\phi_j)$ belongs to the discrete set \mathcal{N} . It is also bounded by D by Fact 4, so we may assume it is constant. Thus $\lambda(\psi_n) = D - \frac{E}{1 + \alpha_{kl}}$ for some E . Now $\lambda(\psi_n)$ is increasing; hence α_{kl} is decreasing. Finally Fact 1 shows that $\alpha_{kl} \in \mathcal{N}$ and therefore is constant for large n . This proves that λ satisfies the DCC.

In order to find the limit points of λ , suppose $\lambda(\psi_n) \rightarrow \lambda \in \lambda$, $\lambda \neq \infty$. By extracting a subsequence and using the DCC we may assume that $\lambda(\psi_n)$ is increasing. The argument above applies again: for n large, we have $\lambda(\psi_n) = D - E(1 + \alpha_{kl})^{-1}$ for some constants $D, E > 0$, where D is an integer. If $\lambda(\psi_n)$ is not stationary, α_{kl} increases to infinity, so that $\lambda = D \in \mathbf{N}$. This shows that the limit points of λ are included in $\mathbf{N} \cup \{\infty\}$. Conversely, we have $\lambda(y^m + x^n) = mn/(m + n)$. Letting $m \rightarrow \infty$ while keeping n constant gives $n \in \lambda$. Letting $m, n \rightarrow \infty$ gives $\infty \in \lambda$. This concludes the proof. \square

7. IDEALS AS MULTIPLIER IDEALS

Proposition 2.5 asserts that the multiplier ideal associated to any tree potential is integrally closed. Our aim is to prove the converse statement, thereby giving a new proof of a theorem by Lipman and Watanabe [LW].

Theorem 7.1. *If J is an integrally closed ideal of R , then $J = \mathcal{J}(g)$ for some tree potential g .*

Remark 7.2. Our proof is constructive and generates a tree potential of the form $g = c g_I$, where I is an integrally closed ideal and $c > 0$. The choices of I and c are not unique.

Remark 7.3. Lipman and Watanabe work on more general rings. They also prove that if $J = I_\nu$ with ν divisorial, then there exists an ideal I with $\mathcal{J}(I) = J$ iff $b(\nu) = 1$. More generally it would be interesting to know what integrally closed ideals are of the form $\mathcal{J}(I)$ for an ideal I .

Proof. Any integrally closed ideal in R is the product of a principal ideal and a primary ideal. In view of Proposition 2.10 we may hence assume that J is primary.

Let us recall some notation from the analysis of tree transforms of ideals in Section 1.3. The tree measure of J on \mathcal{V} is of the form $\rho_J = \sum_1^r n_i b_i \nu_i$. Here $n_i \in \mathbf{N}$ and ν_i is divisorial with generic multiplicity b_i . Write $\alpha_i = \alpha(\nu_i)$ and $A_i = A(\nu_i)$. Set $m = m(J) = \sum_i n_i b_i$ and let \mathcal{T}_J be the support of the tree potential g_J , i.e., the smallest subtree containing ν_m and all the ν_i .




FIGURE 3. Construction of a tree potential with a predetermined multiplier ideal I . The Zariski factorization of I is $I = I_1 I_2 I_3^2 I_4^3$ and ν_i is the Rees valuation of I_i , $1 \leq i \leq 4$. The curves C_{ij} are chosen as curvettes of ν_i and the valuations ν_{ij} are well chosen on the segments $[\nu_i, \nu_{C_{ij}}]$. See the proof of Theorem 7.1.

Pick $\psi_{ij} \in \mathfrak{m}$ irreducible for $1 \leq i \leq r$, $1 \leq j \leq n_i$ such that $m(\psi_{ij}) = b_i$, $\nu_{\psi_{ij}} > \nu_i$ and the $\nu_{\psi_{ij}}$ represent n_i distinct tangent vectors at ν_i . The curves $C_{ij} := \{\psi_{ij} = 0\}$ are then curvettes of ν_i . Define ν_{ij} to be the unique quasimonomial valuation in $]\nu_i, \nu_{\psi_{ij}}[$ whose thinness $A_{ij} = A(\nu_{ij})$ is given by $A_{ij} - A_i = R_i$ for a constant $R_i > 0$ depending only on i to be fixed shortly. See Figure 3. Write $\alpha_{ij} = \alpha(\nu_{ij})$ so that $b_i(\alpha_{ij} - \alpha_i) = R_i$. For a given $\varepsilon > 0$, we define the tree potential g by

$$\Delta g = (1 + \varepsilon) \sum_{i \leq r, j \leq n_i} b_i \nu_{ij}.$$

Let \mathcal{T}_g be the support of g (i.e., \mathcal{T}_J with the segments $[\nu_i, \nu_{ij}]$ attached) and let $c(J) = \inf_\nu A(\nu)/\nu(J)$ be the complex singularity exponent of the ideal J (see Corollary 5.1). In the sequel, we fix $\varepsilon < m(J)^{-1}$. With this choice, we now show that we can pick $R_i > 0$, for $i \leq r$, so that

$$(7.1) \quad \nu_i(J) + A_{ij} - b_i^{-1} < g(\nu_{ij}) < \nu_i(J) + A_{ij}.$$

To see this, notice that

$$(7.2) \quad g(\nu_{ij}) = (1 + \varepsilon)(\nu_i(J) + R_i).$$

Hence $g(\nu_{ij}) - \nu_i(J) - A_{ij} = \varepsilon\nu_i(J) - A_i + \varepsilon R_i =: S_i$. We want to pick $R_i > 0$ so that $S_i \in (-b_i^{-1}, 0)$. Clearly S_i is an affine function of R_i and $S_i \rightarrow \infty$ as $R_i \rightarrow \infty$. Hence it suffices to show that $S_i < 0$ when $R_i = 0$; i.e., that $\varepsilon\nu_i(J) < A_i$. But the tree potential properties of g_J give $\nu_i(J) \leq m\alpha_i$. Hence $\varepsilon\nu_i(J) < \alpha_i < A_i$.

Finally, we note that g is proportional to the tree transform of some integrally closed ideal. Indeed multiplying $\sum b_i\nu_{ij}$ by the least common multiple of all the generic multiplicities $b(\nu_{ij})$ we obtain the tree measure ρ_I of some ideal I . We can thus write $g = c g_I$ for some $c > 0$ and some integrally closed ideal I .

We claim that $\mathcal{J}(g) = J$. Let us first prove that $\mathcal{J}(g) \supset J$. By Proposition 2.4 (ii) it suffices to show that $\nu(J) + A(\nu) > g(\nu)$ for $\nu \in \mathcal{T}_J$ and $\nu = \nu_{ij}$. The latter case is taken care of by (3.1). As for the former, if $\nu \in \mathcal{T}_J$, then $g(\nu) = (1 + \varepsilon)\nu(J)$. Moreover, $\nu(J) \leq m(J)\alpha(\nu) < m(J)A(\nu)$, so

$$\frac{g(\nu)}{\nu(J) + A(\nu)} = \frac{1 + \varepsilon}{1 + A(\nu)/\nu(J)} < \frac{1 + \varepsilon}{1 + m(J)^{-1}} < 1.$$

We now complete the proof by proving that $\mathcal{J}(g) \subset J$. Fix $\psi \in \mathcal{J}(g)$. Then

$$(7.3) \quad \nu(\psi) > g(\nu) - A(\nu)$$

for all $\nu \in \mathcal{V}_{\text{qm}}$. By Section 1.3 we must show that $\nu_i(\psi) \geq \nu_i(J)$ for all $i \leq r$.

For $i \leq r$ and any $j \leq n_i$, (7.1) and (7.3) imply that

$$\nu_{ij}(\psi) > g(\nu_{ij}) - A_{ij} > \nu_i(J) - b_i^{-1}.$$

We let $\mathcal{N} \subset \{1, \dots, r\}$ be the set of i for which there exists j such that ψ has no irreducible factor $\tilde{\phi}$ with $\nu_{\tilde{\phi}} \wedge \nu_{ij} > \nu_i$. After relabeling we have $\mathcal{N} = \{1, \dots, k\}$.

For $1 \leq i \leq k$, we infer immediately that $\nu_i(\psi) = \nu_{ij}(\psi) > \nu_i(J) - b_i^{-1}$. Since ν_i is divisorial we have $\nu_i(\psi) \in b_i^{-1}\mathbf{N}$ (see Section 1.2.9); hence $\nu_i(\psi) \geq \nu_i(J)$.

If $k = r$, the proof is complete. Thus assume $k < r$ and pick i with $k < i \leq r$. For any $j \leq n_i$, ψ admits an irreducible factor ϕ_{ij} with $\nu_{\phi_{ij}} \wedge \nu_{ij} > \nu_i$. The multiplicity of ϕ_{ij} is necessarily a multiple of b_i , say $p_{ij}b_i$.

We now replace ϕ_{ij} by $\psi_{ij}^{p_{ij}}$, where ψ_{ij} are defined above. By doing this replacement whenever $k < i \leq r$ and $1 \leq j \leq n_i$ we obtain from ψ a new $\hat{\phi} \in \mathfrak{m}$, for which $\nu(\hat{\phi}) = \nu(\psi)$ for all $\nu \in \mathcal{T}_J$.

We can then write $\hat{\phi} = \phi' \prod_{k < i \leq r} \psi_i$ where $\psi_i = \prod_j \psi_{ij}$ and $\phi' \in \mathfrak{m}$. Notice that if $\nu \in \mathcal{T}_J$, then $\nu(\psi_i) = b_i \nu \cdot \nu_i$ and $\nu(J) = \sum_1^r n_i b_i \nu \cdot \nu_i$. Thus, for $\nu \in \mathcal{T}_J$ we have

$$(7.4) \quad \nu(\psi) - \nu(J) = \nu(\hat{\phi}) - \nu(J) = \nu(\phi') - \sum_1^k n_i b_i \nu \cdot \nu_i = \nu(\phi') - \nu(J'),$$

where J' is the integrally closed ideal associated to the measure $\sum_1^k n_i b_i \nu_i$ (see Section 1.3). We have seen that $\nu_i(\phi') - \nu_i(J') = \nu_i(\psi) - \nu_i(J) > -b_i^{-1}$ for all $i \leq k$. As above we conclude that in fact $\nu_i(\phi') - \nu_i(J') \geq 0$. As the support of $\rho_{J'}$ is included in the ν_i 's, we infer $\nu(\phi') \geq \nu(J')$ for all $\nu \in \mathcal{V}_{\text{qm}}$. Thus (7.4) gives $\nu(\psi) \geq \nu(J)$ for all $\nu \in \mathcal{T}_J$. Hence $\psi \in J$, which concludes the proof. \square

Remark 7.4. It follows from the proof that we may in fact choose the numbers R_i of the form $R_i = n/b_i$ for $n \in \mathbf{N}^*$. One can then check that our construction coincides with that of Lipman and Watanabe [LW].

8. EQUISINGULARITY

In general we may refer to two objects as *equisingular* if they have the same tree transform. Two ideals are equisingular iff they have the same integral closure. (This is classical, but see [FJ1, Theorem 8.12].) Two psh functions are equisingular iff their pull-backs by any composition of blowups have the same Lelong number at all points on the exceptional divisor; see [FJ2, Proposition 6.2].

Our objective now is to show that equisingularity can be detected at the level of multiplier ideals. This is achieved through the following theorem, which allows us to recover a tree potential from the multiplier ideals of all its constant multiples.

Theorem 8.1. *Suppose h is a potential. For $t > 0$, let h_t be the tree transform of the ideal $\mathcal{J}(th)$. Then $h - t^{-1}A \leq t^{-1}h_t \leq h$, where A denotes thinness. In particular $t^{-1}h_t$ converges to h in \mathcal{P} as $t \rightarrow \infty$.*

Remark 8.2. The sequence $(n^{-1}h_n)_{n=1}^\infty$ can be thought of as an analogue for a tree potential of the Demailly approximating sequence of a psh function.

Corollary 8.3. *Let $h_1, h_2 \in \mathcal{P}$. Then $h_1 = h_2$ iff $\mathcal{J}(th_1) = \mathcal{J}(th_2)$ for all $t \geq 0$.*

Using the results of Sections 3 and 4 we immediately infer two corollaries.

Corollary 8.4. *Let I_1, I_2 be integrally closed ideals. Then $I_1 = I_2$ iff $\mathcal{J}(I_1^n) = \mathcal{J}(I_2^n)$ for all $n \in \mathbf{N}^*$.*

Corollary 8.5. *Let u_1, u_2 be psh functions. Then $\mathcal{J}(tu_1) = \mathcal{J}(tu_2)$ for all $t \geq 0$ iff the tree transforms of u_1 and u_2 coincide.*

Remark 8.6. Corollary 8.5 can be deduced directly from Ohsawa-Takegoshi's theorem. Similarly, Corollary 8.4 is a consequence of Skoda's theorem, which implies $\mathcal{J}(I^n) = I^{n-1}\mathcal{J}(I)$.

Remark 8.7. It is not true that if I_\bullet, J_\bullet are graded systems of ideals with $\mathcal{J}(I_\bullet^c) = \mathcal{J}(J_\bullet^c)$ for all $c > 0$, then $I_\bullet = J_\bullet$. See Example 3.11.

We give a proof of Theorem 8.1 based on the subadditivity property for multiplier ideals proved in [DEL].

Proof of Theorem 8.1. One inequality is elementary: for any $\phi \in \mathcal{J}(th)$ we have $\nu(\phi) > th(\nu) - A(\nu)$; hence $t^{-1}h_t \geq h - A/t$. The other inequality, on the other hand, is quite deep, but it is an immediate consequence of the following result, whose proof is given below. \square

Lemma 8.8. *For any tree potential h , the tree transform of $\mathcal{J}(h)$ is dominated by h . In other words, $g_{\mathcal{J}(h)} \leq h$.*

Remark 8.9. Explicitly, this lemma asserts that for any tree potential h , and any $\nu \in \mathcal{V}_{\text{qm}}$, one can find $\phi \in \mathcal{J}(h)$ so that $\nu(\phi) \leq h(\nu)$. It would be interesting to have a more direct construction of such an element ϕ .

Proof of Lemma 8.8. Let $\rho = \rho_h$ be the measure associated to h . We will gradually build the proof using successively more general measures ρ .

First suppose $h = g_I$ for an integrally closed ideal I . If $\psi \in I$, then $\nu(\psi) \geq \nu(I)$ for all $\nu \in \mathcal{V}$. Hence $\nu(\psi) + A(\nu) > \nu(I)$ for all $\nu \in \mathcal{V}_{\text{qm}}$ and $\rho_\psi\{\nu\} + m(\nu) > \nu_I\{\nu\}$ for all curve valuations ν . By Proposition 2.4 this implies $\psi \in \mathcal{J}(I)$. Hence $I \subset \mathcal{J}(I)$, so Lemma 8.8 is immediate in this case.

We now consider the more general case $h = c g_I$, where I is an integrally closed ideal and $c > 0$ is rational. By the subadditivity theorem of Demailly-Ein-Lazarsfeld [DEL] we have $\mathcal{J}(I^{c^q}) \subset \mathcal{J}(I^c)^q$ for any q . (Our situation is strictly speaking not contained in [DEL]; a more general statement than what we need can be found in [TW].) Therefore $I^{c^q} \subset \mathcal{J}(I^c)^q$ if $c^q \in \mathbf{N}$. Hence $c^q g_I \geq q g_{\mathcal{J}(I^c)}$, which gives $h \geq g_{\mathcal{J}(h)}$.

The next more general case is when the support of ρ is included in a finite tree \mathcal{T} whose ends are quasimonomial valuations. We approximate ρ by atomic measures $\rho_n = \sum_i c_i^n \nu_i^n$. We can suppose ν_i^n are all divisorial valuations, and c_i^n are rational numbers. We can also impose that $h_n = g_{\rho_n}$ decreases towards $g_\rho = h$. To do so, note that by linearity, we may take the support of ρ to be included in a segment $T = [\nu_m, \nu_0]$. The set of divisorial valuations in T coincides with the set of valuations with rational skewness and is hence dense in T . We now have to approximate a concave function on T by a decreasing sequence of piecewise linear functions with rational slopes, and whose second derivatives are measures supported on the rationals. This can be done in an elementary way.

By Section 1.3, $q h_n$ is the tree transform of an integrally closed ideal I if q is the product of the denominators of c_i^n with the generic multiplicities of ν_i^n over all i . By the preceding argument, the tree transform of $\mathcal{J}(h_n)$ is bounded by h_n . As $h \leq h_n$ we infer $g_{\mathcal{J}(h)} \leq g_{\mathcal{J}(h_n)} \leq h_n$. We conclude, by letting $n \rightarrow \infty$, that $g_{\mathcal{J}(h)} \leq h$ as desired.

Finally consider a general tree potential h . First assume that $\rho\{\nu\} < m(\nu)$ for all curve valuations ν . Since ρ has finite mass we may pick $\varepsilon > 0$ such that in fact $\rho\{\nu\} < (1 - \varepsilon)m(\nu)$. Define $\mathcal{T} = \{\nu \in \mathcal{V} : \rho\{\mu \geq \nu\} \geq (1 - \varepsilon)m(\nu)\}$. By Lemma 2.12, \mathcal{T} is a finite subtree of \mathcal{V} and $\mathcal{J}(h_{\mathcal{T}}) = \mathcal{J}(h)$. Our choice of ε means that the ends of \mathcal{T} are quasimonomial. Hence the previous analysis gives $g_{\mathcal{J}(h_{\mathcal{T}})} \leq h_{\mathcal{T}}$. Since $h_{\mathcal{T}} \leq h$ this yields $g_{\mathcal{J}(h)} \leq h$.

In the most general case we can decompose $h = g_\phi + h'$, where $\phi \in R$ (possibly reducible) and h' is a tree potential with associated measure satisfying $\rho'\{\nu\} < m(\nu)$ for all curve valuations ν . By repeated use of Proposition 2.10 we have $\mathcal{J}(h) = \phi \mathcal{J}(h')$. Therefore the previous step gives

$$g_{\mathcal{J}(h)} = g_\phi + g_{\mathcal{J}(h')} \leq g_\phi + h' = h,$$

which concludes the proof. \square

APPENDIX A. SUBADDITIVITY AND SKODA'S THEOREM

In this appendix we prove two fundamental properties of multiplier ideals associated to tree potentials. The proofs appear here as they rely on the corresponding properties of multiplier ideals of formal powers of ideals. It would be interesting to have direct proofs, not using a reduction to the case of powers of ideals.

Proof of Proposition 2.8. Write $h = h_1 + h_2$. Let $\rho_i = \rho_{h_i}$ and $\rho = \rho_h = \rho_1 + \rho_2$ be the measures associated to h_i and h , respectively. As in the proof of Lemma 8.8 we will gradually build the proof using successively more general measures.

First suppose that $h_i = c_i g_{I_i}$, $i = 1, 2$, for ideals I_1, I_2 and $c_1, c_2 > 0$ rational. By Proposition 3.2 our subadditivity statement translates into $\mathcal{J}(I_1^{c_1} \cdot I_2^{c_2}) \subset \mathcal{J}(I_1^{c_1}) \cdot \mathcal{J}(I_2^{c_2})$, which holds by [DEL] (see the proof of Lemma 8.8).

Next suppose the support of ρ is included in a finite tree \mathcal{T} whose ends are quasimonomial valuations. As in the proof of Lemma 8.8 we approximate h_i from above by tree potentials of the form $h_i^n = c_i^n g_{I_i^n}$, where $c_i^n > 0$ are rational and I_i^n are primary ideals. By semicontinuity (Proposition 2.6) we have $\mathcal{J}(h_i^n) = \mathcal{J}(h_i)$ and $\mathcal{J}(h_1^n + h_2^n) = \mathcal{J}(h)$ for large n , completing the proof in this case.

Now consider general tree potentials h_i . First assume that $\rho_i\{\nu\} < m(\nu)$ for all curve valuations ν , $i = 1, 2$. Since ρ_i has finite mass we may pick $\varepsilon > 0$ so that in fact $\rho_i\{\nu\} < (1 - \varepsilon)m(\nu)$. Define $\mathcal{T}_i = \{\nu \in \mathcal{V} : \rho_i\{\nu\} \geq \varepsilon\}$. By Lemma 2.12, \mathcal{T}_i is a finite subtree of \mathcal{V} and $\mathcal{J}((h_i)_{\mathcal{T}_i}) = \mathcal{J}(h_i)$. Our choice of ε means that the ends of \mathcal{T}_i are quasimonomial. Hence the previous analysis gives

$$\mathcal{J}(h_1 + h_2) \subset \mathcal{J}((h_1)_{\mathcal{T}_1} + (h_2)_{\mathcal{T}_2}) \subset \mathcal{J}((h_1)_{\mathcal{T}_1}) \cdot \mathcal{J}((h_2)_{\mathcal{T}_2}) = \mathcal{J}(h_1) \cdot \mathcal{J}(h_2).$$

In the most general case we can decompose $h_i = g_{\phi_i} + h'_i$, where $\phi_i \in R$ (possibly reducible) and h'_i is a tree potential with associated measure satisfying $\rho'_i\{\nu\} < m(\nu)$ for all curve valuations ν . By repeated use of Proposition 2.10 we have $\mathcal{J}(h_i) = \phi_i \mathcal{J}(h'_i)$ and $\mathcal{J}(h) = \phi_1 \phi_2 \mathcal{J}(h'_1 + h'_2)$. Therefore the previous step gives

$$\mathcal{J}(h_1 + h_2) = \phi_1 \phi_2 \mathcal{J}(h'_1 + h'_2) \subset \phi_1 \phi_2 \mathcal{J}(h'_1) \cdot \mathcal{J}(h'_2) = \mathcal{J}(h_1) \cdot \mathcal{J}(h_2),$$

which completes the proof. \square

Proof of Proposition 2.9. The proof is based on a reduction to the case $h = c g_I$ for an integrally closed ideal I and $c > 0$ rational. In the latter case the result is well known (see [La, Theorem 9.6.21]) in view of Proposition 3.2. The reduction goes along exactly the same steps as the proof of subadditivity above. The details are left to the reader. \square

REFERENCES

- [BM] M. Blé and S.K. Mimouni. *Singularités et intégrabilité des fonctions plurisousharmoniques*. Ann. Inst. Fourier, to appear.
- [D] J.-P. Demailly. *Multiplier ideal sheaves and analytic methods in algebraic geometry*. In *School on Vanishing Theorems and Effective Results in Algebraic Geometry* (Trieste 2000), 1–148. ICTP Lect. Notes, vol. 6, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2001. MR1919457 (2003f:32020)
- [DEL] J.-P. Demailly, L. Ein and R. Lazarsfeld. *A subadditivity property of multiplier ideals*. Michigan Math. J. 48 (2000), 137–156. MR1786484 (2002a:14016)
- [DK] J.-P. Demailly and J. Kollár. *Semi-continuity of complex singularity exponents and Kähler-Einstein metrics on Fano orbifolds*. Ann. Sci. École Norm. Sup. (4) 34 (2001), no. 4, 525–556. MR1852009 (2002e:32032)
- [ELS] L. Ein, R. Lazarsfeld and K.E. Smith. *Uniform approximation of Abhyankar valuation ideals in smooth function fields*. Amer. J. Math. 125 (2003), no. 2, 409–440. MR1963690 (2003m:13004)
- [FJ1] C. Favre and M. Jonsson. *The valuative tree*. Lecture Notes in Mathematics, 1853. Springer-Verlag, Berlin, 2004. MR2097722
- [FJ2] C. Favre and M. Jonsson. *Valuative analysis of planar plurisubharmonic functions*. Invent. Math., to appear.
- [I] J. Igusa. *On the first terms of certain asymptotic expansions*. Complex analysis and algebraic geometry, 357–368. Iwanami Shoten, Tokyo, 1977. MR0485881 (58:5680)

- [Ki1] C.-O. Kiselman. *Un nombre de Lelong raffiné*. Séminaire d'analyse complexe et géométrie 1985-1987, pp. 61-70. Faculté des sciences de Tunis et Faculté des Sciences et techniques de Monastir.
- [Ki2] C.-O. Kiselman. *Attenuating the singularities of plurisubharmonic functions*. Ann. Polon. Math. 60 (1994), no. 2, 173–197. MR1301603 (95i:32024)
- [Ku] T. Kuwata. *On log canonical thresholds of reducible plane curves*. Amer. J. Math. 121 (1999), no. 4, 701–721. MR1704476 (2001g:14047)
- [La] R. Lazarsfeld. *Positivity in algebraic geometry II*. Ergebnisse der Math. und ihrer Grenzgebiete (3), vol. 49. Springer-Verlag, Berlin, 2004. MR2095472
- [Li] J. Lipman. *Adjoints of ideals in regular local rings*. Math. Res. Lett. 1 (1994), no. 6, 739–755. With an appendix by S. C. Cutkosky. MR1306018 (95k:13028)
- [LW] J. Lipman and K.-I. Watanabe. *Integrally closed ideals in two-dimensional regular local rings are multiplier ideals*. Math. Res. Lett. 10 (2003), no. 4, 423–434. MR1995782 (2004m:13059)
- [MP] J. McKernan and Y. Prokhorov. *Threefold thresholds*. Manuscripta Math. 114 (2004), 281–304. MR2075967
- [Mi] S. K. Mimouni. *Singularités des fonctions plurisousharmoniques et courants de Liouville*. Thèse de la faculté des sciences de Monastir, Tunisie, 2001.
- [N] A. M. Nadel. *Multiplier ideal sheaves and Kähler-Einstein metrics of positive scalar curvature*. Ann. of Math. (2) 132 (1990), 549–596. MR1078269 (92d:32038)
- [PS] D.-H. Phong and J. Sturm. *On a conjecture of Demailly and Kollár*. Kodaira's issue. Asian J. Math. 4 (2000), no. 1, 221–226. MR1803721 (2002b:32043)
- [S] V. V. Shokurov. *Three-dimensional log perestroikas*. Russian Acad. Sci. Izv. Math. 40 (1993), no. 1, 95–202. MR1162635 (93j:14012)
- [Sp] M. Spivakovsky. *Valuations in function fields of surfaces*. Amer. J. Math. 112 (1990), no. 1, 107–156. MR1037606 (91c:14037)
- [S] H. Skoda. *Sous-ensembles analytiques d'ordre fini ou infini dans \mathbf{C}^n* . Bull. Soc. Math. France 100 (1972), 353–408. MR0352517 (50:5004)
- [TW] S. Takagi and K.-I. Watanabe. *When does the subadditivity theorem for multiplier ideals hold?* Trans. Amer. Math. Soc. 356 (2004), 3951–3961. MR2058513 (2005d:13016)

CNRS, INSTITUT DE MATHÉMATIQUES, EQUIPE GÉOMÉTRIE ET DYNAMIQUE, F-75251 PARIS CEDEX 05, FRANCE

E-mail address: favre@math.jussieu.fr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MICHIGAN 48109-1109

Current address: Department of Mathematics, Royal Institute of Technology, SE-100 44 Stockholm, Sweden

E-mail address: mattiasj@umich.edu

E-mail address: mattiasj@kth.se