

THE ESSENTIALLY TAME LOCAL LANGLANDS CORRESPONDENCE, I

COLIN J. BUSHNELL AND GUY HENNIART

To the memory of Albrecht Fröhlich

1. Let F be a non-Archimedean local field; we assume throughout that the residue field \mathbb{k}_F of F is finite, of characteristic p . Let \mathcal{W}_F be the Weil group of F , formed relative to some separable algebraic closure of F . For each integer $n \geq 1$, let $\mathfrak{S}_n^0(F)$ be the set of equivalence classes of *irreducible* smooth (complex) representations of \mathcal{W}_F of dimension n . Likewise, let $\mathcal{A}_n^0(F)$ be the set of equivalence classes of irreducible *supercuspidal* representations of the group $\mathrm{GL}_n(F)$. The Langlands correspondence gives a bijection

$$(1) \quad {}_F\mathcal{L}_n : \mathfrak{S}_n^0(F) \xrightarrow{\cong} \mathcal{A}_n^0(F)$$

for each n . The existence of the family $\{{}_F\mathcal{L}_n\}$ is established indirectly, by methods relying heavily on geometric or global constructions [21], [14], [15]; explicit information about it, of the sort essential for local calculations, is inconveniently hard to obtain. Indeed, in virtually no case is there a complete, explicit account.

2. In a series of papers, of which this is the first, we describe ${}_F\mathcal{L}_n$ in the “essentially tame” case. For $\sigma \in \mathfrak{S}_n^0(F)$, let $t(\sigma)$ be the number of unramified characters χ of \mathcal{W}_F for which $\chi \otimes \sigma \cong \sigma$. The integer $t(\sigma)$ divides n , and we say that σ is *essentially tame* if p does not divide $n/t(\sigma)$. (This is equivalent to demanding that the restriction of σ to the wild inertia subgroup of \mathcal{W}_F be a sum of characters: see Appendix (A.4) below.) We denote by $\mathfrak{S}_n^{\mathrm{et}}(F)$ the set of essentially tame classes $\sigma \in \mathfrak{S}_n^0(F)$. Analogously, we say that $\pi \in \mathcal{A}_n^0(F)$ is *essentially tame* if p does not divide $n/t(\pi)$, where $t(\pi)$ is the number of unramified characters χ of F^\times such that π is equivalent to the representation

$$\chi\pi : g \longmapsto \chi(\det g)\pi(g).$$

Let $\mathcal{A}_n^{\mathrm{et}}(F)$ denote the set of essentially tame $\pi \in \mathcal{A}_n^0(F)$. The correspondence (1) then restricts to a bijection

$$(2) \quad {}_F\mathcal{L}_n^{\mathrm{et}} : \mathfrak{S}_n^{\mathrm{et}}(F) \xrightarrow{\cong} \mathcal{A}_n^{\mathrm{et}}(F).$$

Our aim is to give a complete and explicit description of the map (2).

Received by the editors March 29, 2004.

2000 *Mathematics Subject Classification.* Primary 22E50.

Key words and phrases. Explicit local Langlands correspondence, base change, automorphic induction, tame lifting, admissible pair.

Much of the work in this programme was carried out while the first-named author was visiting, and partly supported by, l’Université de Paris-Sud. Both authors were also partially supported by the EU network “Arithmetical Algebraic Geometry”.

©2005 American Mathematical Society
 Reverts to public domain 28 years from publication

When $p \nmid n$, we have $\mathfrak{G}_n^{\text{et}}(F) = \mathfrak{G}_n^0(F)$ and $\mathcal{A}_n^{\text{et}}(F) = \mathcal{A}_n^0(F)$. Under this hypothesis (and when F has characteristic zero), Moy [23] (see also Reimann [24]) wrote down a bijection $\mathfrak{G}_n^0(F) \rightarrow \mathcal{A}_n^0(F)$ which, at the time, seemed a good candidate for the Langlands correspondence. We show in this paper that Moy's correspondence is, in its general form, quite close to the Langlands correspondence; in later papers, for example [7], we will see that the two diverge significantly at the level of detail. (However, Moy's map is the Langlands correspondence when n is prime.)

3. One classifies the elements of $\mathfrak{G}_n^{\text{et}}(F)$ using *admissible pairs*. We recall the matter briefly: see the Appendix for more details.

Definition. Let E/F be a finite, tamely ramified field extension and let ξ be a quasicharacter of E^\times . The pair $(E/F, \xi)$ is called *admissible* if it satisfies the following two conditions. Let K range over intermediate fields, $F \subset K \subset E$.

- (1) If ξ factors through the relative norm $N_{E/K}$, then $K = E$.
- (2) If $\xi|_{U_E^1}$ factors through $N_{E/K}$, then E/K is unramified.

Let $P_n(F)$ denote the set of F -isomorphism classes of admissible pairs $(E/F, \xi)$ in which $[E:F] = n$. The map

$$(3) \quad \begin{aligned} P_n(F) &\longrightarrow \mathfrak{G}_n^{\text{et}}(F), \\ (E/F, \xi) &\longmapsto \text{Ind}_{E/F} \xi, \end{aligned}$$

then provides a canonical bijection. (Here, we regard ξ as a quasicharacter of \mathcal{W}_E via class field theory, and denote by $\text{Ind}_{E/F}$ the functor of induction from representations of \mathcal{W}_E to representations of \mathcal{W}_F : see (A.3) below for more details.)

We combine the structure theory for supercuspidal representations [9] with the technique of tame lifting of simple characters [3] to construct a canonical bijection

$$(4) \quad \begin{aligned} P_n(F) &\longrightarrow \mathcal{A}_n^{\text{et}}(F), \\ (E/F, \xi) &\longmapsto {}_F\pi_\xi. \end{aligned}$$

Together, (3) and (4) yield a canonical bijection

$$(5) \quad {}_F\mathcal{N}_n : \mathfrak{G}_n^{\text{et}}(F) \longrightarrow \mathcal{A}_n^{\text{et}}(F),$$

which we call the “naïve correspondence”. Our aim is to compare the maps ${}_F\mathcal{N}_n$, ${}_F\mathcal{L}_n^{\text{et}}$. The main result of this paper is:

Theorem A. Let $(E/F, \xi) \in P_n(F)$, and set $\sigma = \text{Ind}_{E/F} \xi$. There is a tamely ramified character $\mu = {}_F\mu_\xi$ of E^\times such that

$${}_F\mathcal{L}_n^{\text{et}}(\sigma) = {}_F\pi_{\mu_\xi}.$$

Evaluation of the character ${}_F\mu_\xi$ in the general case requires the development of quite elaborate methods: in this paper, we treat it only in the following special case. Let $\delta_{E/F}$ denote the discriminant character of the extension E/F , that is, the determinant of the regular representation $\text{Ind}_{E/F} 1$. We show:

Theorem B. Let $(E/F, \xi) \in P_n(F)$; suppose that n is odd and E/F is totally ramified. We then have

$${}_F\mu_\xi = \delta_{E/F} \circ N_{E/F}.$$

4. Our general method is based on the following fundamental fact. Let K/F be a finite, cyclic extension. We thus have the operations of restriction and induction connecting the representations of the Weil, or Weil-Deligne, groups of K and F . Under the Langlands correspondence, these correspond respectively to *base change* $b_{K/F}$ and *automorphic induction* $A_{K/F}$. When K/F is also tamely ramified, the theory of tame lifting [3], [6] describes how the classification from [9] interacts with these operations.

When we restrict to essentially tame representations, the compatibility with base change and automorphic induction is sufficient to determine the Langlands correspondence uniquely: we outline a proof at the end of §3.

In particular, the epsilon factor of pairs [19], [27] plays no rôle whatsoever. Indeed, it is not presently known whether the preservation of L - and epsilon factors does determine the correspondence when restricted to essentially tame representations.

In the later papers of the series, we shall completely determine the character ${}_F\mu_\xi$ of Theorem A. As we shall first see in [7], it takes a rather complicated form. Some of this is traceable to our construction of the map (4). At a certain critical point, it relies on a choice which, while both natural and canonical in the technical sense, is not the only one available. We have retained it for consistency with the literature, and for its simplicity and directness. On the other hand, some of the complexity in ${}_F\mu_\xi$ seems unavoidable. We will return to this matter in a later paper, with more facts in hand.

5. We place our results in their historical context. We recall that, when $p \nmid n$, we have $\mathfrak{G}_n^{\text{et}}(F) = \mathfrak{G}_n^0(F)$ and $\mathcal{A}_n^{\text{et}}(F) = \mathcal{A}_n^0(F)$. In that case, Howe [18] first constructed the map (4) $P_n(F) \rightarrow \mathcal{A}_n^{\text{et}}(F)$, but he was only able to prove it is injective. Moy [23] then took up the problem and proved, under the hypothesis that F has characteristic zero, that (4) is bijective. The restriction on characteristic zero is essential since his method relied on a description of the admissible dual of D^\times , for a central F -division algebra D of dimension n^2 [11], [20], and on the existence of the Jacquet-Langlands correspondence. At that time, the latter was only available in characteristic zero [25]. We, on the other hand, can proceed much more directly and generally: the construction of the map (4) is an instance of the theory of tame lifting [3], [6] and its bijectivity follows from the classification theorems of [9].

One can modify the naïve correspondence by “tame twisting”, to get bijections $P_n(F) \rightarrow \mathcal{A}_n^{\text{et}}(F)$ of the form $(E/F, \xi) \mapsto {}_F\pi_{\nu_\xi}$, where ν_ξ is a tame character of E^\times varying with ξ . Moy observed that one could choose the system $\xi \mapsto \nu_\xi$ so that the corresponding bijection $\mathfrak{G}_n^{\text{et}}(F) \rightarrow \mathcal{A}_n^{\text{et}}(F)$ satisfies those properties of the Langlands correspondence known at the time he was writing: these mainly concern behaviour relative to L - and epsilon factors of a *single* representation. In totality, these properties are rather weak. They do not imply that Moy’s map is the Langlands correspondence, or even that the correspondence can be obtained at all as a tame twist of the naïve correspondence.

Theorem A finally confirms the accuracy of Moy’s basic intuition, even if the fine details are different. It should be noted that Theorem A is, in effect, an instance of the main theorem (7.1) of [6], a result of considerable depth.

We mention that Bushnell and Fröhlich [2] took a similar approach to the representations of D^\times and came to similar conclusions: the Langlands correspondence seemed to be obtainable from the analogue of the naïve correspondence by a process

of tame twisting, but the precise suggestions in [2] are inaccurate to very much the same degree as Moy's.

We point out the later paper of Reimann [24]. He showed that, after the correction of minor errors and tangential misconceptions, the approaches of [23], [2], could be accommodated within the same formal framework. The concision and accuracy of [24] make it a valuable consolidation of the earlier papers, but it gets no closer to the true correspondence than they.

Note on characteristic. The operations of *base change* and *automorphic induction* are constructed, in [1] and [16] respectively, on the hypothesis that F has characteristic zero. The operation of *tame lifting* (of endo-classes of simple characters) [3] is defined in arbitrary characteristic. The main results of [3], which are basic to everything we do here, relate tame lifting to base change and automorphic induction. They depend on the standard character relations and a short list of other properties concerned with linear independence of supercuspidal characters on certain sets. As we remarked in [3, §16], a theory of base change and automorphic induction in positive characteristic, with these auxiliary properties, would instantly extend the main results of [3] to that case. Such a theory is constructed in [17], together with some useful refinements of the kind worked out in the appendices to [8]. The results of [6] depend only on those of [3], [8] along with certain conductor estimates valid in all characteristics. We therefore anticipate the outcome of [17] and proceed as if the results of [3], [6] hold in positive characteristic. This allows us here to treat all characteristics on an equal footing.

Notation and background. Throughout, F is a non-Archimedean local field; we let \mathfrak{o}_F be the discrete valuation ring in F and \mathfrak{p}_F the maximal ideal of \mathfrak{o}_F . We set $\mathbb{k}_F = \mathfrak{o}_F/\mathfrak{p}_F$ and $p = \text{char } \mathbb{k}_F$. We write $U_F = \mathfrak{o}_F^\times$ and $U_F^n = 1 + \mathfrak{p}_F^n$, $n \geq 1$. We let $v_F : F^\times \rightarrow \mathbb{Z}$ be the canonical additive valuation.

If K/F is a finite field extension, we use similar notation relative to K and we write $N_{K/F}$, $\text{Tr}_{K/F}$ for the relative norm and trace respectively.

We fix a character ψ_F of F , trivial on \mathfrak{p}_F but not on \mathfrak{o}_F .

If $\pi \in \mathcal{A}_n^0(F)$, we denote the contragredient of π by $\tilde{\pi}$ and the central quasicharacter of π by ω_π .

a. Let V be a finite-dimensional F -vector space and write $A = \text{End}_F(V)$, $G = \text{Aut}_F(V)$. We use the notation of [9] for simple strata in A and simple characters in G . Thus, if $[\mathfrak{A}, l, 0, \beta]$ is a simple stratum in A , we have the compact open subgroups $H^1(\beta, \mathfrak{A})$ and $J^i(\beta, \mathfrak{A})$ of G , $i = 0, 1$, along with the set $\mathcal{C}(\mathfrak{A}, \beta, \psi_F)$ of simple characters of $H^1(\beta, \mathfrak{A})$. (We drop ψ_F from the notation for much of the time, as in [9].)

b. We shall require the notion of *endo-equivalence* for simple characters, and also the operation of *tame lifting* for endo-equivalence classes (or endo-classes, for short): this theory is developed in [3], but the summary in [6, §1] may be found helpful. We let $\mathcal{E}(F)$ denote the set of endo-classes of simple characters over F . If $[\mathfrak{A}, l, 0, \beta]$ is a simple stratum in A and $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$, we write $\mathcal{E}_F(\theta)$ for the endo-class of θ . We adjoin to $\mathcal{E}(F)$ a trivial element Θ_0 , which may be regarded as the class of trivial characters of groups $U_{\mathfrak{A}}^1$, where \mathfrak{A} ranges over hereditary \mathfrak{o}_F -orders in matrix algebras over F .

c. If K/F is a finite, tamely ramified field extension, there is a canonical surjective map $R_{K/F} : \mathcal{E}(K) \rightarrow \mathcal{E}(F)$, transitive in the extension K/F . If $\Theta \in \mathcal{E}(F)$, the fibre $R_{K/F}^{-1}(\Theta)$ is finite, and its elements are the K/F -lifts of Θ . In particular, if Θ is the endo-class of $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$, the K/F -lifts of Θ are parametrized by the simple components of the algebra $F[\beta] \otimes_F K$.

d. Let $\pi \in \mathcal{A}_n^0(F)$. We attach to π an endo-class $\Theta(\pi) \in \mathcal{E}(F)$ as follows. If π has a fixed vector for the group $U_{\mathfrak{M}}^1$, $\mathfrak{M} = M_n(\mathfrak{o}_F)$, we set $\Theta(\pi) = \Theta_0$ (one says that “ π has level zero”). Otherwise, π contains a simple character $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$, for some simple stratum $[\mathfrak{A}, n, 0, \beta]$. In this case, we set $\Theta(\pi) = \mathcal{E}_F(\theta)$.

We shall frequently need the following special cases of the main result (7.1) of [6]. Let K/F be a finite, tamely ramified, cyclic extension, and let $\pi \in \mathcal{A}_n^0(F)$ and $\rho \in \mathcal{A}_m^0(K)$. First, if the base change $\pi_K = b_{K/F}(\pi)$ is supercuspidal, then

$$\Theta(\pi) = R_{K/F}(\Theta(\pi_K)).$$

More precisely, $\Theta(\pi_K)$ is the unique K/F -lift of $\Theta(\pi)$. Second, if the automorphically induced representation $\rho^K = A_{K/F}(\rho)$ is supercuspidal, then

$$\Theta(\rho^K) = R_{K/F}(\Theta(\rho)).$$

For the general statement, we refer to [6], although we shall only use the cases above.

1. TAMELY RAMIFIED SIMPLE CHARACTERS

Let $\Theta \in \mathcal{E}(F)$; there is a finite-dimensional F -vector space V , a simple stratum $[\mathfrak{A}, l, 0, \beta]$ in $\text{End}_F(V)$, and a simple character $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$ such that $\Theta = \mathcal{E}_F(\theta)$. The quantities

$$e(\Theta) = e(F[\beta]|F), \quad f(\Theta) = f(F[\beta]|F), \quad \deg \Theta = [F[\beta]:F],$$

depend only on the endo-class Θ , not the choice of representatives θ or β , [3, 8.11].

For the trivial element Θ_0 , we define $e(\Theta_0) = f(\Theta_0) = \deg \Theta_0 = 1$.

Let $\Theta \in \mathcal{E}(F)$; we say that Θ is *tame* or *tamely ramified* if p does not divide $e(\Theta)$. We write $\mathcal{E}^{\text{et}}(F)$ for the set of tame endo-classes $\Theta \in \mathcal{E}(F)$: in particular, $\mathcal{E}^{\text{et}}(F)$ contains the trivial element of $\mathcal{E}(F)$.

The aim of this section is to give an independent description of the set $\mathcal{E}^{\text{et}}(F)$.

1.1. Let K/F be a finite, tamely ramified field extension and let $\Theta \in \mathcal{E}^{\text{et}}(F)$. We say that K/F *splits* Θ if Θ has a K/F -lift Θ_K such that $\deg \Theta_K = 1$.

Proposition 1.1. (1) *Let $\Theta \in \mathcal{E}^{\text{et}}(F)$ and let K/F split Θ ; then $\deg \Theta$ divides $[K:F]$.*

(2) *Let $\Theta \in \mathcal{E}^{\text{et}}(F)$ have degree d . There then exists a tamely ramified extension K/F , of degree d , which splits Θ . Moreover, Θ determines K/F uniquely, up to F -isomorphism.*

Proof. We choose a finite-dimensional F -vector space V together with a simple stratum $[\mathfrak{A}, n, 0, \beta]$ in $\text{End}_F(V)$ and a simple character $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$ such that $\mathcal{E}_F(\theta) = \Theta$. By definition, $\deg \Theta = [F[\beta]:F]$ and $e(\Theta) = e(F[\beta]|F)$. In particular, the field extension $F[\beta]/F$ is tamely ramified.

Let K/F be finite and tamely ramified; we write $K \otimes_F F[\beta] = \prod_i E_i$, where the E_i are fields. Each E_i is a tamely ramified extension of K , and the corresponding

K/F -lift Θ_i of Θ satisfies $\deg \Theta_i = [E_i:K]$. Thus K/F splits Θ if and only if $E_i = K$, for some i . This occurs if and only if there exists an F -embedding of $F[\beta]$ in K , so we have proved (1) and the existence statement of (2). Further, if K/F has degree d and splits Θ , this embedding $F[\beta] \rightarrow K$ must be an isomorphism. \square

Let K/F be tamely ramified and let $\Theta \in \mathcal{E}^{\text{et}}(F)$; we say that K/F is a *splitting field for Θ* if it splits Θ and $[K:F] = \deg \Theta$. By the proposition, any $\Theta \in \mathcal{E}^{\text{et}}(F)$ has a splitting field, and this is unique up to F -isomorphism. Moreover, the proof of the proposition shows:

Corollary 1.1. *Let $\Theta \in \mathcal{E}^{\text{et}}(F)$ and let K/F be a finite, tamely ramified field extension. Then K/F splits Θ if and only if K contains a splitting field for Θ .*

Remark. For the moment, let $\Theta \in \mathcal{E}(F)$ be arbitrary, that is, not necessarily tamely ramified. Choose a simple stratum $[\mathfrak{A}, l, 0, \beta]$ in some $\text{End}_F(V)$ such that $\Theta = \mathcal{E}_F(\theta)$ for some $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$. The same argument as above shows that *the maximal tamely ramified sub-extension of $F[\beta]/F$ is uniquely determined, up to F -isomorphism, by the endo-class Θ* . This is not the case for the wildly ramified part.

1.2. We make a simple observation. Let χ be a non-trivial character of U_F^1 : suppose χ is trivial on U_F^{n+1} but not on U_F^n , for some $n \geq 1$. There then exists $c \in F$, with $v_F(c) = -n$, such that

$$\chi(1+x) = \psi_F(cx), \quad 2v_F(x) \geq n+1.$$

The quadruple $[\mathfrak{o}_F, n, 0, c]$ is a simple stratum in $\text{End}_F(F) = F$ and χ is an element of $\mathcal{C}(\mathfrak{o}_F, c, \psi_F)$. We can therefore form the endo-class $\mathcal{E}_F(\chi) \in \mathcal{E}(F)$. If χ is the trivial character of U_F^1 , then $\mathcal{E}_F(\chi)$ is the trivial endo-class. In this way, we can identify the dual \widehat{U}_F^1 of U_F^1 :

$$\widehat{U}_F^1 = \{\Theta \in \mathcal{E}(F) : \deg \Theta = 1\}.$$

If K/F is a finite tame extension and $\chi \in \widehat{U}_F^1$, the endo-class $\mathcal{E}_F(\chi)$ has a unique K/F -lift, namely $\mathcal{E}_K(\chi \circ N_{K/F})$. Put another way, if we write $\xi = \chi \circ N_{K/F} \in \widehat{U}_K^1$, then

$$R_{K/F} \mathcal{E}_K(\xi) = \mathcal{E}_F(\chi).$$

1.3. We consider pairs $(K/F, \chi)$, where K/F is a finite, tamely ramified field extension and $\chi \in \widehat{U}_K^1$. We call $(K/F, \chi)$ an *admissible 1-pair* if χ does not factor through $N_{K/E}$, for any field E with $F \subset E \subsetneq K$. We write $P^{(1)}(F)$ for the set of isomorphism classes of admissible 1-pairs over F .

Let $(K/F, \chi) \in P^{(1)}(F)$. We can thus form the endo-class $R_{K/F} \mathcal{E}_K(\chi) \in \mathcal{E}(F)$.

Theorem 1.3. *Let $(K/F, \chi) \in P^{(1)}(F)$, and write $\Theta_\chi = R_{K/F} \mathcal{E}_K(\chi)$. Then $\Theta_\chi \in \mathcal{E}^{\text{et}}(F)$ and $\deg \Theta_\chi = [K:F]$. Moreover, the map*

$$(1.3.1) \quad \begin{aligned} P^{(1)}(F) &\longrightarrow \mathcal{E}^{\text{et}}(F), \\ (K/F, \chi) &\longmapsto \Theta_\chi \end{aligned}$$

is a canonical bijection, which is natural with respect to automorphisms of the base field F .

Proof. Let $(K/F, \chi) \in P^{(1)}(F)$; by definition, the endo-class $\Theta_\chi \in \mathcal{E}(F)$ has a K/F -lift of degree 1. Since K/F is tamely ramified, it follows that Θ_χ is tame and K/F splits Θ_χ . In particular, $\deg \Theta_\chi$ divides $[K:F]$. Furthermore, K contains a splitting field E/F for Θ_χ (Corollary 1.1). There is, therefore, a character ϕ of U_E^1 such that $\Theta_\chi = R_{E/F} \mathcal{E}_E(\phi)$ and $\mathcal{E}_K(\chi)$ is the K/F -lift of $\mathcal{E}_E(\phi)$. That is, $\chi = \phi \circ N_{K/E}$; since $(K/F, \chi)$ is admissible, we have $E = K$ and $\deg \Theta_\chi = [E:F] = [K:F]$, as required.

For $i = 1, 2$, let $(K_i/F, \chi_i) \in P^{(1)}(F)$, and assume that $\Theta_{\chi_1} = \Theta_{\chi_2}$. We choose a finite, tamely ramified Galois extension L/F containing both K_i . We set $\chi_i^L = \chi_i \circ N_{L/K_i} \in \widehat{U}_L^1$. By hypothesis, we have

$$R_{L/F} \mathcal{E}_L(\chi_1^L) = R_{L/F} \mathcal{E}_L(\chi_2^L).$$

By [3, 9.13 *et seq*], this implies that the endo-classes $\mathcal{E}_L(\chi_i^L)$, hence also the characters χ_i^L themselves, are conjugate under $\text{Gal}(L/F)$. Since K_i/F is the least subextension of L/F such that χ_i^L factors through N_{L/K_i} (*cf.* (A.1), Lemma (3)), this conjugation carries the pair $(K_1/F, \chi_1)$ to $(K_2/F, \chi_2)$, whence the pairs $(K_i/F, \chi_i)$ are conjugate. In other words, $(K_1/F, \chi_1) \cong (K_2/F, \chi_2)$, and we have shown that the map $P^{(1)}(F) \rightarrow \mathcal{E}^{\text{et}}(F)$ is injective.

Let $\Theta \in \mathcal{E}^{\text{et}}(F)$ have degree n . There is a vector space V of dimension n over F , a simple stratum $[\mathfrak{A}, l, 0, \beta]$ in $\text{End}_F(V)$ and a simple character $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$ with $\Theta = \mathcal{E}_F(\theta)$. We put $E = F[\beta]$, so that $[E:F] = n$. We have $U_E^1 = E^\times \cap H^1(\beta, \mathfrak{A})$, and we form the character $\chi = \theta \mid U_E^1$. As in 1.2, χ is a simple character over E , and (by definition [3, 9.4]) $\mathcal{E}_E(\chi)$ is an E/F -lift of Θ . We first have to show that $(E/F, \chi)$ is an admissible 1-pair. Suppose we have a subfield K , $F \subset K \subset E$, such that χ factors through $N_{E/K}$, say $\chi = \phi \circ N_{E/K}$, for some $\phi \in \widehat{U}_K^1$. Then $\mathcal{E}_K(\phi)$ is a K/F -lift of Θ , of degree 1. That is, K/F splits Θ , whence $n = \deg \Theta$ divides $[K:F]$. By 1.1, we have $K = E$ and $(E/F, \chi) \in P^{(1)}(F)$. Therefore $\Theta = \Theta_\chi$, and our map is surjective. It is therefore bijective, as required.

The operation of tame lifting of endo-classes is natural with respect to automorphisms of the base field F . The same therefore applies to the bijection of the theorem. \square

Let us exhibit the inverse of the map $P^{(1)}(F) \rightarrow \mathcal{E}^{\text{et}}(F)$, as implied by the proof of the theorem. Suppose $\Theta \in \mathcal{E}^{\text{et}}(F)$ is the endo-class of $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$, for a simple stratum $[\mathfrak{A}, l, 0, \beta]$ in $A = \text{End}_F(V)$, for some vector space V . Set $E = F[\beta]$ and let B denote the A -centralizer of E . Thus $\mathfrak{B} = B \cap \mathfrak{A}$ is a hereditary \mathfrak{o}_E -order in B and we have $H^1(\beta, \mathfrak{A}) \cap B^\times = U_{\mathfrak{B}}^1$. The restriction of θ to $U_{\mathfrak{B}}^1$ is of the form $\phi \circ \det_B$, for a uniquely determined character ϕ of U_E^1 . The image of Θ in $P^{(1)}(F)$ is then the F -isomorphism class of $(E/F, \phi)$.

Remark. The base field restriction map $R_{K/F} : \mathcal{E}(K) \rightarrow \mathcal{E}(F)$ does not depend on the underlying choice of character ψ_F (*cf.* [6, Remark §1]). The same, therefore, applies to the map (1.3.1).

1.4. Let $[\mathfrak{A}, l, 0, \beta]$ be a simple stratum in $M_n(F)$, say, and let $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$. Let χ be a character of U_F^1 and consider the character $\theta' = \theta \cdot (\chi \circ \det)$ of $H^1(\beta, \mathfrak{A})$. There is then a simple stratum $[\mathfrak{A}, l', 0, \beta']$, where $\beta' = \beta + a$, for some $a \in F^\times$, such that $H^1(\beta', \mathfrak{A}) = H^1(\beta, \mathfrak{A})$ and $\theta' \in \mathcal{C}(\mathfrak{A}, \beta')$: see [10, Appendix] for details of this construction.

In the present context, let $(K/F, \xi) \in P^{(1)}(F)$, and let χ be a character of U_F^1 . Setting $\xi' = \xi \cdot (\chi \circ N_{K/F})$, the 1-pair $(K/F, \xi')$ is admissible, and

$$(1.4.1) \quad \Theta_{\xi'} = \mathcal{E}_F(\theta_\xi \cdot (\chi \circ \det)),$$

for any simple character θ_ξ such that $\mathcal{E}_F(\theta_\xi) = \Theta_\xi$.

2. ESSENTIALLY TAME SUPERCUSPIDAL REPRESENTATIONS

Let π be an irreducible supercuspidal representation of $G = \text{GL}_n(F)$. As in the introduction, we let $t(\pi)$ denote the number of unramified characters χ of F^\times such that $\chi\pi \cong \pi$. Thus $t(\pi)$ is an integer dividing n ; we say that π is *essentially tame* if p does not divide $n/t(\pi)$. We write $\mathcal{A}_n^{\text{et}}(F)$ for the set of equivalence classes of essentially tame, irreducible supercuspidal representations of $\text{GL}_n(F)$.

The aim of this section is to construct a canonical bijection $P_n(F) \rightarrow \mathcal{A}_n^{\text{et}}(F)$, for each $n \geq 1$.

2.1. We recall briefly the main features of the classification [9] of irreducible supercuspidal representations of $G = \text{GL}_n(F)$, in terms of maximal simple types. Let $\pi \in \mathcal{A}_n^0(F)$. According to [9, (8.4.1)], π contains a unique G -conjugacy class of maximal simple types (J, λ) , in the sense of [9, (5.5.10, 6.2)]. Moreover, if π contains the maximal simple type (J, λ) and \mathbf{J}_λ denotes the G -normalizer of the pair (J, λ) , there is a unique representation A of \mathbf{J}_λ which occurs in π and satisfies $A|_J \cong \lambda$. We then have

$$\pi \cong c\text{-Ind}_{\mathbf{J}_\lambda}^G A.$$

(We refer to pairs of the form (\mathbf{J}_λ, A) as *extended* maximal simple types.)

The definition *loc. cit.* divides the maximal simple types into two subclasses. In the first, dubbed “of level zero”, we have $J = U_{\mathfrak{A}}$, where $\mathfrak{A} \cong M_n(\mathfrak{o}_F)$. The representation λ is the inflation of an irreducible *cuspidal* representation $\bar{\lambda}$ of $U_{\mathfrak{A}}/U_{\mathfrak{A}}^1 \cong \text{GL}_n(\mathbb{k}_F)$. The group \mathbf{J}_λ is $F^\times J$. A representation $\pi \in \mathcal{A}_n^0(F)$ is said to be of level zero if it contains a maximal simple type of level zero.

In the second case, there is a simple stratum $[\mathfrak{A}, l, 0, \beta]$ in $A = M_n(F)$, such that \mathfrak{A} is maximal among $F[\beta]^\times$ -stable hereditary \mathfrak{o}_F -orders in A , and a simple character $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$ which occurs in π : indeed, θ is the unique simple character occurring in π , up to G -conjugacy. The group $J = J^0(\beta, \mathfrak{A})$ is the $U_{\mathfrak{A}}$ -normalizer of θ ; the natural representation λ of J on the θ -isotypic space in π gives a maximal simple type (J, λ) in π . (We recall more of the representation λ below.) The group \mathbf{J}_λ is $F[\beta]^\times J$.

For $\pi \in \mathcal{A}_n^0(F)$, we define an invariant $\Theta(\pi) \in \mathcal{E}(F)$ as follows: $\Theta(\pi)$ is trivial if π is of level zero, while otherwise $\Theta(\pi) = \mathcal{E}_F(\theta)$, where θ is a simple character occurring in π .

Proposition 2.1. *Let $\pi \in \mathcal{A}_n^0(F)$; then π is essentially tame if and only if $\Theta(\pi) \in \mathcal{E}^{\text{et}}(F)$.*

Proof. If π has level zero, then $t(\pi) = n$ while, if π contains a simple character $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$, then $t(\pi) = n/e(F[\beta]|F)$ [9, (6.2.5)]. In other words, $n/t(\pi) = e(\Theta(\pi))$, whence the result follows. □

2.2. It is convenient to deal separately with the first subclass. Let $\mathcal{A}_n^0(F)_0$ denote the set of $\pi \in \mathcal{A}_n^0(F)$ which are of level zero. Let $P_n(F)_0$ denote the set of $(E/F, \xi) \in P_n(F)$ for which $\xi|_{U_E^1} = 1$.

Take $(E/F, \xi) \in P_n(F)_0$. By definition, E/F is unramified. We set $\Sigma = \text{Gal}(E/F)$. The admissibility of the pair $(E/F, \xi)$ is then equivalent to ξ being Σ -regular, in the sense that the conjugates ξ^σ , $\sigma \in \Sigma$, are distinct. The character $\xi|_{U_E}$ is the inflation of a character $\bar{\xi}$ of \mathbb{k}_E^\times . If we identify Σ with $\text{Gal}(\mathbb{k}_E/\mathbb{k}_F)$ by reduction, the admissibility of $(E/F, \xi)$ is again equivalent to $\bar{\xi}$ being Σ -regular.

We now use the Green parametrization [13]: this gives a canonical bijection $\chi \mapsto \bar{\lambda}_\chi$ between the set of Σ -orbits of Σ -regular characters χ of \mathbb{k}_E^\times and the set of equivalence classes of irreducible cuspidal representations $\bar{\lambda}$ of $\text{GL}_n(\mathbb{k}_F)$.

We choose an F -embedding of E in $A = M_n(F)$, and let \mathfrak{A} be the unique E^\times -stable hereditary \mathfrak{o}_F -order in A : we have $\mathfrak{A} \cong M_n(\mathfrak{o}_F)$ and $U_{\mathfrak{A}}/U_{\mathfrak{A}}^1 \cong \text{GL}_n(\mathbb{k}_F)$. The character $\bar{\xi}$ gives a cuspidal representation $\bar{\lambda}_\xi$ of $\text{GL}_n(\mathbb{k}_F)$, which we inflate to a representation λ_ξ of $U_{\mathfrak{A}}$. We extend λ_ξ to a representation A_ξ of $\mathbf{J} = F^\times U_{\mathfrak{A}}$ by deeming that $A_\xi|_{F^\times}$ be a multiple of ξ . We set

$$(2.2.1) \quad {}_F\pi_\xi = c\text{-Ind}_{\mathbf{J}}^G A_\xi.$$

Proposition 2.2. *Let $(E/F, \xi) \in P_n(F)_0$ and define ${}_F\pi_\xi$ as in (2.2.1). Then:*

- (1) ${}_F\pi_\xi \in \mathcal{A}_n^0(F)_0$, and the equivalence class of π depends only on that of the pair $(E/F, \xi)$;
- (2) the map $(E/F, \xi) \mapsto {}_F\pi_\xi$ is a canonical bijection $P_n(F)_0 \rightarrow \mathcal{A}_n^0(F)_0$.

Proof. The first assertion is immediate, so we have a well-defined map $P_n(F)_0 \rightarrow \mathcal{A}_n^0(F)_0$. If we take $\pi \in \mathcal{A}_n^0(F)_0$, it contains a maximal simple type $(\text{GL}_n(\mathfrak{o}_F), \lambda)$ (unique up to conjugacy), where λ is inflated from an irreducible cuspidal representation of $\text{GL}_n(\mathbb{k}_F)$. We reverse the procedure above to construct from π a pair $(E/F, \xi) \in P_n(F)_0$. This gives a well-defined map $\mathcal{A}_n^0(F)_0 \rightarrow P_n(F)_0$, inverse to the first one. \square

2.3. We take an admissible pair $(E/F, \xi)$, of degree n , such that $\xi|_{U_E^1} \neq 1$. We construct from this pair a representation ${}_F\pi_\xi \in \mathcal{A}_n^{\text{et}}(F)$, the equivalence class of which will depend only on the isomorphism class of $(E/F, \xi)$.

Let E'/F be the minimal sub-extension of E/F such that $\xi|_{U_E^1}$ factors through $N_{E'/E}$ (see Appendix (A.1), Lemma (3)), and write $\xi|_{U_E^1} = \phi \circ N_{E'/E}$. We choose a simple stratum $[\mathfrak{A}, l, 0, \beta]$ in $A = M_n(F)$ and a simple character $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$ such that $\mathcal{E}_F(\theta) = R_{E'/F} \mathcal{E}_{E'}(\phi)$. We then have $F[\beta] \cong E'$; we henceforward identify E' with $F[\beta] \subset A$.

We can choose \mathfrak{A} to be maximal among E'^\times -stable hereditary \mathfrak{o}_F -orders in A . That done, the $G = \text{GL}_n(F)$ -conjugacy class of the pair (\mathfrak{A}, θ) is determined by the F -isomorphism class of $(E'/F, \phi)$ (and hence by that of $(E/F, \xi)$). The pair (\mathfrak{A}, θ) determines the groups $J^1 = J^1(\beta, \mathfrak{A})$, $J^0 = J^0(\beta, \mathfrak{A})$ and $\mathbf{J} = E'^\times J^0$: these groups are the normalizers of θ in, respectively, $U_{\mathfrak{A}}^1$, $U_{\mathfrak{A}}$ and G .

Let B denote the A -centralizer of E' . Let η be the unique irreducible representation of J^1 which contains θ [9, (5.1.1)].

Lemma 1. *There exists a unique irreducible representation κ of J such that*

- (1) $\kappa|_{J^1} \cong \eta$;

- (2) κ is intertwined by every element of B^\times ;
- (3) the character $\det \kappa$ has finite, p -power order.

Proof. This follows from [9, (5.2.2)]. □

To proceed further, we have to choose a prime element ϖ_F of F .

Lemma 2. *There is a unique representation $\tilde{\kappa}$ of \mathbf{J} satisfying the following conditions:*

- (1) $\tilde{\kappa} \mid J \cong \kappa$;
- (2) $\varpi_F \in \text{Ker } \tilde{\kappa}$;
- (3) $\det \tilde{\kappa}$ has finite, p -power order.

Proof. We have $\mathbf{J} = E^\times J = E'^\times J$ and $E'^\times \cap J = U_{E'}$. We can therefore extend κ by triviality to the group $\langle \varpi_F, J \rangle$ generated by ϖ_F and J . The quotient $\mathbf{J}/\langle \varpi_F, J \rangle$ is cyclic of order $e(E'|F)$. Since $p \nmid e(E'|F)$, κ admits a unique extension $\tilde{\kappa}$ of the required form. □

For the next step, we use the same prime element ϖ_F to impose a factorization on the quasicharacter ξ .

Lemma 3. *There is a unique character ξ_w of E^\times with the following properties:*

- (1) $\xi_w \mid U_E^1 = \xi \mid U_E^1$;
- (2) $\xi_w(\varpi_F) = 1$;
- (3) ξ_w has finite, p -power order.

The proof is immediate. We now set $\xi_t = \xi_w^{-1}\xi$. The quasicharacter ξ_t is tamely ramified, and the pair $(E/E', \xi_t)$ is admissible.

Let $\mathfrak{B} = \mathfrak{A} \cap B$. Thus \mathfrak{B} is a maximal $\mathfrak{o}_{E'}$ -order in B . We have $J = U_{\mathfrak{B}} J^1$ and $J/J^1 \cong U_{\mathfrak{B}}/U_{\mathfrak{B}}^1 \cong \text{GL}_d(\mathbb{k}_{E'})$, where $d = [E:E']$. We follow the procedure of 2.2 to define, from the admissible pair $(E/E', \xi_t)$, an irreducible representation Λ_t of $E^\times U_{\mathfrak{B}}/U_{\mathfrak{B}}^1 \cong \mathbf{J}/J^1$. We view Λ_t as a representation of \mathbf{J} trivial on J^1 . We put $\Lambda_\xi = \Lambda_t \otimes \tilde{\kappa}$, and note that this definition is independent of the intermediate choice of ϖ_F .

At this point, it becomes useful to condense our notation a little:

Notation. Let $(E/F, \xi) \in P_n(F)$, and let E'/F be the minimal sub-extension of E/F such that $\xi \mid U_E^1$ factors through $N_{E'/E'}$. Thus $\xi \mid U_E^1 = \xi' \circ N_{E'/E'}$, for a uniquely determined character ξ' of $U_{E'}^1$. We put

$$\Theta_\xi = \Theta_{\xi'} = R_{E'/F}(\mathcal{E}_{E'}(\xi')) = R_{E/F}(\mathcal{E}_E(\xi \mid U_E^1)).$$

Proposition 2.3. *The representation ${}_F\pi_\xi = c\text{-Ind}_{\mathbf{J}}^G \Lambda_\xi$ is irreducible, supercuspidal, essentially tame, and it satisfies $\Theta({}_F\pi_\xi) = \Theta_\xi$. The equivalence class of ${}_F\pi_\xi$ depends only on the isomorphism class of the pair $(E/F, \xi)$.*

Proof. Setting $\lambda = \Lambda_\xi \mid J$, the pair (J, λ) is a maximal simple type in G and $(\mathbf{J}, \Lambda_\xi)$ is an extended maximal simple type. Thus $\pi = {}_F\pi_\xi$ lies in $\mathcal{A}_n^0(F)$ and by construction, $\Theta(\pi) = \Theta_\xi \in \mathcal{E}^{\text{et}}(F)$. Thus π is essentially tame. By Theorem 1.3 and the uniqueness of the pair (\mathfrak{A}, θ) described above, different choices of pairs $(E/F, \xi)$ in the same isomorphism class lead to conjugate types (\mathbf{J}, Λ) and hence equivalent representations ${}_F\pi_\xi$. □

Taking Proposition 2.2 into account, the operation $(E/F, \xi) \mapsto {}_F\pi_\xi$ gives a well-defined map $P_n(F) \rightarrow \mathcal{A}_n^{\text{et}}(F)$.

Theorem 2.3. *The map*

$$(2.3.1) \quad \begin{aligned} P_n(F) &\longrightarrow \mathcal{A}_n^{\text{et}}(F), \\ (E/F, \xi) &\longmapsto {}_F\pi_\xi \end{aligned}$$

is a bijection.

Proof. A representation $\pi \in \mathcal{A}_n^{\text{et}}(F)$ contains an extended maximal simple type (\mathbf{J}, λ) , of level zero or based on a simple stratum [9, (8.4.1)]. In the latter case, the stratum is tamely ramified by Proposition 2.1. Any such (\mathbf{J}, λ) arises from an element of $P_n(F)$ as above: all steps in the construction are reversible. Thus the map (2.3.1) is surjective.

For $i = 1, 2$, let $(E_i/F, \xi_i) \in P_n(F)$ and suppose that ${}_F\pi_{\xi_1} = {}_F\pi_{\xi_2} = \pi$, say. Let E'_i/F be the minimal sub-extension of E_i/F such that $\xi_i \mid U_{E_i}^1$ factors through N_{E_i/E'_i} . Let ϕ_i be the character of $U_{E'_i}^1$ such that $\xi_i \mid U_{E_i}^1 = \phi_i \circ N_{E_i/E'_i}$. By Theorem 1.3 and the proposition, the admissible 1-pairs $(E'_i/F, \phi_i)$ are isomorphic, so we may take $E'_1 = E'_2 = E'$ and $\phi_1 = \phi_2$. The extensions E_i/E' are unramified of the same degree, so we may as well assume $E_1 = E_2 = E$, say, and $\xi_1 \mid U_E^1 = \xi_2 \mid U_E^1$.

If we follow through the constructions above, for the pairs $(E/F, \xi_i)$, we arrive at the same stratum $[\mathfrak{A}, l, 0, \beta]$ and the same $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$. Furthermore, if we choose a prime element ϖ_F of F to factorize the ξ_i as in Lemma 3, we get $\xi_{1,w} = \xi_{2,w}$. The corresponding extended simple types (\mathbf{J}, λ_i) , $\mathbf{J} = E'^{\times} J^0(\beta, \mathfrak{A})$, lie in the same representation π and so are G -conjugate. Indeed, they are conjugate under the G -normalizer of θ , which is none other than \mathbf{J} . In other words, $\lambda_1 = \lambda_2$. In the factorizations $\lambda_i = \lambda_{i,t} \otimes \tilde{\kappa}$, the factors $\lambda_{i,t}$ are equivalent, so the admissible pairs $(E/E', \xi_i \xi_{i,w}^{-1})$ are E' -isomorphic. We conclude that the pairs $(E/F, \xi_i)$ are F -isomorphic, as required. \square

Remark 1. The only debatable step in the construction of ${}_F\pi_\xi$ is the definition of the factor $\tilde{\kappa}$ held to correspond to the wildly ramified factor ξ_w of ξ . Other choices of $\tilde{\kappa}$ are possible, and some give variants of (2.3.1) with smoother properties. We shall analyze this phenomenon elsewhere. For the present, the version given above has the advantage of being simple, straightforward, and in need of no preliminary analysis.

Remark 2. Constructions such as (2.3.1) originated in [18], and were further developed in [23]. Slightly different conventions are used in [24], where the relation between the approaches is discussed. Our method is actually identical to that of [24], as one verifies easily, starting from Theorem 1.3. We have subsumed the extensive details of the earlier papers into the general theory of [9] and [3], to obtain the simple and direct proof of bijectivity. This option was not available to those authors who were also confined to the case $p \nmid n$ and F of characteristic zero.

2.4. We collect some simple properties of the bijection (2.3.1).

Proposition 2.4. (1) *The map (2.3.1) is natural with respect to automorphisms of the base field F .*

(2) *If $(K/F, \xi) \in P_n(F)$ and $\pi = {}_F\pi_\xi$, then*

$${}_F\pi_{\xi^{-1}} = \tilde{\pi} \quad \text{and} \quad \omega_\pi = \xi \mid F^\times.$$

(3) Let $(K/F, \xi) \in P_n(F)$ and let χ be a quasicharacter of F^\times . Write $\chi_K = \chi \circ N_{K/F}$. Then $(K/F, \xi \cdot \chi_K) \in P_n(F)$ and

$$F\pi_{\xi \cdot \chi_K} = \chi \cdot F\pi_\xi$$

(4) The map (2.3.1) is natural with respect to (not necessarily continuous) automorphisms of \mathbb{C} , that is,

$$F\pi_{\gamma \circ \xi} = \gamma \circ F\pi_\xi,$$

for $(E/F, \xi) \in P_n(F)$ and $\gamma \in \text{Aut } \mathbb{C}$.

All of these assertions are proved by following through the construction of $F\pi_\xi$, step by step and, for (3), comparing with 1.4.

3. THE LANGLANDS CORRESPONDENCE AND THE NAÏVE CORRESPONDENCE

We make an initial comparison between the constructions of §2 and the Langlands correspondence for essentially tame representations.

3.1. We recall that we have a canonical bijection $P_n(F) \rightarrow \mathfrak{G}_n^{\text{et}}(F)$ given by $(E/F, \xi) \mapsto \text{Ind}_{E/F} \xi$ (A.3). Combining this with Theorem 2.3, we have the first statement of:

Theorem 3.1. *For each $n \geq 1$, the maps*

$$\begin{aligned} P_n(F) &\longrightarrow \mathfrak{G}_n^{\text{et}}(F), & P_n(F) &\longrightarrow \mathcal{A}_n^{\text{et}}(F), \\ (E/F, \xi) &\longmapsto \text{Ind}_{E/F} \xi, & (E/F, \xi) &\longmapsto F\pi_\xi \end{aligned}$$

together induce a bijection

$$\mathcal{N} = {}_F\mathcal{N}_n : \mathfrak{G}_n^{\text{et}}(F) \xrightarrow{\cong} \mathcal{A}_n^{\text{et}}(F).$$

The map \mathcal{N} has the following properties:

- (1) ${}_F\mathcal{N}_1$ is the bijection $\mathfrak{G}_1(F) \rightarrow \mathcal{A}_1(F)$ of local class field theory;
- (2) ${}_F\mathcal{N}_n$ is natural with respect to automorphisms of the base field F ;
- (3) if $\sigma \in \mathfrak{G}_n^{\text{et}}(F)$ and $\pi = \mathcal{N}(\sigma)$, then $\mathcal{N}(\check{\sigma}) = \check{\pi}$;
- (4) if $\sigma \in \mathfrak{G}_n^{\text{et}}(F)$ and $\chi \in \mathfrak{G}_1(F)$, then $\mathcal{N}(\chi \otimes \sigma) = \chi \cdot \mathcal{N}(\sigma)$;
- (5) if $\sigma = \text{Ind}_{E/F} \xi$, for $(E/F, \xi) \in P_n(F)$, and $\pi = \mathcal{N}(\sigma) = F\pi_\xi$, then

$$(3.1.1) \quad \det \sigma = \omega_\pi \delta_{E/F}.$$

Proof. Property (1) is a matter of definition, while the others follow from Proposition 2.4. □

Remark. We observe also that \mathcal{N} commutes with the natural actions of $\text{Aut } \mathbb{C}$.

3.2. Let ${}_F\mathcal{L}_n : \mathfrak{G}_n^0(F) \rightarrow \mathcal{A}_n^0(F)$ be the Langlands correspondence. We have observed that ${}_F\mathcal{L}_n$ satisfies the analogue of Theorem 3.1(4), so it induces a bijection

$${}_F\mathcal{L}_n^{\text{et}} : \mathfrak{G}_n^{\text{et}}(F) \xrightarrow{\cong} \mathcal{A}_n^{\text{et}}(F).$$

We usually write simply ${}_F\mathcal{L}_n^{\text{et}} = \mathcal{L}$.

Let K/F be a cyclic field extension of degree d . We then have the operations of base change $b_{K/F}$ [1], [17], and automorphic induction $A_{K/F}$ [16], [17]. These correspond, via the Langlands correspondence, to the operations of restriction and induction connecting representations of the Weil-Deligne groups of F and K *loc. cit.* Translating this in terms of our constructions, we have:

Proposition 3.2. *The Langlands correspondence induces a bijection*

$$\mathcal{L} = {}_F\mathcal{L}_n^{\text{et}} : \mathfrak{S}_n^{\text{et}}(F) \longrightarrow \mathcal{A}_n^{\text{et}}(F),$$

for each $n \geq 1$. This map satisfies the analogues of (1)–(4) of Theorem 3.1, while instead of (5) we have

$$(3.2.1) \quad \omega_\pi = \det \sigma, \quad \pi = \mathcal{L}(\sigma).$$

In addition:

- (6) Let $(E/F, \xi) \in P_n(F)$ and let K/F be a cyclic sub-extension of E/F . Put $\tau = \text{Ind}_{E/K} \xi$, $\rho = \mathcal{L}(\tau)$, $\sigma = \text{Ind}_{E/F} \xi$ and $\pi = \mathcal{L}(\sigma)$. Then π is automorphically induced by ρ :

$$\pi = A_{K/F} \rho.$$

- (7) Let $(E/F, \xi) \in P_n(F)$, let L/F be a finite, cyclic tamely ramified extension, and suppose that the pair $(EL/L, \xi_L)$ is admissible, where $\xi_L = \xi \circ N_{EL/E}$. Set $\sigma = \text{Ind}_{E/F} \xi$, $\pi = \mathcal{L}(\sigma)$ and $\sigma_L = \text{Ind}_{EL/L} \xi_L = \sigma | \mathcal{W}_L$. Then

$$\mathcal{L}(\sigma_L) = b_{L/F} \pi.$$

We recall [6, (7.1)] that in (6) we have $\Theta(\pi) = R_{K/F} \Theta(\rho)$ while, in (7), $\Theta(b_{L/F} \pi)$ is the unique L/F -lift of $\Theta(\pi)$.

Remark. The list of properties in the proposition is sufficient to determine the family ${}_F\mathcal{L}_n^{\text{et}}$ uniquely. This is not a new observation: it is implicit in [22] and also in our arguments. We sketch a direct proof in 3.5 below.

3.3. We now give our main results, using the notation of 2.3.

Theorem 3.3. *Let $(E/F, \xi) \in P_n(F)$; set $\sigma = \text{Ind}_{E/F} \xi \in \mathfrak{S}_n^{\text{et}}(F)$ and $\pi = \mathcal{L}(\sigma) \in \mathcal{A}_n^{\text{et}}(F)$. Then*

$$\Theta(\pi) = \Theta_\xi = \Theta({}_F\pi_\xi).$$

Proof. Set $\pi' = N(\sigma) = {}_F\pi_\xi$ and abbreviate $R_{E/F}(\xi|U_E^1) = \Theta_\xi$. The relation $\Theta_\xi = \Theta(\pi')$ is then given by Propositions 2.2, 2.3.

To prove that $\Theta(\pi) = \Theta_\xi$, we first suppose that E/F is totally ramified. Thus σ is totally ramified, in the sense that $t(\sigma) = 1$. Let L/F be the unramified extension of F generated by a primitive n -th root of unity and set $\xi_L = \xi \circ N_{LE/E}$. The pair $(EL/L, \xi_L)$ is then admissible, and we put

$$\sigma_L = \text{Ind}_{EL/L} \xi_L = \sigma | \mathcal{W}_L \in \mathfrak{S}_n^{\text{et}}(L).$$

Set $\rho = {}_L\mathcal{L}_n(\sigma_L)$; since the extension EL/L is cyclic, we have $\rho = A_{EL/L} \xi_L$. Since L/F is cyclic, we also have $\rho = b_{L/F} \pi$. We now use [6, (7.1)]. The relation $\rho = A_{EL/L} \xi_L$ implies $\Theta(\rho) = R_{EL/L} \mathcal{E}_{EL}(\xi_L|U_{EL}^1)$, while $\rho = b_{L/F} \pi$ implies $R_{L/F} \Theta(\rho) = \Theta(\pi)$. In all,

$$\Theta(\pi) = R_{L/F} \Theta(\rho) = R_{EL/F} \mathcal{E}_{EL}(\xi_L|U_{EL}^1) = R_{E/F}(\xi|U_E^1) = \Theta_\xi,$$

since the map R is transitive with respect to the base field extension. This proves the theorem in the case where E/F is totally ramified.

In the general case, let K/F be the maximal unramified sub-extension of E/F , put $\tau = \text{Ind}_{E/K} \xi$ and $\rho = \mathcal{L}(\tau)$. Thus $\pi = A_{K/F} \rho$. By the first case, we have $\Theta(\rho) = R_{E/K} \mathcal{E}_E(\xi | U_E^1)$, whence *loc. cit.*

$$\Theta(\pi) = R_{K/F} R_{E/K} \mathcal{E}_E(\xi | U_E^1) = \Theta_\xi.$$

This completes the proof of the theorem. \square

We can re-interpret the theorem as follows:

Corollary 3.3. *Let $(E/F, \xi) \in P_n(F)$ and set $\sigma = \text{Ind}_{E/F} \xi$. There exists a tamely ramified character $\mu = {}_F\mu_\xi$ of E^\times such that $(E/F, \mu\xi)$ is admissible and*

$${}_F\mathcal{L}_n^{\text{et}}(\sigma) = {}_F\pi_{\mu\xi}.$$

The character ${}_F\mu_\xi$ further satisfies ${}_F\mu_\xi | F^\times = \delta_{E/F}$.

Proof. Let us write $\pi = \mathcal{N}(\sigma) = {}_F\pi_\xi$ and $\pi' = \mathcal{L}(\sigma)$. By Theorem 2.3, there is an admissible pair $(E'/F, \xi') \in P_n(F)$ such that $\pi' = {}_F\pi_{\xi'}$. By the theorem, we have

$$\Theta_{\xi'} = \Theta(\pi') = \Theta_\xi = \Theta(\pi).$$

Let K/F be the minimal sub-extension of E/F such that $\xi | U_E^1$ factors through $N_{E/K}$, and let ϕ be the character of U_K^1 for which $\xi | U_E^1 = \phi \circ N_{E/K}$. Similarly define K'/F and ϕ' . The 1-pairs $(K/F, \phi), (K'/F, \phi')$ are admissible; we have

$$\mathbf{R}_{K/F}\mathcal{E}_K(\phi) = \Theta_\xi = \Theta_{\xi'} = \mathbf{R}_{K'/F}\mathcal{E}_{K'}(\phi'),$$

so $(K'/F, \phi')$ is F -isomorphic to $(K/F, \phi)$ (Theorem 1.3). We may as well, therefore, take $K = K'$ and $\phi = \phi'$. The extensions $E/K, E'/K$ are unramified of the same degree $n/[K:F]$, so we are justified in setting $E' = E$. These identifications give $\xi | U_E^1 = \xi' | U_E^1$, so $\xi^{-1}\xi' = \mu$ is tamely ramified, as required. The final assertion follows from comparison of (3.2.1) with (3.1.1). \square

Remark. One can say little about the character ${}_F\mu_\xi$ without further analysis. One sees easily that it has finite order: we show elsewhere that it depends only on $\xi | U_E^1$ and that its order divides 4. Also, we have noted that the correspondence \mathcal{N} is algebraic, in the sense of commuting with $\text{Aut } \mathbb{C}$. However, \mathcal{L} is not algebraic, as is clear from the discussion in, for example, [5, §7]. The function $(E/F, \xi) \mapsto {}_F\mu_\xi$ therefore cannot be algebraic.

3.4. It will be useful to record a variation on the theme of the corollary. For this, we take an admissible pair $(E/F, \xi) \in P_n(F)$ and a cyclic sub-extension K/F of E/F , of degree d . The pair $(E/K, \xi)$ is then admissible, $(E/K, \xi) \in P_m(F)$, where $md = n$, so we can form the representation $\rho = {}_K\pi_\xi \in \mathcal{A}_m^{\text{et}}(K)$.

Proposition 3.4. *Let $\pi = \mathbf{A}_{K/F} \rho$ be the representation of $\text{GL}_n(F)$ automorphically induced by ρ . The representation π is supercuspidal, essentially tame, and there exists a tamely ramified character $\mu = {}_{K/F}\mu_\xi$ of E^\times such that $\pi = {}_F\pi_{\mu\xi}$. The character μ also satisfies $\mu | F^\times = \delta_{K/F}^m$.*

Proof. Let $\Sigma = \text{Gal}(K/F)$; we view E as a subfield of a separable algebraic closure \overline{F}/F and extend each $\sigma \in \Sigma$ to an F -embedding, also denoted σ , of E in \overline{F} . The admissible pairs $(E^\sigma/K, \xi^\sigma), \sigma \in \Sigma$, are then mutually non-isomorphic over K . The construction (cf. Proposition 2.4 (1)) yields ${}_K\pi_{\xi^\sigma} = \rho^\sigma$. The representations $\rho^\sigma, \sigma \in \Sigma$, are therefore mutually inequivalent or, in other words, ρ is Σ -regular. It follows [8, 2.6] that $\pi = \mathbf{A}_{K/F} \rho$ is supercuspidal and [6, (7.1)] $\Theta(\pi) = \mathbf{R}_{K/F} \Theta(\rho)$. Thus π is essentially tame, and of the form ${}_F\pi_{\xi'}$, for some $(E'/F, \xi') \in P_n(F)$.

Furthermore, we have $\Theta(\pi) = \mathbf{R}_{E'/F}\mathcal{E}_{E'}(\xi'|U_{E'}^1) = \mathbf{R}_{E/F}\mathcal{E}_E(\xi|U_E^1)$. The proof concludes in the same manner as that of Corollary 3.3. \square

3.5. We remarked in 3.2 that the family of correspondences ${}_F\mathcal{L}_n^{\text{et}}$ is uniquely determined by the properties listed in Proposition 3.2. We shall never use this fact directly, but we give an outline of the proof since it illuminates our strategy.

We take $\sigma \in \mathfrak{G}_n^{\text{et}}(F)$; we have to construct $\mathcal{L}(\sigma)$ using only properties from the list in section 3.2. We proceed by induction on n . If $n = 1$, then $\mathcal{L}(\sigma)$ is given by class field theory (the analogue for \mathcal{L} of Theorem 3.1(1)). We therefore assume $n > 1$ and we write $\sigma = \text{Ind}_{E/F} \xi$, for some $(E/F, \xi) \in P_n(F)$. Suppose first that E/F admits a non-trivial cyclic sub-extension K/F . Setting $\tau = \text{Ind}_{E/K} \xi$, the representation $\rho = \mathcal{L}(\tau)$ is known, by the induction hypothesis, while $\mathcal{L}(\sigma) = \mathbb{A}_{K/F} \rho$, by Proposition 3.2(6).

We are so reduced to the case where E/F admits no non-trivial cyclic sub-extension. Thus E/F is totally ramified and $n = [E:F]$ is odd. Let ℓ be the least prime divisor of n , and let L/F be the unramified extension generated by a primitive ℓ -th root of unity. Setting $\xi_L = \xi \circ N_{EL/E}$, the pair $(EL/L, \xi_L)$ is admissible; let $\sigma_L = \text{Ind}_{EL/L} \xi_L = \sigma | \mathcal{W}_L$. The extension EL/L surely admits a non-trivial cyclic sub-extension so, by the first case, $\pi_L = \mathcal{L}(\sigma_L)$ has been constructed. Setting $\pi = \mathcal{L}(\sigma)$, we have $\pi_L = \mathfrak{b}_{L/F} \pi$, by Proposition 3.2(7). Let $k = [L:F]$, and let X_k be the group of unramified characters χ of F^\times satisfying $\chi^k = 1$. The set of $\rho \in \mathfrak{A}_n^{\text{et}}(F)$ such that $\mathfrak{b}_{L/F} \rho = \pi_L$ forms a single X_k -orbit. The central quasicharacter of $\chi\rho$ is $\chi^n \omega_\rho$. Since $\omega_\pi = \det \sigma$ (3.2.1) and $(k, n) = 1$, the representation π is uniquely determined by the two conditions $\mathfrak{b}_{L/F} \pi = \pi_L$ and $\omega_\pi = \det \sigma$. \square

4. TOTALLY RAMIFIED REPRESENTATIONS

We make a preliminary calculation of the character ${}_{K/F}\mu_\xi$ of Proposition 3.4 in a special case. We then specialize further, and prove Theorem B of the introduction.

4.1. Let $(E/F, \xi) \in P_n(F)$, and suppose that E/F is *totally ramified*. In particular, $p \nmid n$. For the moment, we set $\pi = {}_F\pi_\xi$. Thus π is *totally ramified* in that $t(\pi) = 1$.

We choose an F -embedding of E in $A = M_n(F)$, and henceforward regard E as a subfield of A . As in section 2.3, the pair $(E/F, \xi)$ gives a simple stratum $[\mathfrak{A}, l, 0, \beta]$ in A with $E = F[\beta]$, and a simple character $\theta \in \mathcal{C}(\mathfrak{A}, \beta)$ such that $\theta(x) = \xi(x)$, for $x \in U_E^1$. We abbreviate $H^1 = H^1(\beta, \mathfrak{A})$, $J^1 = J^1(\beta, \mathfrak{A})$ and $\mathbf{J} = E^\times J^1$. We let η be the unique irreducible representation of J^1 which contains θ .

We consider the finite p -group $\Omega = J^1/\text{Ker } \theta$. The centre of Ω is the cyclic group $\mathcal{Z} = H^1/\text{Ker } \theta$, and Ω/\mathcal{Z} is an elementary abelian p -group. Since θ is stable under conjugation by E^\times , the group Ω inherits an action of E^\times , the subgroup $F^\times U_E^1$ acting trivially. We therefore view Ω as a module over the finite group $\Gamma = E^\times/F^\times U_E^1$; this action of Γ stabilizes η , viewed as a representation of Ω . Since E/F is totally ramified, the group Γ is cyclic.

The group Ω^Γ of Γ -fixed points is exactly \mathcal{Z} [4, 4.1, Lemma 1]. We apply the machinery of the Glauberman correspondence [12] (or see [4, Appendix] for a discussion in the present context).

(4.1.1). (1) *There exists a unique representation $\tilde{\eta}$ of the group $\Gamma \times \Omega$ such that $\tilde{\eta} | \Omega \cong \eta$ and $\det \tilde{\eta} | \Gamma = 1$.*

(2) *There exists a constant $\epsilon_F = \pm 1$ such that $\text{tr } \tilde{\eta}(\gamma u) = \epsilon_F \xi(u)$, for every generator γ of Γ and every $u \in U_E^1$.*

We recall that there is a unique irreducible representation Λ of \mathbf{J} such that $\Lambda \mid H^1$ is a multiple of θ and $\pi = c\text{-Ind } \Lambda$. The construction of Λ in section 2.3 can now be expressed more simply in this case, as follows:

Lemma 4.1. *Defining ϵ_F as in (4.1.1), the representation Λ satisfies*

$$\text{tr } \Lambda(h) = \epsilon_F \xi(h),$$

for every $h \in E^\times$ such that $v_E(h)$ is relatively prime to n . This relation characterizes Λ among representations of \mathbf{J} which extend η .

Proof. Take the representation $\tilde{\eta}$ of (4.1.1) and inflate it to a representation Λ_0 of $E^\times \rtimes \mathcal{Q}$. Let $\tilde{\xi}$ denote the inflation of ξ to a character of $E^\times \rtimes \mathcal{Q}$ and form the representation $\Lambda_1 = \tilde{\xi} \otimes \Lambda_0$ of $E^\times \rtimes \mathcal{Q}$. We inflate Λ_1 to a representation $\tilde{\Lambda}_1$ of $E^\times \rtimes J^1$.

There is a canonical surjection $E^\times \rtimes J^1 \rightarrow \mathbf{J}$ given by $(x, j) \mapsto xj$. The kernel is the group of elements (x, x^{-1}) , $x \in U_E^1$. Since $\tilde{\Lambda}_1$ is trivial on this kernel, it is the inflation of an irreducible representation Λ_2 of \mathbf{J} . The representation Λ_2 satisfies the character relation of the lemma.

The representation Λ_2 agrees with Λ on $F^\times J^1$, so $\Lambda_2 = \chi \otimes \Lambda$, for a character χ of $\mathbf{J}/F^\times J^1 \cong E^\times / F^\times U_E^1$. Comparing the constructions, we have, for $y \in E^\times$,

$$\det \Lambda_2(y) = \xi(y)^{\dim \eta} = \zeta_y \det \Lambda(y),$$

where ζ_y is some root of unity of p -power order. Since $\dim \eta$ is a power of p and χ has order prime to p , we conclude that $\chi = 1$, as required. \square

We consider elements h of $G = \text{GL}_n(F)$ such that $v_F(\det h)$ is relatively prime to n . We call such elements *G-special*.

We will only be concerned with such elements h for which there exists a hereditary \mathfrak{o}_F -order \mathfrak{A} such that $h\mathfrak{A}h^{-1} = \mathfrak{A}$. Under these conditions, the algebra $F[h]$ is a field, totally ramified and of degree n over F .

Proposition 4.1. *Let $(E/F, \xi) \in P_n(F)$. Suppose that E/F is totally ramified and put $\pi = {}_F\pi_\xi$.*

- (1) *Let $h \in G$ be G-special and suppose that $\text{tr } \pi(h) \neq 0$. Then h is G-conjugate to an element of E^\times .*
- (2) *Let $h \in E^\times$ be G-special. Then*

$$\text{tr } \pi(h) = \epsilon_F \sum_{\alpha \in \text{Aut}(E|F)} \xi^\alpha(h).$$

Proof. The condition $\text{tr } \pi(h) \neq 0$ implies that h has a G -conjugate in \mathbf{J} , so h is an elliptic regular element of G . We can therefore apply the Mackey formula [3, (A.14)]

$$(4.1.2) \quad \text{tr } \pi(h) = \sum_{x \in G/\mathbf{J}} \text{tr } \Lambda(x^{-1}hx),$$

where we view $\text{tr } \Lambda$ as a function on G vanishing outside \mathbf{J} . Clearly, if $\text{tr } \pi(h) \neq 0$, then h is G -conjugate to an element of \mathbf{J} . We may therefore take $h = bj$, where $j \in J^1$ and $b \in E^\times$ satisfies $v_E(b) = v_F(\det b) = v_F(\det h)$. In particular, $v_E(b)$ is relatively prime to n . The argument of [3, 15.19] now applies to show that bj is J^1 -conjugate to an element bj' , where $j' \in J^1$ commutes with b . Since $F[b] = E$, this gives $j' \in U_E^1 = E^\times \cap J^1$, and the first assertion is proved.

We now take a G -special element h of E^\times and an element $x \in G$ such that $x^{-1}hx \in \mathbf{J}$. As in the first paragraph of the proof, there exists $j \in J^1$ such that $h' = j^{-1}x^{-1}hxj \in E^\times$. We have $E = F[h'] = F[h]$, so conjugation by xj induces an F -automorphism of the field E or, equivalently, $xj \in N_G(E^\times)$. We have $N_G(E^\times) \cap \mathbf{J} = E^\times(N_G(E^\times) \cap J^1)$. As J^1 is a pro- p group and $\text{Aut}(E|F) = N_G(E^\times)/E^\times$ has order prime to p , the intersection $N_G(E^\times) \cap J^1$ is contained in E^\times , whence the sum (4.1.2) is effectively taken over $\text{Aut}(E|F)$. The result now follows from the lemma. \square

4.2. For the next step, we are given a cyclic sub-extension K/F of E/F , of degree d , with $\Sigma = \text{Gal}(K/F)$. The pair $(E/K, \xi)$ is then admissible and we may form the representation $\rho = {}_K\pi_\xi \in \mathcal{A}_m^{\text{et}}(K)$, where $m = n/d$.

The representation ρ contains an extended maximal simple type $(\mathbf{J}_K, \Lambda_K)$, say, based on a simple stratum in $M_m(K)$. We use this to define a Glauberman sign $\epsilon_K = \pm 1$, as in (4.1.1). This gives us:

Lemma 4.2. *The representation Λ_K satisfies*

$$\text{tr } \Lambda_K(h) = \epsilon_K \xi(h),$$

for all $h \in E^\times$ with $v_E(h)$ relatively prime to n .

4.3. We now change notation, and let $\pi = A_{K/F} \rho$ be the representation of G which is automorphically induced by ρ . By Proposition 3.4, $\pi \in \mathcal{A}_n^{\text{et}}(F)$ and $\pi = {}_F\pi_{\mu\xi}$, for some tamely ramified character $\mu = {}_{K/F}\mu_\xi$ of E^\times . In particular, π contains an extended maximal simple (\mathbf{J}, Λ) , based on the same simple stratum $[\mathfrak{A}, l, 0, \beta]$ as in section 4.1. Replacing ξ by $\mu\xi$ does not change the constant $\epsilon_F = \pm 1$ of (4.1.1), since ϵ_F depends only on the action of E^\times on Ω , which in turn depends only on $\xi|U_E^1$.

We recall the automorphic induction equation. The cyclic extension K/F corresponds, via class field theory, to a cyclic group $Y_{K/F}$ of characters of F^\times . We choose a generator \varkappa of $Y_{K/F}$; following the conventions of [3], we form the \varkappa -twisted character Ξ_π^\varkappa of π , relative to the \varkappa -operator which acts as the identity on the θ -isotypic subspace of π . Writing G_K for the G -centralizer of K^\times , we have [16, 3.11], [17],

$$\Xi_\pi^\varkappa(h) = c \delta(h) \sum_{\sigma \in \Sigma} \text{tr } \rho^\sigma(h), \quad h \in G_K \cap G_{\text{reg}}^{\text{ell}},$$

where $G_{\text{reg}}^{\text{ell}}$ is the set of elliptic regular elements of G . In this formula, δ is a transfer factor: in the conventions of [16], $\delta(h) = \Delta^2(h)/\Delta^1(h)$. The factor c is some non-zero constant.

We prove:

Theorem 4.3. *Let $(E/F, \xi) \in P_n(F)$, and suppose that E/F is totally ramified. Let K/F be a cyclic sub-extension of E/F . The character $\mu = {}_{K/F}\mu_\xi$ satisfies*

$$(4.3.1) \quad \mu(h) = \epsilon_F \epsilon_K c \delta(h),$$

for every $h \in E^\times$ such that $v_E(h)$ is relatively prime to n .

Proof. Let Λ be the unique irreducible representation of $\mathbf{J} = E^\times J^1(\beta, \mathfrak{A})$ which occurs in π and contains the simple character θ . With our definition of Ξ_π^\varkappa , we

have [3, 15.8]:

$$\Xi_\pi^\varkappa(h) = \sum_{x \in G/\mathbf{J}} \varkappa(\det x^{-1}) \operatorname{tr} \Lambda(x^{-1}hx).$$

We evaluate this sum exactly as in Proposition 4.1 to get

$$\Xi_\pi^\varkappa(h) = \epsilon_F \sum_{\alpha \in \operatorname{Aut}(E|F)} \varkappa(\det \alpha) \mu^\alpha \xi^\alpha(h),$$

valid for $h \in E^\times$ of valuation prime to n .

We consider the character values $\operatorname{tr} \rho^\sigma(h)$, for $\sigma \in \Sigma$ and h as before.

Lemma 4.3. *Let $h \in E^\times$ have valuation prime to n , and let $\sigma \in \Sigma$. Then $\operatorname{tr} \rho^\sigma(h) = 0$ unless σ extends to an F -automorphism of E .*

Proof. For each $\sigma \in \Sigma$, we choose $t_\sigma \in G$ such that conjugation by t_σ induces the action of σ on K . Suppose that $\operatorname{tr} \rho^\sigma(h) = \operatorname{tr} \rho(t_\sigma h t_\sigma^{-1}) \neq 0$. As in Proposition 4.1, there exists $x \in G_K$ such that $x t_\sigma h t_\sigma^{-1} x^{-1} \in E^\times$, which implies $x t_\sigma \in N_G(E^\times)$ and hence the result. \square

We can apply the considerations of section 4.2 to the representation ρ^σ in place of ρ . In particular, ρ^σ gives rise to a constant $\epsilon_K^{(\sigma)} = \pm 1$. For h as before and $\sigma \in \Sigma$, we obtain

$$\operatorname{tr} \rho^\sigma(h) = \sum_{\substack{\alpha \in \operatorname{Aut}(E|F), \\ \alpha|K=\sigma}} \epsilon_K^{(\sigma)} \xi^\alpha(h),$$

with the understanding that the sum is zero when the index set is empty. In all, therefore,

$$\epsilon_F \sum_{\alpha \in \operatorname{Aut}(E|F)} \varkappa(\det \alpha) \mu^\alpha \xi^\alpha(h) = c \delta(h) \sum_{\alpha \in \operatorname{Aut}(E|F)} \epsilon_K^{(\alpha|K)} \xi^\alpha(h).$$

This relation continues to hold on replacing h by hu , for $u \in U_E^1$. The definition [16] shows readily that $\delta(hu) = \delta(h)$, so

$$\epsilon_F \sum_{\alpha \in \operatorname{Aut}(E|F)} \varkappa(\det \alpha) \mu^\alpha \xi^\alpha(h) \xi^\alpha(u) = c \delta(h) \sum_{\alpha \in \operatorname{Aut}(E|F)} \epsilon_K^{(\alpha|K)} \xi^\alpha(h) \xi^\alpha(u),$$

for $h \in E^\times$ with $v_E(h)$ prime to n and $u \in U_E^1$. Since $(E/F, \xi)$ is admissible and E/F is totally ramified, the characters ξ^α of U_E^1 are distinct, as α ranges over $\operatorname{Aut}(E|F)$. We therefore multiply the last equation by $\xi(u)^{-1}$ and integrate over U_E^1 to get the result. \square

Remark. The formula (4.3.1) determines the character μ completely. However, we cannot give it an explicit form since the value of the constant c is unknown. Note that the function $\delta(h)$ depends only on n and the extension K/F (modulo certain normalizations independent of ξ). We will show elsewhere that, with our definition of the twisted trace, the constant c depends only on the restriction $\xi|U_E^1$, whence the same is true of μ . We do not use that property in this paper.

4.4. We now specialize to the case where n is odd, and prove Theorem B of the Introduction:

Theorem 4.4. *Let $(E/F, \xi) \in P_n(F)$. Suppose that n is odd and that E/F is totally ramified. We then have*

$$F\mu_\xi = \delta_{E/F} \circ N_{E/F}.$$

The discriminant character $\delta_{E/F}$ is easy to describe: it is given by a Jacobi symbol

$$\delta_{E/F}(x) = \left(\frac{q}{n}\right)^{v_F(x)}, \quad x \in F^\times$$

(see, for example, [2, 10.1.6]).

4.5. The first step in the proof of Theorem 4.4 is:

Proposition 4.5. *Let $(E/F, \xi) \in P_n(F)$, with n odd and E/F totally ramified. Let K/F be a cyclic sub-extension of E/F . The character $K/F\mu_\xi$ is then trivial.*

Proof. Since E/F is totally tamely ramified, there exists a prime element ϖ of E such that $\varpi^n = \varpi_F$ is a prime element of F . We consider the transfer factor $\delta(\varpi^r)$, for integers r relatively prime to n .

Lemma 4.5. *As r ranges over the integers relatively prime to n , the function $r \mapsto \delta(\varpi^r)$ is constant.*

Proof. To compute the transfer factors, we follow the recipes in [16]. We need to fix a generator σ_0 of Σ and an element e_m which, since n is odd, we may set equal to 1. The character relation in Theorem 4.3 shows that the $|\delta(h)|$ is constant on G -special elements of E^\times . In the definition of the transfer factor, $|\delta(h)| = \Delta^1(h)^{-1}$, so it is enough to show that the function $r \mapsto \Delta^2(\varpi^r)$ is constant.

We have $\sigma_0(\varpi^m) = \eta\varpi^m$, for some d -th root of unity η . For $0 \leq i \leq d-1$, the eigenvalues of $\sigma_0^i(\varpi)$ are therefore $\zeta^i\varpi$, where ζ ranges over the m -th roots of η . In the notation of [16], we have $\Delta^2 = \varkappa \circ \tilde{\Delta}$ and

$$\tilde{\Delta}(\varpi) = \varpi^{m^2 d(d-1)/2} \prod_{0 \leq i < j \leq d-1} \prod_{\zeta_1, \zeta_2} (\zeta_1^i - \zeta_2^j),$$

where ζ_1, ζ_2 range independently over the m -th roots of η .

We can describe the double product

$$\Pi = \prod_{0 \leq i < j \leq d-1} \prod_{\zeta_1, \zeta_2} (\zeta_1^i - \zeta_2^j)$$

in another way. Let η_1, η_2 range independently over the n -th roots of unity. Then Π is the product of all differences $(\eta_1 - \eta_2)$, subject to two conditions: $\eta_1^m \neq \eta_2^m$ and exactly one of the possibilities $\pm(\eta_1 - \eta_2)$ occurs.

We apply the same analysis to $\tilde{\Delta}(\varpi^r)$, $(r, n) = 1$, to obtain

$$\tilde{\Delta}(\varpi^r) = \pm \varpi^{r m^2 d(-1)/2} \Pi.$$

Since n is odd and K/F is totally ramified, we have $\varkappa(\varpi_F) = 1$. Thus

$$\varkappa(\varpi^{m^2 d(d-1)/2}) = \varkappa(\varpi_F^{m(d-1)/2}) = 1.$$

Likewise, $\varkappa(-1) = 1$, so $\Delta^2(\varpi^r) = \varkappa(\Pi) = \Delta^2(\varpi)$, as required. □

Returning to the proof of the proposition, we have $\mu(\varpi^r) = c \epsilon_F \epsilon_K \delta(\varpi^r)$, whenever $(r, n) = 1$. The character μ is trivial on U_E^1 and also on F^\times , since $\delta_{K/F} = 1$. In effect, therefore, μ is a character of the group $E^\times / F^\times U_E^1 \cong \mathbb{Z}/n\mathbb{Z}$. The lemma shows that it is constant on the generators of this group and, since n is odd, μ must be trivial. \square

4.6. The proof of Theorem 4.4 relies on an auxiliary result (which does not require our assumption that n is odd).

Theorem 4.6. *Let $(E/F, \xi) \in P_n(F)$, and suppose that E/F is totally ramified. Let L/F be unramified of degree d , with $(d, n) = 1$. Write $\xi_L = \xi \circ N_{EL/E}$. The pair $(EL/L, \xi_L)$ is then admissible and*

$$L\pi_{\xi_L} = b_{L/F}(F\pi_\xi).$$

We shall prove this in the following paragraphs. We first use it to complete the proof of Theorem 4.4.

We proceed by induction on n . Let ℓ be the least prime divisor of n and let K/F be the unique sub-extension of E/F of degree ℓ . By the induction hypothesis, we have

$$\delta_{E/K} \cdot K\pi_\xi = \mathcal{L}(\text{Ind}_{E/K} \xi).$$

The character $\delta_{E/K}$ is unramified and hence of the form $\delta_{E/K} = \phi \circ N_{K/F}$, for some unramified character ϕ of F^\times , having the same order as $\delta_{E/K}$. In particular, $\phi^2 = 1$.

If K/F is cyclic, we have $\delta_{K/F} = 1$ and, by Proposition 4.5, $A_{K/F}(K\pi_\xi) = F\pi_\xi$ whence

$$\phi \cdot F\pi_\xi = A_{K/F}(\delta_{E/K} \cdot K\pi_\xi) = \mathcal{L}(\text{Ind}_{E/F} \xi).$$

However, $\delta_{E/F} = \delta_{K/F}^{n/\ell} \delta_{E/K} \mid F^\times = \phi^\ell = \phi$, as required for Theorem 4.4.

We therefore assume that K/F is not cyclic. We put $\sigma = \text{Ind}_{E/F} \xi$. We let L/F be the unramified extension obtained by adjoining to F a primitive ℓ -th root of unity. Let $k = [L:F]$; since $k < \ell$ and ℓ is the least prime divisor of n , we have $(k, n) = 1$. Applying Theorem 4.6, we get

$$(4.6.1) \quad L\pi_{\xi_L} = b_{L/F}(F\pi_\xi).$$

The extension LK/L is cyclic so, by the first case,

$$L\pi_{\xi_L} = \delta_{EL/L} \cdot \mathcal{L}(\sigma_L),$$

where $\sigma_L = \sigma \mid \mathcal{W}_L = \text{Ind}_{EL/L} \xi_L$. The character $\delta_{EL/L}$ is just $\delta_{E/F} \circ N_{L/F}$, so

$$(4.6.2) \quad L\pi_{\xi_L} = b_{L/F}(\delta_{E/F} \cdot \mathcal{L}(\sigma)).$$

Comparing (4.6.1), (4.6.2), we get $F\pi_\xi = \chi \delta_{E/F} \cdot \mathcal{L}(\sigma)$, for some unramified character χ of F^\times such that $\chi^k = 1$ [8, 2.6]. However, comparing central quasicharacters, we get

$$\xi = \chi^n \delta_{E/F}^n \det \sigma = \chi^n \delta_{E/F}^{n+1} \xi;$$

since n is odd, this reduces to $\chi^n = 1$. We also have $\chi^k = 1$ and $(k, n) = 1$, so $\chi = 1$, as required.

This completes the proof of Theorem 4.4. \square

4.7. We now prove Theorem 4.6. We abbreviate $\pi = {}_F\pi_\xi$, $\pi_L = {}_L\pi_{\xi_L}$ and $\Pi = {}_{b_{L/F}}\pi$. In this notation, we have to show that $\pi_L = \Pi$.

The endo-class $\Theta(\pi_L)$ is given by $\Theta(\pi_L) = R_{EL/L}(\xi_L | U_{E_L}^1)$ whence

$$(4.7.1) \quad R_{L/F}\Theta(\pi_L) = \Theta(\pi);$$

thus $\Theta(\pi_L)$ is an L/F -lift of $\Theta(\pi)$. (Indeed, it is the unique L/F -lift of $\Theta(\pi)$.) On the other hand, this equals $\Theta(\Pi)$, by [6, (7.1)]. The representations Π , π_L have the same central quasicharacter, namely $\omega_\pi \circ N_{L/F}$. Since E/F is totally ramified, we deduce:

Lemma 4.7. *There is an unramified character χ of L^\times such that $\chi^n = 1$ and $\pi_L = \chi\Pi$.*

We have to show that $\chi = 1$.

4.8. We pass to the group $G(L) = GL_n(L)$, and set $\Sigma = \text{Gal}(L/F)$. The pair $(LE/L, \xi_L)$ gives an extended maximal simple type $(\mathbf{J}_L, \Lambda_L)$ in π_L , which we now describe. Write $\mathfrak{A}_L = \mathfrak{A} \otimes_{\mathfrak{o}_F} \mathfrak{o}_L$; the quadruple $[\mathfrak{A}_L, l, 0, \beta]$ is then a simple stratum in $A(L) = M_n(L)$ and, setting $\psi_L = \psi_F \circ \text{Tr}_{L/F}$, there is a simple character $\theta_L \in \mathcal{C}(\mathfrak{A}_L, \beta, \psi_L)$ such that $\mathcal{E}_L(\theta_L)$ is the unique L/F -lift of $\mathcal{E}_F(\theta)$ [3, (11.2) *et seq.*]. We may therefore take $\mathbf{J}_L = EL^\times J^1(\beta, \mathfrak{A}_L)$ and assume that $\Lambda_L | J^1(\beta, \mathfrak{A}_L) = \eta_L$, the unique irreducible representation of $J^1(\beta, \mathfrak{A}_L)$ containing θ_L .

The character θ_L is Σ -stable (remark following [3, (11.3)]), and the same therefore applies to η_L . It follows (*cf.* Lemma 4.1) that Λ_L is Σ -stable.

We choose a generator σ of Σ . The quasicharacter ξ_L of LE^\times is Σ -stable, so we can extend it to a quasicharacter $\tilde{\xi}_L$ of $LE^\times \rtimes \Sigma$ by setting $\tilde{\xi}_L | \Sigma = 1$. Likewise, Λ_L admits an extension to a representation of $\mathbf{J}_L \rtimes \Sigma$.

Lemma 4.8. *Let $\tilde{\Lambda}_L$ be a representation of $\mathbf{J}_L \rtimes \Sigma$ which extends Λ_L . There is a non-zero constant ϵ'_L such that*

$$\text{tr } \tilde{\Lambda}_L(\varpi^r \zeta x \sigma) = \epsilon'_L \tilde{\xi}_L(\varpi^r \zeta x \sigma) = \epsilon'_L \xi_L(\varpi^r \zeta x),$$

for all integers r such that $(r, n) = 1$ and all $\zeta \in \boldsymbol{\mu}_F$, $x \in U_E^1$.

Proof. Set $J_L^1 = J^1(\beta, \mathfrak{A}_L)$ and $H_L^1 = H^1(\beta, \mathfrak{A}_L)$. Observe that the assertion to be proved only concerns the restriction of $\tilde{\Lambda}_L$ to $E^\times J_L^1$.

Write $\tilde{\theta}_L = \tilde{\xi}_L | H_L^1 \rtimes \Sigma$. There is a unique irreducible representation $\tilde{\eta}_L$ of $J_L^1 \rtimes \Sigma$ which restricts, on $H_L^1 \rtimes \Sigma$, to a multiple of $\tilde{\theta}_L$. This representation $\tilde{\eta}_L$ is inflated from a representation of the finite group $J_L^1 \rtimes \Sigma / \text{Ker } \theta_L$ of order dp^s , for some integer s . The group E^\times acts, by conjugation, as a cyclic group of automorphisms of order n , which is prime to dp^s . This action fixes $\tilde{\eta}_L$ and so gives rise to a Glauberman sign $\epsilon_L = \pm 1$. Thus there exists a unique representation $\overline{\Lambda}_L$ of $E^\times J_L^1 \rtimes \Sigma$ such that

$$\text{tr } \overline{\Lambda}_L(\varpi^r \zeta x \sigma) = \epsilon_L \tilde{\xi}_L(\varpi^r \zeta x \sigma) = \epsilon_L \xi_L(\varpi^r \zeta x),$$

for $(r, n) = 1$, $\zeta \in \boldsymbol{\mu}_F$, $x \in U_E^1$.

Finally, we note that $\tilde{\Lambda}_L | E^\times J_L^1 \rtimes \Sigma$ is of the form $\phi \otimes \overline{\Lambda}_L$, for some character ϕ of Σ . The result follows, with $\epsilon'_L = \epsilon_L \phi(\sigma)$. \square

We put

$$\tilde{\pi}_L = c\text{-Ind}_{\mathbf{J}_L \rtimes \Sigma}^{G(L) \rtimes \Sigma} \tilde{\Lambda}_L.$$

This is an irreducible representation of $G(L) \rtimes \Sigma$ extending π_L .

Again we say that an element h of \mathbf{J}_L is $G(L)$ -special if $v_L(\det h)$ is relatively prime to n .

Proposition 4.8. *Let $h \in \mathbf{J}_L$ be $G(L)$ -special. Then*

$$\mathrm{tr} \tilde{\pi}_L(h\sigma) = \epsilon_F \epsilon'_L \mathrm{tr} \pi(\mathcal{N}_\sigma h),$$

where $\mathcal{N}_\sigma h$ is the σ -norm of h in G .

Proof. Let $h \in \mathbf{J}_L$ be $G(L)$ -special. We have $v_F(\det \mathcal{N}_\sigma h) = dv_L(\det h)$; since $(d, n) = 1$, the element $\mathcal{N}_\sigma h$ is G -special.

We write $J_L^1 = J^1(\beta, \mathfrak{A}_L)$. We have $h = jh_1$, where $h_1 \in LE^\times$ and $j \in J_L^1$. The element $h\sigma$ acts on J_L^1 , by conjugation, as an automorphism of finite order relatively prime to p . It follows that $h\sigma$ is J_L^1 -conjugate to an element $j_1 h_1 \sigma$, where $j_1 \in J_L^1$ commutes with $h_1 \sigma$. That is, $h\sigma$ is J_L^1 -conjugate to an element of $LE^\times \sigma$. If $h \in LE^\times$, we have $\mathcal{N}_\sigma h = N_{LE/E}(h)$ so $\mathcal{N}_\sigma(h)$ is an elliptic regular, G -special element of G .

Let $h, h' \in LE^\times$ be $G(L)$ -special, and suppose that $h'\sigma = x^{-1}h\sigma x$, for some $x \in G(L)$. The σ -norms of h, h' are then conjugate by the same element x , whence $N_{LE/E}(h') = x^{-1}N_{LE/E}(h)x$. Each of these norms generates E/F . We conclude that conjugation by x is an L -automorphism of LE , which we now denote by α . This satisfies $\alpha(\varpi) = \zeta\varpi$, for some n -th root of unity ζ in L .

We consider the element $x^{-1}h\sigma x = \alpha(h)x^{-1}\sigma x$. Conjugation by the element $x^{-1}\sigma x\sigma^{-1}$ is an L -automorphism of LE mapping ϖ to $\sigma^{-1}(\zeta^{-1})\zeta\varpi$. By definition, we have $\alpha(h)x^{-1}\sigma x\sigma^{-1} = h'$, so $x^{-1}\sigma x\sigma^{-1} \in \mathbf{J}_L$. The automorphism of LE induced by $x^{-1}\sigma x\sigma^{-1}$ is therefore trivial, so σ commutes with α and therefore $\sigma(\zeta) = \zeta$. It follows that $\alpha \in \mathrm{Aut}(EL/L) \cong \mathrm{Aut}(E|F)$. Therefore, if $h \in LE^\times$ is $G(L)$ -special, we have

$$\begin{aligned} \mathrm{tr} \tilde{\pi}(h\sigma) &= \epsilon'_L \sum_{\alpha \in \mathrm{Aut}(E|F)} \xi_L(\alpha(h)) \\ &= \epsilon'_L \sum_{\alpha \in \mathrm{Aut}(E|F)} \xi(\alpha(N_{LE/E}(h))) \\ &= \epsilon'_F \epsilon_L \mathrm{tr} \pi(\mathcal{N}_\sigma h), \end{aligned}$$

by Proposition 4.1, and the result follows. □

4.9. We return to the representation $\Pi = b_{L/F} \pi = \chi\pi_L$, where χ is unramified and $\chi^n = 1$. We extend Π to a representation $\tilde{\Pi}$ of $G(L) \rtimes \Sigma$; there is then a non-zero constant c such that

$$\mathrm{tr} \tilde{\Pi}(h\sigma) = c \mathrm{tr} \pi(\mathcal{N}_\sigma h),$$

for all $h \in G(L)$ such that $\mathcal{N}_\sigma h$ is a quasi-regular element of G [3, Appendix]. In particular, if $h \in LE^\times$ is $G(L)$ -special, Proposition 4.8 gives

$$\epsilon_F \epsilon'_L \mathrm{tr} \tilde{\pi}_L(h\sigma) = \mathrm{tr} \pi(\mathcal{N}_\sigma h) = c^{-1} \mathrm{tr} \tilde{\Pi}(h\sigma) = c^{-1} \chi(\det h) \mathrm{tr} \tilde{\pi}_L(h\sigma).$$

In other words, there is a constant c' such that

$$\chi(\det h) \mathrm{tr} \tilde{\pi}_L(h\sigma) = c' \mathrm{tr} \tilde{\pi}_L(h\sigma),$$

for all $G(L)$ -special elements h of LE^\times . We now write $\chi = \phi \circ N_{L/F}$, for some unramified character ϕ of F^\times satisfying $\phi^n = 1$ (as we may, since $(d, n) = 1$). Invoking Proposition 4.8 again, there is a constant c'' such that

$$\phi(\det x) \operatorname{tr} \pi(x) = c'' \operatorname{tr} \pi(x),$$

for all G -special $x \in E^\times$ with $v_F(\det x) \equiv 0 \pmod{d}$.

The representation π is totally ramified so, given $a \in \mathbb{Z}$, there exists $g \in G$ with $v_F(\det g) = a$ and $\operatorname{tr} \pi(g) \neq 0$. By Proposition 4.1, if $(a, n) = 1$, we can take $g \in E^\times$. Therefore ϕ must be constant on $(F^\times)^d U_F$, whence $\phi^d = 1$. Since $\phi^n = 1$ and $(n, d) = 1$, we have $\phi = 1$, as required.

This completes the proof of Theorem 4.6. \square

APPENDIX. CHARACTERS OF TAME EXTENSIONS

We gather here some elementary results concerning quasicharacters of finite, tamely ramified extensions of the base field F . While these are, broadly speaking, well known, they are also widely forgotten. They are scattered in literature which is now rather old, often in the wrong degree of generality. It seems more satisfactory, therefore, to collect them in this brief appendix.

For our present purposes, we define a representation $\sigma \in \mathfrak{S}_n^0(F)$ to be *essentially tame* if its restriction to the wild inertia subgroup of \mathcal{W}_F is a sum of characters.

A.1. Let E/F be a finite, Galois, tamely ramified field extension and put $\Gamma = \operatorname{Gal}(E/F)$. The basic fact, on which all depends, is that the Γ -modules U_E^n , $n \geq 1$, are *cohomologically trivial*. The same applies to U_E when E/F is unramified. (See [26] for a comprehensive account of such matters.) Consequently:

- Lemma.** (1) *Let E/F be a finite, Galois, tamely ramified field extension, and let K/F be a sub-extension with $\Delta = \operatorname{Gal}(E/K)$. Let χ be a character of U_E^{n+1} with $n \geq 0$ (or $n \geq -1$ if E/K is unramified). Then χ factors through $N_{E/K}$ if and only if $\chi^\delta = \chi$, for all $\delta \in \Delta$.*
- (2) *Suppose, in (1), that $\chi \mid U_E^{n+1}$ factors through $N_{E/K}$. Let $e = e(E|K)$ and $m = [n/e]$, so that $U_E^{n+1} \cap K = U_K^{m+1} = N_{E/K}(U_E^{n+1})$. There is then a unique character η of U_K^{m+1} such that $\chi \mid U_E^{n+1} = \eta \circ N_{E/K}$.*
- (3) *Let E/F be a finite, tamely ramified field extension and let χ be a character of U_E^n , for some $n \geq 0$. There exists a unique sub-extension E_0/F of E/F such that χ factors through N_{E/E_0} and which is minimal for this property.*

In (3), the field E_0 arises as follows. Let L/F be the normal closure of E/F so that, in particular, L/E is unramified. Set $\chi_L = \chi \circ N_{L/E}$, and let Δ be the subgroup of $\operatorname{Gal}(L/F)$ which fixes $\chi_L \mid U_L^n$. Then $E_0 = L^\Delta$.

A.2. We return to the notion of “admissible pair”, defined in the Introduction. The first property of these pairs is:

Proposition. *Let $(E/F, \xi)$ be an admissible pair. Then $\operatorname{Ind}_{E/F} \xi$ is irreducible and essentially tame.*

Proof. Let L/F be the normal closure of E/F . In particular, L/E is unramified. We put $\Gamma = \operatorname{Gal}(L/F)$, $\Delta = \operatorname{Gal}(L/E)$.

Suppose, for a contradiction, that $\operatorname{Ind}_{E/F} \xi$ is reducible. By Mackey’s criterion, therefore, there exists $g \in \Gamma$, $g \notin \Delta$, such that ξ and ξ^g have the same restriction

to \mathcal{W}_L . This says that the quasicharacter $\xi \circ N_{L/E}$ of L^\times is invariant under the subgroup $\Omega = \langle \Delta, g \rangle$ of Γ generated by Δ and this element g . The restriction of ξ to U_E^1 therefore factors through $N_{E/K}$, where $K = L^\Omega$. Since $(E/F, \xi)$ is admissible, the extension E/K is unramified. Therefore L/K is unramified. The group Ω therefore stabilizes E and, by definition, it stabilizes the quasicharacter $\xi \circ N_{L/E}$ of L^\times . This quasicharacter therefore factors through $N_{L/K}$ and so ξ factors through $N_{E/K}$. Since $(E/F, \xi)$ is admissible, we have $K = E$ and the desired contradiction. The final assertion is immediate. \square

A.3. Let $(E/F, \xi)$ be an admissible pair; the equivalence class of the representation $\sigma = \text{Ind}_{E/F} \xi$ depends only on the F -isomorphism class of $(E/F, \xi)$ and σ is surely essentially tame. Thus we have a canonical map

$$(A.3.1) \quad \begin{aligned} P_n(F) &\longrightarrow \mathfrak{G}_n^{\text{et}}(F), \\ (K/F, \xi) &\longmapsto \text{Ind}_{K/F} \xi, \end{aligned}$$

for every $n \geq 1$.

Theorem. *For every $n \geq 1$, the map (A.3.1) is a bijection.*

Proof. Let $\sigma \in \mathfrak{G}_n^{\text{et}}(F)$ act on the vector space V . Let L/F be a finite, tamely ramified, Galois extension such that $\sigma|_{\mathcal{W}_L}$ is a direct sum of quasicharacters. Let η be a component of $\sigma|_{\mathcal{W}_L}$, viewed as a quasicharacter of L^\times , and let Δ be the subgroup of $\text{Gal}(L/F)$ which fixes $\eta|_{U_L^1}$. Set $K = L^\Delta$. The implied representation τ of \mathcal{W}_K on the isotypic space V^η is irreducible and, by Clifford theory, $\sigma = \text{Ind}_{K/F} \tau$. On the other hand, there is a quasicharacter ϕ of K^\times such that $\phi \circ N_{L/K}$ agrees with η on U_K^1 , so $\tau \cong \phi \otimes \rho$, for some irreducible representation ρ of \mathcal{W}_K trivial on \mathcal{W}_L . Thus ρ can be viewed as an irreducible representation of $\Delta \cong \mathcal{W}_K/\mathcal{W}_L$. The Galois extension L/K is tamely ramified, so ρ is of the form $\rho = \text{Ind}_{E/K} \mu$, where E/K is an unramified extension in L and μ is a tamely ramified quasicharacter of E^\times . We put $\xi = \mu \cdot (\phi \circ N_{E/K})$ to get $\sigma = \text{Ind}_{E/F} \xi$.

We next show that the pair $(E/F, \xi)$ is admissible. Clearly, condition (1) in the definition is satisfied, since $\text{Ind}_{E/F} \xi = \sigma$ is irreducible. Let K' be an extension of F in E and ϕ' a character of $U_{K'}^1$ such that $\phi' \circ N_{E/K'} = \xi|_{U_E^1}$. The group $\text{Gal}(L/K')$ therefore stabilizes $\xi \circ N_{L/E}|_{U_L^1} = \eta|_{U_L^1}$, so $K' \supset K$ whence E/K is unramified. Thus $(E/F, \xi)$ is admissible and the map (A.3.1) is surjective.

Finally, we have to show that (A.3.1) is injective. With σ and $(E/F, \xi)$ as before, suppose we have another admissible pair $(E'/F, \xi')$ such that $\text{Ind}_{E'/F} \xi' = \sigma$. Enlarging L if need be, we can assume that both E and E' are contained in L . The quasicharacters $\xi \circ N_{L/E}$, $\xi' \circ N_{L/E'}$ of L^\times are then conjugate under $\text{Gal}(L/F)$. Replacing $(E'/F, \xi')$ by an F -isomorphic pair, we can assume that $\xi' \circ N_{L/E'} = \xi \circ N_{L/E}$. Let Δ be the subgroup of $\text{Gal}(L/F)$ fixing these quasicharacters on U_L^1 and set $K = L^\Delta$: this is indeed the same as the field K attached to $\sigma = \text{Ind}_{E/F} \xi$ in the first paragraph of the proof. Thus $E' \supset K$ and, by admissibility, E'/K is unramified. The fields E, E' are both unramified extensions of K inside L , and of the same degree $n/[K:F]$. Therefore $E = E'$, and we may assume again that L/F is the Galois closure of $E = E'$ over F . Applying the analysis of the first paragraph to $\sigma = \text{Ind}_{E/F} \xi$, we get

$$\tau = \text{Ind}_{E/K} \xi = \text{Ind}_{E/K} \xi'.$$

Since E/K is cyclic, this implies that the quasicharacters ξ, ξ' of E^\times are conjugate under $\text{Gal}(E/K)$, whence the pairs $(E/F, \xi), (E'/F, \xi')$ are F -isomorphic. Thus (A.3.1) is injective, and we have finished the proof. \square

A.4. Let $\sigma \in \mathfrak{G}_n^0(F)$; we define $T(\sigma)$ to be the group of unramified characters χ of F^\times such that $\chi \otimes \sigma \cong \sigma$. Since $\det(\chi \otimes \sigma) = \chi^n \cdot \det \sigma$, the group $T(\sigma)$ is cyclic of order $t(\sigma)$ dividing n .

Proposition. *Let $(E/F, \xi) \in P_n(F)$ and let $\sigma = \text{Ind}_{E/F} \xi$. Then $t(\sigma) = f(E|F)$.*

Proof. If χ is an unramified character of F^\times , let us abbreviate $\chi \circ N_{E/F} = \chi_E$. Then $\chi \otimes \sigma \cong \text{Ind}_{E/F} \xi'$, where $\xi' = \xi \cdot \chi_E$. If $\chi^{f(E|F)} = 1$, then $\chi_E = 1$, so we deduce that $f(E|F)$ divides $t(\sigma)$.

In the opposite direction, let $\chi \in T(\sigma)$. The pair $(E/F, \chi_E \xi)$ is surely admissible, and so F -isomorphic to $(E/F, \xi)$. That is, there exists $\alpha \in \text{Aut}(E|F)$ such that $\chi_E \xi = \xi^\alpha$. Since χ_E is unramified, this implies that α fixes $\xi | U_E$. If K is the fixed field of α , this implies that $\xi | U_E^1$ factors through $N_{E/K}$, whence E/K is unramified. Now we can deduce that $\xi | U_E$ factors through $N_{E/K}$ as well. Any quasicharacter of E^\times agreeing with ξ on U_E therefore factors through $N_{E/K}$. In particular, this applies to ξ . By admissibility, $\alpha = 1$, so $\chi_E = 1$, and $\chi^{f(E|F)} = 1$, as required. \square

Corollary. *Let $\sigma \in \mathfrak{G}_n^0(F)$. The following conditions are equivalent:*

- (1) σ is essentially tame;
- (2) p does not divide $n/t(\sigma)$.

Proof. If σ is essentially tame, it is of the form $\text{Ind}_{E/F} \xi$, for some $(E/F, \xi) \in P_n(F)$, and so the implication (1) \Rightarrow (2) follows directly from the proposition. Conversely, suppose $p \nmid n/t(\sigma)$. If E/F is unramified of degree $t(\sigma)$, there is an irreducible representation τ of \mathcal{W}_E such that $\sigma \cong \text{Ind}_{E/F} \tau$. The dimension of τ is prime to p so, when we restrict τ to the pro- p group \mathcal{P}_F , it becomes a sum of characters. The same therefore applies to σ , which is therefore essentially tame. \square

REFERENCES

1. J. Arthur and L. Clozel, *Simple algebras, base change, and the advanced theory of the trace formula*, Annals of Math. Studies, vol. 120, Princeton University Press, 1989. MR1007299 (90m:22041)
2. C.J. Bushnell and A. Fröhlich, *Gauss sums and p -adic division algebras*, Lecture Notes in Math., vol. 987, Springer, Berlin, Heidelberg, New York, 1983. MR0701540 (84m:12017)
3. C.J. Bushnell and G. Henniart, *Local tame lifting for $GL(N)$ I: simple characters*, Publ. Math. IHES **83** (1996), 105–233. MR1423022 (98m:11129)
4. ———, *Local tame lifting for $GL(n)$ II: wildly ramified supercuspidals*, Astérisque **254** (1999). MR1685898 (2000d:11147)
5. ———, *Davenport-Hasse relations and an explicit Langlands correspondence*, J. reine angew. Math. **519** (2000), 171–199. MR1739725 (2002e:11171)
6. ———, *Local tame lifting for $GL(n)$ IV: simple characters and base change*, Proc. London Math. Soc. (3) **87** (2003), 337–362. MR1990931 (2004f:22017)
7. ———, *The essentially tame local Langlands correspondence, II: totally ramified representations*, Compositio Mathematica, to appear.
8. C.J. Bushnell, G. Henniart and P.C. Kutzko, *Correspondance de Langlands locale pour GL_n et conducteurs de paires*, Ann. Scient. École Norm. Sup. (4) **31** (1998), 537–560. MR1634095 (99h:22011)
9. C.J. Bushnell and P.C. Kutzko, *The admissible dual of $GL(N)$ via compact open subgroups*, Annals of Math. Studies, vol. 129, Princeton University Press, 1993. MR1204652 (94h:22007)

10. ———, *The admissible dual of $SL(N)$ II*, Proc. London Math. Soc. (3) **68** (1994), 317–379. MR1253507 (94k:22035)
11. L.J. Corwin and R.E. Howe, *Computing characters of tamely ramified p -adic division algebras*, Pacific J. Math. **73** (1977), 461–477. MR0492084 (58:11238)
12. G. Glauberman, *Correspondences of characters for relatively prime operator groups*, Canad. J. Math. **20** (1968), 1465–1488. MR0232866 (38:1189)
13. J.A. Green, *The characters of the finite general linear groups*, Trans. Amer. Math. Soc. **80** (1955), 402–447. MR0072878 (17:345e)
14. M. Harris and R. Taylor, *On the geometry and cohomology of some simple Shimura varieties*, Annals of Math. Studies, vol. 151, Princeton University Press, 2001. MR1876802 (2002m:11050)
15. G. Henniart, *Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p -adique*, Invent. Math. **139** (2000), 439–455. MR1738446 (2001e:11052)
16. G. Henniart and R. Herb, *Automorphic induction for $GL(n)$ (over local non-archimedean fields)*, Duke Math. J. **78** (1995), 131–192. MR1328755 (96i:22038)
17. G. Henniart and B. Lemaire, *Work in progress*.
18. R.E. Howe, *Tamely ramified supercuspidal representations of GL_n* , Pacific J. Math. **73** (1977), 437–460. MR0492087 (58:11241)
19. H. Jacquet, I. Piatetski-Shapiro and J. Shalika, *Rankin-Selberg convolutions*, Amer. J. Math. **105** (1983), 367–483. MR0701565 (85g:11044)
20. H. Koch and E.-W. Zink, *Zur Korrespondenz von Darstellungen der Galoisgruppen und der zentralen Divisionsalgebren über lokalen Körpern (der zahme Fall)*, Math. Nachr. **98** (1980), 83–119. MR0623696 (83h:12025)
21. G. Laumon, M. Rapoport and U. Stuhler, *\mathcal{D} -elliptic sheaves and the Langlands correspondence*, Invent. Math. **113** (1993), 217–338. MR1228127 (96e:11077)
22. C. Mœglin, *Sur la correspondance de Langlands-Kazhdan*, J. Math. Pures et Appl. (9) **69** (1990), 175–226. MR1067450 (91g:11141)
23. A. Moy, *Local constants and the tame Langlands correspondence*, Amer. J. Math. **108** (1986), 863–930. MR0853218 (88b:11081)
24. H. Reimann, *Representations of tamely ramified p -adic division and matrix algebras*, J. Number Theory **38** (1991), 58–105. MR1105671 (92h:11103)
25. J. Rogawski, *Representations of $GL(n)$ and division algebras over a local field*, Duke Math. J. **50** (1983), 161–196. MR0700135 (84j:12018)
26. J.-P. Serre, *Corps locaux*, Hermann, Paris, 1965. MR0150130 (27:133)
27. F. Shahidi, *Fourier transforms of intertwining operators and Plancherel measures for $GL(n)$* , Amer. J. Math. **106** (1984), 67–111. MR0729755 (86b:22031)

DEPARTMENT OF MATHEMATICS, KING'S COLLEGE LONDON, STRAND, LONDON WC2R 2LS,
UNITED KINGDOM

E-mail address: bushnell@mth.kcl.ac.uk

DÉPARTEMENT DE MATHÉMATIQUES & UMR 8628 DU CNRS, BÂTIMENT 425, UNIVERSITÉ DE
PARIS-SUD, 91405 ORSAY CEDEX, FRANCE

E-mail address: Guy.Henniart@math.u-psud.fr