

DIVISIBILITY OF THE STABLE MILLER-MORITA-MUMFORD CLASSES

SOREN GALATIUS, IB MADSEN, AND ULRIKE TILLMANN

1. INTRODUCTION AND RESULTS

Let $\Gamma_{g,b}^s$ denote the mapping class group of a surface of genus g with b ordered boundary components and s marked points. We will suppress s or b when their value is zero. Gluing a disk or a torus with two boundary components to one of the boundary components induces homomorphisms

$$(1.1) \quad \Gamma_{g,b-1}^s \longleftarrow \Gamma_{g,b}^s \longrightarrow \Gamma_{g+1,b}^s.$$

Recall that by Harer-Ivanov's stability theory both homomorphisms induce a homology isomorphism in dimensions $*$ with $2* + 1 < g$; cf. [H2], [I]. Let $\Gamma_\infty := \lim_{g \rightarrow \infty} \Gamma_{g,2}$ be the stable mapping class group.

Mumford in [Mu] introduced certain tautological classes in the cohomology of moduli spaces of Riemann surfaces. Miller [Mi] and Morita [Mo] studied topological analogues. Let $e \in H^2(B\Gamma_{g,b}^1; \mathbb{Z})$ be the Euler class of the central extension

$$(1.2) \quad \mathbb{Z} \longrightarrow \Gamma_{g,b+1} \longrightarrow \Gamma_{g,b}^1$$

which is induced by gluing a disk with a marked point to one of the boundary components. Define

$$\kappa_i := \pi_!(e^{i+1}) \in H^{2i}(B\Gamma_{g,b}; \mathbb{Z})$$

where $\pi_!$ is the Umkehr (or integration along the fibre) map associated to the forgetful map $\Gamma_{g,b}^1 \rightarrow \Gamma_{g,b}$. These correspond under the maps of (1.1) when $i > 0$ and hence define classes in $H^*(B\Gamma_\infty; \mathbb{Z})$. We will only be concerned with these stable classes in this paper.

By the proof of the Mumford conjecture [MW],

$$H^*(B\Gamma_\infty; \mathbb{Q}) \simeq \mathbb{Q}[\kappa_1, \kappa_2, \dots].$$

In contrast, little is known about κ_i in integral cohomology though it follows from [H1] that κ_1 is precisely divisible by 12 (cf. [MT, p. 537]). We write

$$H_{free}^*(B\Gamma_\infty) := H^*(B\Gamma_\infty; \mathbb{Z})/\text{Torsion}$$

for the integral lattice in $H^*(B\Gamma_\infty; \mathbb{Q})$.

Received by the editors October 31, 2004.

2000 *Mathematics Subject Classification*. Primary 57R20, 55P47.

Key words and phrases. Mapping class group, characteristic classes, surface bundles.

The third author was supported by an Advanced Fellowship of the EPSRC.

©2006 American Mathematical Society
 Reverts to public domain 28 years from publication

Theorem 1.1. *Let D_i be the maximal divisor of κ_i in $H_{free}^*(B\Gamma_\infty)$. Then for all $i \geq 1$,*

$$D_{2i} = 2 \quad \text{and} \quad D_{2i-1} = \text{den}\left(\frac{B_i}{2i}\right).$$

Here B_i denotes the i -th Bernoulli number and den is the function that takes a rational number when expressed as a fraction in its lowest terms to its denominator. It is well known, cf. [MSt, Appendix B], that $\text{den}(B_i)$ is the product of all primes p such that $p - 1$ divides $2i$, and that a prime divides $\text{den}(B_i/2i)$ if and only if it divides $\text{den}(B_i)$. So in terms of their p -adic valuation the D_i are determined by the formula

$$(1.3) \quad \nu_p(D_i) = \begin{cases} 1 + \nu_p(i + 1) & \text{if } i + 1 \equiv 0 \pmod{p - 1} \\ 0 & \text{if } i + 1 \not\equiv 0 \pmod{p - 1}, \end{cases}$$

and $D_1 = 2^2 \cdot 3$, $D_3 = 2^3 \cdot 3 \cdot 5$, $D_5 = 2^2 \cdot 3^2 \cdot 7$, \dots

Our Theorem 1.1 is inspired by a conjecture of T. Akita [Ak] which we also prove:

Theorem 1.2. *The element κ_i in $H^{2i}(B\Gamma_\infty; \mathbb{F}_p)$ vanishes if and only if $i + 1 \equiv 0 \pmod{p - 1}$.*

Remark 1.3. The divisor D_i of κ_i in $H_{free}^*(B\Gamma_\infty)$ is not necessarily equal to the maximal divisor $D_i^{\mathbb{Z}}$ of κ_i in integral cohomology $H^*(B\Gamma_\infty; \mathbb{Z})$ but only provides an upper bound for it. However, Theorem 1.2 gives

$$p \text{ divides } D_i^{\mathbb{Z}} \iff p \text{ divides } D_i,$$

which was strengthened by the first author in [G2] to

$$p^2 \text{ divides } D_i^{\mathbb{Z}} \iff p^2 \text{ divides } D_i.$$

It follows that for all even i and for many odd i ($i = 1, 5, 9, 13, \dots$), D_i is indeed equal to $D_i^{\mathbb{Z}}$, and one may expect that $D_i = D_i^{\mathbb{Z}}$ for all $i \geq 1$.

Remark 1.4. The integral lattice $H_{free}^*(B\Gamma_\infty)$ inherits a Hopf algebra structure. The graded module of primitive elements $P(H_{free}^*(B\Gamma_\infty))$ is a copy of \mathbb{Z} in each even degree, and κ_i is a primitive element of $H_{free}^*(B\Gamma_\infty)$. The structure of the Hopf algebra $H_{free}^*(B\Gamma_\infty)$ is not completely understood at present, but we have the following partial results.

Theorem 1.5. *For odd primes p there is an isomorphism of Hopf algebras over the p -local integers $\mathbb{Z}_{(p)}$:*

$$H_{free}^*(B\Gamma_\infty; \mathbb{Z}_{(p)}) \simeq H^*(BU; \mathbb{Z}_{(p)}).$$

This fails for $p = 2$. In fact, the squaring map

$$\xi : H_{free}^2(B\Gamma_\infty) \otimes \mathbb{F}_2 \rightarrow H_{free}^4(B\Gamma_\infty) \otimes \mathbb{F}_2$$

is not injective, so the algebra $H_{free}^(B\Gamma_\infty; \mathbb{Z}_{(2)})$ is not polynomial.*

In outline, the proofs of the above theorems depend on previous results as follows. For Theorem 1.2, the proof of the “if” part in Section 3.3 is a calculation of characteristic classes which relies on the fact that there is a map of infinite loop spaces $\alpha : \mathbb{Z} \times B\Gamma_\infty^+ \rightarrow \Omega^\infty \mathbb{C}P_{-1}^\infty$ (compare [T] and [MT] or Theorems 2.1 and 2.2 below). The “only if” part is implied by Theorem 1.1: if p divides κ_i , then in particular it must divide its reduction to the free part.

For Theorem 1.1, we first establish a lower bound: $D_{2i} \geq 2$ by the “if” part of Theorem 1.2 and $D_{2i-1} \geq \text{den}(B_i/2i)$ by a well-known relation between the κ_i classes and the symplectic characteristic classes for surface bundles (here stated as Theorem 4.2). The main theorem of [MT] provides an upper bound which is tight for i even and precisely twice the lower bound for i odd. To eliminate the indeterminacy of the factor 2, the main theorem of [MW] (Theorem 2.4 below), as well as calculations from [G1] (Proposition 4.4) and a stronger version of the main result of [MT] (given in Theorem 2.2 and proved in Section 5) are used.

Theorem 1.5 is proved in Section 6.

Given the interest in the mapping class groups also outside the topology community we have strived to make this paper as self-contained as possible. In particular we have spelled out some of the more obscure parts of [MT].

2. SPECTRUM COHOMOLOGY AND EARLIER RESULTS

2.1. Spectra and spectrum cohomology. Let $E = \{E_n, \epsilon_n\}$ be a CW-spectrum¹ in the sense of [A1]: E_n is a sequence of pointed CW-complexes and $\epsilon_n : SE_n \rightarrow E_{n+1}$ a (pointed) isomorphism onto a subcomplex, where $S(-)$ denotes suspension. The associated infinite loop space is the direct limit

$$\Omega^\infty E = \text{colim } \Omega^n E_n$$

of the n -th loop space of E_n ; the limit is taken over the adjoint maps $\epsilon'_n : E_n \rightarrow \Omega E_{n+1}$.

The k -th homotopy group of E is defined to be the direct limit of $\pi_{n+k}(E_n)$. It is equal to the k -th homotopy group of the space $\Omega^\infty E$. In particular, the group of components of $\Omega^\infty E$ is the direct limit of $\pi_n(E_n)$. For $\alpha \in \pi_0(\Omega^\infty E)$ we let $\Omega_\alpha^\infty E$ be the component determined by α . In particular we write $\Omega_0^\infty E$ for the component of the zero element.

The homology and cohomology groups of E are

$$H^k(E) = \lim_{n \rightarrow \infty} \tilde{H}^{k+n}(E_n), \quad H_k(E) = \text{colim}_{n \rightarrow \infty} \tilde{H}_{k+n}(E_n),$$

where the limits are induced from the maps ϵ_n together with the suspension isomorphisms.² In contrast to homotopy groups the cohomology groups of a spectrum are usually much simpler than the cohomology groups of $\Omega^\infty E$.

The evident evaluation map from $S^n \Omega^n E_n$ to E_n induces maps

$$(2.1) \quad \sigma^* : H^*(E) \longrightarrow \tilde{H}^*(\Omega_0^\infty E), \quad \sigma_* : \tilde{H}_*(\Omega_0^\infty E) \longrightarrow H_*(E).$$

If we use field coefficients in the cohomology groups, then $H^*(\Omega_0^\infty E)$ is a connected Hopf algebra and the image of σ^* is contained in the graded vector space $PH^*(\Omega_0^\infty E)$ of primitive elements. We shall be particularly concerned with the torsion free integral homology and cohomology groups

$$H_{free}^*(\Omega_0^\infty E) = H^*(\Omega_0^\infty E; \mathbb{Z})/\text{Torsion}, \quad H_*^{free}(\Omega_0^\infty E) = H_*(\Omega_0^\infty E; \mathbb{Z})/\text{Torsion}.$$

¹If one does not assume the spaces to be CW-complexes, then one should assume that ϵ_n is a closed cofibration.

²The k -th spectrum cohomology of E is normally defined as the group of homotopy classes of degree k spectrum maps from E to the Eilenberg-Mac Lane spectrum $K(\mathbb{Z})$. This coincides with the above formula for $H^k(E)$ whenever for all k the inverse system $\lim_{n \rightarrow \infty} \tilde{H}^{k+n}(E_n)$ satisfies the Mittag-Löffler condition (see [A1]). In particular this will be the case when the structure maps $\epsilon : SE_n \rightarrow E_{n+1}$ are $c(n)$ -connected for some function $c(n)$ with $c(n) \rightarrow \infty$ as $n \rightarrow \infty$.

They are lattices in $H^*(\Omega_0^\infty E; \mathbb{Q})$ and $H_*(\Omega_0^\infty E; \mathbb{Q})$ and are dual Hopf algebras. Moreover, the image of σ^* is contained in the module of primitive elements

$$\sigma^* : H_{free}^*(E) \longrightarrow P(H_{free}^*(\Omega_0^\infty E)),$$

and dually σ_* factors over the indecomposable elements of $H_*^{free}(\Omega_0^\infty E)$.

Given a pointed space X we have the associated suspension spectrum $S^\infty X$ whose n -th term is $S^n X$ with infinite loop space $\Omega^\infty S^\infty X$. There is an obvious inclusion $i : X \rightarrow \Omega^\infty S^\infty X$ inducing a splitting of σ^* (and σ_*):

$$(2.2) \quad H^*(S^\infty X) \xrightarrow{\sigma^*} \tilde{H}^*(\Omega_0^\infty S^\infty X) \xrightarrow{i^*} \tilde{H}^*(X)$$

is the suspension isomorphism. $\Omega^\infty S^\infty X$ is the free infinite loop space on X and satisfies the universal property that any pointed map from X to some infinite loop space Y can be extended in a unique way up to homotopy to a map of infinite loop spaces from $\Omega^\infty S^\infty X$ to Y .

The spectra of most relevance to us are $\mathbb{C}P_{-1}^\infty$ and the suspension spectrum $S^\infty \mathbb{C}P_+^\infty$ of $\mathbb{C}P^\infty \sqcup \{+\}$. We recall the definition of the former. There are two complex vector bundles over the complex projective n -space $\mathbb{C}P^n$, namely the tautological line bundle L_n and its n -dimensional complement L_n^\perp in $\mathbb{C}P^n \times \mathbb{C}^{n+1}$. Its Thom space (or one point compactification) is denoted by $\text{Th}(L_n^\perp)$. Since the restriction of L_n^\perp to $\mathbb{C}P^{n-1} \subset \mathbb{C}P^n$ is equal to $L_{n-1}^\perp \oplus \mathbb{C}$, where \mathbb{C} denotes the trivial line bundle over $\mathbb{C}P^{n-1}$, we get a map

$$\epsilon : S^2 \text{Th}(L_{n-1}^\perp) \longrightarrow \text{Th}(L_n^\perp).$$

The spectrum $\mathbb{C}P_{-1}^\infty$ has

$$(\mathbb{C}P_{-1}^\infty)_{2n} = \text{Th}(L_{n-1}^\perp), \quad (\mathbb{C}P_{-1}^\infty)_{2n+1} = S \text{Th}(L_{n-1}^\perp)$$

and the structure map ϵ_{2n+1} is given by the above ϵ . The associated infinite loop space is

$$\Omega^\infty \mathbb{C}P_{-1}^\infty = \text{colim}_{n \rightarrow \infty} \Omega^{2n} \text{Th}(L_{n-1}^\perp).$$

The inclusion of L_{n-1}^\perp into $L_{n-1}^\perp \oplus L_{n-1} = \mathbb{C}P^{n-1} \times \mathbb{C}^n$ via the zero section of L_{n-1} induces a map from $\text{Th}(L_{n-1}^\perp)$ into $S^{2n}(\mathbb{C}P_+^{n-1})$ and hence a map

$$\omega : \Omega^\infty \mathbb{C}P_{-1}^\infty \longrightarrow \Omega^\infty S^\infty(\mathbb{C}P_+^\infty).$$

This map fits into a fibration sequence

$$(2.3) \quad \Omega^\infty \mathbb{C}P_{-1}^\infty \xrightarrow{\omega} \Omega^\infty S^\infty(\mathbb{C}P_+^\infty) \xrightarrow{\partial} \Omega^\infty S^{\infty-1}$$

where the right-hand term is the direct limit of $\Omega^n S^{n-1}$ [R]. Indeed the inclusion of a fibre $\mathbb{C}^n \rightarrow L_n^\perp$ induces a map $S^{2n} \rightarrow \text{Th}(L_n^\perp)$ and gives rise to a cofibre sequence of spectra $S^\infty(S^{-2}) \rightarrow \mathbb{C}P_{-1}^\infty \rightarrow S^\infty \mathbb{C}P_+^\infty \rightarrow S^\infty(S^{-1})$. (2.3) is the associated fibration sequence of infinite loop spaces.

The component groups of (2.3) are

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\pi_0(\omega)} \mathbb{Z} \xrightarrow{\pi_0(\partial)} \mathbb{Z}/2 \longrightarrow 0,$$

so $\pi_0(\omega)$ is multiplication by ± 2 , depending on the choice of generators. There is a canonical splitting of infinite loop spaces

$$(2.4) \quad \Omega^\infty S^\infty(\mathbb{C}P_+^\infty) \simeq \Omega^\infty S^\infty(\mathbb{C}P^\infty) \times \Omega^\infty S^\infty.$$

We fix the generator of $\pi_0\Omega^\infty S^\infty(\mathbb{C}P_+^\infty)$ to be the element that maps to $+1$ under the isomorphisms

$$\pi_0(\Omega^\infty S^\infty(\mathbb{C}P_+^\infty)) \xrightarrow{\pi_0(c)} \pi_0(\Omega^\infty S^\infty) \xrightarrow{\text{degree}} \mathbb{Z},$$

where c collapses $\mathbb{C}P^\infty$ to the non-basepoint of S^0 . We fix the generator of $\pi_0(\Omega^\infty \mathbb{C}P_{-1}^\infty)$ so that $\pi_0(\omega)$ is multiplication by -2 .

2.2. Review of results used. Our divisibility result of Theorem 1.1 is based upon the following three theorems.

Theorem 2.1 ([T]). *The spaces $\mathbb{Z} \times B\Gamma_\infty^+$ and $B\Gamma_\infty^+ (= \{0\} \times B\Gamma_\infty^+)$ are infinite loop spaces.*

Here the superscript $(+)$ denotes Quillen’s plus construction; cf. [B]. The product structure can be described as follows. We may view $\Gamma_{g,2}$ as the mapping class group of surfaces with one incoming and one outgoing boundary component. Gluing the incoming boundary component of one surface to the outgoing component of the other defines a map

$$\Gamma_{g,2} \times \Gamma_{h,2} \longrightarrow \Gamma_{g+h,2}$$

and a corresponding map of classifying spaces that makes the disjoint union $\bigsqcup B\Gamma_{g,2}$ over all $g \geq 0$ into a topological monoid. Consider the map

$$\bigsqcup_{g \geq 0} B\Gamma_{g,2} \longrightarrow \mathbb{Z} \times B\Gamma_\infty^+$$

that sends $B\Gamma_{g,2}$ into the component $\{g\} \times B\Gamma_\infty^+$ by the stabilization map (1.1) followed by the map into the plus construction. The infinite loop space structure on $\mathbb{Z} \times B\Gamma_\infty^+$ is compatible with the monoidal structure on $\bigsqcup B\Gamma_{g,2}$, and the induced map

$$\Omega B(\bigsqcup_{g \geq 0} B\Gamma_{g,2}) \longrightarrow \mathbb{Z} \times B\Gamma_\infty^+$$

is a homotopy equivalence of loop spaces. We refer to [T] for details.

To state the next result, for each prime p we pick a positive integer $k = k(p)$ so that $-k$ reduces to a generator of the units $(\mathbb{Z}/p^2)^\times$ when p is odd. We pick $k = 5$ when $p = 2$. Write ψ^{-k} for the self-map of $\mathbb{C}P^\infty$ that multiplies by $-k$ on the second cohomology group. Composing with the inclusion into $\Omega^\infty S^\infty(\mathbb{C}P^\infty)$ and using the loop sum we have a map

$$1 + k\psi^{-k} : \mathbb{C}P^\infty \longrightarrow \Omega^\infty S^\infty(\mathbb{C}P^\infty),$$

and, using the universal property of free infinite loop spaces, a unique extension to a self-map of $\Omega^\infty S^\infty(\mathbb{C}P^\infty)$, again denoted $1 + k\psi^{-k}$.

Theorem 2.2. *There are infinite loop maps*

$$\alpha : \mathbb{Z} \times B\Gamma_\infty^+ \longrightarrow \Omega^\infty \mathbb{C}P_{-1}^\infty, \quad \mu_p : \Omega^\infty S^\infty(\mathbb{C}P_+^\infty) \longrightarrow (\mathbb{Z} \times B\Gamma_\infty^+)_p^\wedge$$

such that the composition $\omega \circ \alpha \circ \mu_p$ and the self-map

$$\begin{pmatrix} 1 + k\psi^{-k} & 0 \\ 0 & -2 \end{pmatrix} : \Omega^\infty S^\infty \mathbb{C}P^\infty \times \Omega^\infty S^\infty \longrightarrow \Omega^\infty S^\infty \mathbb{C}P^\infty \times \Omega^\infty S^\infty$$

become homotopic after p -adic completion.

This is an improvement on the main theorem of [MT], where the map in the lower left corner had been left undetermined. For our calculations in Section 4.3 we need this map however to be zero. A proof of Theorem 2.2 is given in the final Section 5.

Remark 2.3. The reader is referred to [BK] for the notion of p -adic completion (also called \mathbb{F}_p -completion). For connected, compact CW-complexes X and infinite loop spaces $\Omega^\infty E$ of finite type one has

$$[X, (\Omega^\infty E)_p^\wedge] = [X, \Omega^\infty E] \otimes \mathbb{Z}_p, \quad H^*((\Omega^\infty E)_p^\wedge; \mathbb{Z}) = H^*(\Omega^\infty E; \mathbb{Z}_p).$$

Furthermore, note that the homotopy class of the map α in Theorem 2.2 is uniquely determined by its composition with $\mathbb{Z} \times B\Gamma_\infty \rightarrow \mathbb{Z} \times B\Gamma_\infty^+$. Indeed since $\Omega^\infty \mathbb{C}P_{-1}^\infty$ is an infinite loop space the induced map

$$[\mathbb{Z} \times B\Gamma_\infty^+, \Omega^\infty \mathbb{C}P_{-1}^\infty] \longrightarrow [\mathbb{Z} \times B\Gamma_\infty, \Omega^\infty \mathbb{C}P_{-1}^\infty]$$

is an isomorphism. This is a standard property of the plus construction; cf. [B].

Before we give a detailed description of α in the next section, we state here the third result.

Theorem 2.4 ([MW]). *The map α is a homotopy equivalence.*

3. CHARACTERISTIC CLASSES OF SURFACE BUNDLES

3.1. Universal surface bundles. The methods used in this and the surrounding papers do not use the mapping class groups directly but rather the topological groups of orientation-preserving diffeomorphisms of surfaces. We briefly review the correspondence.

Let $F_{g,b}$ be a connected surface of genus g with b boundary circles. We write $\text{Diff}(F_{g,b}; \partial)$ for the topological group of orientation-preserving diffeomorphisms that keep (a neighborhood of) the boundary pointwise fixed. For $g \geq 2$, results from [EE] and [ES] yield

$$B\Gamma_{g,b} \simeq B\text{Diff}(F_{g,b}; \partial)$$

so that $B\Gamma_{g,b}$ classifies diffeomorphism classes of smooth fibre bundles $\pi : E \rightarrow X$ with fibre $F_{g,b}$ and standard boundary behavior:

$$\partial E = X \times \bigsqcup_1^b S^1, \quad \pi|_{\partial E} = \text{proj}_X.$$

Similarly,

$$B\Gamma_{g,b}^s \simeq B\text{Diff}(F_{g,b}; \partial \sqcup \{x_1, \dots, x_s\})$$

where x_1, \dots, x_s are distinct interior points of $F_{g,b}$. Take $s = 1$. Since $\text{Diff}(F_{g,b}; \partial)$ acts transitively on the interior of $F_{g,b}$,

$$E(F_{g,b}) := E\text{Diff}(F_{g,b}; \partial) \times_{\text{Diff}(F_{g,b}; \partial)} F_{g,b} \simeq B\text{Diff}(F_{g,b}; \partial \sqcup \{x\}) \simeq B\Gamma_{g,b}^1.$$

The forgetful map $\pi : B\Gamma_{g,b}^1 \rightarrow B\Gamma_{g,b}$ corresponds to the universal smooth $F_{g,b}$ bundle

$$(3.1) \quad F_{g,b} \longrightarrow E(F_{g,b}) \longrightarrow B\text{Diff}(F_{g,b}; \partial).$$

The central extension (1.2) is classified by “the differential at x ”,

$$\text{Diff}(F_{g,b}; \partial \sqcup \{x\}) \longrightarrow \text{GL}^+(T_x F_{g,b}) \simeq \text{SO}(2).$$

Hence the circle bundle induced from (1.2) by applying the classifying space functor corresponds to the circle bundle of the vertical tangent bundle associated with (3.1).

3.2. The map α and the kappa classes. The map of infinite loop spaces $\alpha : \mathbb{Z} \times B\Gamma_\infty^+ \rightarrow \Omega^\infty \mathbb{C}P_{-1}^\infty$ constructed in section 2 of [MT] restricts to a map $\alpha_{g,2} : B\Gamma_{g,2} \rightarrow \Omega_g^\infty \mathbb{C}P_{-1}^\infty$ that is homotopic to the composition

$$(3.2) \quad \alpha_{g,2} : B\Gamma_{g,2} \longrightarrow B\Gamma_{g+1} \xrightarrow{\alpha_{g+1}} \Omega_g^\infty \mathbb{C}P_{-1}^\infty,$$

where the left-hand map is induced from gluing the two parametrized boundary circles together. These maps are up to homotopy compatible with the monoidal structure on $\bigsqcup B\Gamma_{g,2}$.

We next recall a description of α_{g+1} which is well-suited for identifying the kappa classes. Let $\pi : E \rightarrow X$ be a smooth surface bundle with closed fiber F . Thus $E = P \times_{\text{Diff}(F)} F$ where P is a principal $\text{Diff}(F)$ bundle over X . We do not assume that X is smooth or finite dimensional, only that X is paracompact (or a CW-complex).

We denote by $\text{Emb}(F, \mathbb{R}^n)$ the space of smooth embeddings in the C^∞ -topology, and let \mathbb{R}^∞ and $\text{Emb}(F, \mathbb{R}^\infty)$ be the colimits of \mathbb{R}^n and $\text{Emb}(F, \mathbb{R}^n)$, respectively. We shall consider fiberwise embeddings $\iota : E \rightarrow X \times \mathbb{R}^\infty$, that is, fiberwise maps such that each $\iota_x : E_x \rightarrow \{x\} \times \mathbb{R}^\infty$ is an embedding and such that the adjoint $\text{Diff}(F)$ -equivariant map $P \rightarrow \text{Emb}(F, \mathbb{R}^\infty)$ is continuous. Such an ι is equivalent to a section of $P \times_{\text{Diff}(F)} \text{Emb}(F, \mathbb{R}^\infty)$. Note that $\text{Emb}(F, \mathbb{R}^\infty)$ is contractible so that such a section always exists.

An embedding $\iota_x : F \hookrightarrow \mathbb{R}^{n+2}$ extends to a map from the normal bundle $N^n \iota_x = \{(p, v) | v \perp T_p F\}$ into \mathbb{R}^{n+2} by sending (p, v) to $p + v$. (Here we have identified F with its image under ι_x .) We call the embedding ι_x *fat* if this map restricts to an embedding of the unit disk bundle $D(N^n \iota_x)$. The subspace of fat embeddings $\text{Emb}^f(F, \mathbb{R}^\infty) \subset \text{Emb}(F, \mathbb{R}^\infty)$ is contractible, since the inclusion is a homotopy equivalence by the tubular neighborhood theorem and since $\text{Emb}(F, \mathbb{R}^\infty)$ is contractible by Whitney’s embedding theorem. A fibrewise fat embedding $\iota : E \rightarrow X \times \mathbb{R}^\infty$ is then a section of the fibre bundle $P \times_{\text{Diff}(F)} \text{Emb}^f(F, \mathbb{R}^\infty)$.

Suppose first that $\iota : E \rightarrow X \times \mathbb{R}^{n+2}$ is a fibrewise fat embedding of codimension n . The Pontryagin–Thom construction associates a “collapse” map onto the Thom space of the fibrewise normal bundle,

$$c_{\pi, \iota} : X_+ \wedge S^{n+2} \longrightarrow D(N_\pi^n \iota) / S(N_\pi^n \iota) = \text{Th}(N_\pi^n \iota).$$

We are particularly interested in its adjoint map $X \rightarrow \Omega^{n+2} \text{Th}(N_\pi^n \iota)$.

Let $G(2, n)$ be the Grassmann manifold of oriented 2-dimensional subspaces of \mathbb{R}^{n+2} , and let U_n and U_n^\perp be the two complementary universal bundles over it of dimension 2 and n , respectively. The fat embedding ι induces bundle maps

$$T_\pi E \longrightarrow U_n, \quad N_\pi^n \iota \longrightarrow U_n^\perp$$

and a commutative diagram

$$(3.3) \quad \begin{array}{ccccc} X_+ \wedge S^{n+2} & \xrightarrow{c_{\pi, \iota}} & \text{Th}(N_\pi^n \iota) & \xrightarrow{s} & \text{Th}(T_\pi E \oplus N_\pi^n \iota) \\ \parallel & & \downarrow & & \downarrow \\ X_+ \wedge S^{n+2} & \xrightarrow{c_n} & \text{Th}(U_n^\perp) & \xrightarrow{s} & \text{Th}(U_n \oplus U_n^\perp). \end{array}$$

In the general case of a fiberwise fat embedding $\iota : E \rightarrow X \times \mathbb{R}^\infty$, the base space X is the colimit of the subspaces

$$X_n := \{x \in X \mid \iota_x(E_x) \subset \{x\} \times \mathbb{R}^{n+2}\},$$

and the diagram

$$\begin{CD} (X_n)_+ \wedge S^{n+2} @>>> \text{Th}(U_n^\perp) \\ @VVV @VVV \\ (X_{n+1})_+ \wedge S^{n+3} @>>> \text{Th}(U_{n+1}^\perp) \end{CD}$$

is commutative since $U_{n+1}^\perp|_{G(2,n)} = U_n^\perp$. Taking adjoints we get

$$\alpha_{\pi,\iota} : X \longrightarrow \text{colim}_{n \rightarrow \infty} \Omega^{n+2} \text{Th}(U_n^\perp).$$

Since $\text{Emb}^f(F, \mathbb{R}^\infty)$ is contractible, all sections of $P \times_{\text{Diff}(F)} \text{Emb}^f(F, \mathbb{R}^\infty)$ are homotopic, and consequently the homotopy class $[\alpha_{\pi,\iota}]$ is independent of the choice of ι . We will therefore from now on suppress the subscript ι .

Realification gives a $(2n - 1)$ -connected map from $\mathbb{C}P^n$ into the oriented Grassmannian $G(2, 2n)$ covered by a bundle map $L_n^\perp \rightarrow U_{2n}^\perp$. Thus $G(2, \infty) \simeq \mathbb{C}P^\infty$ and

$$\Omega^\infty \mathbb{C}P_{-1}^\infty = \text{colim} \Omega^{2n+2} \text{Th}(L_n^\perp) \xrightarrow{\simeq} \text{colim} \Omega^{2n+2} \text{Th}(U_{2n}^\perp)$$

is a homotopy equivalence. Altogether we have a well-defined homotopy class

$$\alpha_\pi : X \longrightarrow \Omega^\infty \mathbb{C}P_{-1}^\infty.$$

For $X = B\text{Diff}(F_{g+1}) \simeq B\Gamma_{g+1}$ this is the map α_{g+1} of (3.2).

Let us check that the image of α_{g+1} , and hence the image of $\alpha_{g,2}$, lie in the g -component of $\Omega^\infty \mathbb{C}P_{-1}^\infty$, or equivalently that the composition

$$\text{proj} \circ \omega \circ \alpha_{g+1} : B\Gamma_{g+1} \longrightarrow \Omega^\infty S^\infty(\mathbb{C}P_+^\infty) \longrightarrow \Omega^\infty S^\infty$$

lands in the $-2g$ component (with identification of components chosen at the end of Section 2.1). Consider (3.3) with X a single point and $E = F_{g+1}$. The bottom row in (3.3) is thus $s \circ c_n : S^{n+2} \rightarrow G(2, 2n)_+ \wedge S^{n+2}$, and we need to compute the degree of the composition of this map with the projection onto S^{n+2} . This degree is given by the evaluation of the pullback of the generator of $H^{n+2}(S^{n+2})$ on the fundamental class $[S^{n+2}]$. Under the projection the fundamental class is pulled back to the Thom class of the trivial bundle $U_n \oplus U_n^\perp$. Writing λ_U for the Thom class of the vector bundle U , the degree is thus given by

$$\begin{aligned} \langle c_\pi^* s^*(\lambda_{U_n} \cdot \lambda_{U_n^\perp}), [S^{2n+2}] \rangle &= \langle c_\pi^*(e(TF_{g+1}) \cdot \lambda_{U_n^\perp}), [S^{2n+2}] \rangle \\ &= \langle e(TF_{g+1}), [F_{g+1}] \rangle = -2g, \end{aligned}$$

as claimed.

Next we compute the maps $\alpha_{g,2}$ and $\alpha_{g+1,2}$ under the map $B\Gamma_{g,2} \rightarrow B\Gamma_{g+1,2}$ induced from gluing a torus with two boundary circles $F_{1,2}$ to $F_{g,2}$. Considering $F_{1,2}$ as a fibre bundle over a point the construction above gives an element $[1] \in \Omega_1^\infty \mathbb{C}P_{-1}^\infty$. Loop sum with $[1]$ in $\Omega^\infty \mathbb{C}P_{-1}^\infty$ translates the g -component into the

$(g + 1)$ -component and

$$(3.4) \quad \begin{array}{ccc} B\Gamma_{g,2} & \xrightarrow{\alpha_{g,2}} & \Omega_g^\infty \mathbb{C}P_{-1}^\infty \\ \downarrow & & \downarrow * [1] \\ B\Gamma_{g+1,2} & \xrightarrow{\alpha_{g+1,2}} & \Omega_{g+1}^\infty \mathbb{C}P_{-1}^\infty \end{array}$$

is homotopy commutative. To see this observe that the left vertical map is multiplication by the basepoint of $B\Gamma_{1,2}$ in the monoid $\bigsqcup_{g \geq 0} B\Gamma_{g,2}$ and the right vertical map is multiplication by its image in $\Omega_1^\infty \mathbb{C}P_{-1}^\infty$. Homotopy commutativity of diagram (3.4) now follows because the $\alpha_{g,2}$ induce a map of monoids up to homotopy. This is because α is a map of infinite loop spaces, α restricts to $\alpha_{g,2}$ on $B\Gamma_{g,2}$, and the infinite loop space structure on $\mathbb{Z} \times B\Gamma_\infty^+$ is compatible with the monoidal structure. (Alternatively, homotopy commutativity of (3.4) follows from a calculation of pretransfers; cf. [G2].)

Let $\tilde{\alpha}$ denote the restriction of α in Theorem 2.2 to the zero component. Restricted to $B\Gamma_{g,2}$ it is homotopic to $\tilde{\alpha}_{g,2} = (*[-g]) \circ \alpha_{g,2}$. We can now relate the kappa classes to spectrum cohomology. Consider

$$B\Gamma_{g,2} \xrightarrow{\tilde{\alpha}_{g,2}} \Omega_0^\infty \mathbb{C}P_{-1}^\infty \xrightarrow{\omega} \Omega_0^\infty S^\infty(\mathbb{C}P_+^\infty)$$

and recall the cohomology suspension from Section 2.1:

$$\sigma^* : H^{2i}(\mathbb{C}P^\infty; \mathbb{Z}) \simeq H^{2i}(S^\infty \mathbb{C}P_+^\infty) \longrightarrow H^{2i}(\Omega_0^\infty S^\infty(\mathbb{C}P_+^\infty)).$$

Theorem 3.1. *The Miller-Morita-Mumford class κ_i is equal to $(\omega \circ \tilde{\alpha})^*(\sigma^* e^i)$, where $e \in H^2(\mathbb{C}P^\infty; \mathbb{Z})$ is the Euler class of the canonical line bundle.*

Proof. Let $\pi : E \rightarrow X$ be a smooth fibre bundle with fibre F_{g+1} , classified by $f_\pi : X \rightarrow B\Gamma_{g+1}$. By definition

$$f_\pi^*(\kappa_i) = \pi_!(e(T_\pi E)^{i+1}) \in H^{2i}(X; \mathbb{Z}),$$

where $\pi_!$ is the composition of the Thom isomorphism and the Pontrjagin-Thom collapse map

$$\tilde{H}^{2i+2}(E; \mathbb{Z}) \xrightarrow{\simeq} \tilde{H}^{2i+2n+2}(\text{Th}(N_\pi^i); \mathbb{Z}) \xrightarrow{c_\pi^*} H^{2i+2n+2}(S^{2n+2} \wedge X_+; \mathbb{Z})$$

followed by the $(2n + 2)$ -nd desuspension; here the notation is as in (3.3). Let $x \in H^k(\mathbb{C}P^N; \mathbb{Z}) \simeq H^k(G(2, 2N); \mathbb{Z})$ for $N \gg k$. The $(2n + 2)$ -fold suspension $\Sigma^{2n+2}(x) \in H^{k+2n+2}(S^{2n+2} \wedge \mathbb{C}P^N; \mathbb{Z})$ is x times the Thom class of the trivial $2n + 2$ real bundle $L_n^\perp \oplus L_n$. Thus

$$s^*(\Sigma^{2n+2}(x)) = s^*(\lambda_{L_n^\perp} \cdot \lambda_{L_n} \cdot x) = \lambda_{L_n^\perp} \cdot e(L_n) \cdot x.$$

To interpret these formulas, recall that the cohomology of the Thom space of a vector bundle is a module over the cohomology of the base space, and a map of bundles (such as s) induces a map of modules. Furthermore, as elements in the cohomology of the Thom space of the trivial bundle, $\lambda_{L_n^\perp}$ and λ_{L_n} are pulled back along the bundle projections π_1 and π_2 of $L_n^\perp \oplus L_n$ onto the first and second summands. Hence, as $\pi_1 \circ s$ is the identity, we have $s^*(\lambda_{L_n^\perp}) = \lambda_{L_n^\perp}$, and as $\pi_2 \circ s$ factors through the base space, $s^*(\lambda_{L_n}) = 1 \cdot e(L_n)$. Finally, take $x = e^i$ in the above formula and use the commutativity of (3.3) to complete the proof. \square

3.3. One part of Akita's conjecture. For our next theorem we need the relation between Steenrod operations and characteristic classes of vector bundles. Recall the i -th Steenrod operation:

$$\begin{aligned} P^i &: H^k(X; \mathbb{F}_p) \longrightarrow H^{k+2i(p-1)}(X; \mathbb{F}_p), \quad p \text{ odd,} \\ Sq^i &: H^k(X, \mathbb{F}_2) \longrightarrow H^{k+i}(X, \mathbb{F}_2). \end{aligned}$$

Let E be an oriented vector bundle over X and λ_U its cohomology Thom class. One defines $v_i(E) \in H^{2i(p-1)}(X; \mathbb{F}_p)$, respectively $v_i(E) \in H^i(X; \mathbb{F}_2)$, by

$$P^i(\lambda_E) = v_i(E)\lambda_E, \quad Sq^i(\lambda_E) = v_i(E)\lambda_E.$$

For $p = 2$, these are the Stiefel-Whitney classes, and for p odd they were first defined by Wu; cf. [MSt]. For an oriented 2-plane bundle (or complex line bundle) L ,

$$\begin{aligned} v_1(L) &= e(L)^{p-1} && \text{for } p \text{ odd,} \\ v_2(L) &= e(L) \quad \text{and} \quad v_1(L) = 0 && \text{for } p = 2. \end{aligned}$$

Moreover, the total class

$$v(E) = 1 + v_1(E) + v_2(E) + \cdots \in H^*(X; \mathbb{F}_p)$$

takes direct sums of oriented vector bundles into (graded) products.

Theorem 3.2. *The modulo p reduction of $\kappa_i \in H^{2i}(B\Gamma_\infty; \mathbb{F}_p)$ is zero when $i + 1 \equiv 0 \pmod{p - 1}$.*

*Proof.*³ Let $H(\mathbb{Z}, k)$ denote the Eilenberg-MacLane space with non-trivial homotopy \mathbb{Z} in dimension k . The Thom class $\lambda_n = \lambda_{L_n^\perp}$ is represented by a map from $\text{Th}(L_n^\perp)$ to $H(\mathbb{Z}, 2n)$. We let Y_{2n+2} be its homotopy fibre, so that there is a fibration sequence

$$Y_{2n+2} \xrightarrow{j_{2n+2}} \text{Th}(L_n^\perp) \xrightarrow{\lambda_n} H(\mathbb{Z}, 2n).$$

The spaces Y_{2n+2} are the $(2n + 2)$ -nd terms of a spectrum Y and the j_{2n+2} define a map $j : Y \rightarrow \mathbb{C}P_{-1}^\infty$ of spectra. Since $\Omega^{2n+2}H(\mathbb{Z}, 2n)$ has vanishing homotopy groups,

$$(3.5) \quad \Omega^\infty Y \xrightarrow{\Omega^\infty j} \Omega^\infty \mathbb{C}P_{-1}^\infty$$

is a (weak) homotopy equivalence. Thus the bottom vertical map in the commutative diagram

$$\begin{array}{ccc} H^{2i}(\mathbb{C}P_{-1}^\infty; \mathbb{F}_p) & \xrightarrow{j^*} & H^{2i}(Y, \mathbb{F}_p) \\ \sigma^* \downarrow & & \sigma^* \downarrow \\ H^{2i}(\Omega_0^\infty \mathbb{C}P_{-1}^\infty; \mathbb{F}_p) & \xrightarrow{(\Omega^\infty j)^*} & H^{2i}(\Omega^\infty Y; \mathbb{F}_p) \end{array}$$

is an isomorphism. The proof of Theorem 3.1 shows that

$$\tilde{\alpha}^* \sigma^*(e^{i+1}\lambda_{L^+}) = \kappa_i,$$

so it suffices to prove that j^* vanishes when $i + 1 \equiv 0 \pmod{p - 1}$. Equivalently, we must show that $e^{i+1}\lambda_n$ is in the image of

$$\lambda_n^* : H^*(H(\mathbb{Z}, 2n); \mathbb{F}_p) \longrightarrow H^*(\text{Th}(L_n^\perp); \mathbb{F}_p)$$

³We thank John Rognes for this proof; it replaces a more cumbersome earlier argument.

in the stated dimensions. This is implied by

$$v(L_n^\perp) = v(L_n)^{-1} = \begin{cases} (1 + e^{p-1})^{-1}, & p > 2, \\ (1 + e)^{-1}, & p = 2. \end{cases}$$

Since $P^i(\lambda_n) \in \text{image}(\lambda_n^*)$ the result follows. □

Remark 3.3. The relation $P^i(\lambda_{L^\perp}) = \kappa_{i(p-1)-1}\lambda_{L^\perp}$ used above is further exploited in [G2] to define secondary classes μ_i with $p\mu_i = \kappa_{i(p-1)-1}$ in cohomology with $\mathbb{Z}/p^2\mathbb{Z}$ coefficients.

Theorem 3.2 proves half of Akita's, Theorem 1.2. The other half is implied by Theorem 1.1, which is proved below.

4. PROOF OF THEOREM 1.1

4.1. Segal's splitting. The inclusion of CP^∞ in BU that represents the reduced canonical line bundle (of virtual dimension zero) extends to a map of infinite loop spaces

$$l : \Omega^\infty S^\infty(CP^\infty) \longrightarrow BU$$

by Bott periodicity and the universal property of the free infinite loop space functor $\Omega^\infty S^\infty$. Graeme Segal [S] proved that this map has a splitting. In [C] Michael Crabb gives a construction of this map as the S^1 -equivariant J -homomorphism.

Theorem 4.1 ([S], [C]). *The map l has a left inverse up to homotopy. In the resulting decomposition*

$$\Omega^\infty S^\infty(CP^\infty) \simeq BU \times \text{Fib}(l)$$

the homotopy fiber $\text{Fib}(l)$ has vanishing rational cohomology.

In particular, this gives an identification of Hopf algebras

$$(4.1) \quad H_{free}^*(\Omega^\infty S^\infty(CP^\infty)) \simeq H^*(BU; \mathbb{Z})$$

induced by l^* . The graded module of primitive elements of the right-hand side is a copy of \mathbb{Z} in each even degree generated by the integral Chern character class $s_i = i!ch_i$. Since $l \circ i$ represents the (reduced) line bundle and $ch_i(L) = \frac{1}{i!}e^i$, (4.1) implies that

$$i^* : H_{free}^*(\Omega^\infty S^\infty(CP^\infty)) \longrightarrow H^*(CP^\infty; \mathbb{Z})$$

sends the primitive generator s_i to e^i . Note from (2.2) that we also have that

$$H^*(CP^\infty; \mathbb{Z}) = H^*(S^\infty CP^\infty; \mathbb{Z}) \xrightarrow{\sigma^*} P(H_{free}^*(\Omega^\infty S^\infty(CP^\infty)))$$

maps e^i to s_i . Therefore, Theorem 3.1 translates into

$$(4.2) \quad \kappa_i = (l \circ \omega \circ \alpha)^*(s_i).$$

The map ω in (2.3) induces an isomorphism on rational cohomology because the cohomology of $\Omega^\infty S^{\infty-1}$ is well known to be all torsion. The isomorphisms

$$H^*(BU; \mathbb{Q}) \xrightarrow{l^*} H^*(\Omega_0^\infty S^\infty(CP_+^\infty); \mathbb{Q}) \xrightarrow{\omega^*} H^*(\Omega^\infty CP_{-1}^\infty; \mathbb{Q})$$

together with Theorem 2.4 thus give an isomorphism

$$H^*(B\Gamma_\infty; \mathbb{Q}) \simeq H^*(BU; \mathbb{Q})$$

of Hopf algebras. Hence $P(H_{free}^*(B\Gamma_\infty))$ is a copy of \mathbb{Z} in each even degree. We choose a generator

$$\tau_i \in P(H_{free}^{2i}(B\Gamma_\infty))$$

such that $\kappa_i = D_i\tau_i$ for some positive number D_i . The next section gives close upper and lower bounds for D_i .

4.2. A lower and an upper bound. Recall the definition of the symplectic characteristic classes for surface bundles. The action of $\Gamma_{g,1}$ on $H^1(F_{g,1}; \mathbb{Z}) = \mathbb{Z}^{2g} = H^1(F_g; \mathbb{Z})$ induces the standard symplectic representation, and hence a representation of $\Gamma_{g,2}$ via the map $\Gamma_{g,2} \rightarrow \Gamma_{g,1}$ induced by gluing a disk to one of the boundary components. We may let $g \rightarrow \infty$ and obtain

$$B\Gamma_\infty \longrightarrow BSp(\mathbb{Z}).$$

This map can be composed with the map into $BSp(\mathbb{R}) \simeq BU$ so that we have a map

$$\eta : B\Gamma_\infty \longrightarrow BU.$$

Theorem 4.2 ([Mo], [Mu]). *In $H^*(B\Gamma_\infty; \mathbb{Q})$ one has the relation*

$$\eta^*(s_{2i-1}) = (-1)^i \binom{B_i}{2i} \kappa_{2i-1}.$$

We are now in a position to prove

Theorem 4.3. *For all $i \geq 1$, $D_{2i} = 2$ and*

$$D_{2i-1} = \text{den}\left(\frac{B_i}{2i}\right) \quad \text{or} \quad D_{2i-1} = 2 \text{den}\left(\frac{B_i}{2i}\right).$$

Proof. As s_{2i-1} and κ_{2i-1} are integral classes, Theorem 4.2 implies immediately that modulo torsion $\text{den}(B_i/2i)$ divides κ_{2i-1} . By Theorem 3.2 (for $p = 2$) we also know that 2 divides κ_{2i} . This establishes the lower bounds for all $i \geq 1$. The upper bounds are a consequence of Theorem 2.2 as we explain now.

As in Section 2.2, let k be a positive integer such that $-k$ generates $(\mathbb{Z}/p^2)^\times$ for odd p and let $k = 5$ when $p = 2$. Then by Theorem 2.2 there is a factorization

$$1 + k\psi^{-k} : \mathbb{C}P^\infty \xrightarrow{\mu_p} (B\Gamma_\infty^+)_p^\wedge \xrightarrow{\omega \circ \alpha} \Omega_0^\infty S^\infty(\mathbb{C}P_+^\infty)_p^\wedge \xrightarrow{\text{proj}} \Omega^\infty S^\infty(\mathbb{C}P^\infty)_p^\wedge.$$

Since $\Omega^\infty S^\infty(\mathbb{C}P^\infty)$ is of finite type,

$$H_{free}^*(\Omega^\infty S^\infty(\mathbb{C}P^\infty)_p^\wedge) = H_{free}^*(\Omega^\infty S^\infty(\mathbb{C}P^\infty)) \otimes \mathbb{Z}_p.$$

It follows from (4.2) that $\mu_p^*(\kappa_i)$ is the image of the primitive generator s_i under

$$(1 + k\psi^{-k})^* : PH_{free}^{2i}(\Omega^\infty S^\infty(\mathbb{C}P^\infty)) \otimes \mathbb{Z}_p \longrightarrow H^{2i}(\mathbb{C}P^\infty; \mathbb{Z}_p).$$

Both groups are copies of \mathbb{Z}_p and $(1 + k\psi^{-k})^*$ in dimension $2j$ is multiplication by $1 + k(-k)^j = 1 - (-k)^{j+1}$. Therefore, $\nu_p(D_j) \leq \nu_p(1 - (-k)^{j+1})$.

We have the following well-known table of p -adic valuations (see, for example, lemma 2.12 in [A2]):

$$\nu_p(1 - (-k)^s) = \begin{cases} 1 + \nu_p(s) & \text{if } s \equiv 0 \pmod{p-1}, p \text{ odd,} \\ 0 & \text{if } s \not\equiv 0 \pmod{p-1}, p \text{ odd,} \end{cases}$$

$$\nu_2(1 - (-k)^s) = \begin{cases} 2 + \nu_2(s) & \text{if } s \equiv 0 \pmod{2}, \\ 1 & \text{if } s \not\equiv 0 \pmod{2}. \end{cases}$$

When $j = 2i$, for p odd, $\nu_p(1 - (-k)^{j+1})$ is zero and for $p = 2$ it is equal to 1. This gives $D_{2i} \leq 2$. When $j = 2i - 1$, for p odd we have $\nu_p(1 - (-k)^{j+1}) = 1 + \nu_p(2i)$ if $2i$ divides $p - 1$ and zero otherwise. This is precisely $\nu_p(\text{den}(B_i/2i))$ (compare (1.3)). If $p = 2$, however, $\nu_2(1 - (-k)^{j+1}) = 2 + \nu_2(2i)$, which is one more than $\nu_2(\text{den}(B_i/2i))$. This gives $D_{2i-1} \leq 2 \text{den}(B_i/2i)$. \square

4.3. The final factor of 2. Theorem 4.3 leaves us with an indeterminacy of a factor of 2 in the odd case, and Theorem 1.1 will follow immediately from Theorem 4.5 below. The proof requires several extra results: the improvement of the main theorem from [MT] as stated in Theorem 2.2 (for $p = 2$), Theorem 2.4, as well as part of theorem 1.3 from [G1], which we state as

Theorem 4.4. $H_*(\Omega_0^\infty \mathbb{C}P_{-1}^\infty; \mathbb{F}_2) \xrightarrow{\omega_*} H_*(\Omega_0^\infty S^\infty(\mathbb{C}P_+^\infty); \mathbb{F}_2)$ is injective.

Theorem 4.5. $\nu_2(D_{2i-1}) = 1 + \nu_2(2i)$.

Proof: As in the proof of Theorem 4.3 we have

$$1 - (-5)^{j+1} : P(H_{free}^{2j}(\Omega^\infty S^\infty(\mathbb{C}P_+^\infty))) \otimes \mathbb{Z}_2 \xrightarrow{(\omega \circ \alpha)^*} P(H_{free}^{2j}(B\Gamma_\infty^+)) \otimes \mathbb{Z}_2$$

$$\xrightarrow{\mu_2^*} H^{2j}(\mathbb{C}P^\infty; \mathbb{Z}_2).$$

All groups are copies of \mathbb{Z}_2 , $\kappa_j = D_j \tau_j$ with τ_j a generator and $\kappa_j = (\omega \circ \alpha)^*(s_j)$, where s_j is the generator of the left term. Suppose that $\nu_2(D_j) = 2 + \nu_2(j + 1)$ for some $j = 2i - 1$. Then $\mu_2^*(\tau_j)$ would be a generator and dually

$$\mu_{2*} : H_{2j}(\mathbb{C}P^\infty; \mathbb{F}_2) \longrightarrow H_{2j}^{free}(B\Gamma_\infty^+) \otimes \mathbb{F}_2$$

and hence

$$\mu_{2*} : H_{2j}(\mathbb{C}P^\infty; \mathbb{F}_2) \longrightarrow H_{2j}(B\Gamma_\infty^+) \otimes \mathbb{F}_2$$

would be non-zero. Now apply Theorem 2.4 and Theorem 4.4 to conclude that

$$H_{4i-2}(\mathbb{C}P^\infty; \mathbb{F}_2) \xrightarrow{(\omega \circ \alpha \circ \mu_2)^*} H_{4i-2}(\Omega_0^\infty S^\infty(\mathbb{C}P_+^\infty); \mathbb{F}_2)$$

would be non-zero. But this leads to a contradiction as we now argue.

Indeed, by Theorem 2.2 the above map $(\omega \circ \alpha \circ \mu_2)_*$ is $(1 + 5\psi^{-5}, 0)_*$. The self-map ψ^{-5} of $\mathbb{C}P^\infty$ induces the identity on $H_*(\mathbb{C}P^\infty; \mathbb{F}_2)$. Hence $1 + 5\psi^{-5}$ induces

the same map on \mathbb{F}_2 -homology as six times the canonical inclusion map:

$$6i : \mathbb{C}P^\infty \longrightarrow \Omega^\infty S^\infty(\mathbb{C}P^\infty).$$

But $(6i)_* = 0$ on \mathbb{F}_2 -homology in degrees $4i - 2$, giving the desired contradiction. Indeed, $6i$ is the composition of $2i$ and multiplication by 3 in the loop sum sense in $\Omega^\infty S^\infty(\mathbb{C}P^\infty)$. $2i$ in turn is the composition of the diagonal map on $\mathbb{C}P^\infty$ composed with $i \times i$ and loop sum. The diagonal map sends the generator a_j of the $2j$ -th homology group to $\Sigma_{s+t=j} a_s \otimes a_t$. Loop sum replaces tensor product by Pontryagin product, which here is commutative. Thus in \mathbb{F}_2 -homology for j odd, $2i$ is zero. \square

5. PROOF OF THEOREM 2.2

We offer two proofs of Theorem 2.2. The first one is in the spirit of [MT] and provides an improvement on the geometric construction of μ_p given there. The second proof was suggested by the referee. It is short, purely homotopy theoretic, but it does not provide any insight in the map μ_p from $\Omega^\infty S^\infty \mathbb{C}P_+^\infty$ to $B\Gamma_\infty^+$. The map μ_p , constructed from explicit cyclic ramified covers of $\mathbb{C}P^1$, provides an important link between the algebraic approach and the topological approach to the mapping class group.

5.1. A constructive proof. In order to prove that the map $\Omega^\infty S^\infty \mathbb{C}P^\infty \rightarrow \Omega^\infty S^\infty$ induced by $\omega \circ \alpha \circ \mu_p$ is zero after p -adic completion, we will have to review the construction of μ_p . μ_p depends on p and the choice of k ; here k can be any integer which is a unit modulo p^n for all $n \geq 1$. We take the opportunity to give a variant of the proof for the main theorem in [MT] from sections 3.1-3.3. For the fact that α is a map of infinite loop spaces we refer to section 2 of [MT].

We first describe Riemann surfaces Σ with holomorphic actions of the q -th roots of unity $\mu_q \subset \mathbb{C}^\times$. This gives maps $B\mu_q \rightarrow B\text{Diff}(\Sigma)$.

Consider a divisor $D = \sum n_i p_i$ of $\mathbb{C}P^1$ with support $A = \{p_0, p_1, \dots, p_k\}$, $n_i \in \mathbb{N}$ and $n_0 + n_1 + \dots + n_k \equiv 0 \pmod{q}$. Assume for simplicity that $\gcd(q, n_i) = 1$ for $i = 0, 1, \dots, k$. Let Σ_D be the branched cover associated with the Galois extension

$$\mathbb{C}(z) \hookrightarrow \mathbb{C}(z)[T]/(F(T)), \quad F(T) = T^q - \prod_{i=0}^k (z - p_i)^{n_i}$$

(see e.g. [F], chap. 1.8). The Galois group is the group μ_q . The surface Σ_D has a holomorphic action of μ_q with orbit space $\mathbb{C}P^1$. The induced map $\pi : \Sigma_D \rightarrow \mathbb{C}P^1$ is holomorphic, branched over A , and $\pi^{-1}(p_i)$ is a single point for each i (since we assumed $\gcd(n_i, q) = 1$).

Thus the μ_q action on Σ_D is free outside A , and A is fixed pointwise by all elements in μ_q .

Let γ_i be a small loop in $\mathbb{C}P^1$ around p_i . The fundamental group $\mathbb{C}P^1 \setminus A$ is the free group of rank k generated by $\gamma_0, \dots, \gamma_k$ with the single relation $\prod \gamma_i = 1$. The covering $\Sigma_d \setminus A \rightarrow \mathbb{C}P^1 \setminus A$ is classified by the map from $\pi_1(\mathbb{C}P^1 \setminus A)$ to μ_q that sends γ_i to $e^{2\pi i n_i / q}$. The complex tangent line $T_{p_i} \Sigma_D$ at p_i is a μ_q representation; $u \in \mu_q$ multiplies by $u^{\bar{n}_i}$ where $\bar{n}_i \in \mathbb{Z}/q$ is the multiplicative inverse of n_i .

If D and D' are two divisors and q divides their difference $D - D'$, then there is a biholomorphic map between Σ_D and $\Sigma_{D'}$ that is equivariant w.r.t. the μ_q action. Thus it is only the class of D in $\tilde{H}_0(A; \mathbb{Z}/q)$ that matters.

Recall from Section 3.2 that given a surface bundle $\pi : E \rightarrow X$ there is a diagram

$$(5.1) \quad \begin{array}{ccccc} X & \longrightarrow & \Omega^\infty(\text{Th}(N_\pi E)) & \longrightarrow & \Omega^\infty S^\infty(E_+) \\ \parallel & & \downarrow & & T_\pi \downarrow \\ X & \xrightarrow{\alpha_{p_i}} & \Omega^\infty \mathbb{C}P^\infty_{-1} & \xrightarrow{\omega} & \Omega^\infty S^\infty(\mathbb{C}P^\infty_+) \end{array}$$

where $\Omega^\infty(\text{Th}(N_\pi E)) = \text{colim } \Omega^{n+2}(N_\pi^n e_n)$ for suitable fat fibrewise embeddings $e_n : E \rightarrow X \times \mathbb{R}^{n+2}$. The upper horizontal composition is the Becker-Gottlieb transfer map $t_E = t_\pi : X \rightarrow \Omega^\infty S^\infty(E_+)$.

We need the following properties of the Becker-Gottlieb transfer for smooth manifold bundles with compact fiber and compact Lie structure group:

(A1) Let $f : E \rightarrow E'$ be a fiberwise homotopy equivalence. Then

$$t_{E'} = \Omega^\infty S^\infty(f_+) \circ t_E \in [X, \Omega^\infty S^\infty(E'_+)].$$

(A2) Suppose

$$\begin{array}{ccc} E_{12} & \xrightarrow{j_1} & E_1 \\ j_2 \downarrow & & i_1 \downarrow \\ E_2 & \xrightarrow{i_2} & E \end{array}$$

is fiberwise homotopy co-Cartesian. In $[X, \Omega^\infty S^\infty(E_+)]$,

$$t_E = \Omega^\infty S^\infty(i_{1+}) \circ t_{E_1} + \Omega^\infty S^\infty(i_{2+}) \circ t_{E_2} - \Omega^\infty S^\infty(i_{12+}) \circ t_{E_{12}},$$

where $i_{12} = j_1 \circ i_1 = j_2 \circ i_2$.

(A3) If the tangent bundle along fibers $T_\pi E$ admits a non-zero section, then $t_E \in [X, \Omega^\infty S^\infty(E_+)]$ is the zero element.

The proof of (A1) and (A2) can be found in [LMS], pp. 189–190 or in [BS]. Property (A3) is much simpler. It follows because

$$\text{Th}(N_\pi E) \longrightarrow \text{Th}(N_\pi E \oplus T_\pi E)$$

is homotopic to the constant map at ∞ whenever $T_\pi E$ has an everywhere non-zero section.

Let $\Sigma = \Sigma_D$ be the μ_q -surface constructed above. We shall study the transfer of the associated smooth surface bundle

$$\pi : E\mu_q \times_{\mu_q} \Sigma \longrightarrow B\mu_q.$$

To shorten notation we write

$$t_\Sigma : B\mu_q \longrightarrow \Omega^\infty(E_+), \quad E = E\mu_q \times_{\mu_q} \Sigma$$

for the associated transfer.

Let $D = \sum n_i p_i$ and $q \in \mathbb{N}$ satisfy $A = \text{supp}(D) = \{p_0, \dots, p_k\}$, $n_0 + \dots + n_k \equiv 0$, $\text{gcd}(q, n_i) = 1$ for $i = 0, \dots, k$. For each i , the inclusion of $E\mu_q \times_{\mu_q} \{p_i\} \subset E$ induces a map

$$\hat{p}_i : B\mu_q \longrightarrow E \longrightarrow \Omega^\infty S^\infty(E_+).$$




FIGURE 1.

The principal μ_q bundle $E\mu_q \rightarrow B\mu_q$ induces a transfer from $B\mu_q$ to $\Omega^\infty S^\infty(E\mu_{q+}) \simeq \Omega^\infty S^\infty$, and hence

$$\hat{t}_q : B\mu_q \longrightarrow \Omega^\infty S^\infty \longrightarrow \Omega^\infty S^\infty(E_+)$$

upon choosing a point of E .

Lemma 5.1. *The transfer t_Σ is equal to $\sum \hat{p}_i + (1 - k)\hat{t}_q$ in $[B\mu_q, \Omega^\infty S^\infty(E_+)]$.*

Proof. We make a cell decomposition of $S^2 = \mathbb{C}P^1$ with two 0-cells $\{0, \infty\}$, $k + 1$ 1-cells I_i and $k + 1$ 2-cells D_i such that $p_i \in \text{int } D_i$. (See Figure 1.)

There are obvious co-Cartesian diagrams

$$\begin{array}{ccc} \bigsqcup \partial I_i & \longrightarrow & \bigsqcup I_i & & \bigsqcup \partial D_i & \longrightarrow & \bigsqcup D_i \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \{0, \infty\} & \longrightarrow & G & & G & \longrightarrow & S^2 \end{array}$$

where G denotes the 1-skeleton. This cell structure lifts to a cell structure of Σ :

$$(5.2) \quad \begin{array}{ccc} \bigsqcup \partial \tilde{I}_i & \longrightarrow & \bigsqcup \tilde{I}_i & & \bigsqcup \partial \tilde{D}_i & \longrightarrow & \bigsqcup \tilde{D}_i \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \{0, \infty\}^\sim & \longrightarrow & \tilde{G} & & \tilde{G} & \longrightarrow & \Sigma \end{array}$$

with $\tilde{G} = \pi^{-1}(G)$, etc. We can apply the functor $E\mu_q \times_{\mu_q} (-)$ to (5.2) and use (A1–A3) to evaluate $t_{\tilde{G}}$ and t_Σ . First by (A1) and (A2),

$$t_{\tilde{G}} = (1 + k)\hat{t}_q + 2\hat{t}_q - 2(k + 1)\hat{t}_q = (1 - k)\hat{t}_q,$$

since $\tilde{I}_i = \mu_q \times I_i$, $\partial \tilde{I}_i = \mu_q \times \partial I_i$ and $\{0, \infty\}^\sim = \mu_q \times \{0, \infty\}$. Second, the inclusion of p_i in \tilde{D}_i is a homotopy equivalence, so $t_{\tilde{D}_i} = \hat{p}_i$. Moreover, $E\mu_q \times_{\mu_q} \partial \tilde{D}_i = E\mu_q \times S^1$ has trivial vertical tangent bundle, so $t_{\partial \tilde{D}_i}$ is homotopically constant. One more application of (A1–A2) completes the proof. \square

The tangent representation $T_{p_i}\Sigma$ is given by multiplication with $e^{2\pi i\bar{n}_i/q}$, so an application of Lemma 5.1 gives

Corollary 5.2. *The homotopy class of*

$$B\mu_q \longrightarrow \Omega^\infty \mathbb{C}P_{-1}^\infty \longrightarrow \Omega^\infty S^\infty(\mathbb{C}P_+^\infty)$$

is $\sum \psi^{\bar{n}_i} + (1 - k)\hat{t}_q$. Here $\psi^{\bar{n}_i}$ is the composition

$$\psi^{\bar{n}_i} : B\mu_q \longrightarrow \mathbb{C}P^\infty \longrightarrow \Omega^\infty S^\infty(\mathbb{C}P_+^\infty)$$

with the left-hand map induced from the group homomorphism $\mu_q \rightarrow S^1$ that sends u to $u^{\bar{n}_i}$.

We are now ready to complete the proof of Theorem 2.2. As in section 3.3 of [MT] we let $q = p^n$ be a prime power and consider the divisor

$$D = p_0 + mp_1 + \cdots + mp_k, \quad m \equiv -1/k \pmod{p^n}.$$

We use the notation

$$F(n) = \Sigma_D, \quad C_{p^n} = \mu_{p^n}, \quad \tau_n = \hat{t}_{p^n}$$

and consider diagram (5.1) with

$$X = B\text{Diff}(F(n)), \quad E = E\text{Diff}(F(n)) \times_{\text{Diff}(F(n))} F(n).$$

Composing with the map $BC_{p^n} \rightarrow B\text{Diff}(F(n))$ induced by the C_{p^n} action on $F(n)$, and using that $B\Gamma_{g(n)} \simeq B\text{Diff}(F(n))$ with $g(n) = \frac{1}{2}(p^n - 1)(k - 1)$ we get the diagram

$$\begin{array}{ccccc} B\Gamma_{g(n)} & \xrightarrow{\alpha_n} & \Omega_{g(n)-1}^\infty \mathbb{C}P_{-1}^\infty & \xrightarrow{T} & \Omega_0^\infty \mathbb{C}P_{-1}^\infty \\ \mu_n \uparrow & & \omega \downarrow & & \omega \downarrow \\ BC_{p^n} & \longrightarrow & \Omega_{2-2g(n)}^\infty S^\infty(\mathbb{C}P_+^\infty) & \xrightarrow{T} & \Omega_0^\infty S^\infty(\mathbb{C}P_+^\infty). \end{array}$$

The right-hand horizontal maps are translations of the indicated component into the zero component. The lower horizontal composition is by Corollary 5.2 equal to

$$(1 + k\psi^{-k}, \tilde{\tau}_n) \in [BC_{p^n}, \Omega^\infty S^\infty(\mathbb{C}P^\infty) \times \Omega_0^\infty S^\infty]$$

with $\tilde{\tau}_n = T \circ \tau_n$.

Lemma 5.3. *Let $i_{n-1} : BC_{p^{n-1}} \rightarrow BC_{p^n}$ be the map associated with $C_{p^{n-1}} \subset C_{p^n}$. Then*

$$[\tau_n \circ i_{n-1}] = p[\tau_{n-1}] \in [BC_{p^{n-1}}, \Omega^\infty S^\infty].$$

Proof. Let E be a contractible space with a free action of C_{p^n} for all n , e.g. the union of odd-dimensional spheres $E = \bigcup_{m \geq 1} S^{2m-1}$. Consider the diagram

$$\begin{array}{ccccc} EC_{p^n} & \xrightarrow{\pi_{n-1}} & BC_{p^{n-1}} & \xrightarrow{i_{n-1}} & BC_{p^n} \\ \uparrow & & \uparrow & & \uparrow i_{n-1} \\ \bigsqcup_1^p EC_{p^n} & \longrightarrow & \bigsqcup_1^p BC_{p^{n-1}} & \longrightarrow & BC_{p^{n-1}} \end{array}$$

where $BC_{p^{n-1}}$ and BC_{p^n} are the orbit spaces $E/C_{p^{n-1}}$ and E/C_{p^n} and i_{n-1} is represented by the obvious quotient map. The lower sequence in the above diagram is the pull-back of the upper sequence, and $i_{n-1} \circ \pi_{n-1} = \pi_n$. The transfer of a composition is the composition of transfers, and transfers are natural for pull-backs.

Thus $[\tau_n \circ i_{n-1}]$ is the transfer of the lower sequence composed with the fold map $\bigsqcup_1^p E \rightarrow E$. This comes out to be $p[\tau_{n-1}]$. \square

The Ivanov-Harer stability theorems imply that the map of plus constructions,

$$B\Gamma_{g(n)-1,2}^+ \longrightarrow B\Gamma_{g(n)}^+,$$

is b_n -connected with $b_n = \lfloor \frac{g(n)-1}{2} \rfloor$. Thus

$$[BC_{p^n}^{(b_n)}, B\Gamma_{g(n)-1,2}^+] \simeq [BC_{p^n}^{(b_n)}, B\Gamma_{g(n)}^+],$$

where the superscript indicates the b_n -skeleton. Let $\mu_{n,2} : BC_{p^n}^{(b_n)} \rightarrow B\Gamma_{g(n)-1,2}^+$ correspond to μ_n so that

$$(5.3) \quad T \circ \omega \circ \alpha_{n,2} \circ \mu_{n,2} \simeq (1 + k\psi^{-k}, (1 - k)\tilde{\tau}_n)$$

in $[BC_{p^n}^{(b_n)}, \Omega_0^\infty S^\infty(\mathbb{C}P_+^\infty)]$. Replacing $\mu_{n,2}$ by the composition $\mu_{n+l,2} \circ i_{n+l-1} \circ \dots \circ i_n$ on the left side of (5.3), by Lemma 5.3, the second component on the right-hand side has to be replaced by $p^l(1 - k)\tilde{\tau}_n$. This second component is an element in $[BC_{p^n}^{(b_n)}, (\Omega_0^\infty S^\infty)_p^\wedge]$ which is a finite p -group. Hence, for some l large enough, $p^l(1 - k)\tilde{\tau}_n$ is homotopic to zero.

Consider the subset G_n of $[BC_{p^n}^{(b_n)}, (B\Gamma_{\infty,2}^+)_p^\wedge]$ of elements that satisfy (5.3) with the second component on the right-hand side actually zero. As we have argued it is non-empty. Furthermore, any two elements in G_n differ by a map into the fibre of ω . The set $[BC_{p^n}^{(b_n)}, \Omega_0^\infty S^{\infty-2}]$ is however finite and hence G_n is finite. By Tychonov's theorem, the inverse limit of the G_n is therefore non-empty. We pick $\tilde{\mu}_p \in \varprojlim G_n$. Since $\text{colim}_{n \rightarrow \infty} BC_{p^n}^{(b_n)} = BC_{p^\infty}$ has p -adic completion homotopy equivalent to $(\mathbb{C}P^\infty)_p^\wedge$, the map $\tilde{\mu}_p$ extends to a map

$$\tilde{\mu}_p : \Omega^\infty S^\infty(\mathbb{C}P^\infty) \longrightarrow (B\Gamma_{\infty,2}^+)_p^\wedge$$

and

$$T \circ \omega \circ \alpha \circ \mu_p \simeq (1 + k\psi^k, 0).$$

This proves that the first column in the matrix described in Theorem 2.2 is $(1 + k\psi^{-k}, 0)$. The second column corresponds to the homotopy class of

$$(5.4) \quad \Omega^\infty S^\infty \xrightarrow{\mu} \mathbb{Z} \times B\Gamma_{\infty,2}^+ \xrightarrow{\alpha} \Omega^\infty \mathbb{C}P_{-1}^\infty \xrightarrow{\omega} \Omega^\infty S^\infty(\mathbb{C}P_+^\infty).$$

An infinite loop map with source $\Omega^\infty S^\infty$ is determined by its restriction to $S^0 = \{-1, +1\} \hookrightarrow \Omega^\infty S^\infty$. The μ above maps the basepoint $+1$ of S^0 into $(0, *)$ and the non-basepoint into $(1, *)$. The composition (5.4) maps the basepoint into the basepoint of $\Omega_0^\infty S^\infty(\mathbb{C}P^\infty)$ and the non-basepoint into the basepoint of $\Omega_{-2}^\infty S^\infty(\mathbb{C}P_+^\infty)$. This shows that the second column of μ of the matrix in Theorem 2.2 is $(0, -2)$ as claimed. \square

5.2. The referee's proof. By Theorem 2.4, we may identify $\mathbb{Z} \times B\Gamma_\infty^+$ with $\Omega^\infty \mathbb{C}P_{-1}^\infty$ via the map α . Hence it suffices to show that the self-map

$$(1 + k\psi^{-k}, -2) : \Omega^\infty S^\infty(\mathbb{C}P^\infty) \times \Omega^\infty S^\infty \longrightarrow \Omega^\infty S^\infty(\mathbb{C}P^\infty) \times \Omega^\infty S^\infty$$

lifts through

$$\omega : \Omega^\infty \mathbb{C}P_{-1}^\infty \rightarrow \Omega^\infty S^\infty(\mathbb{C}P_+^\infty) \simeq \Omega^\infty S^\infty(\mathbb{C}P^\infty) \times \Omega^\infty S^\infty$$

after p -completion. By the discussion in section 2.1, the factor of -2 on the summand $\Omega^\infty S^\infty$ can be lifted by sending the non-basepoint of S^0 to the basepoint of

the 1-component in $\Omega^\infty \mathbb{C}P_{-1}^\infty$, and extending this map to the uniquely determined infinite loop space map from $\Omega^\infty S^\infty$. Hence, by the fibration sequence (2.3), we are left to show that after p -completion the composite map

$$(5.5) \quad \Omega^\infty S^\infty(\mathbb{C}P^\infty) \xrightarrow{1+k\psi^k} \Omega^\infty S^\infty(\mathbb{C}P^\infty) \xrightarrow{\partial} \Omega^\infty S^{\infty-1}$$

is null-homotopic.

We recall that ∂ is the S^1 -transfer map; cf. [R]. Stefan Stolz gave the following, alternative description of the S^1 -transfer map. Let $l : \mathbb{C}P^\infty \rightarrow BU$ be the map that represents the reduced canonical line bundle, $\beta : BU \rightarrow \Omega U$ be the Bott map, and $J : U \rightarrow \Omega^\infty S^\infty$ be the J -homomorphism that associates to a unitary transformation the induced map of spheres.

Lemma 5.4 ([St]). *The map ∂ is homotopic to the infinite loop space map induced by the composite map*

$$\mathbb{C}P^\infty \xrightarrow{l} BU \xrightarrow{\beta} \Omega U \xrightarrow{\Omega J} \Omega^\infty S^{\infty-1}.$$

The maps ψ^{-k} give rise to the Adams operations $\psi^{-k} : BU \rightarrow BU$. These operations are stable in the sense that they satisfy the identity

$$\Omega^2(\psi^{-k}) \circ \beta \simeq \beta \circ (-k\psi^{-k}).$$

Here we identify $U \simeq \Omega BU$. Hence, we have a commutative diagram

$$\begin{array}{ccc} \mathbb{C}P^\infty & \xrightarrow{1+k\psi^{-k}} & \Omega^\infty S^\infty(\mathbb{C}P^\infty) \\ \downarrow l & & \downarrow l \\ BU & \xrightarrow{1+k\psi^{-k}} & BU \\ \downarrow \beta & & \downarrow \beta \\ \Omega^2 BU & \xrightarrow{\Omega^2(1-\psi^{-k})} & \Omega^2 BU \end{array} \quad \begin{array}{c} \nearrow \partial \\ \nearrow \Omega^2 BJ \end{array} \rightarrow \Omega^\infty S^{\infty-1}.$$

By the affirmed Adams conjecture, after inverting $-k$ and hence after p -completion, $BJ \circ (1 - \psi^{-k})$ is null-homotopic. Thus the bottom row defines a null-homotopic map after p -completion, which proves (5.5). □

6. PROOF OF THEOREM 1.5

It follows from [MS, Theorem 7.8] that for odd primes p there is a splitting of p -localized spaces

$$(\Omega_0^\infty \mathbb{C}P_{-1}^\infty)_{(p)} \simeq T_{(p)} \times BU_{(p)},$$

where $T_{(p)}$ has torsion homotopy and homology groups. Therefore we have

$$H_{free}^*(\Omega_0^\infty \mathbb{C}P_{-1}^\infty; \mathbb{Z}_{(p)}) \simeq H_{free}^*(BU; \mathbb{Z}_{(p)}) = H^*(BU; \mathbb{Z}_{(p)}).$$

For $p = 2$ we use the fact that $H_*(\Omega^\infty \mathbb{C}P_{-1}^\infty; \mathbb{F}_2)$ is a sub-Hopf-algebra of $H_*(\Omega^\infty S^\infty \mathbb{C}P_+^\infty; \mathbb{F}_2)$ by theorem 1.3 of [G1]. We need explicit additive generators in low dimensions. Let $a_i \in H_{2i}(\Omega^\infty S_1^\infty \mathbb{C}P_+^\infty; \mathbb{F}_2)$ be the non-zero element in the image of $\mathbb{C}P^\infty \rightarrow \Omega^\infty S_1^\infty \mathbb{C}P_+^\infty$ into the 1-component of $\Omega^\infty S^\infty \mathbb{C}P_+^\infty$ and let

$$Q^s : H_*(\Omega^\infty S^\infty \mathbb{C}P_+^\infty; \mathbb{F}_2) \rightarrow H_{*+s}(\Omega^\infty S^\infty \mathbb{C}P_+^\infty; \mathbb{F}_2)$$

be the s -th homology operation for infinite loop spaces. The following elements form an \mathbb{F}_2 -basis for $H_*(\Omega_0^\infty \mathbb{C}P_{-1}^\infty; \mathbb{F}_2)$:

$$\begin{aligned} x_2 &= (Q^1 a_0)^2 a_0^{-4}, \\ x_3 &= (Q^3 a_0) a_0^{-2} + (Q^2 Q^1 a_0) a_0^{-4} + (Q^2 a_0)(Q^1 a_0) a_0^{-4} + (Q^1 a_0)^3 a_0^{-6}, \\ x_4 &= (a_1)^2 a_0^{-2}, \quad y_4 = (Q^2 a_0)^2 a_0^{-4}, \quad x_2^2, \end{aligned}$$

in positive degrees less than 5. The rational homology of $\Omega_0^\infty \mathbb{C}P_{-1}^\infty$ and BU agree, so

$$H_2^{free}(\Omega_0^\infty \mathbb{C}P_{-1}^\infty) = \mathbb{Z}, \quad H_3^{free}(\Omega_0^\infty \mathbb{C}P_{-1}^\infty) = 0, \quad H_4^{free}(\Omega_0^\infty \mathbb{C}P_{-1}^\infty) = \mathbb{Z} \oplus \mathbb{Z}.$$

We now employ the Bockstein spectral sequence. For an infinite loop space X , it is a singly graded spectral sequence of Hopf algebras with

$$E_*^1 = H_*(X; \mathbb{F}_2), \quad E_*^\infty = H_*^{free}(X) \otimes \mathbb{F}_2.$$

The differential d^1 is the standard Bockstein [Br]. The elements listed above for $X = \Omega_0^\infty \mathbb{C}P_{-1}^\infty$ all survive to E_*^2 . The general formula for d^2 applied to an even square [M, Proposition 1.5] gives

$$d^2 y_4 = x_3 + (Q^3 a_0) a_0^{-2} + (Q^1 a_0)^3 a_0^{-6} = x_3$$

in $E_*^2(\Omega_0^\infty S_0^\infty \mathbb{C}P_+^\infty)$, and $d^2 x_4 = d^2 x_2^2 = 0$. On the other hand, the E_*^∞ -term is \mathbb{F}_2 , 0, and $\mathbb{F}_2 \oplus \mathbb{F}_2$ in degrees 2, 3 and 4, respectively. It follows that

$$E_2^\infty(\Omega_0^\infty \mathbb{C}P_{-1}^\infty) = \mathbb{F}_2 \langle x_2 \rangle, \quad E_4^\infty(\Omega_0^\infty \mathbb{C}P_{-1}^\infty) = \mathbb{F}_2 \langle x_4 \rangle \oplus \mathbb{F}_2 \langle x_2^2 \rangle.$$

Both x_2^2 and x_4 are primitive elements (since that is the case in $\Omega_0^\infty S_0^\infty \mathbb{C}P_+^\infty$), so the reduced diagonal from E_4^∞ to $E_2^\infty \otimes E_2^\infty$ is zero. Dually the squaring map is zero on $H_{free}^2(\Omega_0^\infty \mathbb{C}P_{-1}^\infty) \otimes \mathbb{F}_2$, so $H_{free}^*(\Omega_0^\infty \mathbb{C}P_{-1}^\infty; \mathbb{Z}_{(2)})$ is not polynomial. \square

Remark. The non-zero differential $d^2 : E_4^2 \rightarrow E_3^2$ proves that the two-torsion in $H_3(B\Gamma_\infty)$ is $\mathbb{Z}/4\mathbb{Z}$.

REFERENCES

- [A1] J.F. Adams, *Lectures on generalized cohomology, category theory, homology theory and their applications III*, LNM vol. 99, Springer, 1969. MR0251716 (40:4943)
- [A2] J.F. Adams, On the groups $J(X)$ II, *Topology*, vol. 3, 1965, pp. 137–171. MR0198468 (33:6626)
- [Ak] T. Akita, *Nilpotency and triviality of mod p Morita-Mumford classes of mapping class groups of surfaces*, Nagoya Math. J., vol. 165, 2002, pp. 1–22. MR1892095 (2002k:55033)
- [BS] J.C. Becker, R.E. Schultz, *Axioms for transfers and traces*, *Math. Z.*, vol. 227, 1988, pp. 583–605. MR1621939 (99f:55021)
- [B] J. Berrick, *An approach to algebraic K-theory*, Research Notes in Mathematics, vol. 56, Pitman (Advanced Publishing Program), Boston, 1982. MR0649409 (84g:18028)
- [BK] A.K. Bousfield, D.M. Kan, *Homotopy Limits, Completions and Localizations*, LNM vol. 304, Springer LNM, 1972. MR0365573 (51:1825)
- [Br] W. Browder, *Torsion in H-spaces*, vol. 74, *Ann. of Math.*, 1961, pp. 24–51. MR0124891 (23:A2201)
- [C] M. Crabb, *$\mathbb{Z}/2$ -Homotopy Theory*, LMS Lecture Note Series, vol. 44, Cambridge University Press, 1980. MR0591680 (83m:55010)
- [EE] C.J. Earle, J. Eells, *A fibre bundle description of Teichmüller theory*, *J. Diff. Geom.*, vol. 3, 1969, pp. 19–43. MR0276999 (43:2737a)
- [ES] C.J. Earle, A. Schatz, *Teichmüller theory for surfaces with boundary*, *J. Diff. Geom.*, vol. 4, 1970, pp. 169–185. MR0277000 (43:2737b)
- [F] O. Forster, *Lectures on Riemann surfaces*, Graduate Texts in Mathematics, vol. 81, Springer, New York-Berlin, 1991. MR1185074 (93h:30061)

- [G1] S. Galatius, *Mod p homology of the stable mapping class group*, *Topology*, vol. 43, 2004, pp. 1105–1132. MR2079997 (2006a:57020)
- [G2] S. Galatius, *Secondary characteristic classes of surface bundles*, preprint, math.AT/0402226.
- [H1] J.L. Harer, *The second homology group of the mapping class group of an orientable surface*, *Invent. Math.*, vol. 72, 1983, pp. 221–239. MR0700769 (84g:57006)
- [H2] J.L. Harer, *Stability of the homology of the mapping class groups of orientable surfaces*, *Annals Math.*, vol. 121, 1985, pp. 215–249. MR0786348 (87f:57009)
- [I] N.V. Ivanov, *Stabilization of the homology of Teichmüller modular groups*, Original: *Algebra i Analiz*, vol. 1, 1990, pp. 675–691, Leningrad Math. J. MR1015128 (91g:57010)
- [LMS] G. Lewis, J. P. May, M. Steinberger, *Equivariant stable homotopy theory*, LNM vol. 1213, Springer, 1986. MR0866482 (88e:55002)
- [M] I. Madsen, *Higher Torsion in SG and BSG* , vol. 143, *Math. Z.*, 1975, pp. 55–80. MR0375307 (51:11503)
- [MS] I. Madsen, C. Schlichtkrull, *The circle transfer and K -theory*, *AMS Contemporary Math.*, vol. 258, 2000, pp. 307–328. MR1778114 (2001e:55021)
- [MT] I. Madsen, U. Tillmann, *The stable mapping class group and $Q(\mathbb{C}P^\infty)$* , vol. 145, *Invent. Math.*, 2001, pp. 509–544. MR1856399 (2002h:55011)
- [MSt] J. Milnor, J. Stasheff, *Characteristic Classes*, Study, vol. 76, Princeton University Press, 1974. MR0440554 (55:13428)
- [MW] I. Madsen, M. Weiss, *The stable moduli space of Riemann surfaces: Mumford’s conjecture*, math.AT/0212321.
- [Mi] E.Y. Miller, *The homology of the mapping class group*, vol. 24, *J. Diff. Geom.*, 1986, pp. 1–14. MR0857372 (88b:32051)
- [Mo] S. Morita, *Characteristic classes of surface bundles*, vol. 90, *Invent. Math.*, 1987, pp. 551–577. MR0914849 (89e:57022)
- [Mu] D. Mumford, *Towards an enumerative geometry of the moduli space of curves in Arithmetic and Geometry* (M. Artin and J. Tate, editors), *Progr. Math.*, vol. 36, Birkhäuser, 1983, pp. 271–328. MR0717614 (85j:14046)
- [R] D. Ravenel, *The Segal conjecture for cyclic groups and its consequences*, vol. 106, *Amer. J. Math.*, 1984, pp. 415–446. MR0737779 (85g:55015)
- [S] G. Segal, *The stable homotopy of complex projective space*, vol. 24, *Quart. J. Math. Oxford*, 1973, pp. 1–5. MR0319183 (47:7729)
- [St] S. Stolz, *Relationships between transfer and J -homomorphisms*, vol. 125, *Bonner Mathematische Zeitschriften*, Universität Bonn, 1980. MR0598816 (82g:57012)
- [T] U. Tillmann, *On the homotopy of the stable mapping class group*, vol. 130, *Invent. Math.*, 1997, pp. 257–275. MR1474157 (99k:57036)

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CALIFORNIA 94305
E-mail address: galatius@math.stanford.edu

MATEMATISK INSTITUT, AARHUS UNIVERSITET, 8000 AARHUS C, DENMARK
E-mail address: imadsen@imf.au.dk

MATHEMATICAL INSTITUTE, 24-29 ST. GILES STREET, OXFORD OX1 3LB, UNITED KINGDOM
E-mail address: tillmann@maths.ox.ac.uk