

COFINALITY SPECTRUM THEOREMS IN MODEL THEORY, SET THEORY, AND GENERAL TOPOLOGY

M. MALLIARIS AND S. SHELAH

CONTENTS

1. Introduction	237
1.1. List of main theorems	242
2. Cofinality Spectrum Problems	243
2.1. A motivating example	243
2.2. Main definitions	245
2.3. Key examples	249
2.4. Basic tools	250
3. The Function lcf Is Well Defined Strictly Below $\min\{\mathfrak{p}_s^+, \mathfrak{t}_s\}$	250
4. Cofinality Spectrum Problems Are Locally Saturated	254
5. Cofinality Spectrum Problems Have Sufficient Peano Arithmetic	255
6. There Are No Symmetric Cuts Strictly Below $\min\{\mathfrak{p}_s^+, \mathfrak{t}_s\}$	260
7. If $\text{lcf}(\aleph_0, \mathfrak{s}) > \aleph_1$, Then $\text{lcf}(\aleph_0, \mathfrak{s}) \geq \min\{\mathfrak{p}_s^+, \mathfrak{t}_s\}$	262
8. There Are No Asymmetric Cuts Strictly Below $\min\{\mathfrak{p}_s^+, \mathfrak{t}_s\}$	265
9. $\mathcal{C}(\mathfrak{s}, \mathfrak{t}_s) = \emptyset$	271
10. A New Characterization of Good Regular Ultrafilters	272
11. SOP_2 Implies Maximality in Keisler’s Order	278
12. If \mathcal{D} Is Good for Some Nonsimple Theory, Then \mathcal{D} Is Flexible	282
13. There Is a Minimum Nonsimple Class in Keisler’s Order	285
14. $\mathfrak{p} = \mathfrak{t}$	289
Acknowledgments	295
References	295

1. INTRODUCTION

We connect and solve two fundamental open problems in quite different areas: the model-theoretic problem of the maximality of SOP_2 theories in Keisler’s order, and the problem from general topology/set theory of whether $\mathfrak{p} = \mathfrak{t}$, as well as some natural set-theoretic questions about cuts in regular ultrapowers of linear order.

Received by the editors August 27, 2012 and, in revised form, January 14, 2014 and January 11, 2015.

2010 *Mathematics Subject Classification*. Primary 03C20, 03C45, 03E17; Secondary 03E05.

The first author was partially supported by NSF grant DMS-1001666, by Shelah’s NSF grant DMS-1101597, by a Gödel research fellowship, and by a Sloan fellowship.

The second author was partially supported by the Israel Science Foundation grants 710/07 and 1053/11. This is paper 998 in Shelah’s list of publications.

Let us begin with the simpler of the two problems to state: whether $\mathfrak{p} = \mathfrak{t}$. Cantor proved in 1874 that the continuum is uncountable, i.e., $\aleph_0 < 2^{\aleph_0}$ [6]. The study of cardinal invariants or characteristics of the continuum illuminates this gap by studying connections between cardinals measuring the continuum which arise from different perspectives: combinatorics, algebra, topology, and measure theory. Although there are many cardinal invariants and many open questions about them (see, e.g., the surveys of van Douwen [8], Vaughan [49], and Blass [5]), the problem of whether $\mathfrak{p} = \mathfrak{t}$ is the oldest and so holds a place of honor. (Moreover, usually if such an equality was not obviously true it was consistently false, by forcing.)

Before reviewing the history, we give the easily stated definition.

Definition 1.1. (See, e.g., [8].) *We define several properties which may hold of a family $D \subseteq [\mathbb{N}]^{\aleph_0}$, i.e., a family of infinite sets of natural numbers. Let $A \subseteq^* B$ mean that $\{x : x \in A, x \notin B\}$ is finite.*

- *D has a pseudo-intersection if there is an infinite $A \subseteq \mathbb{N}$ such that for all $B \in D, A \subseteq^* B$.*
- *D has the s.f.i.p. (strong finite intersection property) if every nonempty finite subfamily has infinite intersection.*
- *D is called a tower if it is well ordered by \supseteq^* and has no infinite pseudo-intersection.*

Then,

$$\mathfrak{p} = \min\{|\mathcal{F}| : \mathcal{F} \subseteq [\mathbb{N}]^{\aleph_0} \text{ has the s.f.i.p. but has no infinite pseudo-intersection}\},$$

$$\mathfrak{t} = \min\{|\mathcal{T}| : \mathcal{T} \subseteq [\mathbb{N}]^{\aleph_0} \text{ is a tower}\}.$$

Clearly, both \mathfrak{p} and \mathfrak{t} are at least \aleph_0 and no more than 2^{\aleph_0} . It is easy to see that $\mathfrak{p} \leq \mathfrak{t}$, since a tower has the s.f.i.p. By a 1934 theorem of Hausdorff $\aleph_1 \leq \mathfrak{p}$ [13]. In 1948 [37], Rothberger proved (in our terminology) that $\mathfrak{p} = \aleph_1$ implies $\mathfrak{p} = \mathfrak{t}$, which begs the question of whether $\mathfrak{p} = \mathfrak{t}$.

Problem 1. *Is $\mathfrak{p} = \mathfrak{t}$?*

Problem 1 appears throughout the literature. Van Douwen presents six primary cardinal invariants (of a model of set theory), $\mathfrak{a}, \mathfrak{b}, \mathfrak{d}, \mathfrak{p}, \mathfrak{s}$, and \mathfrak{t} ; he attributes $\mathfrak{b}, \mathfrak{p}, \mathfrak{t}$ to Rothberger in 1939 and 1948, \mathfrak{d} to Katetov in 1960, \mathfrak{a} to Hechler in 1972 and Solomon in 1977, and \mathfrak{s} to Booth in 1974 (see [8], p. 123). Vaughan’s [49] Problem 1.1 includes the only inequalities about van Douwen’s six cardinals which remained open in 1990; it is noted there that “we believe (a)” (i.e., whether $\mathfrak{p} < \mathfrak{t}$ is consistent with Zermelo-Fraenkel set theory with the axiom of choice (ZFC)) “is the most interesting.” Following Shelah’s solution of Vaughan’s 1.1(b) in [43] (showing it was independent), Problem 1 is therefore both the oldest and the only remaining open inequality about van Douwen’s cardinals.

There has been much work on \mathfrak{p} and \mathfrak{t} , for example: Bell [4] proved that \mathfrak{p} is the first cardinal μ for which $MA_\mu(\sigma\text{-centered})$ fails; in topological language, this asserts that no separable compact Hausdorff space can be covered by fewer than \mathfrak{p} nowhere dense sets. Szymański proved that \mathfrak{p} is regular. Piotrowski and Szymański [34] proved that $\mathfrak{t} \leq \text{add}(\mathcal{M})$, where \mathcal{M} is the ideal of meager sets, and $\text{add}(I)$ denotes the smallest number of sets in an ideal I with the union not in I . Shelah proved in [44] that if $\mathfrak{p} < \mathfrak{t}$, then there is a so-called peculiar cut in ${}^\omega\omega$ (see Section 14, Theorem G below); we will leverage this result in the present work.

In Section 14 of this paper, we apply the methods developed in earlier sections to answer Problem 1.

Theorem. (Theorem 14.1) $\mathfrak{p} = \mathfrak{t}$.

In the context of the general framework we build (of cofinality spectrum problems), this answer is natural, but it is a priori very surprising. Given the length of time this problem had remained open, the expectation was an independence result.

We now describe the second problem, which concerns a criterion for maximality in Keisler's 1967 order on theories. In this discussion, all theories are complete and countable. If \mathcal{D} is an ultrafilter on λ , let us say that \mathcal{D} *saturates* M if the ultrapower M^λ/\mathcal{D} is λ^+ -saturated.¹ When \mathcal{D} is a regular ultrafilter on λ and M, N are both models of the same complete, countable first-order theory T , then \mathcal{D} saturates M if and only if (iff) \mathcal{D} saturates N , so we may simply say that \mathcal{D} saturates T .

Keisler's order is the relation: $T_1 \preceq T_2$ iff for any infinite cardinal λ and any regular ultrafilter \mathcal{D} on λ ; if \mathcal{D} saturates T_2 , then \mathcal{D} saturates T_1 . Keisler's order is a pre-order on theories, and a partial order on the equivalence classes. It was recognized early on that this order gave a powerful way of comparing the complexity of any two theories, even in different languages. Determining the structure of this order is a large-scale, and largely open, problem in model theory. It is not known, for instance, whether the order is linear, or whether it is absolute, although it has long been thought to have finitely many classes and be linearly ordered. Its structure on the stable theories is known [40], and the first dividing line in Keisler's order among the unstable theories has just been discovered [28].

It has been known since 1967 that the order has a maximum class, whose model-theoretic identity has remained elusive. Around 1963, Keisler proved the existence of a strong family of ultrafilters, the *good* regular ultrafilters: see [14], or Definition 10.8 below.² In defining his order, Keisler connected these filters to model theory by showing that on the one hand, good regular ultrafilters saturate any theory, and on the other, there exist theories complex enough to be able to code any failure of goodness as an omitted type in the ultrapower. In other words, there exist T such that \mathcal{D} saturates T iff \mathcal{D} is good. This shows the existence of a maximum class in Keisler's order, which can be characterized set-theoretically as the set of complete countable first-order theories which are saturated only by those regular \mathcal{D} which are good.

A first surprise was that complexity in the sense of coding is not an essential model-theoretic feature of theories in the maximum Keisler class: in 1978, Shelah [40] proved that any theory of linear order, or more precisely with the strict order property, belongs to the maximum class. Later, this was weakened to the strong order property SOP_3 , which retains many features of linear order. For a long time, it was not clear whether to expect that any theories *without* the obvious features of linear order might belong to the maximum class. Attention focused on SOP_2 , a kind of maximally inconsistent tree: we say that a theory T has SOP_2

¹Saturation is a fullness condition. Given a model M and $A \subseteq \text{dom}(M)$, the n -types over A are the maximal consistent sets of formulas in n free variables with parameters from A . A type $p(\bar{x})$ is realized in M if there is \bar{a} such that $M \models \varphi(\bar{a})$ for each $\varphi \in p$; otherwise it is omitted. A model is called κ -saturated if all types, or equivalently all 1-types, over all subsets of M of size $< \kappa$ are realized.

²Keisler's proof assumed the generalized continuum hypothesis (GCH), which was later eliminated by Kunen.

if for some formula³ $\varphi(\bar{x}, \bar{y})$, we may, in some sufficiently saturated model of T , find a set of parameters for φ indexed by the nodes of a binary tree such that the set of instances of φ along any path through the tree is consistent, whereas any two instances assigned to incomparable nodes are inconsistent. This occurs easily in models of linear order, but also in the generic triangle-free graph. (These properties are discussed further in Section 10.) The second major problem this paper addresses is the following.

Problem 2. *Does SOP_2 imply maximality in Keisler's order?*

Problem 2 does not settle the identity of the maximum class; why, then, is it so significant? For one, it moves the possible boundary of the maximum class onto what we believe to be a dividing line, in the sense of classification theory. Moreover, we believe the following conjecture.

Conjecture 1.2. *SOP_2 characterizes maximality in Keisler's order.*

Evidence for this conjecture is discussed in 11.12 below. In Section 11 below we give a positive answer to Problem 2.

Theorem. (Theorem 11.11) *Let T be a theory with SOP_2 . Then T is maximal in Keisler's order.*

To prove Theorem 11.11 we develop a general theory of cofinality spectrum problems, a main contribution of the paper (which we will return to presently). Applying a theorem of Malliaris [24, 25], we then show the following theorem.

Theorem. (Theorem 13.1) *There is a minimum class among the nonsimple theories in Keisler's order, which contains the theory T_{feq} of a parametrized family of independent equivalence relations.*

In order to state the paper's remaining central theorem, we now outline some features of our approach which connects and solves both Problems 1 and 2 above. In this discussion, we use \mathcal{D} to denote a regular ultrafilter on λ .

By *cut* we will mean an unfilled pre-cut (in set-theoretic literature cuts in reduced powers are often referred to as *gaps*). Define the cut spectrum of the ultrafilter \mathcal{D} to be

$$\mathcal{C}(\mathcal{D}) = \{(\kappa_1, \kappa_2) : \kappa_1, \kappa_2 \text{ regular, } \kappa_1 + \kappa_2 \leq \lambda, (\mathbb{N}, <)^\lambda / \mathcal{D} \text{ has a } (\kappa_1, \kappa_2)\text{-cut}\}.$$

Since any theory of linear order is in the maximal Keisler class, we have the following fact.

Fact 1.3. *Let \mathcal{D} be a regular ultrafilter on λ . Then \mathcal{D} is λ^+ -good iff $\mathcal{C}(\mathcal{D}) = \emptyset$.*

By a tree $(\mathcal{T}, \trianglelefteq)$ we will mean a set \mathcal{T} of sequences partially ordered by initial segment \trianglelefteq such that the set of predecessors of any element of \mathcal{T} is well-ordered. Say that \mathcal{D} has κ -treetops if for any tree $(\mathcal{T}, \trianglelefteq)$ and any infinite regular cardinal $\gamma < \kappa$, any strictly increasing γ -indexed sequence in $(\mathcal{T}, \trianglelefteq)^\lambda / \mathcal{D}$ has an upper bound. In Section 11 below, we prove that if \mathcal{D} saturates some theory with SOP_2 , then \mathcal{D} has λ^+ -treetops. Then a positive answer to Problem 2 would follow from a positive answer to Problem 3.

Problem 3. *Suppose \mathcal{D} has λ^+ -treetops. Is $\mathcal{C}(\mathcal{D}) = \emptyset$?*

³For transparency we simplify: we should really speak about pairs (T, φ) .

This is because a positive answer to Problem 3 would establish that any ultrafilter able to saturate some theory with SOP_2 is necessarily good.

To further compare $\mathcal{C}(\mathcal{D})$ with the treetops of \mathcal{D} , let us define $\mathfrak{p}_{\mathcal{D}}$ to be the least κ such that there exist κ_1, κ_2 with $\kappa_1 + \kappa_2 = \kappa$ and $(\kappa_1, \kappa_2) \in \mathcal{C}(\mathcal{D})$. Likewise, define $\mathfrak{t}_{\mathcal{D}}$ to be the greatest κ such that \mathcal{D} has κ -treetops. Below $\mathfrak{p}_{\mathcal{D}}$ there are no cuts, then, and below $\mathfrak{t}_{\mathcal{D}}$ infinite paths through trees have upper bounds. The names $\mathfrak{p}_{\mathcal{D}}$, $\mathfrak{t}_{\mathcal{D}}$ are chosen to be suggestive of the cardinal invariants of Section 14, and note that Problem 3 is equivalent to asking the question of Problem 3.

Problem 4. *Can it happen that $\mathfrak{p}_{\mathcal{D}} < \mathfrak{t}_{\mathcal{D}}$?*

This analogy between Problems 1 and 2 becomes formal as follows. Preliminary investigation, e.g., the claim in Section 2.1 below, shows that the hypothesis of treetops has nontrivial consequences for the possible cuts in ultrapowers of linear orders. Moreover, such proofs may be carried out using only a few essential features of ultrapowers: the fact that ultrapower of $(\mathbb{N}, <)$ is pseudo-finite (it remains a discrete linear order in which any nonempty, bounded, definable subset has a first and a last element), and the fact that ultrapowers commute with reducts (meaning here that we may expand a given model of linear order to have available certain trees of functions from the order to itself, which the proof may then manipulate). Section 2.1 below gives further details.

In Section 2, we define the true general context: cofinality spectrum problems, a central definition of the paper. These are given essentially as the data \mathbf{s} of an elementary pair of models $M \preceq M_1$, equipped with a suitably closed set of formulas Δ which define distinguished pseudo-finite linear orders in M_1 , and which admit expansion to the elementary pair $M^+ \preceq M_1^+$ where this expanded language has available uniform definitions for certain trees of functions from each distinguished linear order to itself (trees under the partial order given by the initial segment). For a given cofinality spectrum problem \mathbf{s} , let $\text{Or}(\mathbf{s})$ denote the set of distinguished orders and $\text{Tr}(\mathbf{s})$ the set of associated trees. Let $\mathfrak{t}_{\mathbf{s}}$ be the minimal regular cardinal τ such that some tree in $\text{Tr}(\mathbf{s})$ has a strictly increasing τ -sequence with no upper bound. When $\lambda < \mathfrak{t}_{\mathbf{s}}$, we say \mathbf{s} has λ^+ -treetops. Let $\mathcal{C}^{\text{ct}}(\mathbf{s})$ be the set of pairs of regular cardinals (κ_1, κ_2) which appear as the cofinalities of a cut in some distinguished linear order of \mathbf{s} .

In this context, the problem of comparing linear orders and trees has become Question 1.4.

Question 1.4 (Central question). *Let \mathbf{s} be a cofinality spectrum problem. What are the possible values of*

$$\mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}}) = \{(\kappa_1, \kappa_2) : (\kappa_1, \kappa_2) \in \mathcal{C}^{\text{ct}}(\mathbf{s}), \kappa_1 + \kappa_2 < \mathfrak{t}_{\mathbf{s}}\} \quad ?$$

In particular, is $\mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}}) = \emptyset$? That is, for a cofinality spectrum problem \mathbf{s} , can it happen that $\mathfrak{p}_{\mathbf{s}} < \mathfrak{t}_{\mathbf{s}}$? As the definition of cofinality spectrum problems includes that of regular ultrapowers, this generalizes Problem 4. Thus, the maximality of SOP_2 in Keisler's order would follow from proving that for any cofinality spectrum problem \mathbf{s} , $\mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}}) = \emptyset$.

In Section 14, it is proved that *if $\mathfrak{p} < \mathfrak{t}$, then there exists a cofinality spectrum problem in which $\mathfrak{p}_{\mathbf{s}} \leq \mathfrak{p} < \mathfrak{t} \leq \mathfrak{t}_{\mathbf{s}}$, where $\mathfrak{p}, \mathfrak{t}$ are the cardinal invariants from Definition 1.1. (In fact, $\mathfrak{p}_{\mathbf{s}} = \mathfrak{p}$ and $\mathfrak{t}_{\mathbf{s}} = \mathfrak{t}$, though this is not used.) This cofinality spectrum problem is built using $M = M^+ = (\mathcal{H}(\omega_1), \epsilon)$ with \mathbf{G} a generic ultrafilter*

for the forcing notion $([\omega]^{\aleph_0}, \supseteq^*)$ and $M_1 = M_1^+$ the generic ultrapower M^ω/\mathbf{G} . By definition of \mathfrak{p}_s , this says that if $\mathfrak{p} < \mathfrak{t}$, there is a cofinality spectrum problem \mathfrak{s} for which $\mathcal{C}(\mathfrak{s}, \mathfrak{t}_s) \neq \emptyset$. Thus, in order to prove $\mathfrak{p} = \mathfrak{t}$, it would suffice to prove, in ZFC, that $\mathcal{C}(\mathfrak{s}, \mathfrak{t}_s) = \emptyset$ for all cofinality spectrum problems \mathfrak{s} .

These reductions underline the importance of Question 1.4, whose solution is the core of the paper. The main steps are as follows, and the proofs are primarily model-theoretic. In Section 2, we introduce the right level of abstraction. In Section 3, we prove that for any cardinal $\kappa < \min\{\mathfrak{p}_s^+, \mathfrak{t}_s\}$ there is at most one κ' such that $(\kappa, \kappa') \in \mathcal{C}(\mathfrak{s}, \mathfrak{t}_s)$. In other words, the lower cofinality (or coinitality) function is well defined. In Section 4, we prove that cofinality spectrum problems have a certain amount of local saturation; here, local types are types of elements in one of the distinguished linear orders. In Section 5, we show that any cofinality spectrum problem contains a certain amount of Peano arithmetic, sufficient to carry out Gödel coding. In Section 6, we rule out symmetric cuts in $\mathcal{C}(\mathfrak{s}, \mathfrak{t}_s)$. In Section 7, a warm-up to Section 8, we show that if the lower cofinality of \aleph_0 is not \aleph_1 , then it is at least $\min\{\mathfrak{p}_s^+, \mathfrak{t}_s\}$. In the key Section 8, we rule out *all* asymmetric cuts in $\mathcal{C}(\mathfrak{s}, \mathfrak{t}_s)$. In Section 9, we give one of the paper’s main theorems: for any cofinality spectrum problem \mathfrak{s} , $\mathcal{C}(\mathfrak{s}, \mathfrak{t}_s) = \emptyset$.

Sections 10–13 develop the consequences for regular ultrapowers and Keisler’s order. Section 14 addresses \mathfrak{p} and \mathfrak{t} .

Our work here also gives a new characterization of Keisler’s good ultrafilters, Theorem 10.25 quoted below. Cofinality spectrum problems include other central examples, such as Peano arithmetic, as developed further in our paper [33].

There we give evidence that $NSOP_2$ is a dividing line, including characterizing maximality in a relative of Keisler’s order. In other continuations of the present work, we prove Keisler’s order is not so simple [32] and try to understand simple unstable theories [31].

* * * * *

1.1. List of main theorems. The main results of the paper are the following.

Main Theorem. (Theorem 9.1) *Let \mathfrak{s} be a cofinality spectrum problem. Then $\mathcal{C}(\mathfrak{s}, \mathfrak{t}_s) = \emptyset$.*

Theorem. (Theorem 11.11) *Let T be a theory with SOP_2 . Then T is maximal in Keisler’s order.*

Theorem. (Theorem 10.25) *For a regular ultrafilter \mathcal{D} on λ , the following are equivalent: (a) \mathcal{D} has λ^+ -treetops, (b) $\kappa \leq \lambda$ implies $(\kappa, \kappa) \notin \mathcal{C}(\mathcal{D})$, (c) $\mathcal{C}(\mathcal{D}) = \emptyset$, (d) \mathcal{D} is λ^+ -good.*

Theorem. (Theorem 13.1) *There is a minimum class among the nonsimple theories in Keisler’s order, which contains the theory T_{feq} of a parametrized family of independent equivalence relations.*

Theorem. (Theorem 14.1) $\mathfrak{p} = \mathfrak{t}$.

Our methods give various results about the structure of regular ultrapowers. For instance, combining our work here with the work of Malliaris [22, 24], we prove that any ultrafilter which saturates some nonlow or nonsimple theory must be flexible; see Conclusion 12.16 below.

The sections on regular ultrafilters and on cardinal invariants of the continuum may be read independently of each other, while Sections 2–9 are prerequisites for both.

In a work in preparation we intend to deal also with the following questions: Do cofinality spectrum problems require high theories (like bounded Peano arithmetic, in Section 5 below)? Last elements? Successor? Do the theorems on the cofinality spectrum [i.e., 3.1 (thus, 3.4, 3.5, 3.6, 3.8, 7.1–7.4, 8.1 and also 2.15, 3.1, 3.1, 3.7)] really need the same assumptions? What happens with so-called $\mathfrak{p}_\lambda, \mathfrak{t}_\lambda$? This work in preparation will also show that the lower cofinality is well defined in a more general setting; see 14.12 below. We also intend to deal with characterizing the maximum class in the related order \leq_* , leveraging the work of [9, 45] mentioned above and the results here.

2. COFINALITY SPECTRUM PROBLEMS

In this section we give some central definitions of the paper.

Convention 2.1 (Conventions on notation).

- (1) $\ell(\bar{y})$ is the length of a tuple of variables \bar{y} . This length exists in the ambient model of set theory. It is always a standard ordinal and is usually finite.
- (2) Definable means with parameters, unless otherwise stated.
- (3) We say that a linear order is pseudo-finite if it is discrete and every non-empty, bounded, definable subset has a first and a last element.
- (4) When T is a theory or M a model, we follow model-theoretic convention and write $\tau(T), \tau(M), \tau_T, \tau_M$, etc., to indicate the ambient vocabulary or signature.
- (5) Given an ultrapower M^I/\mathcal{D} , we fix in advance some lifting $M^I/\mathcal{D} \rightarrow M^I$. Then for any element a of the ultrapower and $t \in I$, we write $a[t]$ to denote the projection to the t th coordinate of this fixed representation.

2.1. A motivating example. Let \mathcal{D} be a regular ultrafilter on λ . Let $\mathcal{C}(\mathcal{D})$ and λ^+ -treetops be as defined in the Introduction immediately before and after Fact 1.3. We begin with a proof to illustrate the effect of treetops on $\mathcal{C}(\mathcal{D})$, which will serve to motivate the definition of cofinality spectrum problems. Namely, with Problems 3 and 4 from the Introduction in mind, we rule out symmetric cuts.

Lemma 2.2. *Suppose \mathcal{D} is a regular ultrafilter on λ with λ^+ -treetops, and $\kappa < \lambda^+$ is regular. Then $\mathcal{C}(\mathcal{D})$ has no (κ, κ) -cuts, i.e., $(\kappa, \kappa) \notin \mathcal{C}(\mathcal{D})$.*

Proof. Let $M = (\mathbb{N}, <)$ and $M_1 = M^\lambda/\mathcal{D}$. Note that by the first fundamental theorem of ultraproducts, Los’ theorem, any nonempty bounded definable subset of the domain of M_1 has a first and a last, i.e., least and greatest, element.

Assume for a contradiction that in M_1 there is $(\bar{a}, \bar{b}) = (\langle a_\alpha : \alpha < \kappa \rangle, \langle b_\alpha : \alpha < \kappa \rangle)$ such that for all $\beta < \alpha < \kappa$, $M_1 \models a_\beta < a_\alpha < b_\alpha < b_\beta$ but there is no $c \in M_1$ such that $a_\alpha < c < b_\alpha$ for all $\alpha < \kappa$.

Let (\mathcal{T}, \leq) be the tree whose elements are finite sequences of pairs of natural numbers, partially ordered by initial segment. Expand M to a model M^+ in which the tree (\mathcal{T}, \leq) is definable, for instance, by letting $M^+ = (\mathcal{H}(\omega_1), \epsilon)$ and identifying $\mathbb{N} = \text{dom}(M)$ with ω in M^+ . In this expansion, what is important is that

- (a) \mathbb{N} is a definable set, \mathcal{T} is a definable set, and \leq is a definable relation.
- (b) Elements of \mathcal{T} are functions from an initial segment of \mathbb{N} into $\mathbb{N} \times \mathbb{N}$.

- (c) The following are definable, uniformly, for each $a \in \mathcal{T}$:
- (i) the length function $\text{lg}(a)$,
 - (ii) the function giving $\max(\text{dom}(a))$, i.e. $\text{lg}(a) - 1$,
 - (iii) for each $n \leq \max(\text{dom}(a))$, the evaluation function $a(n)$,
 - (iv) for each $n \leq \max(\text{dom}(a))$, the projection functions $a(n, 0)$ and $a(n, 1)$, where these denote the two coordinates of $a(n)$.

The second fundamental theorem of ultraproducts, Theorem A below, states that ultraproducts commute with reducts. Having chosen M^+ , then, let M_1^+ be the induced expansion of M_1 . Then, e.g., elements of $\mathcal{T}^{M_1^+}$ are functions from an initial segment of the nonstandard integers into pairs of nonstandard integers, and (a)–(c) remain definable in M_1^+ . The length function lg will then often take a nonstandard length. (Throughout the paper, we will use lg to denote a possibly nonstandard length function definable in the structure under consideration. In contrast, we use $\ell(\bar{y})$ following Convention 2.1.)

Let $\varphi(x)$ be a formula expressing $x \in \mathcal{T}$ and $n < m \leq \max(\text{dom}(x))$ implies $x(n, 0) < x(m, 0) < x(m, 1) < x(n, 1)$. In M_1^+ , φ defines an infinite subtree of $\mathcal{T}^{M_1^+}$, which we denote in the rest of the proof⁴ as \mathcal{T}_* . Note that if $M_1^+ \models \varphi(c)$ and $n \leq \max(\text{dom}(c))$, then $M_1^+ \models \varphi(c \upharpoonright_n)$.

By induction on $\alpha < \kappa$ we now choose elements c_α of \mathcal{T}_* and n_α of $\mathbb{N}^{M_1^+}$ such that

- (1) for all $\beta < \alpha < \kappa$, $M_1^+ \models c_\beta \trianglelefteq c_\alpha$,
- (2) for all $\alpha < \kappa$, $n_\alpha = \max(\text{dom}(c_\alpha))$,
- (3) for all $\alpha < \kappa$, $c_\alpha(n_\alpha, 0) = a_\alpha$, and $c_\alpha(n_\alpha, 1) = b_\alpha$.

For the base case, let $c_0 = \langle (a_0, b_0) \rangle$.

When $\alpha = \beta + 1$, having defined c_β , let $c_\alpha = c_\beta \wedge \langle (a_\alpha, b_\alpha) \rangle$. Then $c_\alpha \in \mathcal{T}_*$ by definition of the sequences \bar{a}, \bar{b} . Let $n_\alpha = n_\beta + 1$.

When α is a limit ordinal, by the hypothesis of treetops there is $c_* \in \mathcal{T}_*$ such that $\beta < \alpha$ implies $c_\beta \trianglelefteq c_*$. Let $n_* = \max(\text{dom}(c_*))$. By definition of \mathcal{T}_* and of \trianglelefteq , it will be the case that $\beta < \alpha$ implies $c_\beta(n_\beta, 0) = c_*(n_\beta, 0) < c_*(n_*, 0) < c_*(n_*, 1) < c_*(n_\beta, 1) = c_\beta(n_\beta, 1)$, but it may also be the case that $a_\alpha < c_*(n_*, 0) < c_*(n_*, 1) < b_\alpha$. However, the set

$$\{n \leq n_* : c_*(n, 0) < a_\alpha \wedge b_\alpha < c_*(n, 1)\}$$

is a nonempty bounded definable subset of nonstandard integers, so it has a maximal element; call it m_* . Note that necessarily $c_\beta \trianglelefteq c_* \upharpoonright_{m_*}$ for each $\beta < \alpha$. Now $c_\alpha = c_* \upharpoonright_{m_*} \wedge \langle (a_\alpha, b_\alpha) \rangle$ realizes φ and, letting also $n_\alpha = m_*$, this completes the inductive construction.

Now the path $\bar{c} = \langle c_\alpha : \alpha < \kappa \rangle$ in \mathcal{T}_* corresponds, by construction, to a cofinal sequence in our original cut. By hypothesis of treetops, there is an upper bound $c_* \in \mathcal{T}_*$ so $\alpha < \kappa$ implies $c_\alpha \trianglelefteq c_*$. Let $n_* = \max(\text{dom}(c_*))$. Then for each $\alpha < \kappa$, by definition of φ ,

$$a_\alpha = c_\alpha(n_\alpha, 0) = c_*(n_\alpha, 0) < c_*(n_*, 0) < c_*(n_*, 1) < c_*(n_\alpha, 1) = c_\alpha(n_\alpha, 1) = b_\alpha,$$

realizing the cut (\bar{a}, \bar{b}) . This contradiction shows (\bar{a}, \bar{b}) cannot exist. □

⁴Why not simply begin with the definition for \mathcal{T}_* ? The present approach generalizes more easily to later sections, where we will distinguish certain families of basic trees and then work, as necessary, within their definable subtrees.

Lemma 2.2 hints at the power of the treetops hypothesis with respect to saturation of the underlying order. It suggests that *in regular ultrapowers*, one might productively analyze the existence or nonexistence of other families of cuts in $\mathcal{C}(\mathcal{D})$ by considering progressively more sophisticated trees. However, on closer inspection, the use of regular ultrapowers does not appear essential. Rather, the key features of that setting were the following:

- (1) Los' theorem, used to show $M \preceq M_1$, and that the set \mathbb{N}^{M_1} behaved in a pseudo-finite way: it remained a definable, discrete linear order in which any bounded, nonempty definable subset had a first and a last element.
- (2) Ultrapowers commute with reducts, invoked to study a given order in M, M_1 by uniformly (simultaneously) expanding both models to have available a suitable tree of sequences from the order to itself. Note that even after the expansion, any bounded, nonempty definable subset of $\mathbb{N}^{M_1^+}$ has a first and a last element.

The setting of cofinality spectrum problems, a fundamental definition of this paper, can be seen as a suitable abstraction of this setting (i.e., studying cuts in linear orders under the hypothesis of the existence of paths through related families of trees) which retains the key features just described. Although Definitions 2.3 and 2.5 have wider application, regular ultrapowers will be an important context for the second half of the paper.

2.2. Main definitions. We now give several main definitions of the paper.

First, given a family of formulas defining linear orders, we specify which trees we would like to have available.⁵ The conditions are of two kinds: on the one hand, we require that Δ have a certain form and satisfy some closure conditions, and on the other hand, we require that certain sets derived from Δ are definable.

Definition 2.3. (ESTT, enough set theory for trees) *Let M_1 be a model and Δ a nonempty set of formulas in the language of M_1 . We say that (M_1, Δ) has enough set theory for trees when the following conditions are true.*

- (1) Δ consists of first-order formulas $\varphi(\bar{x}, \bar{y}; \bar{z})$, with $\ell(\bar{x}) = \ell(\bar{y})$.
- (2) For each $\varphi \in \Delta$ and each parameter $\bar{c} \in {}^{\ell(\bar{z})}M_1$, $\varphi(\bar{x}, \bar{y}, \bar{c})$ defines a discrete linear order on $\{\bar{a} : M_1 \models \varphi(\bar{a}, \bar{a}, \bar{c})\}$ with a first and a last element.
- (3) The family of all linear orders defined in this way will be denoted $\text{Or}(\Delta, M_1)$. Specifically, each $\mathbf{a} \in \text{Or}(\Delta, M_1)$ is a tuple $(X_{\mathbf{a}}, \leq_{\mathbf{a}}, \varphi_{\mathbf{a}}, \bar{c}_{\mathbf{a}}, d_{\mathbf{a}})$, where
 - (a) $X_{\mathbf{a}}$ denotes the underlying set $\{\bar{a} : M_1 \models \varphi_{\mathbf{a}}(\bar{a}, \bar{a}, \bar{c}_{\mathbf{a}})\}$,
 - (b) $\bar{x} \leq_{\mathbf{a}} \bar{y}$ abbreviates the formula $\varphi_{\mathbf{a}}(\bar{x}, \bar{y}, \bar{c}_{\mathbf{a}})$,
 - (c) $d_{\mathbf{a}} \in X_{\mathbf{a}}$ is a bound for the length of elements in the associated tree; it is often, but not always, $\max X_{\mathbf{a}}$. If $d_{\mathbf{a}}$ is finite, we call \mathbf{a} trivial.
- (4) For each $\mathbf{a} \in \text{Or}(\mathbf{a})$, $(X_{\mathbf{a}}, \leq_{\mathbf{a}})$ is pseudo-finite, meaning that any bounded, nonempty, M_1 -definable subset has a $\leq_{\mathbf{a}}$ -greatest and a $\leq_{\mathbf{a}}$ -least element.
- (5) For each pair \mathbf{a} and \mathbf{b} in $\text{Or}(\Delta, M_1)$, there is $\mathbf{c} \in \text{Or}(\Delta, M_1)$ such that
 - (a) there exists an M_1 -definable bijection $\text{Pr} : X_{\mathbf{a}} \times X_{\mathbf{b}} \rightarrow X_{\mathbf{c}}$ such that the coordinate projections are M_1 -definable.
 - (b) if $d_{\mathbf{a}}$ is not finite in $X_{\mathbf{a}}$ and $d_{\mathbf{b}}$ is not finite in $X_{\mathbf{b}}$, then also $d_{\mathbf{c}}$ is not finite in $X_{\mathbf{c}}$.

⁵Although in Section 2.1 we used that ultrapowers commute with reducts, giving us access to any reasonable tree, it is not necessary to have all or even many trees available. Here we ask for uniform definitions for the necessary families of trees.

(6) For some nontrivial $\mathbf{a} \in \text{Or}(\Delta, M_1)$, there is $\mathbf{c} \in \text{Or}(\Delta, M_1)$ such that $X_{\mathbf{c}} = \text{Pr}(X_{\mathbf{a}} \times X_{\mathbf{a}})$ and the ordering $\leq_{\mathbf{c}}$ satisfies

$$M_1 \models (\forall x \in X_{\mathbf{a}})(\exists y \in X_{\mathbf{c}})(\forall x_1, x_2 \in X_{\mathbf{a}})(\max\{x_1, x_2\} \leq_{\mathbf{a}} x \iff \text{Pr}(x_1, x_2) \leq_{\mathbf{c}} y).$$

(7) To the family of distinguished orders, we associate a family of trees, as follows. For each formula $\varphi(\bar{x}, \bar{y}; \bar{z})$ in Δ there are formulas ψ_0, ψ_1, ψ_2 of the language of M_1 such that for any $\mathbf{a} \in \text{Or}(\mathbf{s})$ with $\varphi_{\mathbf{a}} = \varphi$

(a) $\psi_0(\bar{x}, \bar{c}_{\mathbf{a}})$ defines a set, denoted $\mathcal{T}_{\mathbf{a}}$, of functions from $X_{\mathbf{a}}$ to $X_{\mathbf{a}}$.

(b) $\psi_1(\bar{x}, \bar{y}, \bar{c})$ defines a function $\text{lg}_{\mathbf{a}} : \mathcal{T}_{\mathbf{a}} \rightarrow X_{\mathbf{a}}$ satisfying

(i) for all $b \in \mathcal{T}_{\mathbf{a}}$, $\text{lg}_{\mathbf{a}}(b) \leq_{\mathbf{a}} d_{\mathbf{a}}$.

(ii) for all $b \in \mathcal{T}_{\mathbf{a}}$, $\text{lg}_{\mathbf{a}}(b) = \max(\text{dom}(b))$.

(c) $\psi_2(\bar{x}, \bar{y}, \bar{c})$ defines a function from $\{(b, a) : b \in \mathcal{T}_{\mathbf{a}}, a \in X_{\mathbf{a}}, a <_{\mathbf{a}} \text{lg}_{\mathbf{a}}(b)\}$ into $X_{\mathbf{a}}$ whose value is called $\text{val}_{\mathbf{a}}(b, c)$, and abbreviated $b(a)$.

(i) if $c \in \mathcal{T}_{\mathbf{a}}$, $\text{lg}_{\mathbf{a}}(c) < d_{\mathbf{a}}$, and $a \in X_{\mathbf{a}}$, then $c \frown \langle a \rangle$ exists; i.e., there is $c' \in \mathcal{T}_{\mathbf{a}}$ such that $\text{lg}_{\mathbf{a}}(c') = \text{lg}_{\mathbf{a}}(c) + 1$, $c'(\text{lg}_{\mathbf{a}}(c)) = a$, and

$$(\forall a <_{\mathbf{a}} \text{lg}_{\mathbf{a}}(c))(c(a) = c'(a)).$$

(ii) $\psi_0(\bar{x}, \bar{c})$ implies that if $b_1 \neq b_2 \in \mathcal{T}_{\mathbf{a}}$, $\text{lg}_{\mathbf{a}}(b_1) = \text{lg}_{\mathbf{a}}(b_2)$, then for some $n <_{\mathbf{a}} \text{lg}_{\mathbf{a}}(b_1)$, $b_1(n) \neq b_2(n)$.

(d) $\psi_3(\bar{x}, \bar{y}, \bar{c})$ defines the partial order $\leq_{\mathbf{a}}$ on $\mathcal{T}_{\mathbf{a}}$ given by initial segment, that is, such that $b_1 \leq_{\mathbf{a}} b_2$ implies

(i) for all $b, c \in \mathcal{T}_{\mathbf{a}}$, $b \leq c$ implies $\text{lg}_{\mathbf{a}}(b) \leq_{\mathbf{a}} \text{lg}_{\mathbf{a}}(c)$.

(ii) $\text{lg}_{\mathbf{a}}(b_1) \leq_{\mathbf{a}} \text{lg}_{\mathbf{a}}(b_2)$.

(iii) $(\forall a <_{\mathbf{a}} \text{lg}_{\mathbf{a}}(b_1))(b_2(a) = b_1(a))$.

The family of all $\mathcal{T}_{\mathbf{a}}$ defined this way will be denoted $\text{Tr}(\Delta, M_1)$. We refer to elements of this family as trees.

In 2.3(2), we ask for both first and last elements. The first element is important: we repeatedly use that these orders are pseudo-finite. The last element is technical; we could alter this but would need to make other changes to ensure that the derived trees are not too large. Also, the condition in (6) should be understood as saying that while we need Cartesian products to exist, it is largely unimportant what exactly the order on these products is (as long as it is discrete, pseudo-finite, and so forth). It is sufficient that one such order behaves well, like the usual Gödel pairing function. We could have alternately asked that (6) hold for all products.

Convention 2.4. Many proofs will consider elements of a given tree $\mathcal{T}_{\mathbf{a}}$ or a given linear order $X_{\mathbf{a}}$. Although these orders or trees may be defined by some $\varphi(\bar{x}, \dots)$ with $\ell(\bar{x}) > 1$, we will often omit overlines on their elements; this should not cause confusion as the provenance of such elements is clear.

Definition 2.5. Say that $(M, M_1, M^+, M_1^+, T, \Delta)$ is a cofinality spectrum problem when

(1) $M \preceq M_1$.

(2) $T \supseteq \text{Th}(M)$ is a theory in a possibly larger vocabulary.

(3) Δ is a set of formulas in the language of M ; i.e., we are interested in studying the orders of $\mathcal{L}(M) = \mathcal{L}(M_1)$ in the presence of the additional structure of $\mathcal{L}(M^+) = \mathcal{L}(M_1^+)$.

(4) M^+, M_1^+ expand M, M_1 , respectively, so that $M^+ \preceq M_1^+ \models T$ and (M_1^+, Δ) has enough set theory for trees.

- (5) We may refer to the components of \mathbf{s} as $M^{\mathbf{s}}$, $\Delta^{\mathbf{s}}$, etc., for definiteness. When $T = Th(M)$, $M = M^+$, $M_1 = M_1^+$, or Δ is the set of all formulas $\varphi(x, y, \bar{z})$ in the language of T which satisfy 2.3(2)–(4), these may be omitted.

In what follows, we work almost exclusively with M^+, M_1^+ but our results are for M, M_1 . We will make the following conventions in the remainder of the paper.

Convention 2.6 (Conventions on cofinality spectrum problems).

- (1) Extending Definition 2.3(3)(c) we say that \mathbf{s} is trivial if every $\mathbf{a} \in \text{Or}(\mathbf{s})$ is trivial (note that this need not mean all the orders are finite, just that the choice of tree-height bound $d_{\mathbf{a}}$ is in each instance finite). We will assume that all cofinality spectrums we work with are nontrivial, though sometimes this is repeated for emphasis.
- (2) Likewise, by “ $\mathbf{a} \in \text{Or}(\mathbf{s})$ ” we will mean “ $\mathbf{a} \in \text{Or}(\mathbf{s})$ and \mathbf{a} is nontrivial,” unless otherwise indicated.
- (3) By definable we shall mean definable in the larger expanded model, i.e., in M_1^+ , possibly with parameters, unless otherwise stated.
- (4) We will present Cartesian products without explicitly mentioning the pairing functions, writing, e.g., “let $\mathbf{a} \in \text{Or}(\mathbf{s})$ and let \mathbf{b} be such that $X_{\mathbf{b}} = X_{\mathbf{a}} \times X_{\mathbf{a}}$.”
- (5) If $\mathbf{c}, \mathbf{a}_0, \dots, \mathbf{a}_k \in \text{Or}(\mathbf{s})$ are such that $X_{\mathbf{c}} = X_{\mathbf{a}_0} \times \dots \times X_{\mathbf{a}_k}$, and $c \in \mathcal{T}_{\mathbf{c}}$ and $n \in X_{\mathbf{c}}$, $n < \text{lg}(c)$ (thus, $c(n)$ is well defined), then we will write $c(n, i)$ to mean the i th coordinate of $c(n)$.

Definition 2.7. When $\mathbf{s}_1, \mathbf{s}_2$ are cofinality spectrum problems, write $\mathbf{s}_1 \leq \mathbf{s}_2$ to mean

- $M^{\mathbf{s}_1} = M^{\mathbf{s}_2}$, $M_1^{\mathbf{s}_1} = M_1^{\mathbf{s}_2}$.
- $\tau(M^{+, \mathbf{s}_1}) \subseteq \tau(M^{+, \mathbf{s}_2})$, i.e., the vocabulary may be larger, and likewise $T^{+, \mathbf{s}_1} \subseteq T^{+, \mathbf{s}_2}$.
- $(M^{+, \mathbf{s}_2} \upharpoonright_{\tau(M^{+, \mathbf{s}_1})}) \cong M^{+, \mathbf{s}_1}$.
- $(M_1^{+, \mathbf{s}_2} \upharpoonright_{\tau(M^{+, \mathbf{s}_1})}) \cong M_1^{+, \mathbf{s}_1}$.
- $\Delta^{\mathbf{s}_1} \subseteq \Delta^{\mathbf{s}_2}$.

We will study properties of orders and trees arising in cofinality spectrum problems, as we now describe. We follow the model-theoretic terminology, writing cuts instead of gaps. For clarity, see Definition 2.8.

Definition 2.8. (Cuts, pre-cuts, and representations of cuts) Let \mathbf{s} be a cofinality spectrum problem, $\mathbf{a} \in \text{Or}(\mathbf{s})$. When κ_1, κ_2 are regular, define

- (1) A (κ_1, κ_2) -cut in $X_{\mathbf{a}}$, i.e., in $(X_{\mathbf{a}}, \leq_{\mathbf{a}})$, is given by a pair of sets (C_1, C_2) such that
 - (a) $C_1 \cap C_2 = \emptyset$,
 - (b) C_1 is downward closed, C_2 is upward closed,
 - (c) $(\forall x \in C_1)(\forall y \in C_2)(x <_{\mathbf{a}} y)$,
 - (d) $C_1 \cup C_2 = X_{\mathbf{a}}$.

- (2) If (C_1, C_2) satisfies conditions (a), (b), (c) for being a cut, it is called a pre-cut, meaning that possibly $(\exists c)(C_1 < c < C_2)$.
- (3) By a (κ_1, κ_2) -representation of a pre-cut (C_1, C_2) in $X_{\mathbf{a}}$ we mean a pair of sequences $(\langle a_i : i < \kappa_1 \rangle, \langle b_j : j < \kappa_2 \rangle)$ of elements of $X_{\mathbf{a}}$ such that
 - (a) κ_1, κ_2 are regular cardinals,
 - (b) $\langle a_i : i < \kappa_1 \rangle$ is strictly $<_{\mathbf{a}}$ -increasing and cofinal in C_1 ,
 - (c) $\langle b_j : j < \kappa_2 \rangle$ is strictly $<_{\mathbf{a}}$ -decreasing and coinital in C_2 .

When the (pre-)cut (C_1, C_2) has a (κ_1, κ_2) -representation, we say that it is a (κ_1, κ_2) -(pre-)cut. When there is no danger of confusion, we may informally identify cuts or pre-cuts with one of their representations.

Definition 2.9. For a cofinality spectrum problem \mathbf{s} we define the following:

- (1) $\text{Or}(\mathbf{s}) = \text{Or}(\Delta^{\mathbf{s}}, M_1^{\mathbf{s}})$.
- (2) $\mathcal{C}^{\text{ct}}(\mathbf{s}) = \{(\kappa_1, \kappa_2) : \text{for some } \mathbf{a} \in \text{Or}(\mathbf{s}, M_1), (X_{\mathbf{a}}, \leq_{\mathbf{a}}) \text{ has a } (\kappa_1, \kappa_2)\text{-cut}\}$.
- (3) $\text{Tr}(\mathbf{s}) = \{\mathcal{T}_{\mathbf{a}} : \mathbf{a} \in \text{Or}(\mathbf{s})\} = \text{Tr}(\Delta^{\mathbf{s}}, M_1^{\mathbf{s}})$.
- (4) $\mathcal{C}^{\text{ttp}}(\mathbf{s}) = \{\kappa : \kappa \geq \aleph_0, \mathbf{a} \in \text{Or}(\mathbf{s}), \text{ and there is in the tree } \mathcal{T}_{\mathbf{a}} \text{ a strictly increasing sequence of cofinality } \kappa \text{ with no upper bound}\}$.
- (5) Let $\mathbf{t}_{\mathbf{s}}$ be $\min \mathcal{C}^{\text{ttp}}(\mathbf{s})$ and let $\mathbf{p}_{\mathbf{s}}$ be $\min\{\kappa : (\kappa_1, \kappa_2) \in \mathcal{C}^{\text{ct}}(\mathbf{s}) \text{ and } \kappa = \kappa_1 + \kappa_2\}$.

Our main focus in this paper will be $\mathcal{C}(\mathbf{s}, \mathbf{t}_{\mathbf{s}})$ where this means

- (6) For λ an infinite cardinal, write

$$\mathcal{C}(\mathbf{s}, \lambda) = \{(\kappa_1, \kappa_2) : \kappa_1 + \kappa_2 < \lambda, (\kappa_1, \kappa_2) \in \mathcal{C}^{\text{ct}}(\mathbf{s})\}.$$

Note that by definition of $\mathcal{C}^{\text{ct}}(\mathbf{s})$ and $\mathcal{C}^{\text{ttp}}(\mathbf{s})$, both $\mathbf{t}_{\mathbf{s}}$ and $\mathbf{p}_{\mathbf{s}}$ are regular. The following is a central definition of the paper.

Definition 2.10. (Treetops) Let \mathbf{s} be a cofinality spectrum problem and $\mathbf{t}_{\mathbf{s}}$ be given by 2.9(5). When $\lambda \leq \mathbf{t}_{\mathbf{s}}$ we say that \mathbf{s} has λ -treetops. Our main focus will be the case $\lambda = \mu^+$ for some $\mu < \mathbf{t}_{\mathbf{s}}$.

The name *treetops* reflects the definition of $\mathbf{t}_{\mathbf{s}}$: when $\kappa = \text{cf}(\kappa) < \mathbf{t}_{\mathbf{s}}$, $\mathbf{a} \in \text{Or}(\mathbf{s})$ and thus $\mathcal{T}_{\mathbf{a}} \in \text{Tr}(\mathbf{s})$, 2.9(5), any strictly increasing κ -sequence of elements of $\mathcal{T}_{\mathbf{a}}$ has an upper bound in $\mathcal{T}_{\mathbf{a}}$.

Definition 2.11. Let \mathbf{s} be a cofinality spectrum problem, $\mathbf{a} \in \text{Or}(\mathbf{s})$.

- (0) Write $0_{\mathbf{a}}$ for the $\leq_{\mathbf{a}}$ -least element of $X_{\mathbf{a}}$.
- (1) For any natural number k and any $a \in X_{\mathbf{a}}$, let $S_{\mathbf{a}}^k(a)$ denote the k th successor of a in the discrete linear order $\leq_{\mathbf{a}}$, if defined, and likewise let $S_{\mathbf{a}}^{-k}(a)$ denote the k th predecessor of a , if defined. We will generally write $S^k(a)$, $S^{-k}(a)$ when \mathbf{a} is clear from context.
- (2) Say that $c \in \mathcal{T}_{\mathbf{a}}$ is below the ceiling if $S^k(\text{lg}(c)) <_{\mathbf{a}} d_{\mathbf{a}}$ for all $k < \omega$, i.e., if these successors exist and the statements are true.

We retain the notation $S^k(\dots)$ rather than abbreviating to $+k$ in Definition 2.11 because addition will be introduced formally later on.

To conclude this section, let us emphasize that Definition 2.3 is a choice of abstraction, suitable for our purposes but also relatively strong. In future work, it will be useful to consider various substitutions and weakenings. Toward this, we include some alternatives in 2.12 and 2.13 below.

Discussion 2.12. (Alternate version A: one-to-one) *In Definition 2.3 one could drop the bound $d_{\mathbf{a}}$ and retain an implicit bound on the length of sequences in $\mathcal{T}_{\mathbf{a}}$ by requiring no repetition in the range. Formally, one would modify 2.3 as follows: drop $d_{\mathbf{a}}$ from the definition of $\text{Or}(\Delta, M_1)$, add condition (1) below, and replace (7)(b)(iv) by (2) below.*

- (1) $a_1 <_{\mathbf{a}} a_2 <_{\mathbf{a}} \text{lg}_{\mathbf{a}}(b)$ implies $(b(a_1)) \neq (b(a_2))$.
- (2) if $b \in \mathcal{T}_{\mathbf{a}}$, $a \in X_{\mathbf{a}} \setminus \{\text{val}_{\mathbf{a}}(b(a')) : a' < \text{lg}_{\mathbf{a}}(b)\}$, and there is still room to concatenate, i.e.,

$$|X_{\mathbf{a}} \setminus (\{a\} \cup \{\text{val}_{\mathbf{a}}(b(a')) : a' < \text{lg}_{\mathbf{a}}(b)\})| \geq 2,$$

then $b \hat{\ } \langle a \rangle$ exists; i.e., there is b' such that $b \preceq_{\mathbf{a}} b'$, $\text{lg}_{\mathbf{a}}(b') = \text{lg}_{\mathbf{a}}(b) + 1$, and $\text{val}(b'(\text{lg}(b))) = a$.

Discussion 2.13. (Alternate version B: Allowing other orders) *In Definition 2.3 one could rename the current set $\text{Or}(\mathbf{s})$ as $\text{Psf-ord}(\mathbf{s})$, the set of pseudo-finite orders, and make the following changes. First, allow other kinds of linear orders in $\text{Or}(\mathbf{s})$ (e.g., dense linear orders, or any definable linear order). Second, change the requirements on trees so that any tree is a set of sequences from some $X_{\mathbf{a}}$ into some $X_{\mathbf{b}}$ where $\mathbf{a} \in \text{Psf-ord}(\mathbf{s})$ and $\mathbf{b} \in \text{Or}(\mathbf{s})$.*

2.3. Key examples. A first motivating example is that of Section 2.1. Consider $M \models (\mathbb{N}, <)$. In this case, there are a set of \mathcal{L} -formulas $\Delta \supseteq \{x \leq y\}$, an expanded language \mathcal{L}^+ , and an \mathcal{L}^+ -theory $T \supseteq \text{Th}(M)$ such that $(M, M_1, M^+, M_1^+, T, \Delta)$ is a cofinality spectrum problem. For instance, we may take $T = \text{Th}(\mathcal{H}(\omega_1), \epsilon)$ and identify \mathbb{N} with ω . The reader interested primarily in our model-theoretic conclusions for regular ultrapowers may wish to look first at Section 11, where the main definitions are specialized to the case of regular ultrapowers.

A second motivating example involves pairs of models which admit expansions. Suppose that $M \preceq M_1$, and so that this example is nontrivial, that M includes an infinite definable linear order. M, M_1 can be, respectively, expanded to models M^+, M_1^+ in a larger signature $\tau_{M^+} \supseteq \tau_M$ such that

- (1) $M^+ \preceq M_1^+$,
- (2) $M = (M^+ \upharpoonright P^{M^+}) \upharpoonright \tau_M$,
- (3) $M_1 = (M_1^+ \upharpoonright P^{M_1^+}) \upharpoonright \tau_M$,
- (4) $M^+ = (\mathcal{H}(\chi), \in, P^{M^+}, R^{M^+})_{R \in \tau(M)}$ for some χ , where $M \in \mathcal{H}(\chi)$.

As ultrapowers commute with reducts, this example generalizes the first one.

A third motivating example will be defined in Section 14, but is fundamental to one of our applications so we mention it here. Let $M = (\mathcal{H}(\aleph_1), \in)$, let $\mathbf{Q} = ([\mathbb{N}]^{\aleph_0}, \supseteq^*)$, let \mathbf{G} be a generic subset of \mathbf{Q} , and let $N = M^\omega / \mathbf{G}$ be the generic ultrapower. Then $M \preceq N$. It will be shown in Section 14 that there is a natural cofinality spectrum problem \mathbf{s} with $M = M_{\mathbf{s}}^+$, $N = M_{1, \mathbf{s}}^+$.

A fourth motivating example involves arithmetic with bounded induction. Let M be a model of Peano arithmetic with bounded induction, meaning that we have induction only for formulas all of whose quantifiers are bounded. Choose M_1 so that $M \preceq M_1$. Let $T = \text{Th}(M)$ and let Δ be a set which includes $\{(\exists w)(x + w = y)\}$ and its finite products. So the elements of $\text{Or}(\Delta, M_1)$ are (finite Cartesian products of) initial segments of the domain of $M_1^{\mathbf{s}}$, and the fact that this may be regarded as a cofinality spectrum problem follows from the fact that Gödel coding can be carried out in this context, meaning that we may speak about sequences—which

are not too long—of elements of $X_{\mathbf{a}}$ and thus about trees. The naturalness of this example appears in Section 5 below.

2.4. Basic tools. We now develop some consequences of Definition 2.5. Recall that definable means with parameters in M_1^+ , and that *below the ceiling* was defined in Definition 2.11 above.

Claim 2.14. (Treetops for definable subtrees) *Let \mathbf{s} be a cofinality spectrum problem, $M_1^+ = M_1^{+,s}$. Let $\mathbf{a} \in \text{Or}(\mathbf{s})$, so $\mathcal{T}_{\mathbf{a}} \in \text{Tr}(\mathbf{s})$. Let φ be a formula, possibly with parameters, in M_1^+ , and let $(\mathcal{T}, \trianglelefteq_{\mathbf{a}})$ be the subtree of $(\mathcal{T}_{\mathbf{a}}, \trianglelefteq_{\mathbf{a}})$ defined by φ in M_1^+ . Let $\langle c_\alpha : \alpha < \kappa \rangle$ be a $\trianglelefteq_{\mathbf{a}}$ -increasing sequence of elements of \mathcal{T} , $\kappa = \text{cf}(\kappa) < \mathfrak{t}_{\mathbf{s}}$. Then there is $c_* \in \mathcal{T}$ such that for all $\alpha < \kappa$, $c_\alpha \trianglelefteq_{\mathbf{a}} c_*$.*

Proof. By definition of $\mathfrak{t}_{\mathbf{s}}$, there is an element $c \in \mathcal{T}_{\mathbf{a}}$, not necessarily in \mathcal{T} , such that for all $\alpha < \kappa$, $c_\alpha \trianglelefteq_{\mathbf{a}} c$. The set $\{\text{lg}(c') : c' \trianglelefteq_{\mathbf{a}} c \text{ and } c' \in \mathcal{T}\}$ is a nonempty definable subset of $X_{\mathbf{a}}$, and hence contains a last member a_* . Let c_* be such that $c_* \trianglelefteq c$ and $\text{lg}(c_*) = a_*$, i.e., $c_* = c \upharpoonright_{a_*}$. \square

Lemma 2.15. (Treetops below the ceiling) *Let \mathbf{s} be a cofinality spectrum problem, $\mathbf{a} \in \text{Or}(\mathbf{s})$, $\kappa < \min\{\mathfrak{p}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}}\}$. Let $\mathcal{T} \subseteq \mathcal{T}_{\mathbf{a}}$ be a definable subtree and $\bar{c} = \langle c_\alpha : \alpha < \kappa \rangle$ a strictly $\trianglelefteq_{\mathbf{a}}$ -increasing sequence of elements of \mathcal{T} . Then there exists $c_{**} \in \mathcal{T}$ such that $\alpha < \kappa$ implies $c_\alpha \trianglelefteq c_{**}$ and c_{**} is below the ceiling.*

Proof. Let $c_* \in \mathcal{T}$ be such that $\alpha < \beta$ implies $c_\alpha \trianglelefteq c_*$, as given by Claim 2.14. As we assumed the sequence \bar{c} is strictly increasing, for each $\alpha < \kappa$ the element c_α must be below the ceiling. If c_* is also below the ceiling, we finish. Otherwise,

$$(\{\text{lg}(c_\beta) : \beta < \alpha\}, \{S^{-k}(\text{lg}(c_*)) : k < \omega\})$$

describes a pre-cut in $X_{\mathbf{a}}$. It cannot be a cut, as then $(\aleph_0, \kappa) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$, contradicting the definition of $\mathfrak{p}_{\mathbf{s}}$. Choose $a \in X_{\mathbf{a}}$ realizing this pre-cut, and let $c_{**} = c_* \upharpoonright_a$. \square

Normally, $\mathfrak{p}_{\mathbf{s}} \leq \mathfrak{t}_{\mathbf{s}}$ though we shall not need this for the main theorems of the paper, instead keeping track of each separately.

3. THE FUNCTION lcf IS WELL DEFINED STRICTLY BELOW $\min\{\mathfrak{p}_{\mathbf{s}}^+, \mathfrak{t}_{\mathbf{s}}\}$

In the next six sections, we give a series of constructions which show how to translate certain conditions on realization of pre-cuts in linear order into conditions on existence of paths through trees, leading to the central theorem of Section 9.

We generally write, e.g., M_1^+ rather than $M_1^{+,s}$, but this should not cause confusion; the components of a cofinality spectrum problem are always understood to depend on a background \mathbf{s} fixed at the beginning of a proof.

Lemma 3.1. *Let \mathbf{s} be a cofinality spectrum problem. If $\mathbf{a} \in \text{Or}(\mathbf{s})$ is nontrivial, then for each infinite regular $\kappa \leq \mathfrak{p}_{\mathbf{s}}$,*

- (1) *there is a strictly decreasing κ -indexed sequence $\bar{a} = \langle a_\alpha : \alpha < \kappa \rangle$ of elements of $X_{\mathbf{a}}$ such that*

$$(\{S^k(0_{\mathbf{a}}) : k < \omega\}, \{a_\alpha : \alpha < \kappa\})$$

represents a pre-cut (and possibly a cut) in $X_{\mathbf{a}}$.

(2) there is a strictly increasing κ -indexed sequence $\bar{a} = \langle a_\alpha : \alpha < \kappa \rangle$ of elements of $X_{\mathbf{a}}$ such that

$$(\{a_\alpha : \alpha < \kappa\}, \{S^{-k}(d_{\mathbf{a}}) : k < \omega\})$$

represents a pre-cut (and possibly a cut) in $X_{\mathbf{a}}$.

- (3) thus, there is at least one infinite regular θ such that $(\kappa, \theta) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$, witnessed by a (κ, θ) -cut in $X_{\mathbf{a}}$.
- (4) thus, there is at least one infinite regular θ' such that $(\theta', \kappa) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$, witnessed by a (θ', κ) -cut in $X_{\mathbf{a}}$.

Proof. (1) By induction on $\alpha < \kappa$ we choose elements $a_\alpha \in X_{\mathbf{a}}$ such that

- (a) for each $\alpha < \kappa$ and each $k < \omega$, $S^k(0_{\mathbf{a}}) <_{\mathbf{a}} a_\alpha$,
- (b) $\beta < \alpha$ implies $a_\alpha <_{\mathbf{a}} a_\beta$.

For $\alpha = 0$, let a_0 be the last element of $X_{\mathbf{a}}$. As \mathbf{a} is nontrivial, condition (a) is satisfied. For $\alpha = \beta + 1$, let $a_\alpha = S^{-1}(a_\beta)$, recalling that any nonempty definable subset of $X_{\mathbf{a}}$ has a greatest element so the predecessor of any element not equal to $0_{\mathbf{a}}$ is well defined. As (a) holds for β by inductive hypothesis, it will remain true for $\beta + 1$. For limit α , by inductive hypothesis,

$$(\{S^k(0_{\mathbf{a}}) : k < \omega\}, \{a_\beta : \beta < \alpha\})$$

is a pre-cut. However, it cannot be a cut, as then we would have $(\aleph_0, \text{cf}(\alpha)) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$ with $|\alpha| + \aleph_0 < |\kappa| \leq \mathfrak{p}_{\mathbf{s}}$, contradicting the definition of $\mathfrak{p}_{\mathbf{s}}$. Let a_α be any element realizing this pre-cut. This completes the construction of the sequence and thus the proof.

(2) The argument is parallel to (1), going up instead of down.

(3) Let \bar{a} be a κ -indexed strictly increasing $\leq_{\mathbf{a}}$ -monotonic sequence of elements of $X_{\mathbf{a}}$ given by (2). By construction, $B = \{b \in X_{\mathbf{a}} : \alpha < \kappa \text{ implies } a_\alpha <_{\mathbf{a}} b\} \neq \emptyset$. Let θ be the cofinality of B considered with the reverse order. It cannot be the case that for some $b \in B$,

$$(\{a_\alpha : \alpha < \kappa\}, \{b\})$$

represents a cut, since \bar{a} is strictly increasing, thus $\alpha < \kappa$ implies $a_\alpha <_{\mathbf{a}} S^{-1}(b) <_{\mathbf{a}} b$. So θ is an infinite (regular) cardinal, and $(\kappa, \theta) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$.

(4) The argument is parallel using (1). □

Of course, in Lemma 3.1, the cardinals θ, θ' may be quite large. In what follows, we will be interested in whether they must always be at least the size of $\mathfrak{t}_{\mathbf{s}}$.

Theorem 3.1. (Uniqueness) *Let \mathbf{s} be a cofinality spectrum problem. Then for each regular $\kappa \leq \mathfrak{p}_{\mathbf{s}}$, $\kappa < \mathfrak{t}_{\mathbf{s}}$,*

- (1) there is one and only one λ such that $(\kappa, \lambda) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$.
- (2) $(\kappa, \lambda) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$ iff $(\lambda, \kappa) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$.

Proof. Fix κ satisfying the hypotheses of the theorem. We will prove that whenever we are given

- $\mathbf{a}, \mathbf{b} \in \text{Or}(\mathbf{s})$,
- $(\langle a_\alpha^1 : \alpha < \kappa \rangle, \langle b_\epsilon^1 : \epsilon < \theta_1 \rangle)$ representing a (κ, θ_1) -cut in $(X_{\mathbf{a}}, <_{\mathbf{a}})$,
- $(\langle b_\epsilon^2 : \epsilon < \theta_2 \rangle, \langle a_\alpha^2 : \alpha < \kappa \rangle)$ representing a (θ_2, κ) -cut in $(X_{\mathbf{b}}, <_{\mathbf{b}})$,

then $\theta_1 = \theta_2$. The proof will proceed essentially by threading together the κ -sides of the cuts.

To see why this will suffice for the theorem, note first that Lemma 3.1 guarantees the existence of *some* θ_1, θ_2 such that $(\kappa, \theta_1) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$ and $(\theta_2, \kappa) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$. The statement in the previous paragraph will establish that $\theta_1 = \theta_2$. It will then follow by transitivity of equality that if $(\kappa, \theta), (\kappa, \theta') \in \mathcal{C}^{\text{ct}}(\mathbf{s})$, then $\theta = \theta'$, and likewise if $(\theta, \kappa), (\theta', \kappa) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$, then $\theta = \theta'$.

Let $\mathbf{c} \in \text{Or}(\mathbf{s})$ be such that $X_{\mathbf{c}} = X_{\mathbf{a}} \times X_{\mathbf{b}}$. Thus, for any $x \in \mathcal{T}_{\mathbf{c}}$ and any $n \leq \max(\text{dom}(x))$, $x(n)$ is a pair $(x(n, 1), x(n, 2))$, with $x(n, 1) \in X_{\mathbf{a}}$, and $x(n, 2) \in X_{\mathbf{b}}$. Consider the definable subtree $\mathcal{T}_0 \subseteq \mathcal{T}_{\mathbf{c}}$ consisting of all $x \in \mathcal{T}_{\mathbf{c}}$ such that x is strictly increasing in the first coordinate and strictly decreasing in the second; i.e.,

$$n' <_{\mathbf{c}} n <_{\mathbf{c}} \text{lg}(x) \text{ implies } (x(n', 1) <_{\mathbf{a}} x(n, 1)) \wedge (x(n, 2) <_{\mathbf{b}} x(n', 2)).$$

Keeping in mind the representations of cuts fixed at the beginning of the proof, we now choose c_{α}, n_{α} by induction on $\alpha < \kappa$, such that

- (1) $c_{\alpha} \in \mathcal{T}_0$ and $n_{\alpha} \in X_{\mathbf{c}}$,
- (2) $\beta < \alpha$ implies $M_1^+ \models c_{\beta} \trianglelefteq_{\mathbf{c}} c_{\alpha}$,
- (3) c_{α} is below the ceiling, Definition 2.11,
- (4) $n_{\alpha} = \text{lg}(c_{\alpha}) - 1$, so $\max(\text{dom}(c_{\alpha}))$ is well defined (and c_{α} is not the empty sequence),
- (5) $c_{\alpha}(n_{\alpha}, \ell) = a_{\alpha}^{\ell}$ for $\ell = 1, 2$.

For $\alpha = 0$: let $c_0 = \langle a_0^1, a_0^2 \rangle$, and let $n_0 = 0_{\mathbf{a}}$.

For $\alpha = \beta + 1$: since c_{β} is below the ceiling, concatenation is possible. So by conditions (4) and (5) of the inductive hypothesis, we may concatenate $\langle a_{\beta+1}^1, a_{\beta+1}^2 \rangle$ onto the existing sequence, and let $n_{\alpha} = n_{\beta} + 1$.

For $\alpha < \kappa$ limit: As $\text{cf}(\alpha) < \min\{\mathfrak{p}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}}\}$, apply Lemma 2.15 to choose $c \in \mathcal{T}_0$ such that $\beta < \alpha$ implies $M_1^+ \models c_{\beta} \trianglelefteq c$, and c is below the ceiling. Then the set

$$\{n : n <_{\mathbf{c}} \text{lg}(c), M_1^+ \models (c(n, 1) <_{\mathbf{a}} a_{\alpha}^1) \wedge (a_{\alpha}^2 <_{\mathbf{b}} c(n, 2))\}$$

is definable, bounded, and nonempty in $X_{\mathbf{c}}$, so it has a maximal element n_* . By (b) above and the choice of c as an upper bound, necessarily for all $\beta < \alpha$

$$M_1^+ \models (a_{\beta}^1 <_{\mathbf{a}} c(n_*, 1)) \wedge (c(n_*, 2) <_{\mathbf{b}} a_{\beta}^2).$$

As c and thus all of its initial segments are below the ceiling, concatenation is possible. Define c_{α} to be $(c|_{n_*+1}) \frown \langle a_{\alpha}^1, a_{\alpha}^2 \rangle$. Let $n_{\alpha} = n_* + 1$. By construction, c_{α} will remain strictly monotonic in all coordinates and will $\trianglelefteq_{\mathbf{c}}$ -extend the existing sequence, as desired. This completes the inductive choice of the sequence.

As $\kappa < \mathfrak{t}_{\mathbf{s}}$, by Claim 2.14 there is $c \in \mathcal{T}_0$ so that $\alpha < \kappa$ implies $c_{\alpha} \trianglelefteq_{\mathbf{a}} c$. Let $n_{**} = \text{lg}(c) - 1$, so $n_{**} \in X_{\mathbf{c}}$. For $\ell = 0, 1, 2$ and each $\epsilon < \theta_{\ell}$, recalling the b_{ϵ}^i from our original choice of cuts, define

$$n_{\epsilon, 1} = \max\{n \leq_{\mathbf{c}} n_{**} : c(n, 1) <_{\mathbf{a}} b_{\epsilon}^1\},$$

$$n_{\epsilon, 2} = \max\{n \leq_{\mathbf{c}} n_{**} : b_{\epsilon}^2 <_{\mathbf{b}} c(n, 2)\}.$$

Clearly, $\alpha < \kappa$ implies $n_{\alpha} <_{\mathbf{c}} n_{\epsilon, \ell}$ for $\ell = 1, 2$. By the choice of sequences witnessing the original cuts, for $\ell = 1, 2$ we have that

$$\langle \langle n_{\alpha} : \alpha < \kappa \rangle, \langle n_{\epsilon, \ell} : \epsilon < \theta_{\ell} \rangle \rangle$$

represents a cut in $X_{\mathbf{c}}$. (Clearly it is a pre-cut; if it were filled, say, by i_* , then the elements $c(i_*, 1)$ and $c(i_*, 2)$ would realize our two original cuts, a contradiction.) As we assumed θ_1, θ_2 were each regular, clearly $\theta_1 = \theta_2$.

This completes the proof. □

In light of Theorem 3, the following will be well defined.

Definition 3.2 (The lower cofinality $\text{lcf}(\kappa, \mathbf{s})$). *Let \mathbf{s} be a cofinality spectrum problem. For regular $\kappa \leq \mathfrak{p}_{\mathbf{s}}$, $\kappa < \mathfrak{t}_{\mathbf{s}}$, we define $\text{lcf}(\kappa, \mathbf{s})$ to be the unique θ such that $(\kappa, \theta) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$.*

Remark 3.3. *Lemma 2.15 could be proved under the hypothesis that $\text{lcf}(\aleph_0, \mathbf{s})$ is large, as could Lemma 3.1.*

The function $\kappa \mapsto \text{lcf}(\kappa, \mathbf{s})$ remains interesting even after the main theorems of this paper.

Corollary 3.4. *Let \mathbf{s} be a cofinality spectrum problem and γ a regular cardinal, $\gamma \leq \mathfrak{p}_{\mathbf{s}}$, $\gamma < \mathfrak{t}_{\mathbf{s}}$. Then the following are equivalent:*

- (1) $\text{lcf}(\gamma, \mathbf{s}) = \gamma'$,
- (2) $(\gamma, \gamma') \in \mathcal{C}^{\text{ct}}(\mathbf{s})$,
- (3) $(\gamma', \gamma) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$.

Corollary 3.5. *Let \mathbf{s} be a cofinality spectrum problem and let κ, θ be regular cardinals with $\kappa \leq \mathfrak{p}_{\mathbf{s}}$, $\kappa < \mathfrak{t}_{\mathbf{s}}$. In order to show that $(\kappa, \theta) \notin \mathcal{C}^{\text{ct}}(\mathbf{s})$, it is sufficient to show that for some nontrivial $\mathbf{a} \in \text{Or}(\mathbf{s})$, $X_{\mathbf{a}}$ has no (κ, θ) -cut.*

Corollary 3.5 shows that each of the orders is in some sense a reflection of the structure of the whole cofinality spectrum problem, at least for the cardinals we consider. As a result, when proving results about $\mathcal{C}^{\text{ct}}(\mathbf{s})$ we are free to work in the nontrivial \mathbf{a} which appears most suited to the given proof. Moreover, see Conclusion 3.6.

Conclusion 3.6. *Let \mathbf{s} be a cofinality spectrum problem. In light of 3.1 we may, without loss of generality, study $\mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}})$ by looking at*

$$\{(\kappa_1, \kappa_2) : (\kappa_1, \kappa_2) \in \mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}}), \kappa_1 \leq \kappa_2\}.$$

The proof of Theorem 3.1 has the following very useful corollary.

Corollary 3.7 (Definable monotonic maps exist). *Let \mathbf{s} be a cofinality spectrum problem, $\mathbf{a}, \mathbf{b} \in \text{Or}(\mathbf{s})$, and $\kappa = \text{cf}(\kappa) \leq \mathfrak{p}_{\mathbf{s}}$, $\kappa < \mathfrak{t}_{\mathbf{s}}$. Let $\bar{a} = \langle a_\alpha : \alpha < \kappa \rangle$ be a strictly $\langle \mathbf{a} \rangle$ -monotonic sequence of elements of $X_{\mathbf{a}}$, and let $\bar{b} = \langle b_\alpha : \alpha < \kappa \rangle$ be a strictly $\langle \mathbf{b} \rangle$ -monotonic sequence of elements of $X_{\mathbf{b}}$. Then in M_1^+ there is a definable monotonic partial injection f from $X_{\mathbf{a}}$ to $X_{\mathbf{b}}$ whose domain includes $\{a_\alpha : \alpha < \kappa\}$ and such that $f(a_\alpha) = b_\alpha$ for all $\alpha < \kappa$.*

Proof. Without loss of generality, \bar{a} is increasing and \bar{b} is decreasing (otherwise carry out this proof twice, and compose the functions).

Carry out the proof of Theorem 3.1, substituting \bar{a} here for $\langle a_\alpha^1 : \alpha < \kappa \rangle$ there, and \bar{b} here for $\langle a_\alpha^2 : \alpha < \kappa \rangle$ there. Let $\bar{c} = \langle c_\alpha : \alpha < \kappa \rangle$ be the corresponding path through the definable tree \mathcal{T}_0 constructed in the proof of Theorem 3.1, and let $c \in \mathcal{T}_0$ be an upper bound for this path in the tree, as there. Then let $f : \bar{a} \rightarrow \bar{b}$ be the function whose graph is $\{(c(n, 0), c(n, \ell)) : n <_{\mathbf{c}} \text{lg}(c)\}$. Clearly f is definable, and the hypothesis of monotonicity in each coordinate from the proof of Theorem 3.1 guarantees that f is an injection, which takes a_α to b_α for all $\alpha < \kappa$. \square

A note to model theorists: Recall that by a characterization of Morley and Vaught the saturated models are exactly those which are both homogeneous and universal. Regular ultrapowers are universal for models of cardinality no larger

than the size of the index set, and thus failures of saturation come from failures of homogeneity. Saturation of regular ultrapowers can therefore be gauged by the absence or existence of internal structure-preserving maps between small subsets of the ultrapower. A recurrent theme of this paper, illustrated by Corollary 3.7, is how trees assist in the building of such maps.

Corollary 3.8. *Let $\mathbf{s}_1, \mathbf{s}_2$ be cofinality spectrum problems and suppose that $M^{\mathbf{s}_1} = M^{\mathbf{s}_2}, M_1^{+, \mathbf{s}_1} = M_1^{+, \mathbf{s}_2}$. If $\text{Or}(\mathbf{s}_1) \cap \text{Or}(\mathbf{s}_2)$ contains a nontrivial \mathbf{a} (thus $\Delta^{\mathbf{s}_1} \cap \Delta^{\mathbf{s}_2} \neq \emptyset$), then for all regular κ with $\kappa \leq \mathfrak{p}_{\mathbf{s}}, \kappa < \mathfrak{t}_{\mathbf{s}}$,*

$$\text{lcf}(\kappa, \mathbf{s}_1) = \text{lcf}(\kappa, \mathbf{s}_2).$$

Moreover, the same conclusion holds if $\mathbf{s} \leq \mathbf{s}'$.

Thus, without loss of generality, when computing $\mathcal{C}^{\text{ct}}(\mathbf{s})$ for such κ we may work in a larger language (provided $M^{\mathbf{s}}, M_1^{\mathbf{s}}$ admit the corresponding expansion and remain an elementary pair in the larger language) and/or consider a larger set of formulas Δ , provided that it meets the closure conditions of Definition 2.3.

Corollary 3.9. *Given a cofinality spectrum problem \mathbf{s} , we may assume $\text{Or}(\mathbf{s})$ is closed under definable subsets of $X_{\mathbf{a}}$; i.e., whenever $\mathbf{a} \in \text{Or}(\mathbf{s}), \psi(x)$ a formula in the language of M_1 such that $\psi(x) \vdash x \in X_{\mathbf{a}}$, there is $\mathbf{b} \in \text{Or}(\mathbf{s})$ with $\leq_{\mathbf{b}} = \leq_{\mathbf{a}}$ and*

$$X_{\mathbf{b}} = \{a \in X_{\mathbf{a}} : M_1 \models \psi(a)\}.$$

For definiteness, we specify that $d_{\mathbf{b}} = \min\{d_{\mathbf{a}}, \max\{x : \psi(x)\}\}$.

Proof. Let $\Delta' \supseteq \Delta^{\mathbf{s}}$ be the set of $\varphi'(x, y, \bar{z}')$ such that for some $\varphi(x, y, \bar{z}_1) \in \Delta$ and $\psi(x, \bar{z}_2) \in \tau(M^{\mathbf{s}})$ we have that $\bar{z}' = \bar{z}_1 \hat{\ } \bar{z}_2$ and $\varphi' = \varphi \wedge \psi$. Now apply Corollary 3.8 in the case where $\mathbf{s}' = (M, M_1, M^+, M_1^+, T, \Delta')$ (i.e., only the last component changes). Clearly, \mathbf{s}' remains nontrivial. \square

In fact, for our proofs below, we use this only to assume that for $\mathbf{a} \in \text{Or}(\mathbf{s})$ and $a \in X_{\mathbf{a}}$, the order $(\{x : x \in X_{\mathbf{a}}, x \leq a\}, <_{\mathbf{a}})$ may be regarded as an element of $\text{Or}(\mathbf{s})$. This would be a perfectly reasonable condition to add to the definition of ESTT, though we prefer to derive it.

4. COFINALITY SPECTRUM PROBLEMS ARE LOCALLY SATURATED

In this section we prove that cofinality spectrum problems have a certain amount of local saturation. Here “local” means partial types which are finitely satisfiable in one of the distinguished linear orders.

Definition 4.1. *Let \mathbf{s} be a cofinality spectrum problem and λ a regular cardinal. Let $p = p(x_0, \dots, x_{n-1})$ be a consistent partial type with parameters in M_1^+ . We say that p is an Or-type over M_1^+ if p is a consistent partial type in M_1^+ and for some $\mathbf{a}_0, \dots, \mathbf{a}_{n-1} \in \text{Or}(\mathbf{s})$, we have that*

$$p \vdash \bigwedge_{i < n} x_i \in X_{\mathbf{a}_i}.$$

We say simply that M_1^+ is λ -Or-saturated if every Or-type over M_1^+ over a set of size $< \lambda$ is realized in M_1^+ . Finally, we say that \mathbf{s} is λ -Or-saturated if M_1^+ is.

As we asked that $\text{Or}(\mathbf{s})$ be closed under small Cartesian products, without loss of generality we prove Theorem 4.1 assuming that $p = p(x)$ where $p \vdash x \in X_{\mathbf{a}}$ for some $\mathbf{a} \in \text{Or}(\mathbf{s})$. (Note p may not be a 1-type, for instance, if the formula defining $X_{\mathbf{a}}$ has arity greater than 1.)

Theorem 4.1. *Let \mathbf{s} be a cofinality spectrum problem. If $\kappa < \min\{\mathbf{p}_{\mathbf{s}}, \mathbf{t}_{\mathbf{s}}\}$, then \mathbf{s} is κ^+ -Or-saturated.*

Proof. We prove the claim by induction on $\kappa < \min\{\mathbf{p}_{\mathbf{s}}, \mathbf{t}_{\mathbf{s}}\}$. Suppose that $\kappa = \aleph_0$ or that the theorem holds for all infinite cardinals $\rho < \kappa$. Suppose we are given $\mathbf{a} \in \text{Or}(\mathbf{s})$ and $p = \{\varphi_i(x, \bar{a}_i) : i < \kappa\}$ which is finitely satisfiable in $X_{\mathbf{a}}$. By a second (call it internal) induction on $\alpha \leq \kappa$, we choose $c_\alpha \in \mathcal{T}_{\mathbf{a}}$ and $n_\alpha \in X_{\mathbf{a}}$ such that

- (1) $n_\alpha = \text{lg}(c_\alpha) - 1$.
- (2) $\beta < \alpha$ implies $c_\beta \trianglelefteq c_\alpha$.
- (3) if $i < \beta \leq \alpha$ and $n_\beta \leq_{\mathbf{a}} n \leq_{\mathbf{a}} n_\alpha$, then $M_1 \models \varphi_i(c_\alpha(n), \bar{a}_i)$.
- (4) c_α is below the ceiling.

For $\alpha = 0$ this is trivial. For $\alpha = \beta + 1$, by the external inductive hypothesis (or by the definition of a consistent type if $\kappa = \aleph_0$), let d realize $\{\varphi_i(x, \bar{a}_i) : i \leq \beta\}$, since $|\alpha| < \kappa$. By the internal inductive hypothesis (4), we may concatenate, so let $c_\alpha = c_\beta \hat{\ } \langle d \rangle$, $n_\alpha = n_\beta + 1$. For α limit $\leq \kappa$, $\text{cf}(\alpha) < \min\{\mathbf{p}_{\mathbf{s}}, \mathbf{t}_{\mathbf{s}}\}$ so by Lemma 2.15 there is $c_* \in \mathcal{T}$ such that $\beta < \alpha$ implies $c_\beta \trianglelefteq c_*$ and c_* is below the ceiling. Let $n_* = \text{lg}(c_*) - 1$. Now we correct the value at the limit by restricting to a suitable initial segment which preserves item (3). That is, for each $i < \alpha$, define

$$n(i) = \max\{n \leq_{\mathbf{a}} n_* : \models \varphi_i(c_*(n_*), \bar{a}_i) \text{ for all } m \text{ such that } n_i <_{\mathbf{a}} m \leq_{\mathbf{a}} n\}.$$

As this is a bounded nonempty subset of $X_{\mathbf{a}}$, $n(i)$ exists for each $i < \alpha$. By the internal inductive hypothesis (3), $n(i)_{\mathbf{a}} > n_\beta$ for each $i, \beta < \alpha$. Thus, $(\{n_\beta : \beta < \alpha\}, \{n(i) : i < \alpha\})$ represents a pre-cut in $X_{\mathbf{a}}$. Let γ be the reverse cofinality of the set $\{n(i) : i < \alpha\}$, i.e., its cofinality under the reverse order. Necessarily $\gamma \leq |\alpha| \leq \kappa < \mathbf{p}_{\mathbf{s}}$. If $(\text{cf}(\alpha), \gamma) \in \mathcal{C}(\mathbf{s}, \mathbf{t}_{\mathbf{s}})$, we would contradict the definition of $\mathbf{p}_{\mathbf{s}}$. Thus, there is an element n_{**} such that for all $i < \alpha$ and $\gamma < \alpha$, $n_\gamma <_{\mathbf{a}} n_{**} <_{\mathbf{a}} n(i)$. Let $c_\alpha = c \upharpoonright_{n_{**}}$ and let $n_\alpha = \text{lg}(c_\alpha) - 1$. Then by construction,

$$i < \alpha \text{ implies } M_1 \models \varphi_i(c_\alpha(n_\alpha), \bar{a}_i)$$

as desired, completing the limit step. As the limit case was also proved for $\alpha = \kappa$, $c_\kappa(n_\kappa)$ realizes the type p , which completes the proof. \square

Note that Theorem 4.1 does not require that the type consist of instances of formulas from some finite set (a usual definition of local); however, this is a kind of local saturation in the sense that we use $p(x) \vdash x \in X_{\mathbf{a}}$ for some suitable \mathbf{a} .

5. COFINALITY SPECTRUM PROBLEMS HAVE SUFFICIENT PEANO ARITHMETIC

In this section two things are accomplished. First, we build a certain amount of arithmetic, which shows the naturalness of the corresponding example from Section 2.3 and makes available addition and multiplication which will be useful in later proofs. Second, we show that for certain $\mathbf{a} \in \text{Or}(\mathbf{s})$, we may regard all definable subtrees $\mathcal{T} \subseteq \mathcal{T}_{\mathbf{a}} \in \text{Tr}(\mathbf{s})$ as definable subsets of some $X_{\mathbf{b}}$, $\mathbf{b} \in \text{Or}(\mathbf{s})$. Thus, in our future constructions we will have available more powerful trees, e.g., of sequences of finite tuples some of whose coordinates belong to some $X_{\mathbf{a}}$ and some of whose

coordinates effectively belong to a definable tree (thus are internal partial functions on, or subsets of, some $X_{\mathbf{a}}$, $\mathbf{a} \in \text{Or}(\mathbf{s})$). This could have been guaranteed simply by making stronger assumptions in Definition 2.3, which would remain true in our cases of main interest. Thus, the reader interested primarily in models of set theory or in regular ultrapowers may wish to simply read Convention 5.1, Conclusion 5.8, Convention 5.16 and continue. However, the analysis here shows that the necessary structure already arises from our more basic assumptions.

To begin, given \mathbf{s} and $\mathbf{a} \in \text{Or}(\mathbf{s})$, we introduce notation for the relative cardinality of definable sets $A \subseteq X_{\mathbf{a}}$, $B \subseteq X_{\mathbf{b}}$ as computed by the model M_1^+ under consideration.

Convention 5.1. (An internal notion of cardinality) *Let \mathbf{s} be a cofinality spectrum problem and $\mathbf{a}, \mathbf{b} \in \text{Or}(\mathbf{s})$. Let $\mathbf{c} = \mathbf{a} \times \mathbf{b}$ and let $\text{Par}(\mathbf{a}, \mathbf{b})$ be the definable subtree of $\mathcal{T}_{\mathbf{c}}$ given by $(\varphi(\mathcal{T}_{\mathbf{c}}), \trianglelefteq_{\mathbf{c}})$, where $\varphi(x)$ says that $\{(x(c, 0), x(c, 1)) : c <_{\mathbf{c}} \text{lg}(x)\}$ is the graph of a partial one-to-one function from $X_{\mathbf{a}}$ to $X_{\mathbf{b}}$. We will adopt the following convention. Whenever A, B are given as definable subsets of $X_{\mathbf{a}}, X_{\mathbf{b}}$, respectively, we write*

$$|A| \leq^{\mathbf{s}} |B|$$

to mean “there exists $x \in \text{Par}(\mathbf{a}, \mathbf{b})$ such that $A \subseteq \{x(n, 0) : n <_{\mathbf{c}} \text{lg}(x)\}$ and $\{x(m, 1) : m <_{\mathbf{c}} \text{lg}(x)\} \subseteq B$.” Likewise, we write

$$|A| <^{\mathbf{s}} |B|$$

to mean “ $(|A| \leq^{\mathbf{s}} |B|) \wedge \neg(|B| \leq^{\mathbf{s}} |A|)$ ”; i.e., $|A| \leq^{\mathbf{s}} |B|$ and there does not exist $x \in \text{Par}(\mathbf{a})$ which is an injection from B into A .

Discussion 5.2. *Does the internal notion of cardinality just described behave more like (the cardinality of) the natural numbers or infinite sets? For our present purposes, it behaves like pseudo-finite numbers. However, the same definition applies in very different contexts with different results, for instance, in weakenings of cofinality spectrum problems studied in [30], where the order is dense.*

We now show that a sufficient amount of Gödel coding is available in any cofinality spectrum problem (CSP). More precisely, we would like to find $\mathbf{a} \in \text{Or}(\mathbf{s})$ such that for some other $\mathbf{b} \in \text{Or}(\mathbf{s})$ extending \mathbf{a} , the set of Gödel codes of partial functions from $X_{\mathbf{a}}$ to $X_{\mathbf{a}}$ may be viewed as a definable subset of $X_{\mathbf{b}}$. Informally, the issue is that $X_{\mathbf{a}}$ may need to be quite small relative to $X_{\mathbf{b}}$; we would like not only for statements of basic arithmetic and bounded induction to hold in $X_{\mathbf{b}}$ of elements in $X_{\mathbf{a}}$, but crucially, for all functions necessary for Gödel coding of $X_{\mathbf{a}}$ -sequences—which, a priori, may be fast growing—to remain available.

Lemma 5.3. *Let \mathbf{s} be a CSP and $\mathbf{b} \in \text{Or}(\mathbf{s})$. Then there is $\mathbf{a} \in \text{Or}(\mathbf{s})$ such that $X_{\mathbf{a}}$ is an initial segment of $X_{\mathbf{b}}$ and all Gödel codes for elements of $\mathcal{T}_{\mathbf{a}}$ belong to $X_{\mathbf{b}}$. In particular, we may identify $\mathcal{T}_{\mathbf{a}}$ with a definable subset of $X_{\mathbf{b}}$.*

Proof. Let $\mathbf{b} \in \text{Or}(\mathbf{s})$ be given. We define a series of formulas in the language of M_1^+ . The variables range over elements of $X_{\mathbf{b}}$ unless otherwise indicated.

First, we define (the graph of a partial function corresponding to) addition to be the set of (x, y, z) such that there is an element η of $\mathcal{T}_{\mathbf{b}}$, i.e., a sequence of elements of $X_{\mathbf{b}}$, of length y whose first element is x , whose last element is z , and which increments by one. Formally, $\varphi_+(x, y, z) = (\exists \eta \in \mathcal{T}_{\mathbf{b}})(\text{lg}(\eta) = y \wedge \eta(0) = x \wedge \eta(y - 1) = z \wedge (\forall i)(i < \text{lg}(\eta) \rightarrow \eta(S(i)) = S(\eta(i)))$. Note that we may do

this uniformly across $\mathbf{b} \in \text{Or}(\mathbf{s})$ if we ask that φ_+ also accept a fourth parameter, the defining parameter $c_{\mathbf{b}}$ for $X_{\mathbf{b}}$, and require that the formula $\varphi_+(x, y, z)$ imply $x, y, z \in X_{\mathbf{b}}$ and that $(\forall i)$ abbreviate $(\forall i \in X_{\mathbf{b}})$. We will assume this to be true and will omit it here and in the remaining formulas of this proof for readability.

Analogously, we may define multiplication $\varphi_{\times}(x, y, z)$ by substituting $(i < \text{lg}(\eta) \rightarrow \eta(S(i)) = \eta(i) + x)$ in the appropriate place, i.e., requiring that the sequence increment by x . Define exponentiation $\varphi_{\text{exp}}(x, y, z)$ by substituting $(i < \text{lg}(\eta) \rightarrow \eta(S(i)) = \eta(i) \times x)$ in the appropriate place, i.e., requiring that the sequence increment by a factor of x .

Note that $\varphi_+(x, y, z)$, $\varphi_{\times}(x, y, z)$, and $\varphi_{\text{exp}}(x, y, z)$ are graphs of partial functions, which we will refer to informally as $x + y$, $x \times y$ and x^y . Let $\theta(x)$ assert that $x + y$, $x \times y$, and x^y exist for all $y < x$.

Clearly, with addition and multiplication we can define “ x is a prime.” Let $\varphi_4(x, y)$ assert that x is the y th prime. Let the formula $\varphi_5(x, n, m)$ assert that x is divisible by the n th prime precisely m times, by asserting the existence of $\eta \in \mathcal{T}_{\mathbf{b}}$ of length m whose first element is x , whose subsequent elements decrease by a factor of the n th prime, and whose last element has no more such factors.

Finally, let the formula $\varphi_6(x, \eta)$ assert that $x \in X_{\mathbf{b}}$ is a Gödel code for η , by asserting the existence of $\eta \neq \emptyset$ such that $\eta \in \mathcal{T}_{\mathbf{b}}$, $x > 2$, and for all $i < \text{lg}(\eta)$, x is divisible by the i th prime precisely $\eta(i) + 1$ times.

To conclude, let $\psi(w)$ be the formula

$$(\forall x < w)\theta(x) \wedge (\forall \eta \in \mathcal{T}_{\mathbf{b}})((\text{lg}(\eta) < w \wedge (\forall i < \text{lg}(\eta))(\eta(i) < w)) \rightarrow (\exists x)(\varphi_6(x, \eta))),$$

which asserts that Gödel codes exist for all functions from $X_{\mathbf{b}} \upharpoonright_w$ to itself. Then $\psi(w)$ holds of $0_{\mathbf{b}}$, and of all its finite successors, so it holds also on some non-standard number a_* . Let \mathbf{a} be the order given by $X_{\mathbf{a}} = (X_{\mathbf{b}} \upharpoonright_{a_*}, \leq_{\mathbf{b}})$, and $d_{\mathbf{a}} = \min\{d_{\mathbf{b}}, \max(X_{\mathbf{a}})\}$. Then $\mathbf{a} \in \text{Or}(\mathbf{s})$ by Corollary 3.9 and inherits nontriviality from \mathbf{b} . Thus \mathbf{a}, \mathbf{b} are as desired.

The “in particular” clause in the statement of the Lemma is clear as this was done in a (uniformly) definable way. □

Definition 5.4. (Covers) *Let \mathbf{s} be a cofinality spectrum problem and $\mathbf{a} \in \text{Or}(\mathbf{s})$.*

- (1) *Say that $\mathbf{b} \in \text{Or}(\mathbf{s})$ is a cover for \mathbf{a} if all Gödel codes for elements of $\mathcal{T}_{\mathbf{a}}$ belong to $X_{\mathbf{b}}$. The usual case is when $X_{\mathbf{a}}$ is an initial segment of $X_{\mathbf{b}}$, so is itself an element of Or by 3.9.*
- (2) *We define k -coverable by induction on $k < \omega$.*
 - (a) *Say that \mathbf{a} is 0-coverable if $\mathbf{a} \in \text{Or}(\mathbf{s})$ is nontrivial.*
 - (b) *Say that \mathbf{a} is $k + 1$ -coverable if there exists $\mathbf{b} \in \text{Or}(\mathbf{s})$ such that \mathbf{b} is a cover for \mathbf{a} and is itself k -coverable.*
- (3) *Say that \mathbf{a} is coverable if it is 1-coverable; this will be our main case.*

Claim 5.5. *Let \mathbf{s} be a cofinality spectrum problem. For each $k < \omega$ the set*

$$\{\mathbf{a} \in \text{Or}(\mathbf{s}) : \mathbf{a} \text{ is } k\text{-coverable}\}$$

is not empty.

Proof. We prove this by induction on $k < \omega$. In the case where $k = 0$ this is trivially true as Δ is nonempty and $\text{Or}(\mathbf{s})$ is assumed to be nontrivial. Suppose then that $\mathbf{b} \in \text{Or}(\mathbf{s})$ is k -coverable; we would like to find $\mathbf{a} \in \text{Or}(\mathbf{s})$ which is $k + 1$ -coverable, which we can do by applying Lemma 5.3. □

It will be useful to have the analogous result for Cartesian products.

Definition 5.6. *Say that $\mathbf{a} \in \text{Or}(\mathbf{s})$ is coverable as a pair (by $\mathbf{d} \in \text{Or}(\mathbf{s})$) when*

- (a) *there is $\mathbf{c} \in \text{Or}(\mathbf{s})$ such that $X_{\mathbf{c}} = X_{\mathbf{a}} \times X_{\mathbf{a}}$ and 2.3(6) holds of \mathbf{a}, \mathbf{c} ,*
- (b) *\mathbf{c} is coverable (by \mathbf{d}).*

Corollary 5.7. *There exists $\mathbf{a} \in \text{Or}(\mathbf{s})$ which is coverable as a pair, say by $\mathbf{d} \in \text{Or}(\mathbf{s})$, and moreover this implies*

- (1) *There is $a \in X_{\mathbf{a}}$, not a finite successor of $0_{\mathbf{a}}$, such that the Gödel codes for functions from $[0, a]_{\mathbf{a}}$ to $[0, a]_{\mathbf{a}} \times [0, a]_{\mathbf{a}}$ may be identified with a definable subset of $X_{\mathbf{d}}$.*
- (2) *\mathbf{a} is coverable by \mathbf{d} .*

When \mathbf{a} is coverable as a pair, we may informally abbreviate condition (1) by saying that some given M_1^+ -definable function may be thought of as an element of $\mathcal{T}_{\mathbf{a} \times \mathbf{a}}$.

Proof. By Claim 5.5 (or just Lemma 5.3) and the fact that Definition 5.6(a) requires that the pairing function be well behaved. □

When constructing trees in the continuation of the paper, we will generally build trees of pseudo-finite sequences of n -tuples ($n < 10$). The components of these tuples will belong to some $X_{\mathbf{a}}$ for $\mathbf{a} \in \text{Or}(\mathbf{s})$, they will be definable partial functions with domain and range contained in some $X_{\mathbf{a}}$ which is k -coverable or coverable as a pair (i.e., they will be elements of certain trees), or they will be domains of such functions, as has been justified by the work in this section.

Conclusion 5.8. (More powerful trees) *Let $\mathbf{a} \in \text{Or}(\mathbf{s})$ be coverable. We may in our future constructions consider as a definable subtree of an element of $\text{Tr}(\mathbf{s})$ any tree of sequences of finite tuples some of whose coordinates range over elements of a definable subtree of $\mathcal{T}_{\mathbf{a}}$. Without loss of generality, we also allow some coordinates to be the domain or range of such a function.*

Proof. By Corollary 3.5, to study $\mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}})$ we may freely choose the order we work in from among the nontrivial elements of $\text{Or}(\mathbf{s})$. By Claim 5.5 we may assume \mathbf{a} is coverable (or k -coverable if necessary), so elements of $\mathcal{T}_{\mathbf{a}}$ may be considered as elements of a definable subset of some $X_{\mathbf{b}}$, $\mathbf{b} \in \text{Or}(\mathbf{s})$. Then the desired tree will be a definable subtree of $\mathcal{T}_{\mathbf{b}}$ or of $\mathcal{T}_{\mathbf{c}}$ where \mathbf{c} is a finite Cartesian product of \mathbf{b} .

For the last line, the use of domains and ranges of such functions is entirely a matter of presentation. We could easily in each case use a different defining formula for the tree which did not list the domain as a separate component of the tuple. For more complex arguments, one could alternatively formalize a suitable internal notion of power set. □

We now justify the naturalness of models of Peano arithmetic with bounded induction as an example of a cofinality spectrum problem. We use exponentiation in the sense of the relation from Lemma 5.3 (which satisfies all of the usual properties of the graph of exponentiation except for the sentence $(\forall x \forall y \exists z) \varphi_{exp}(x, y, z)$).

Definition 5.9. (Weak powers) *Let \mathbf{s} be a cofinality spectrum problem and $\mathbf{a} \in \text{Or}(\mathbf{s})$. Let $e_{\mathbf{a}}$ denote the last element of $X_{\mathbf{a}}$. Say that $a \in X_{\mathbf{a}}$ has weak powers if a is not a finite successor of $0_{\mathbf{a}}$, there exists $z \in X_{\mathbf{a}}$ such that $(a^a = z)$, and for each $\ell < \omega$,*

$$a^a <_{\mathbf{a}} S^{-\ell}(e_{\mathbf{a}}).$$

Remark 5.10. Note that Definition 5.9 implies that $a^k <_{\mathbf{a}} S^{-\ell}(e_{\mathbf{a}})$ for each $\ell, k < \omega$, where a^k means $a^{S^k(0_{\mathbf{a}})}$.

Lemma 5.11. Let \mathbf{s} be a cofinality spectrum problem, $\mathbf{a} \in \text{Or}(\mathbf{s})$, and suppose $a \in X_{\mathbf{a}}$ has weak powers. Then there are definable functions $+_{\mathbf{a}}, \times_{\mathbf{a}}$ such that identifying k with $S^k(0_{\mathbf{a}})$, the model

$$N_{\mathbf{a}}^* := \left(\bigcup_{k < \omega} [0_{\mathbf{a}}, a^k]_{\mathbf{a}}; +_{\mathbf{a}}, \times_{\mathbf{a}}, S_{\mathbf{a}}^1, \leq_{\mathbf{a}}, 0_{\mathbf{a}} \right)$$

may be regarded as a model of $I\Delta_0$, arithmetic with bounded induction, meaning that the arithmetic operations are well defined on the domain of $N_{\mathbf{a}}^*$ and for any bounded formula $\varphi(x)$ in the language of arithmetic which implies $x \in X_{\mathbf{a}}$, the following holds in M_1 :

$$(\forall x \leq a_*) (\forall \bar{y}) (\varphi(0_{\mathbf{a}}, \bar{y}) \wedge (\forall z \leq x) (\varphi(z, \bar{y}) \rightarrow \varphi(S(z), \bar{y})) \rightarrow (\forall z \leq x) \varphi(z, \bar{y})).$$

Remark 5.12. Clearly, the domain of $N_{\mathbf{a}}^*$ is not definable in M_1^+ .

Proof. (of Lemma 5.11) First, we define $+_{\mathbf{a}}, \times_{\mathbf{a}}$ as partial functions from $X_{\mathbf{a}} \times X_{\mathbf{a}} \rightarrow X_{\mathbf{a}}$ and check that they behave as desired. We could use the formulas from Lemma 5.3, but it is interesting to do this another way, based only on 5.1. Let $\text{Int}_{\mathbf{a}} = \{[e_1, e_2] : e_1 <_{\mathbf{a}} e_2, e_1, e_2 \in X_{\mathbf{a}}\}$. Formally, elements of $\text{Int}_{\mathbf{a}}$ are presented as pairs but we think of them as closed and open intervals. Let $E_{\mathbf{a}}$ be the definable equivalence relation on $\text{Int}_{\mathbf{a}}$ given by having the same cardinality in the sense of \mathbf{s} (i.e., writing $X = \{x : e_1 \leq_{\mathbf{a}} x <_{\mathbf{a}} e_2\}$, $Y = \{y : e'_1 \leq_{\mathbf{a}} y <_{\mathbf{a}} e'_2\}$, we have that $|X| \leq^{\mathbf{s}} |Y|$ and $|Y| \leq^{\mathbf{s}} |X|$).

Denote by $0_{\mathbf{a}}$ the $\leq_{\mathbf{a}}$ -minimal element of $X_{\mathbf{a}}$. For addition, define

$$+_{\mathbf{a}}(x, y) = z \text{ if } (\exists w \in X_{\mathbf{a}}) (w \leq_{\mathbf{a}} z \wedge [0_{\mathbf{a}}, x) E_{\mathbf{a}} [0_{\mathbf{a}}, w) \wedge [0_{\mathbf{a}}, y) E_{\mathbf{a}} [w, z)).$$

We will write $1 = 1_{\mathbf{a}}$ for the successor of $0_{\mathbf{a}}$ in the discrete linear order $\leq_{\mathbf{a}}$. Then clearly

$$S^1(w) = +_{\mathbf{a}}(w, 1_{\mathbf{a}}).$$

For multiplication, define

$$\begin{aligned} &\times_{\mathbf{a}}(x, y) = z \text{ if} \\ &(\exists f \in \text{Par}(\mathbf{a})) (\text{dom}(f) = [0, x]_{\mathbf{a}} \wedge (\forall w <_{\mathbf{a}} x = \text{lg}(f)) ([f(w), f(w+1)) E_{\mathbf{a}} [0, y)]). \end{aligned}$$

Clearly $+_{\mathbf{a}}, \times_{\mathbf{a}}$ act like addition and multiplication except for the fact that not all pairs of elements of $X_{\mathbf{a}}$ will have a sum or product (if they are too large). This shows $N_{\mathbf{a}}^*$ is a model of Peano arithmetic without induction in which every nonzero element has a predecessor.

Recalling that $\text{dom}(N_{\mathbf{a}}^*) = \bigcup_{k < \omega} [0_{\mathbf{a}}, a^k]_{\mathbf{a}}$, we verify bounded induction on $[0_{\mathbf{a}}, a^k]$ for each relevant formula and each k . Let $k < \omega$ be given and let $a_* = a^k$. For any bounded formula $\psi(x)$ in the language of arithmetic, possibly with parameters, which implies $x \in X_{\mathbf{a}}$, if $M_1^+ \models \psi(0_{\mathbf{a}})$, then the set $\{x \in X_{\mathbf{a}} : (\forall z \leq_{\mathbf{a}} x) \psi(z)\}$ is a definable nonempty subset of $X_{\mathbf{a}}$, and so has a greatest element n_{ψ} . If $n_{\psi} \leq a_*$, $(\forall z \leq n_{\psi}) (\psi(z) \rightarrow \psi(S(z)))$ is well defined, so it is either true in M_1^+ (in which case we contradict $n_{\psi} \leq a_*$) or false in M_1^+ (in which case this instance of induction is trivially true).

This completes the proof. □

Definition 5.13. Say that $\varphi(x)$ is a weakly initial formula for \mathbf{a} if it is a formula in the language of M_1^+ , possibly with parameters, which implies $x \in X_{\mathbf{a}}$ and which holds on $S^k(0_{\mathbf{a}})$ for all $k < \omega$.

Claim 5.14. Let \mathbf{s} be a cofinality spectrum problem, $\mathbf{a} \in \text{Or}(\mathbf{s})$, $\varphi(x)$ a weakly initial formula for \mathbf{a} .

- (1) There is a nonstandard $a \in X_{\mathbf{a}}$ such that $M_1^+ \models \varphi(a)$.
- (2) We may also require that $a < S^{-\ell}(e_{\mathbf{a}})$ for all $\ell < \omega$.

Proof. For part (1), the set $A = \{b \in X_{\mathbf{a}} : (\forall c \leq_{\mathbf{a}} b)\varphi(c)\}$ contains all the standard elements of $X_{\mathbf{a}}$ and is definable, so it has a last element a_* , which is necessarily nonstandard. For part (2), let a be any element realizing the countable pre-cut described by the standard elements on the left and $\{S^{-\ell}(a_*) : \ell < \omega\}$ on the right. □

Alternately, Theorem 4.1 shows that in Claim 5.14 we may replace φ with a consistent partial type $\Sigma(x)$ closed under conjunction, each of whose formulas (thus, finite subtypes) is weakly initial for \mathbf{a} and such that $|\Sigma| < \min\{\mathfrak{p}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}}\}$.

Conclusion 5.15. Let \mathbf{s} be a cofinality spectrum problem, $\mathbf{a} \in \text{Or}(\mathbf{s})$. There are $a \in X_{\mathbf{a}}$ which have weak powers in the sense of 5.9, i.e., for which the hypotheses of Lemma 5.11 are satisfied.

Proof. Let $\varphi_{exp}(x, y, z) = "x^y = z"$ be the formula from Lemma 5.3, let $\varphi(x) = \exists y\psi(x, x, y)$, and apply Claim 5.14 to $\varphi(x)$. □

Convention 5.16. In what follows, whenever the context of a background $X_{\mathbf{a}}$ is clear, we freely use the partial functions $+$, \times , and exponentiation from Lemma 5.11 (or Lemma 5.3), as well as 0 , 1 , $+k$, $-k$, internal cardinality $|\cdot|$ in the sense of 5.1 (really a means of internally comparing two sets), successor $S^{\ell}, S^{-\ell}$ from 2.11. We also adopt the more usual notation $x + y$, $x \times y$.

6. THERE ARE NO SYMMETRIC CUTS STRICTLY BELOW $\min\{\mathfrak{p}_{\mathbf{s}}^+, \mathfrak{t}_{\mathbf{s}}\}$

With the results of Sections 4 and 5 (local saturation, basic arithmetic) in hand, we return to the main line of our investigation: the cofinality spectrum $\mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}})$. To Uniqueness, Theorem 3.1, we now add Lemma 6.1.

Lemma 6.1. Let \mathbf{s} be a cofinality spectrum problem. Then for all regular κ such that $\kappa \leq \mathfrak{p}_{\mathbf{s}}$, $\kappa < \mathfrak{t}_{\mathbf{s}}$, we have that $(\kappa, \kappa) \notin \mathcal{C}^{ct}(\mathbf{s})$.

Proof. Without loss of generality, we fix $\mathbf{a} \in \text{Or}(\mathbf{s})$ which is coverable, say by \mathbf{b} . We prove the lemma by induction on κ . Arriving to κ , assume for a contradiction that $(\langle a_{\alpha} : \alpha < \kappa \rangle, \langle b_{\alpha} : \alpha < \kappa \rangle)$ represents a (κ, κ) -cut in $X_{\mathbf{a}}$. Fix a definable partial injection g from $X_{\mathbf{b}} \times X_{\mathbf{b}}$ to $X_{\mathbf{b}}$ whose domain includes $X_{\mathbf{a}} \times X_{\mathbf{a}}$. (Such a function always exists as $\mathcal{T}_{\mathbf{a}}$ may be identified with a definable subset of $X_{\mathbf{b}}$, and ordered pairs of elements of $X_{\mathbf{a}}$ may be represented as elements of $\mathcal{T}_{\mathbf{a}}$ of length 2. For definiteness, we may assume $X_{\mathbf{a}} \subseteq X_{\mathbf{b}}$ and use $(a, b) \mapsto (a + b + 1)^2 + a$, which will be defined on all pairs of elements from $X_{\mathbf{a}}$ by the choice of $X_{\mathbf{b}}$.) Define \mathcal{T} to be the set of $x \in \mathcal{T}_{\mathbf{b}}$ such that

- (a) $x(n)$, when defined, is $g(a, b)$ for some $a, b \in X_{\mathbf{a}}$. In slight abuse of notation, we will denote a by $x(n, 0)$ and b by $x(n, 1)$.
- (b) $n_1 <_{\mathbf{b}} n_2 <_{\mathbf{b}} \text{lg}(x)$ implies $x(n_1, 0) <_{\mathbf{a}} x(n_2, 0) < x(n_2, 1) <_{\mathbf{a}} x(n_1, 1)$.

Just as in the proof of Theorem 3.1, we choose a sequence c_α, n_α by induction on $\alpha < \kappa$ so that

- (1) $c_\alpha \in \mathcal{T}$ and $n_\alpha = \text{lg}(c_\alpha)$.
- (2) $\beta < \alpha$ implies $c_\beta \trianglelefteq c_\alpha$.
- (3) c_α is below the ceiling.
- (4) $c_\alpha(n_\alpha) = g(a_\alpha, b_\alpha)$.

For limit $\alpha < \kappa$, Lemma 2.15 applies, so we may carry out the induction. Having defined $\langle c_\alpha : \alpha < \kappa \rangle$, as $\kappa < \mathfrak{t}_s$ we apply Claim 2.14 to choose $c \in \mathcal{T}$ such that $\alpha < \kappa$ implies $c_\alpha \trianglelefteq c$. Let $\mathbf{n} = \text{lg}(c) - 1$. Then by clause (b) of the definition of \mathcal{T} , we have that for all $\alpha < \kappa$,

$$a_\alpha = c(n_\alpha, 0) <_{\mathbf{a}} c(\mathbf{n}, 0) <_{\mathbf{a}} c(\mathbf{n}, 1) <_{\mathbf{a}} c(n_\alpha, 1) = b_\alpha.$$

This contradicts the assumption that $(\langle a_\alpha : \alpha < \kappa \rangle, \langle b_\alpha : \alpha < \kappa \rangle)$ represents a (κ, κ) -cut in $X_{\mathbf{a}}$. □

Note that Lemma 6.1 generalizes Lemma 2.2.

Lemma 6.2. *Let \mathbf{s} be a cofinality spectrum problem and κ a regular cardinal. If $\kappa = \mathfrak{t}_s$, then there is a definable linear order which has a (κ, κ) -cut.*

Remark 6.3. *It is not asserted that this definable linear order is an element of $\text{Or}(\mathbf{s})$.*

Proof of Lemma 6.2. By Definition 2.10, if $\kappa = \mathfrak{t}_s$, then there is some $\mathcal{T}_{\mathbf{a}} \in \text{Tr}(\mathbf{s})$ which contains a strictly $\trianglelefteq_{\mathbf{a}}$ -increasing sequence $\langle c_i : i < \kappa \rangle$ with no upper bound.

We construct a linear order by collapsing the tree so that the presence or absence of κ^+ -treetops (upper bounds of linearly ordered sequences of cofinality κ) corresponds to realization or omission of symmetric (κ, κ) -cuts. Note that as $X_{\mathbf{a}}$ is linearly ordered by $\leq_{\mathbf{a}}$, we have available a definable linear ordering on the immediate successors of any given $c \in \mathcal{T}_{\mathbf{a}}$. To simplify notation, write $\text{cis}(c_i, c_j)$ for the common initial segment of c_i, c_j .

Fix two distinct elements of $X_{\mathbf{a}}$; without loss of generality we use $0_{\mathbf{a}}, 1_{\mathbf{a}}$, called $0, 1$, so $0 <_{\mathbf{a}} 1$. Let X be the set $\mathcal{T}_{\mathbf{a}} \times \{0, 1\}$. Let $<_X$ be the linear order on X defined as follows:

- If $c = d$, then $(c, 0) <_X (c, 1)$.
- If $c \trianglelefteq_{\mathbf{a}} d$ and $c \neq d$, then $(c, 0) <_X (d, 0) <_X (d, 1) <_X (c, 1)$.
- If c, d are \trianglelefteq -incomparable, then let $e \in \mathcal{T}_{\mathbf{a}}, n_c, n_d \in X_{\mathbf{a}}$ be such that $e = \text{cis}(c, d)$, $e \frown n_c \trianglelefteq_{\mathbf{a}} c$, and $e \frown n_d \trianglelefteq_{\mathbf{a}} d$. Necessarily $n_c \neq n_d$ by definition of e , so for $s, t \in \{0, 1\}$ we define

$$(c, s) <_X (d, t) \iff n_c <_{\mathbf{a}} n_d.$$

(Informally speaking, each node separates into a set of matched parentheses enclosing the cone above it.) Then $<_X$ is definable linear order on X with a first and a last element.

Recalling the definition of the sequence $\langle c_i : i < \kappa \rangle$ from the beginning of the proof, it follows that

$$(\langle (c_i, 0) : i < \kappa \rangle, \langle (c_i, 1) : i < \kappa \rangle)$$

represents a (κ, κ) -cut in $(X, <_X)$. This is because it is a definable linear order on a definable set in M_1^+ and every element misses the cut at some initial stage. This completes the proof. □

Lemma 6.2 shows that if we were to, e.g., extend our definition of cuts in $\mathcal{C}^{\text{ct}}(\mathbf{s})$ to include the one in Lemma 6.2 (or to show that Δ may be suitably extended, so 3.8 applies), this would give a characterization of $\mathfrak{t}_{\mathbf{s}}$: if $\kappa = \mathfrak{t}_{\mathbf{s}}$, then $\mathfrak{t}_{\mathbf{s}} = \min\{\kappa : (\kappa, \kappa) \in \mathcal{C}^{\text{ct}}(\mathbf{s})\}$; thus also $\mathfrak{p}_{\mathbf{s}} \leq \mathfrak{t}_{\mathbf{s}}$ by definition of $\mathfrak{p}_{\mathbf{s}}$. This is not necessary for our present arguments; it is established in certain cases, e.g., 10.24 below. The full characterization is proved under similar additional hypotheses (which hold in all the main examples of the present paper) in the authors' paper [33].

7. IF $\text{lcf}(\aleph_0, \mathbf{s}) > \aleph_1$, THEN $\text{lcf}(\aleph_0, \mathbf{s}) \geq \min\{\mathfrak{p}_{\mathbf{s}}^+, \mathfrak{t}_{\mathbf{s}}\}$

As a warm-up to Section 8, in the present section we will prove Theorem 7.1: for a cofinality spectrum problem \mathbf{s} , if $\text{lcf}(\aleph_0, \mathbf{s}) \neq \aleph_1$, then either $\text{lcf}(\aleph_0, \mathbf{s}) > \mathfrak{p}_{\mathbf{s}}$ or $\text{lcf}(\aleph_0, \mathbf{s}) \geq \mathfrak{t}_{\mathbf{s}}$.

Convention 7.1. For \mathbf{s} a cofinality spectrum problem, $\mathbf{a} \in \text{Or}(\mathbf{s})$, $a_1 <_{\mathbf{a}} a_2$ from $X_{\mathbf{a}}$ we write $(a_1, a_2)_{\mathbf{a}}$ for

$$\{x \in X_{\mathbf{a}} : a_1 <_{\mathbf{a}} x <_{\mathbf{a}} a_2\} \subseteq X_{\mathbf{a}}$$

and analogously for closed and half-open intervals.

Convention 7.2. Various subsequent proofs will involve the same three types of sequences: we are given a cut represented by (\bar{d}, \bar{e}) and an additional sequence of constants \bar{a} . We standardize indexing as follows:

- $\bar{d} = \langle d_{\epsilon} : \epsilon < \dots \rangle$;
- $\bar{e} = \langle e_{\alpha} : \alpha < \dots \rangle$;
- $\bar{a} = \langle a_i : i < \dots \rangle$.

The proofs also involve building sequences $\langle c_{\alpha} : \alpha < \dots \rangle$ of elements in a given tree, where the definition of c_{α} generally depends on e_{α} , as the index suggests.

We begin with a preliminary claim. Our main case is $\kappa = \aleph_0$, but it will be useful to have the general statement.

Claim 7.3. Let \mathbf{s} be a cofinality spectrum problem. Suppose that we are given

- (1) $\mathbf{a} \in \text{Or}(\mathbf{s})$, which is coverable as a pair by \mathbf{b} ,
- (2) $e <_{\mathbf{a}} e' \in X_{\mathbf{a}}$ infinitely far apart,
- (3) $\kappa^+ < \min\{\mathfrak{p}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}}\}$,
- (4) $\{a_i : i < \kappa^+\}$ a set of pairwise distinct elements of $X_{\mathbf{a}}$,
- (5) $\gamma = \text{cf}(\gamma) < \min\{\mathfrak{p}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}}\}$,
- (6) $\langle f_{\beta} : \beta < \gamma \rangle$ a sequence of definable partial injections from $X_{\mathbf{a}}$ into $X_{\mathbf{a}}$, such that for all $\beta < \gamma$,

$$\text{dom}(f_{\beta}) \supseteq \{a_i : i < \kappa^+\}.$$

Then there exists a definable partial injection f such that

- (a) $a_i \in \text{dom}(f)$ for all $i < \kappa^+$,
- (b) $\text{dom}(f) \subseteq \text{dom}(f_{\beta})$ for all $\beta < \gamma$,
- (c) $\text{range}(f) \subseteq (e, e')_{\mathbf{a}}$.

Moreover, we may identify f with an element of $X_{\mathbf{b}}$.

Proof. We have assumed \mathbf{a} is coverable as a pair by \mathbf{b} , Definition 5.6, which assumes we fix an appropriate $\mathbf{c} \in \text{Or}(\mathbf{s})$ such that $X_{\mathbf{c}} = X_{\mathbf{a}} \times X_{\mathbf{a}}$ and \mathbf{b} covers \mathbf{c} . Let $\mathcal{T} \subseteq \mathcal{T}_{\mathbf{c}}$ be the definable subtree of elements x such that $\text{range}(x)$ is the graph of a partial injection from $X_{\mathbf{a}}$ into $(e, e')_{\mathbf{a}}$, which we denote g_x . Let p be the partial type of

an element $x \in \mathcal{T}$ whose associated g_x satisfies the conditions (a)–(c) on f given in the statement of the Claim. By hypothesis, elements of \mathcal{T} may be identified with elements of $X_{\mathbf{b}}$. Thus, to show that p is a partial Or-type in M_1^+ of size $\kappa^+ + |\gamma| < \min\{\mathfrak{p}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}}\}$, it will suffice to show that p is finitely satisfiable in \mathcal{T} .

Consider an arbitrary but fixed finite subset $p_* \subseteq p$ involving only $\{f_\beta : \beta \in \sigma\}$ and $\{a_i : i \in \tau\}$ for some fixed $\sigma \in [\gamma]^{<\aleph_0}$ and $\tau \in [\kappa^+]^{<\aleph_0}$. By assumption (6) of the Claim, $\{a_i : i \in \tau\} \subseteq \bigcap_{\beta \in \sigma} \text{dom}(f_\beta)$. List this set as $\{a_{i_k} : k < |\tau|\}$. By assumption (2), there exist $|\tau|$ distinct elements $\{b_k : k < |\tau|\}$ of the interval $(e, e')_{\mathbf{a}}$. Let $c_* = \langle a_0, b_0 \rangle \frown \cdots \frown \langle a_{i_{|\tau|-1}}, b_{|\tau|-1} \rangle$. Note that this function will be an element of \mathcal{T} by closure under concatenation and the hypothesis that \mathbf{a} is nontrivial (thus also \mathbf{c} is nontrivial, by choice of \mathbf{c} in 5.6). Thus, c_* realizes p_* . This completes the proof that p is finitely satisfiable in \mathcal{T} .

By Or-saturation, Theorem 4.1 above, p is realized by some c . Let g_c be the associated definable partial function described in the first paragraph of the proof. Then $f = g_c$ is as required. \square

Note that in the proof just given, we did not require the functions f_β to be uniformly definable nor to belong to $\mathcal{T}_{\mathbf{c}}$.

Lemma 7.4. *Let \mathbf{s} be a cofinality spectrum problem and $\aleph_1 < \lambda = \text{cf}(\lambda) \leq \mathfrak{p}_{\mathbf{s}}$, $\lambda < \mathfrak{t}_{\mathbf{s}}$. If*

$$\text{lcf}(\aleph_1, \mathbf{s}) \geq \aleph_2 \quad \text{and} \quad \text{lcf}(\aleph_0, \mathbf{s}) \geq \lambda,$$

then $\text{lcf}(\aleph_0, \mathbf{s}) > \lambda$.

Proof. Assume for a contradiction that $\text{lcf}(\aleph_0, \mathbf{s}) = \lambda$. Note that the hypotheses of the claim guarantee that $\aleph_1 < \min\{\mathfrak{p}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}}\}$. Since we may choose any order in which to work, let $\mathbf{a} \in \text{Or}(\mathbf{s})$ be coverable as a pair and let $\mathbf{b} \in \text{Or}(\mathbf{s})$ be a cover for it. Without loss of generality, we may choose

$$\langle \langle d_\epsilon : \epsilon < \omega \rangle, \langle e_\alpha : \alpha < \lambda \rangle \rangle$$

representing an (\aleph_0, λ) -cut in $X_{\mathbf{a}}$ such that d_0 is the $\leq_{\mathbf{a}}$ -least element of $X_{\mathbf{a}}$ and $d_{\epsilon+1} = S^1(d_\epsilon)$ for $\epsilon < \omega$, whereas the elements of \bar{e} are infinitely far apart: $S^k(e_{\alpha+1}) < e_\alpha$ for each $\alpha < \lambda$ and $k < \omega$. This can be done as follows. Let $\langle d_\epsilon : \epsilon < \omega \rangle$ be as specified; by choice of $X_{\mathbf{a}}$, $X := X_{\mathbf{a}} \setminus \{d_\epsilon : \epsilon < \omega\} \neq \emptyset$ so by uniqueness there is a sequence $\bar{e} = \langle e_\alpha : \alpha < \lambda \rangle$ coinital in X . If necessary, replace \bar{e} with $\langle e_{\omega \cdot \alpha} : \alpha < \lambda \rangle$ (ordinal product) to achieve the spacing.

To complete the preliminaries, choose a sequence $\langle a_i : i < \aleph_1 \rangle$ of distinct elements of $X_{\mathbf{a}}$, which exists by Claim 3.1 since a fortiori $\aleph_1 \leq \mathfrak{p}_{\mathbf{s}}$.

Let $\mathbf{d} \in \text{Or}(\mathbf{s})$ be such that $X_{\mathbf{d}} = X_{\mathbf{a}} \times X_{\mathbf{a}} \times X_{\mathbf{b}} \times X_{\mathbf{b}}$. Let $\mathcal{T}_{\mathbf{d}}$ be the subtree of $\mathcal{T}_{\mathbf{d}}$ consisting of elements x such that

- (a) $x(n, 0) <_{\mathbf{a}} x(n, 1)$ are from $X_{\mathbf{a}}$;
- (b) $x(n, 2)$ is a subset of $X_{\mathbf{a}}$, equal to $\text{dom}(x(n, 3))$;
- (c) $x(n, 3)$ is a one-to-one function from $x(n, 2)$ into the interval $(x(n, 0), x(n, 1))_{\mathbf{a}}$;
- (d) $n_1 <_{\mathbf{d}} n_2$ in $\text{dom}(x)$ implies $x(n_2, 1) <_{\mathbf{a}} x(n_1, 0)$ and $x(n_2, 2) \subsetneq x(n_1, 2)$.

Recalling the choice of $\langle e_\alpha : \alpha < \lambda \rangle$, we will now choose $c_\alpha \in \mathcal{T}_{\mathbf{d}}$, $n_\alpha \in X_{\mathbf{c}}$ by induction on $\alpha < \lambda$ such that

- (i) $\beta < \alpha$ implies $c_\beta \trianglelefteq c_\alpha$;
- (ii) $n_\alpha = \max(\text{dom}(c_\alpha))$;
- (iii) each c_α is below the ceiling;

- (iv) $e_{\alpha+1} \leq_{\mathbf{a}} c_{\alpha}(n_{\alpha}, 0) <_{\mathbf{a}} c_{\alpha}(n_{\alpha}, 1) \leq_{\mathbf{a}} e_{\alpha}$;
- (v) $a_i \in c_{\alpha}(n_{\alpha}, 2)$ for each $i < \aleph_1$.

Let us carry out the induction:

For $\alpha = 0$: Apply Claim 7.3 with $\kappa^+ = \aleph_1$, $\gamma = 0$, $e = e_1$, $e' = e_0$ to obtain a suitable partial injection g , which can be coded as an element of $X_{\mathbf{b}}$, and whose domain contains $\{a_i : i < \aleph_1\}$. Let $c_0 = \langle e_1, e_0, \text{dom}(g), g \rangle$, and let $n_0 = 0$.

For $\alpha = \beta + 1$: Likewise, apply Claim 7.3 with $\kappa^+ = \aleph_1$, $\gamma = 1$, $f_0 = c_{\beta}(n_{\beta}, 3)$, $e = e_{\alpha+1}$, $e' = e_{\alpha}$ to obtain a definable partial injection $g \in X_{\mathbf{b}}$ whose domain includes $\{a_i : i < \aleph_1\}$ and whose range is included in the interval $(e_{\alpha+1}, e_{\alpha})$. Let $c_{\alpha} = c_{\beta} \hat{\ } \langle e_{\alpha}, e_{\beta}, \text{dom}(g), g \rangle$. Let $n_{\alpha} = n_{\beta} + 1$.

For $\alpha < \lambda$ limit: As $\text{cf}(\alpha) \leq \alpha < \lambda$ thus $\text{cf}(\alpha) < \min\{\mathfrak{p}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}}\}$, by Lemma 2.15 there is $c \in \mathcal{T}_2$ such that $\beta < \alpha$ implies $c_{\beta} \preceq c$ and c is below the ceiling. Then

$$\{n : n <_{\mathbf{c}} \text{lg}(c), M_1^+ \models e_{\alpha} \leq_{\mathbf{a}} c(n, 0)\}$$

is a bounded nonempty subset of $X_{\mathbf{c}}$ which contains $n_{\beta} = \max(\text{dom}(c_{\beta}))$ for all $\beta < \alpha$. Let n_* be its maximal element (necessarily $n_* > n_{\gamma}$ for all $\gamma \leq \beta$). Once more, let g be the definable partial injection given by Claim 7.3 when we use

- (1) $e_{\alpha+1}, e_{\alpha}$ for e, e' in 7.3(2);
- (2) $\kappa^+ = \aleph_1$ in 7.3(3);
- (3) $\{a_i : i < \aleph_1\}$ in 7.3(4);
- (4) $\text{cf}(\alpha)$ for γ in 7.3(5), noting that $\text{cf}(\alpha) < \lambda$ thus $\text{cf}(\alpha) < \min\{\mathfrak{p}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}}\}$;
- (5) $\langle c(n_{\zeta}, 3) : \zeta \in X \rangle$ for the sequence of functions in 7.3(6), where X is any sequence cofinal in α .

Let $c_{\alpha} = \langle e_{\alpha+1}, e_{\alpha}, \text{dom}(g), g \rangle$ and let $n_{\alpha+1} = \max(\text{dom}(c_{\alpha}))$.

This completes the inductive construction of the sequence $\langle c_{\alpha} : \alpha < \lambda \rangle$. As $\lambda < \mathfrak{t}_{\mathbf{s}}$, by Claim 2.14 there is $c \in \mathcal{T}_2$ such that $c_{\alpha} \preceq c$ for all $\alpha < \lambda$. Let $m = \max(\text{dom}(c))$. By the choice of (\bar{d}, \bar{e}) , necessarily both $c(m, 0) <_{\mathbf{a}} c(m, 1)$ are elements of the sequence \bar{d} . For each $i < \aleph_1$, the set of $m' \leq_{\mathbf{c}} m$ such that $M_1^+ \models a_i \in c(m', 2)$ is definable, bounded, and nonempty, so let $m_i \leq_{\mathbf{c}} m$ be its maximal element. By our construction, $n_{\alpha} \leq_{\mathbf{c}} m_i$ for each $i < \aleph_1$ and each $\alpha < \lambda$. Thus, for each $i < \aleph_1$, also $c(m_i, 0) <_{\mathbf{a}} c(m_i, 1)$ are elements of the countable sequence \bar{d} . By the pigeonhole principle, there is $d_* \in \bar{d}$ such that $W := \{i < \aleph_1 : c(m_i, 1) = d_*\}$ is uncountable. Note that by our choice of \bar{d} , the element d_* is finite, so in particular the set $[0, d_*]_{\mathbf{a}}$ of predecessors of d_* is finite. There are two cases. If there exists an uncountable (or just infinite) $W' \subseteq W$ on which the function $i \mapsto m_i$ is constant, let m_* denote this constant value. In this case $c(m_*, 3)$ is a definable injection whose domain includes $\{a_i : i \in W'\}$ and whose range is contained in the finite set of predecessors of d_* , a clear contradiction. Otherwise, we may find an infinite $W' \subseteq W$ on which the function $i \mapsto m_i$ is one-to-one. Let g be the function given by $a_i \mapsto c(m_i, 3)(a_i)$. Then g is one-to-one on $\{a_i : i \in W'\}$ because $k <_{\mathbf{c}} l$ implies $c(l, 0) <_{\mathbf{a}} c(l, 1) <_{\mathbf{a}} c(k, 0) <_{\mathbf{a}} c(k, 1)$. Moreover, $g(a_i) \leq_{\mathbf{a}} d_*$ for all $i \in W'$. Let W'' be any subset of W' of cardinality $d_* + 2$. Then $g \upharpoonright \{a_i : i \in W''\}$ is a definable injection of a finite set of size $d_* + 2$ into a finite set of size $d_* + 1$, a clear contradiction. This completes the proof. \square

We arrive to the main result of the section, Theorem 7.1, whose statement suggests that to show $(\aleph_0, \lambda) \notin \mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}})$ the hardest cut to rule out is the cut (\aleph_0, \aleph_1) . This intuition will be borne out in later cases.

Theorem 7.1. *Let \mathbf{s} be a cofinality spectrum problem. If $\text{lcf}(\aleph_0, \mathbf{s}) \neq \aleph_1$, then either $\text{lcf}(\aleph_0, \mathbf{s}) > \mathfrak{p}_{\mathbf{s}}$ or $\text{lcf}(\aleph_0, \mathbf{s}) \geq \mathfrak{t}_{\mathbf{s}}$.*

Proof. We will prove that if $\text{lcf}(\aleph_1, \mathbf{s}) \geq \aleph_2$, then $\text{lcf}(\aleph_0, \mathbf{s}) > \lambda$ for all regular λ such that $\aleph_1 < \lambda \leq \mathfrak{p}_{\mathbf{s}}$ and $\lambda < \mathfrak{t}_{\mathbf{s}}$. This is equivalent to the statement of the theorem because $\text{lcf}(\aleph_1, \mathbf{s}) \geq \aleph_2$ precisely when $(\aleph_0, \aleph_1) \notin \mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}})$, recalling Theorem 3.1 (Uniqueness) and Lemma 6.1 (Anti-Symmetry).

We prove the claim by induction on $\lambda \geq \aleph_2$. Recall that by Corollary 3.4, for regular cardinals $\gamma_1, \gamma_2 < \mathfrak{t}_{\mathbf{s}}$, $\text{lcf}(\gamma_1, \mathbf{s}) = \gamma_2$ iff $(\gamma_1, \gamma_2) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$ iff $\text{lcf}(\gamma_2, \mathbf{s}) = \gamma_1$.

Base case: $\lambda = \aleph_2$. First, $(\aleph_0, \aleph_0) \notin \mathcal{C}^{\text{ct}}(\mathbf{s})$ by Lemma 6.1. Second, by hypothesis, $\text{lcf}(\aleph_1, \mathbf{s}) \geq \aleph_2$. So by uniqueness, Theorem 3.1, there is $\kappa \geq \aleph_2$ such that $(\aleph_1, \kappa) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$ and thus $(\aleph_0, \aleph_1) \notin \mathcal{C}^{\text{ct}}(\mathbf{s})$. Thus by Corollary 3.4, $\text{lcf}(\aleph_0, \mathbf{s}) \geq \aleph_2$. We then apply Lemma 7.4 to conclude $\text{lcf}(\aleph_0, \mathbf{s}) > \aleph_2$.

Inductive case. In both the successor and limit stages, by inductive hypothesis, $\text{lcf}(\aleph_0, \mathbf{s}) > \kappa$ for all $\kappa < \lambda$. Thus $\text{lcf}(\aleph_0, \mathbf{s}) \geq \lambda$. By Lemma 7.4, $\text{lcf}(\aleph_0, \mathbf{s}) > \lambda$ as desired. □

8. THERE ARE NO ASYMMETRIC CUTS STRICTLY BELOW $\min\{\mathfrak{p}_{\mathbf{s}}^+, \mathfrak{t}_{\mathbf{s}}\}$

In this section we substantially generalize the results of the previous section to prove Theorem 8.1 on asymmetric cuts. We assume $\mathfrak{p}_{\mathbf{s}} < \mathfrak{t}_{\mathbf{s}}$ (though we will state where this is used), as our background goal is to show that $\mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}}) = \emptyset$, which is clearly true if $\mathfrak{p}_{\mathbf{s}} \geq \mathfrak{t}_{\mathbf{s}}$ by definition of $\mathfrak{p}_{\mathbf{s}}$. We also use upgraded trees, following 5.8, without further comment.

Discussion 8.1. (Strategy for 8.1) Theorem 7.1 above, an early prototype of Theorem 8.1, proved that under certain conditions $\text{lcf}(\aleph_0) > \lambda$. The key inductive step in the proof involved assuming, for a contradiction, that equality held. We fixed a representative (\bar{d}, \bar{e}) of an (\aleph_0, λ) -cut in some $X_{\mathbf{a}}$, where the elements of \bar{e} were infinitely far apart and the elements of \bar{d} were all finite, and fixed a distinguished uncountable set $\{a_i : i < \aleph_1\}$. We considered a tree \mathcal{T} where $x \in \mathcal{T}$ and $n \leq \max(\text{dom}(x))$ meant that $x(n)$ gave essentially the data of a (uniformly) definable partial injection along with its domain, which included $\{a_i : i < \aleph_1\}$, and an interval of $X_{\mathbf{a}}$ containing its range. As $n \leq \max(\text{dom}(x))$ grew, the domains of these partial functions formed a decreasing sequence and the intervals bounding the range were pairwise disjoint and moved toward the left. We built a path $\langle c_\alpha : \alpha < \lambda \rangle$ through this tree such that evaluated on the maximal element of its domain, c_α gave an injection into the interval $(e_{\alpha+1}, e_\alpha)$. Informally, we were carrying the uncountable set $\{a_i : i < \aleph_1\}$ left along the right-hand side of the cut. By treetops, this sequence $\langle c_\alpha : \alpha < \lambda \rangle$ had an upper bound c . Since (\bar{d}, \bar{e}) was a cut, the elements $\{a_i : i < \aleph_1\}$ each overspilled into the domains of partial injections into intervals bounded on the right by elements of \bar{d} . From this one could find a partial injection of some large set into a finite set for the contradiction.

The main result of this section, Theorem 8.1, significantly generalizes that construction to show that $\text{lcf}(\kappa) > \lambda$ whenever $\kappa < \lambda \leq \mathfrak{p}_{\mathbf{s}} < \mathfrak{t}_{\mathbf{s}}$. A first issue in this more general case is that when $\kappa > \aleph_0$, it is not a priori sufficient to carry a set of size κ^+ into a κ -indexed sequence to obtain a contradiction. After all, elements of the κ -indexed sequence may be far apart (consider, e.g., the diagonal embedding of κ in a regular ultrapower of $(\kappa, <)$). Our control on size will need to come from internal cardinality in the sense of 5.1. A second issue arises in the case where

$\kappa^+ = \lambda$, the most subtle case of all. (Note that in 7.1, we had to assume the case (\aleph_0, \aleph_1) did not occur.) Our mechanism there for carrying the set of size κ^+ through limit stages, 7.3, required $\kappa^+ < \mathfrak{p}_s$ in order to apply Or-saturation (also $\kappa^+ < \mathfrak{t}_s$, but recall we are assuming $\mathfrak{p}_s < \mathfrak{t}_s$). Clearly, this is not satisfied when $\kappa^+ = \lambda = \mathfrak{p}_s$. In Theorem 8.1, this is solved by gradually growing the size of the set we carry (called there $\{y_{\beta+1} : \beta < \alpha \cap \kappa^+\}$) so that it has size $\leq \kappa$ at each inductive stage α when $\kappa^+ = \lambda$. It will grow all the way to κ^+ if $\kappa^+ < \lambda$, but then it is not a problem. The construction of the present section is built to coordinate these various requirements.

We begin by fine-tuning the construction of cuts. The idea of Claim 8.2 is that what was essential about (\bar{d}, \bar{e}) in Lemma 7.4 was that the elements of \bar{d} increased in internal cardinality in the sense of 5.1 and those of \bar{e} were sufficiently spaced. Here we show how to specify a minimum spacing for elements in the sequence representing the cut without losing the fact that (from the point of view of s) the cardinality genuinely grows. Recall that addition and multiplication are available following Section 5.

Claim 8.2. *Let s be a cofinality spectrum problem, $\mathfrak{a} \in \text{Or}(s)$, $\kappa \leq \mathfrak{p}_s$, $(\kappa, \lambda) \in \mathcal{C}(s, \mathfrak{t}_s)$. Let $f : X_{\mathfrak{a}} \rightarrow X_{\mathfrak{a}}$ be multiplication by 2. Then we may choose sequences $\langle d_\epsilon : \epsilon < \kappa \rangle$, $\langle e_\alpha : \alpha < \lambda \rangle$ of elements of $X_{\mathfrak{a}}$ such that*

$$(\langle d_\epsilon : \epsilon < \kappa \rangle, \langle e_\alpha : \alpha < \lambda \rangle)$$

represents a (κ, λ) -cut, and moreover

- (1) $\alpha < \lambda$ implies $f(e_{\alpha+1}) <_{\mathfrak{a}} e_\alpha$;
- (2) for each $\epsilon < \kappa$ there is $\delta = \delta(\epsilon)$, $\epsilon < \delta < \kappa$ such that

$$M_1^+ \models |d_\epsilon| <^s |d_\delta|,$$

where $|d| := |\{x \in X_{\mathfrak{a}} : x <_{\mathfrak{a}} d\}|$ and cardinality is meant the sense of 5.1.

Remark 8.3. *As is clear, the proof holds for more general functions f . Recall that 5.1 describes an internal relation comparing certain definable sets by means of partial functions available in specific trees.*

Proof of Claim 8.2. Note that $(\kappa, \lambda) \in \mathcal{C}(s, \mathfrak{t}_s)$ implies $\kappa + \lambda < \mathfrak{t}_s$ by definition. Let \mathcal{T}_4 be the definable subtree of $\mathcal{T}_{\mathfrak{a}}$ consisting of elements x such that

- (i) $n_1 <_{\mathfrak{a}} n_2 <_{\mathfrak{a}} \text{lg}(x)$ implies $f(x(n_1)) <_{\mathfrak{a}} x(n_2)$;
- (ii) $n_1 <_{\mathfrak{a}} n_2 <_{\mathfrak{a}} \text{lg}(x)$ implies $|x(n_1)| <^s |x(n_2)|$ in the sense of 5.1.

Note that this tree is nonempty and contains arbitrarily long finite branches. For $a \in X_{\mathfrak{a}}$, say that *cardinality f grows above a* to mean that for each $n < \omega$ there are $x_0, \dots, x_n \in X_{\mathfrak{a}}$ such that $x_0 = a$, $f(x_\ell) <_{\mathfrak{a}} x_{\ell+1}$ for $\ell < n$, and

$$|a| <^s |x_1| <^s \dots <^s |x_n|.$$

Clearly this holds for all finite successors of $0_{\mathfrak{a}}$. By induction on $\epsilon < \kappa$ we choose $c_\epsilon \in \mathcal{T}_4$, $n_\epsilon = \text{lg}(c_\epsilon) - 1 \in X_{\mathfrak{a}}$, and $d_\epsilon \in X_{\mathfrak{a}}$ so that

- (a) $\beta < \epsilon$ implies $c_\beta \preceq c_\epsilon$;
- (b) $c_\epsilon(n_\epsilon) = d_\epsilon$;
- (c) c_ϵ is below the ceiling;
- (d) for each $n \leq_{\mathfrak{a}} n_\epsilon$, cardinality f grows above $c_\epsilon(n)$.

The proof is similar to Theorem 3.1 (Uniqueness), with an adjustment for (d); however, we will give the argument for completeness.

For $\epsilon = 0$: let $c_0 = \langle 0_{\mathbf{a}} \rangle$ and let $n_0 = 0_{\mathbf{a}}$. As \mathbf{a} is nontrivial (so contains all the finite successors of $0_{\mathbf{a}}$) and f is multiplication by 2, (d) is satisfied.

For $\epsilon = \beta + 1$: since c_β is below the ceiling, concatenation is possible. By inductive hypothesis (d), we may choose $d_\epsilon \in X_{\mathbf{a}}$ such that $f(c_\beta(n_\beta)) <_{\mathbf{a}} d_\epsilon$, $|f(c_\beta(n_\beta))| <^{\mathbf{s}} |d_\epsilon|$, and cardinality f grows above d_ϵ . Let $c_\epsilon = c_\beta \hat{\ } \langle d_\epsilon \rangle$, and let $n_\epsilon = n_\beta + 1$, and the inductive hypotheses will be preserved.

For $\epsilon < \kappa$ limit: As $\text{cf}(\epsilon) < \min\{\mathfrak{p}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}}\}$, apply Lemma 2.15 to obtain $c \in \mathcal{T}_4$ such that $\beta < \epsilon$ implies $M_1^+ \Vdash c_\beta \sqsubseteq c$, and c is below the ceiling. Let $d_\infty = c(\text{lg}(c) - 1)$. The key point is to ensure (d). As $c \in \mathcal{T}_4$, (ii) holds. For each $k < \omega$, let $m_k = \max\{n <_{\mathbf{a}} \text{lg}(c) : c(n) \leq_{\mathbf{a}} f^{-k}(d_\infty)\}$. Recalling (i) in the definition of \mathcal{T}_4 and the fact that $\langle c_\beta : \beta < \epsilon \rangle$ is a strictly increasing sequence below c , each m_k is well defined and

$$(\langle n_\beta : \beta < \epsilon \rangle, \langle m_k : k < \omega \rangle)$$

is a pre-cut in $X_{\mathbf{a}}$ and cannot be a cut without contradicting the definition of $\mathfrak{p}_{\mathbf{s}}$. Let $n_{**} \in X_{\mathbf{a}}$ realize this pre-cut. Let $c_\epsilon = c \upharpoonright_{n_{**}+1}$, $n_\epsilon = n_{**}$ and $d_\epsilon = c_\epsilon(n_\epsilon)$. Clearly c_ϵ satisfies (d). This completes the inductive choice of the sequence.

Once the sequence has been constructed, $\kappa < \mathfrak{t}_{\mathbf{s}}$ implies that $\langle c_\epsilon : \epsilon < \kappa \rangle$ has an upper bound c in \mathcal{T}_4 by Claim 2.14. We first choose a cut in the domain of c , as follows. Consider the sequence $\langle n_\epsilon := \text{lg}(c_\epsilon) - 1 : \epsilon < \kappa \rangle$ in $X_{\mathbf{a}}$. By Theorem 3.1 (Uniqueness) and the assumption that $(\kappa, \lambda) \in \mathcal{C}^{\text{ct}}(\mathbf{s})$, there is a sequence $\langle m_\alpha : \alpha < \lambda \rangle$ of elements of $\{n \in X_{\mathbf{a}} : n <_{\mathbf{a}} \text{lg}(c)\}$ such that $(\langle n_\epsilon : \epsilon < \kappa \rangle, \langle m_\alpha : \alpha < \lambda \rangle)$ represents a (κ, λ) -cut.

Finally, let the sequence $\langle e_\alpha : \alpha < \lambda \rangle$ in $X_{\mathbf{a}}$ given by $e_\alpha = c(m_\alpha)$. Then $\langle d_\epsilon : \epsilon < \kappa \rangle, \langle e_\alpha : \alpha < \lambda \rangle$ satisfy our requirements, by the definition of \mathcal{T}_4 . \square

Fact 8.4. *There exists a symmetric function $g : \kappa^+ \times \kappa^+ \rightarrow \kappa$ such that*

$$W \in [\kappa^+]^{\kappa^+} \text{ implies } \sup(\text{range}(g \upharpoonright W \times W)) = \kappa.$$

For instance, a symmetric function g has the desired property if whenever $\alpha < \alpha' < \beta$, $g(\alpha, \beta) \neq g(\alpha', \beta)$.

We now prove the main result of this section.

Theorem 8.1. *Let \mathbf{s} be a cofinality spectrum problem. Suppose that κ, λ are regular and $\kappa < \lambda = \mathfrak{p}_{\mathbf{s}} < \mathfrak{t}_{\mathbf{s}}$. Then $(\kappa, \lambda) \notin \mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}})$.*

Proof. We suppose for a contradiction that $\text{lcf}(\kappa) = \lambda$. Note that necessarily $\kappa + \lambda$ is minimal among $(\kappa, \lambda) \in \mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}})$ by definition of $\mathfrak{p}_{\mathbf{s}}$. Let $\mathbf{a} \in \text{Or}(\mathbf{s})$ be coverable as a pair by \mathbf{a}' in the sense of 5.7 (in this proof we will write $\mathbf{a} \times \mathbf{a}$ to refer to the \mathbf{c} of that definition). Let

$$(\langle d_\epsilon : \epsilon < \kappa \rangle, \langle e_\alpha : \alpha < \lambda \rangle)$$

be a representation of a (κ, λ) -cut in $X_{\mathbf{a}}$ given by Claim 8.2. To complete the preliminaries, fix $g : \kappa^+ \times \kappa^+ \rightarrow \kappa$ given by Fact 8.4. This is an outside function, which will help in the proof.

We now define an appropriate tree. Let $\mathbf{b} \in \text{Or}(\mathbf{s})$ be such that $X_{\mathbf{b}} = X_{\mathbf{a}} \times X_{\mathbf{a}} \times X_{\mathbf{a}} \times X_{\mathbf{a}'} \times X_{\mathbf{a}'}$. We will work in the definable subtree \mathcal{T}_6 of $\mathcal{T}_{\mathbf{b}}$ consisting of elements $x \in \mathcal{T}_{\mathbf{b}}$ such that

- (a) $n_1 <_{\mathbf{b}} n_2 <_{\mathbf{b}} \text{lg}(x)$ implies
 - $x(n_2, 0) \leq_{\mathbf{a}} x(n_2, 1) <_{\mathbf{a}} x(n_2, 2) <_{\mathbf{a}} x(n_1, 0)$;
- (b) $x(n, 3)$ is a nonempty subset of $X_{\mathbf{a}}$ and $|x(n, 3)| \leq^{\mathbf{s}} |X_{\mathbf{a}}|/2$ in the sense of 5.1, i.e., its cardinality is $\leq^{\mathbf{s}}$ that of its complement;
- (c) $x(n, 4)$ is a symmetric 2-place function with domain $x(n, 3) \times x(n, 3)$ and range $\subseteq X_{\mathbf{a}}$, which we call a distance estimate function;
- (d) $x(n, 5)$ is a one-to-one function from $x(n, 3)$ to the interval $(x(n, 1), x(n, 2))_{\mathbf{a}}$ such that

$$a \neq b \in x(n, 3) \text{ implies } |x(n, 4)(a, b)| \leq^{\mathbf{s}} |x(n, 5)(a) - x(n, 5)(b)|.$$

(Recall that for any $z \in X_{\mathbf{a}}$, the internal cardinality $|z|$ is identified with $|[0, z]_{\mathbf{a}}|$ following 8.2(2) above.)

- (e) if $n_1 <_{\mathbf{b}} n_2 <_{\mathbf{b}} \text{lg}(x)$ and a, b in $X_{\mathbf{a}}$ are such that

$$(\forall m) ((n_1 \leq_{\mathbf{b}} m \leq_{\mathbf{b}} n_2) \rightarrow \{a, b\} \subseteq x(m, 3)),$$

then $x(n_1, 4)(a, b) = x(n_2, 4)(a, b)$.

This completes the description of \mathcal{T}_6 . (Informally, given $n \leq \max(\text{dom}(x)) \in \mathcal{T}_6$, $x(n, 0)$ is a marker moving left toward the cut and $x(n, 5)$ is an injection from $x(n, 3)$ into the interval $(x(n, 1), x(n, 2))_{\mathbf{a}}$. The new point is a distance estimate function $x(n, 4)$: we ask $x(n, 5)$ to respect the internal spacing given by this function, and we ask that as n grows, the distance estimate remains unchanged on pairs of elements which remain continuously in the sequence of domains $x(n, 3)$.)

In the next, core part of the proof, we will choose $c_{\alpha} \in \mathcal{T}_6$, $n_{\alpha} = \max(\text{dom}(c_{\alpha}))$ by induction on $\alpha < \lambda$. When α is a successor we will also choose y_{α} . Our inductive hypothesis is as follows:

- For all $\alpha < \lambda$, we ensure that
 - (1) $\beta < \alpha$ implies $c_{\beta} \trianglelefteq c_{\alpha}$.
 - (2) $\beta < \alpha$ implies
 - $e_{\alpha+1} \leq_{\mathbf{a}} c_{\alpha}(n_{\alpha}, 0) <_{\mathbf{a}} c_{\alpha}(n_{\alpha}, 1) <_{\mathbf{a}} c_{\alpha}(n_{\alpha}, 2) <_{\mathbf{a}} e_{\beta+1}$,
 - and if $\alpha = \beta + 1$, then in addition $c(n_{\alpha}, 0) = e_{\alpha+1}$.
 - (3) For all $\gamma < \min\{\alpha, \kappa^+\}$,
 - (a) $y_{\gamma+1} \in c_{\alpha}(n_{\alpha}, 3)$;
 - (b) $(\forall m)[n_{\gamma+1} \leq_{\mathbf{b}} m \leq_{\mathbf{b}} n_{\alpha} \rightarrow y_{\gamma+1} \in c_{\alpha}(m, 3)]$;
 - (c) for all $\zeta + 1 < \gamma + 1$ and for all m such that $n_{\gamma+1} \leq_{\mathbf{b}} m \leq_{\mathbf{b}} n_{\alpha}$,
 - $x(m, 4)(y_{\zeta+1}, y_{\gamma+1}) = d_{g(\zeta+1, \gamma+1)}$.
- When $\alpha = \beta + 1 < \kappa^+$, then in addition we choose $y_{\alpha} = y_{\beta+1}$ so that the following are true:
 - (4) $y_{\beta+1} \in X_{\mathbf{a}} \setminus \{y_{\gamma+1} : \gamma < \beta\}$;
 - (5) $y_{\beta+1} \in c_{\alpha}(n_{\alpha}, 3)$;
 - (6) $y_{\beta+1} \notin c_{\beta}(n_{\beta}, 3)$;
 - (7) $|c_{\alpha}(n_{\alpha}, 3)| \leq^{\mathbf{s}} |X_{\mathbf{a}}|/2$;
 - (8) for all $\gamma + 1 < \beta + 1$ and all n such that $n_{\gamma+1} \leq_{\mathbf{b}} n \leq_{\mathbf{b}} n_{\alpha}$,

$$x(n, 4)(y_{\gamma+1}, y_{\beta+1}) = d_{g(\gamma+1, \beta+1)}.$$

Note that $d_{g(\beta,\gamma)}$ functions here as a constant; g , set at the beginning of the proof, is an outside function not mentioned by the definition of \mathcal{T}_6 . Condition (7) is part of the definition of \mathcal{T}_6 , repeated for clarity.

This completes the statement of the inductive hypothesis; let us now carry out the induction.

For the case $\alpha = 0$: Trivial.

For the case $\alpha = \beta + 1$, when in addition $\alpha < \kappa^+$: If $\alpha = \beta + 1 < \kappa^+$, then we first define $y_\alpha = y_{\beta+1}$. By inductive hypothesis, $M_1^+ \models c_\beta \in \mathcal{T}_6$. Thus by (b) in the definition of \mathcal{T}_6 , $X_{\mathbf{a}} \setminus c_\beta(n_\beta, 3) \neq \emptyset$. Choose $y_{\beta+1} \in X_{\mathbf{a}} \setminus c_\beta(n_\beta, 3)$. Note that we choose $c_\alpha(n_\alpha, 3)$ below and will ensure there that it remains small enough; in particular, it is irrelevant whether $c_\beta(n_\beta, 3) \cup \{y_{\beta+1}\}$ has size no larger than its complement in $X_{\mathbf{a}}$.

Then conditions (4) and (6) of the inductive hypothesis hold, so by condition (e) of the definition of \mathcal{T}_6 we will be allowed to freely choose the value of $c_\alpha(n_\alpha, 4)$ on any pair which includes $y_{\beta+1}$. Label this (**) for later reference.

Having defined $y_{\beta+1}$, continue as in the general successor step:

For the case $\alpha = \beta + 1$ for arbitrary $\alpha < \lambda$: We now assume that $y_{\beta+1}$ has been chosen for all $\beta < \min\{\alpha, \kappa^+\}$ and continue the proof assuming only $\alpha = \beta + 1 < \lambda$.

The key point at this step is to define $c_\alpha(n_\alpha, \ell)$ for $\ell < 6$. The nontrivial cases are $\ell = 3, 4, 5$. We will do this by showing it may be expressed as a consistent partial Or-type, and then applying Theorem 4.1, local saturation. Recall that we are assuming $\mathfrak{p}_{\mathbf{s}} < \mathfrak{t}_{\mathbf{s}}$.

Let $p(x_0, x_1, x_2, x_3, x_4, x_5)$ be the partial type stating the following. (Note that x_0, x_1, x_2 are determined by x_3, x_4, x_5 . The e_α are from the background cut.)

- (p.1) x_0, x_1, x_2 are elements of $X_{\mathbf{a}}$ and x_3, x_4, x_5 are elements of $X_{\mathbf{a}'}$.
- (p.2) $x_0 = e_{\alpha+1} \leq_{\mathbf{a}} x_1 <_{\mathbf{a}} x_2 \leq_{\mathbf{a}} e_\alpha$.
- (p.3) $x_3 = \text{dom}(x_4) \subseteq X_{\mathbf{a}}$ and $|x_3| \leq^{\mathbf{s}} |X_{\mathbf{a}}|/2$.
- (p.4) x_4 is a definable symmetric 2-place function from x_3 to $X_{\mathbf{a}}$.
- (p.5) x_5 is a definable injection from x_3 into the interval $(e_{\alpha+1}, e_\alpha)_{\mathbf{a}}$ such that

$$a \neq b \in x_3 \rightarrow |x_4(a, b)| \leq^{\mathbf{s}} |x_5(a) - x_5(b)|.$$

- (p.6) $x_1 = \min(\text{range}(x_5)), x_2 = \max(\text{range}(x_5))$.
- (p.7) if $a, b \in c_\beta(n_\beta, 3) \cap x_3$ then $x_4(a, b) = c_\beta(n_\beta, 4)(a, b)$.

For $\gamma < \min\{\alpha, \kappa^+\}$ we add

(p.8) $_\gamma$
$$y_{\gamma+1} \in x_3.$$

For $\zeta < \gamma < \min\{\alpha, \kappa^+\}$ we add

(p.9) $_{\zeta,\gamma}$
$$x_4(y_{\zeta+1}, y_{\gamma+1}) = d_{g(\zeta+1,\gamma+1)}.$$

(Note: (p.9) is legitimate by the observation (**) from the case “ $\alpha = \beta + 1, \alpha < \kappa^+$ ” when the pair includes $y_{\beta+1}$, and by (1) of the inductive hypothesis for all other pairs of y 's.)

To show that p is an Or-type in the sense of 4.1–4.1, note that first, p depends on the parameters $\{e_\alpha, e_{\alpha+1}, c_\beta\} \cup \{y_{\gamma+1} : \gamma < \min\{\alpha, \kappa^+\}\}$. Recall from the statement of the Lemma that $\lambda = \mathfrak{p}_{\mathbf{s}}$, and thus $|\alpha| < \mathfrak{p}_{\mathbf{s}}$. It remains to show that p is finitely satisfiable in $X_{\mathbf{b}}$ (i.e., consistent). Since we choose a domain x_3 , a distance estimate function x_4 , and a partial injection x_5 simultaneously, the true constraints come

from the schemata (p.8) and (p.9) which require that certain elements are in the domain, thus certain previously set distance estimates must be respected, and, if applicable, certain new distances set.

Let us check finite satisfiability (by compactness this will suffice). Let a nonempty finite subset $\Gamma \subseteq \alpha \cap \kappa^+$ be given. Let $p_0 \subseteq p$ be finite and such that p_0 implies (p.1)–(p.7), p_0 implies (p.8) $_\gamma$ for each $\gamma \in \Gamma$, and p_0 implies (p.9) $_{\zeta, \gamma}$ for each $\zeta, \gamma \in \Gamma$, $\zeta < \gamma$. We will prove that p_0 is satisfiable.

We define b_3, b_4, b_5, b_1, b_2 as follows:

- (i) Let $b_3 = \{y_{\gamma+1} : \gamma \in \Gamma\}$.
- (ii) Let b_4 be the symmetric 2-place function on b_3 defined by

$$(y_{\zeta+1}, y_{\gamma+1}) \mapsto d_{g(\zeta+1, \gamma+1)}$$

for $\zeta, \gamma \in \Gamma$. (Note that this function exists and may be identified with an element of $\mathcal{T}_{\mathbf{a} \times \mathbf{a}}$, thus of $X_{\mathbf{a}'}$, by the choice of \mathbf{a}, \mathbf{a}' satisfying 5.7, and the fact that finite sequences in this tree are closed under concatenating an additional element, since \mathbf{a}' is nontrivial.)

- (iii) Let $d = \max\{d_{g(\zeta+1, \gamma+1)} : \zeta \neq \gamma \in \Gamma\}$. Note that for all $\alpha < \lambda$, $d < e_\alpha$, by choice of the cuts at the beginning of the proof.
- (iv) Let $\{\gamma_0, \dots, \gamma_n\}$ enumerate Γ in increasing order, without repetition. Let b_5 be the definable partial injection given by

$$y_{\gamma_i+1} \mapsto e_{\alpha+1} +_{\mathbf{a}} 1 +_{\mathbf{a}} i \cdot d$$

for $i = 0, \dots, n$. By the choice of \bar{e} , $\max(\text{range}(b_5)) <_{\mathbf{a}} e_\alpha$. (The parenthetical remark from (ii) applies here also.)

- (v) Let b_0, b_1, b_2 be defined from b_3, b_4, b_5 by conditions (p.1), (p.2), (p.6) of the definition of p .

Let us verify that $(b_0, \dots, b_5) \models p_0$. Condition (p.1) is obvious. For Conditions (p.2) and (p.6), note that by (iv) $\max(\text{range}(b_5)) <_{\mathbf{a}} e_\alpha$. Condition (p.3) is obvious as b_3 is finite and $X_{\mathbf{a}}$ is not. Conditions (p.4), (p.5), and (p.6) are immediate from the definitions of the elements b_i . For $\zeta < \gamma \leq \beta$, Conditions (p.8) $_\gamma$ and (p.9) $_{\zeta, \gamma}$ are also immediate. What about Condition (p.7)? (Note that on our proposed finite fragment, $c_\beta(n_\beta, 3) \cap b_3 \subseteq b_3$.) By inductive hypothesis [(1), (5)], we have that

$$c_\beta(n_\beta, 3) \cap b_3 = \{y_{\gamma+1} : \gamma \in \Gamma, \gamma \neq \beta\}.$$

Thus, when $\zeta < \gamma < \beta$, Condition (p.7) for $a = y_{\zeta+1}, b = y_{\gamma+1}$ is ensured by Condition (p.9) $_{\zeta, \gamma}$, and when $\zeta < \beta$, Condition (p.7) $a = y_{\zeta+1}, b = y_{\beta+1}$ is trivially true since $y_{\beta+1} \notin c_\beta(n_\beta, 3)$.

This completes the proof that p_0 is realized, and *thus* that p is an Or-type in the sense of Definition 4.1 over a set of size $< \mathfrak{p}_s$.

By Theorem 4.1, p has a realization $\langle b_i^* : i < 6 \rangle$. By inductive hypothesis, we may concatenate. Let $n_\alpha = n_\beta + 1$ and let $c_\alpha = c_\beta \hat{\ } \langle b_0^*, \dots, b_5^* \rangle$. This completes the successor step.

For the case of α limit: Since $\text{cf}(\alpha) < \lambda \leq \mathfrak{p}_s \leq \mathfrak{t}_s$, by Lemma 2.15 there is $c \in \mathcal{T}_6$ such that $\beta < \alpha$ implies $c_\beta \preceq c$ and c is below the ceiling. The key point is to adjust this by choosing a suitable initial segment c_α of c so that Conditions (1), (2), and (3)(a) of the inductive hypothesis will be satisfied. (Conditions (3)(b) and 3(c) will then follow by definition of \mathcal{T}_6 .) First, let

$$n_* = \max\{n : n <_{\mathbf{b}} \text{lg}(c), M_1^+ \models (e_\alpha <_{\mathbf{a}} c(n, 0))\}.$$

Necessarily for all $\beta < \alpha$, $\text{lg}(c_\beta) <_{\mathbf{b}} n_*$. Second, for each $\beta < \max\{\alpha, \kappa^+\}$, let

$$n(\beta) = \max\{n \leq_{\mathbf{b}} n_* : y_{\beta+1} \in c(n, 3)\}.$$

For each $\gamma < \beta < \alpha$, inductive hypothesis (3) for β implies $y_{\gamma+1} \in c_\beta(n_\beta, 3)$. In other words,

$$(\{n_\beta : \beta < \alpha \cap \kappa^+\}, \{n(\beta) : \beta < \alpha \cap \kappa^+\})$$

is a pre-cut in $X_{\mathbf{b}}$, and thus a (κ_1, κ_2) -pre-cut for some regular $\kappa_1, \kappa_2 \leq |\alpha \cap \kappa^+|$. It cannot be a cut, as then $\kappa_1 + \kappa_2 \leq |\alpha| < \lambda = \mathfrak{p}_{\mathbf{s}}$ contradicting the definition of $\mathfrak{p}_{\mathbf{s}}$. So we may choose $n_{**} \leq_{\mathbf{b}} n_*$ such that for all $\gamma, \beta < \alpha \cap \kappa^+$,

$$n_\gamma <_{\mathbf{b}} n_{**} <_{\mathbf{b}} n_\beta.$$

Let $n_\alpha = n_{**}$, and let $c_\alpha = c \upharpoonright_{n_\alpha}$. This completes the limit case, and thus the inductive construction of the sequence $\langle c_\alpha, n_\alpha : \alpha < \lambda \rangle$.

We arrive at the final stage of the proof. Having built our sequence $\langle c_\alpha : \alpha < \kappa \rangle$, as $\lambda = \mathfrak{p}_{\mathbf{s}} < \mathfrak{t}_{\mathbf{s}}$ we may choose $c_\lambda \in \mathcal{T}_{\mathbf{6}}$ such that for all $\alpha < \lambda$, $c_\alpha \leq c_\lambda$. Let $n_\lambda = \max(\text{dom}(c_\lambda))$. By construction, $\langle n_\alpha : \alpha < \lambda \rangle$ is an increasing sequence of elements of $X_{\mathbf{b}}$ below n_λ . By Theorem 3.1 (Uniqueness), there is a sequence $\langle m_\epsilon : \epsilon < \kappa \rangle$ such that

$$(\langle n_\alpha : \alpha < \lambda \rangle, \langle m_\epsilon : \epsilon < \kappa \rangle)$$

represents a (λ, κ) -cut in $X_{\mathbf{b}}$. Without loss of generality, for some increasing $\zeta : \kappa \rightarrow \kappa$,

$$d_{\zeta(\epsilon)} <_{\mathbf{a}} c_\lambda(m_\epsilon, 0) <_{\mathbf{a}} d_{\zeta(\epsilon+1)}.$$

Now for each $\beta < \kappa^+$ and each α_1, α_2 with $\beta < \alpha_1 < \alpha_2 < \lambda$, the set

$$X_\beta := \{n : n \leq_{\mathbf{b}} n_\lambda, (\forall n') (n_{\beta+1} \leq_{\mathbf{b}} n' \leq_{\mathbf{b}} n \rightarrow y_{\beta+1} \in c_\lambda(n', 3))\}$$

is a subset of $X_{\mathbf{b}}$ which includes the interval $[n_{\alpha_1}, n_{\alpha_2}]_{\mathbf{b}}$, recalling the notation $n_\gamma = \max(\text{dom}(c_\gamma))$. Thus for some $\epsilon(\beta) < \kappa$,

$$[n_{\beta+1}, m_{\epsilon(\beta)}]_{\mathbf{b}} \subseteq X_\beta.$$

As $\beta < \kappa^+$ was arbitrary, there is $\epsilon_* < \kappa$ such that

$$W = \{\beta < \kappa^+ : \epsilon(\beta) = \epsilon_*\}$$

has cardinality κ^+ . For simplicity of notation, let $F := c_\lambda(m_{\epsilon_*}, 4)$. By Condition (d) in the definition of $\mathcal{T}_{\mathbf{6}}$ and the inductive hypothesis, for every $\beta \neq \gamma \in W$ we have that $F(y_\gamma, y_\beta) = d_{g(\gamma, \beta)}$. However, by the choice of \bar{d} , the choice of the outside function g and the fact that $|W| = \kappa^+$, there exist $\gamma, \beta \in W$ such that $M_1^+ \models |d_{\zeta(\epsilon_*)+1}| <^{\mathbf{s}} |d_{g(\gamma, \beta)}|$. This contradiction completes the proof. \square

9. $\mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}}) = \emptyset$

With the results of the previous sections in hand, we now prove the paper's fundamental result.

Central Theorem 9.1. *Let \mathbf{s} be a cofinality spectrum problem. Then $\mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}}) = \emptyset$.*

Proof. There are two cases.

Case 1. $\mathfrak{p}_{\mathbf{s}} < \mathfrak{t}_{\mathbf{s}}$. Let κ, λ be such that $\kappa + \lambda = \mathfrak{p}_{\mathbf{s}}$ and $(\kappa, \lambda) \in \mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}})$. By Claim 3.1, Conclusion 3.6, and Lemma 6.1, $\kappa \neq \lambda$, and we may assume $\kappa < \lambda = \mathfrak{p}_{\mathbf{s}}$. Then the hypotheses of Theorem 8.1 are satisfied, so $(\kappa, \lambda) \notin \mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}})$, contradiction. This shows Case 1 cannot occur.

Case 2. $\mathfrak{t}_{\mathbf{s}} \leq \mathfrak{p}_{\mathbf{s}}$, so by definition of $\mathfrak{p}_{\mathbf{s}}$, $\mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}}) = \emptyset$. This proves the theorem. \square

10. A NEW CHARACTERIZATION OF GOOD REGULAR ULTRAFILTERS

We now apply cofinality spectrum problems to study regular ultrapowers and Keisler's order. We focus here on the definitions and history most relevant to our present proofs; further details can be found in [26]. In the present section, we give the necessary background and then prove two theorems: Theorem 10.1, the analogue of Theorem 9.1 for regular ultrapowers, and Main Theorem 10.25, a new characterization of Keisler's notion of goodness.

Convention 10.1 (Conventions on ultrapowers).

- (1) For transparency we assume all languages are countable.
- (2) For transparency we assume all theories T are complete.
- (3) We use \mathcal{D} to denote a regular ultrafilter on the index set I (see 10.2), and λ to denote $|I|$.
- (4) Small means of cardinality $\leq |I|$.
- (5) We use notation of the form $a[t]$ following Convention 2.1(6).

Definition 10.2. Say that the filter \mathcal{D} on I is κ -regular when there is a collection $\bar{X} = \{X_i : i < \kappa\} \subseteq \mathcal{D}$ such that for each $t \in I$,

$$|\{i < \kappa : t \in X_i\}| < \aleph_0.$$

Such a collection is called a κ -regularizing family. Call \mathcal{D} regular when it is $|I|$ -regular.

Regularity is a kind of strong incompleteness: a κ -regularizing family is such that for any $\sigma \subseteq \kappa$, if $|\sigma| \geq \aleph_0$, then $\bigcap \{X_i : i \in \sigma\} = \emptyset$.

Definition 10.3. When \mathcal{D} is an ultrafilter on I , M a model, say that \mathcal{D} saturates M to mean M^I/\mathcal{D} is $|I|^+$ -saturated.

Fact 10.4 (Keisler [16]). If \mathcal{D} is a regular ultrafilter on I , then whenever $M \equiv N$, \mathcal{D} saturates M iff \mathcal{D} saturates N .

In particular, when \mathcal{D} is regular, the following is well defined (recalling that all languages are countable).

Definition 10.5. When \mathcal{D} is regular, say that \mathcal{D} saturates T to mean \mathcal{D} saturates M for some, equivalently every, $M \models T$.

Keisler in 1967 proposed that regular ultrafilters could be used to investigate the relative complexity of first-order theories.

Definition 10.6. (Keisler [16]) Let T_1, T_2 be complete, countable first order theories. Say that $T_1 \leq T_2$ if for any infinite cardinal λ and any regular ultrafilter \mathcal{D} on λ , if \mathcal{D} saturates T_2 , then \mathcal{D} saturates T_1 .

In other words, $T_1 \leq T_2$ if for all infinite λ , all $M_1 \models T_1, M_2 \models T_2$, and any \mathcal{D} a regular ultrafilter on λ , if M_2^λ/\mathcal{D} is λ^+ -saturated, then M_1^λ/\mathcal{D} must be λ^+ -saturated.

Determining the structure of Keisler's order is a far-reaching problem. It was studied by Keisler, then by Shelah in Chapter VI of [40], then in a series of papers by Malliaris [22], [23], [24], [25], and very recently by both authors jointly in [26], [27], [28], [29]. We refer the interested reader to the introduction of [26] for motivation and [29] for a survey of what is known. Here, we focus on the part of

the classification problem relevant to our current work (on the maximal class, and on nonsimple theories). Toward this let us give several theorems and definitions. The two fundamental theorems of ultraproducts we use are Łos' theorem and its corollary.

Theorem A. (Ultrapowers commute with reducts) *Let τ, τ' denote vocabularies. Let M be an τ' -structure, $\tau \subseteq \tau'$, \mathcal{D} an ultrafilter on $\lambda \geq \aleph_0$, $N = M^\lambda/\mathcal{D}$. Then*

$$(M^\lambda/\mathcal{D})|_\tau = (M|_\tau)^\lambda/\mathcal{D}.$$

In light of Theorem A, it is useful to consider structure on the ultrapower coming from the index models in some possibly expanded language.

Definition 10.7. (Internal, i.e., induced) *Let $N = M^\lambda/\mathcal{D}$ be an ultrapower. Say that a relation or function X on N is internal, also called induced, if we may expand the language by adding a new symbol Y of the same arity as X , and choose for each $t \in \lambda$ an interpretation Y^{M_t} of Y , so that $Y^N \equiv X \pmod{\mathcal{D}}$, where $Y^N = \{a \in {}^\lambda M : \{t < \lambda : a[t] \in Y^{M_t}\} \in \mathcal{D}\}$.*

Good ultrafilters are a useful family of ultrafilters introduced by Keisler. For any infinite λ , λ^+ -good ultrafilters on λ exist by a theorem of Kunen [20], extending a theorem of Keisler which assumed GCH.

Definition 10.8. (Good filters, Keisler [14]) *Say that the filter \mathcal{D} on I , $|I| = \lambda$ is κ^+ -good if every $f : \mathcal{P}_{\aleph_0}(\kappa) \rightarrow \mathcal{D}$ has a multiplicative refinement; i.e., there is $f' : \mathcal{P}_{\aleph_0}(\kappa) \rightarrow \mathcal{D}$ such that*

- (1) $u \in [\kappa]^{<\aleph_0}$ implies $f'(u) \subseteq f(u)$;
- (2) $u, v \in [\kappa]^{<\aleph_0}$ implies $f'(u) \cap f'(v) = f'(u \cup v)$.

In this paper we assume all good ultrafilters are \aleph_1 -incomplete, and thus regular. We say that a filter is good if it is $|I|^+$ -good.

Keisler proved that Keisler's order has a maximum class, and that this class had a set-theoretic characterization.

Theorem B. (Keisler [16]) *There is a maximum class in Keisler's order, which consists precisely of those theories T such that for $M \models T$ and \mathcal{D} a regular ultrafilter on λ , M^λ/\mathcal{D} is λ^+ -saturated iff \mathcal{D} is λ^+ -good.*

Conclusion 10.9. *Let X be any property of regular ultrafilters. Suppose it can be shown that for some countable first-order theory T and model $M \models T$, the condition " \mathcal{D} has property X " is necessary for M^I/\mathcal{D} to be $|I|^+$ -saturated. Then a consequence of Theorem B is that any (regular) good ultrafilter \mathcal{D} on I must have property X .*

The question of the model-theoretic identity of this maximum class has remained elusive. The importance of this question comes, in part, from its being an outside definition of a class of unstable theories.

The connection in Theorem B between realizing types and multiplicative refinements is based on the following definition and Fact 10.11.

Definition 10.10. *Let $N = M^I/\mathcal{D}$ be a regular ultrapower and $p = \{\psi(x; a_i) : i < \mu\}$ a ψ -type in N of cardinality $\mu \leq |I|$.*

- (1) *The Łos map $d_0 : \mathcal{P}_{\aleph_0}(\mu) \rightarrow \mathcal{D}$ is given by*

$$u \in \mathcal{P}_{\aleph_0}(\mu) \mapsto \{s \in I : M \models \exists x \bigwedge \{\psi(x; a_j) : j \in u\}\}.$$

- (2) A distribution for p is a map $d : \mathcal{P}_{\aleph_0}(p) \rightarrow \mathcal{D}$ such that
 - (a) d refines the Los map;
 - (b) the image of d is a μ -regularizing set for \mathcal{D} ;
 - (c) without loss of generality d is monotonic, i.e., $u \subseteq v$ implies $d(u) \supseteq d(v)$.

Fact 10.11 (See, e.g., [26] Section 1.2). *For a small type p in an ultrapower $N = M^I/\mathcal{D}$, the following are equivalent: (a) p is realized, (b) some distribution of p has a multiplicative refinement.*

If Keisler’s order gives a measure of complexity of theories, a first surprise was the nature of the complexity which maximality suggests.

Theorem C. (Sufficient conditions for maximality, Shelah)

- (1) Any theory with the strict order property, e.g., $(\mathbb{Q}, <)$, is maximum in Keisler’s order [40]. VI.2.

(And considerably later)

- (2) SOP_3 , a weakening of the strict order property, is sufficient for maximality [42].

We will not work directly with SOP_3 , a weaker order, in this paper; a definition can be found in Shelah and Usvyatsov [45] Fact 1.3.

Since 1996, it has been open whether the boundary of the Keisler-maximal class lies at SOP_3 or whether it could be pushed up to SOP_2 , Definition 11.1 below (or beyond). A major technical obstacle has been the lack of a framework within which to compare orders and trees. Note that there are also nontrivial model-theoretic questions about whether SOP_2 may imply SOP_3 on the level of theories, but our analysis is able to avoid this.

Discussion 10.12. (SOP_2 and SOP_3) *It is known that SOP_2 is weaker than SOP_3 on the level of formulas, but what about on the level of theories? This remains open and interesting. Our analysis ultimately circumvents this problem by showing that already SOP_2 suffices for maximality from the point of view of Keisler’s order.*

Before addressing the question of the maximality of SOP_2 in Keisler’s order, we formally connect regular ultrapowers to CSPs. Namely, we prove that the results on cofinality spectrum problems developed above may be applied to the study of cuts in regular ultrapowers of linear order. First, we specialize the definitions of treetops 2.10, lower cofinality 3.2, and $\mathcal{C}(\mathbf{s}, \mathbf{t}_{\mathbf{s}})$ 2.9) to the context of regular ultrapowers. (Generally, what bounds results in this section is not the size of the index set but that of the regularizing family.) As \mathcal{D} is regular, the saturation of M will not matter.

Definition 10.13. (Treetops) *Let \mathcal{D} be an ultrafilter on I and κ a regular cardinal.*

- (1) We say that \mathcal{D} has κ -treetops when for any κ -saturated model M which interprets a tree $(\mathcal{T}_M, \trianglelefteq)$, $N = M^I/\mathcal{D}$, $\gamma = \text{cf}(\gamma) < \kappa$, and for any \trianglelefteq -increasing sequence $\langle a_i : i < \gamma \rangle$ in $(\mathcal{T}_N, \trianglelefteq_N)$ there is $a \in \mathcal{T}_N$ such that $i < \kappa$ implies $a_i \trianglelefteq a$.
- (2) We say that \mathcal{D} has $< \kappa$ -treetops if \mathcal{D} has θ -treetops whenever $\theta = \text{cf}(\theta) < \kappa$.
- (3) We say that \mathcal{D} has $\leq \kappa$ -treetops if \mathcal{D} has $(< \kappa^+)$ -treetops.

Our main case is λ^+ -treetops where \mathcal{D} is a regular ultrafilter on I , $|I| = \lambda$.

Definition 10.14. For \mathcal{D} an ultrafilter on I we define

$$\mathcal{C}(\mathcal{D}) = \left\{ (\kappa_1, \kappa_2) \in (\text{Reg} \cap |I|^+) \times (\text{Reg} \cap |I|^+) : (\omega, <)^I / \mathcal{D} \text{ has a } (\kappa_1, \kappa_2)\text{-cut} \right\}.$$

The paper’s central question from the Introduction thus becomes the following.

Question 10.15. Let \mathcal{D} be a regular ultrafilter on I with $|I|^+$ -treetops. What are the possible values of $\mathcal{C}(\mathcal{D})$?

Corollary 10.16. (Of the maximality of strict order, Theorem C above) Let \mathcal{D} be a regular ultrafilter on I , $|I| = \lambda$. Then $\mathcal{C}(\mathcal{D}) = \emptyset$ iff \mathcal{D} is λ^+ -good.

Claim 10.17. Let \mathcal{D} be a regular ultrafilter on I , $|I| = \lambda$. For any $n < \omega$, write $<_n$ for the order on ω restricted to n , i.e., to $\{0, \dots, n - 1\}$. Then there exists a sequence $\bar{n} = \bar{n}(\mathcal{D}) = \langle n_t : t \in I \rangle \in {}^I \omega$ such that for all regular cardinals κ_1, κ_2 with $\kappa_1 + \kappa_2 \leq \lambda$, the following are equivalent:

- (1) $(\kappa_1, \kappa_2) \in \mathcal{C}(\mathcal{D})$, i.e., $(\omega, <)^I / \mathcal{D}$ has a (κ_1, κ_2) -cut.
- (2) $\prod_t (n_t, <_{n_t}) / \mathcal{D}$ has a (κ_1, κ_2) -cut.

Proof. Note that it suffices to show (1) \rightarrow (2).

Without loss of generality, consider $M = (\omega, <)^I / \mathcal{D}$ and $M_1 = M^I / \mathcal{D}$. Let $\{X_i : i < \lambda\}$ be a regularizing family, 10.2 above. For $t \in I$, let $n_t = |\{i < \lambda : t \in X_i\}| + 1$. We verify that $\langle n_t : t \in I \rangle$ works. Let $(\langle a_\alpha : \alpha < \kappa_1 \rangle, \langle b_\beta : \beta < \kappa_2 \rangle)$ represent a (κ_1, κ_2) -cut in $(\omega, <)^\lambda / \mathcal{D}$. As $\kappa_1 + \kappa_2 \leq \lambda$, there is a map $d : \kappa_1 \times \{0\} \cup \kappa_2 \times \{1\} \rightarrow \mathcal{D}$ such that for each $t \in I$, $|\{x \in \text{dom}(d) : t \in d(x)\}| < n_t$. For each $t \in I$, let $X_t = \{a_\alpha[t] : t \in d((\alpha, 0))\} \cup \{b_\beta[t] : t \in d((\beta, 1))\}$, which is a (linearly ordered) subset of $(\omega, <)^M$ with fewer than n_t elements. Let $<_{X_t}$ denote the restriction of the linear order on ω to X_t . Then we may choose at each index t an order preserving injection $h_t : (X_t, <_{X_t}) \rightarrow (n_t, <_{n_t})$ whose image is an interval. Let h be the internal function whose projection to t is h_t . Then by Los’ theorem and the requirement that the range be an interval, we have that $(\langle h(a_\alpha) : \alpha < \kappa_1 \rangle, \langle h(b_\beta) : \beta < \kappa_2 \rangle)$ represents a (κ_1, κ_2) -cut in $\prod_t (n_t, <_{n_t}) / \mathcal{D}$. This completes the proof. \square

Definition 10.18. Let \mathcal{D} be a regular ultrafilter on I , M a model extending $(\omega, <)$. If $\langle n_t : t \in I \rangle \in {}^I \omega$ is a sequence satisfying the conclusion of 10.17 for \mathcal{D} and $(X, <_X) \subseteq M^I / \mathcal{D}$ is given by

$$(X, <_X) = \prod_t (n_t, <_{n_t}) / \mathcal{D},$$

then we say $(X, <_X)$ captures pseudo-finite cuts. [Clearly, this depends on the background data of I, \mathcal{D}, M .]

The next Claims 10.19, 10.21, and 10.22 justify regarding regular ultrapowers extending the theory of linear order [see Claim 10.21] as cofinality spectrum problems, and show that the specialized definitions for $\mathcal{C}(\mathcal{D})$ and treetops accurately reflect the properties of this background CSP.

Claim 10.19. (Regular ultrapowers as CSPs) Let \mathcal{D} be a regular ultrafilter on I , $|I| = \lambda$. Let M expand $(\omega, <)$ and let $M_1 = M^I / \mathcal{D}$.

Then there exist expansions M^+, M_1^+ of M, M_1 , respectively, such that $M_1^+ = (M^+)^I / \mathcal{D}$ and a set of formulas $\Delta \supseteq \{x < y < z\}$ of the language of M such that

- (1) $\mathbf{s} = (M, M_1, M^+, M_1^+, \text{Th}(M^+), \Delta)$ is a cofinality spectrum problem, and
- (2) some nontrivial $\mathbf{a} \in \text{Or}(\mathbf{s})$ captures pseudo-finite cuts in the sense of 10.18.

Proof. As ultrapowers commute with reducts, for (1) choose any expansion M^+ of M which will code sufficient set theory for trees in the sense of 2.3, e.g., the complete expansion, or an expansion to a model of $(\mathcal{H}(\chi), \in)$ for some sufficiently large χ . Let $M_1^+ = (M^+)^I/\mathcal{D}$. For (2), let $\langle n_t : t \in I \rangle$ be given by 10.17. By construction, the linear order $\prod_t(n_t, <_{n_t})/\mathcal{D}$ is Δ -definable in M_1 and captures pseudo-finite cuts. It will correspond to some nontrivial $\mathbf{a} \in \text{Or}(\mathbf{s})$ provided we choose $d_{\mathbf{a}}$ to not be a natural number. \square

Definition 10.20. *Call any \mathbf{s} satisfying the conclusion of 10.19 a cofinality spectrum problem associated to \mathcal{D} .*

In the next series of claims we verify that cuts and trees behave as expected.

Claim 10.21. *Let $\mathcal{D}, I, \lambda, M, M_1$ be as in 10.19 and let \mathbf{s} be a cofinality spectrum problem given by that Claim. For $\kappa_1 + \kappa_2 \leq \lambda$, κ_1, κ_2 regular, the following are equivalent:*

- (1) *There is a (κ_1, κ_2) -cut in some M_1^+ -definable linearly ordered set.*
- (2) *$(\kappa_1, \kappa_2) \in \mathcal{C}(\mathbf{s}, |I|^+)$.*
- (3) *$(\kappa_1, \kappa_2) \in \mathcal{C}(\mathcal{D})$.*

Proof. (3) \rightarrow (1): By Claim 10.19(2).

(1) \rightarrow (2): It suffices to show that

Subclaim. For any definable linear order $(Y, <_Y)$ in M_1^+ [i.e., both Y and $<_Y$ are M_1^+ -definable but this order is not necessarily in $\text{Or}(\mathbf{s})$] and any discrete $A \subseteq Y$, $|A| \leq \lambda$ (e.g., a representation of a cut) there exist a nontrivial $\mathbf{a} \in \text{Or}(\mathbf{s})$ and an internal partial injection f such that

- (a) $A \subseteq \text{dom}(f)$, $\text{range}(f) \subseteq X_{\mathbf{a}}$;
- (b) f is order preserving, i.e. for all $a, b \in \text{dom}(f)$, $a \leq_Y b$ iff $f(a) \leq_{\mathbf{a}} f(b)$;
- (c) $\text{range}(f)$ is an interval in $(X_{\mathbf{a}}, \leq_{\mathbf{a}})$.

The proof is almost exactly the same as that of 10.17, using $(Y, <_Y)$ here instead of the representation of the cut there, and letting $\mathbf{a} \in \text{Or}(\mathbf{s})$ be given by 10.19(2). (The point: by regularity, any discrete linearly ordered set $|A| \leq \lambda$ in the ultrapower may be considered as a subset of some internal, pseudo-finite linear order.)

(2) \rightarrow (3): Assume (2), so there are regular cardinals κ_1, κ_2 with $\kappa_1 + \kappa_2 \leq \lambda$ and some nontrivial $\mathbf{b} \in \text{Or}(\mathbf{s})$ such that $X_{\mathbf{b}}$ contains a (κ_1, κ_2) -cut. To conclude that $(\kappa_1, \kappa_2) \in \mathcal{C}(\mathcal{D})$, let \mathbf{a} be given by 10.19(2). By Observation 3.5 (“any $\mathbf{a} \in \text{Or}(\mathbf{s})$ will work”), also $(X_{\mathbf{a}}, <_{\mathbf{a}})$ has a (κ_1, κ_2) -cut, and thus also $(\omega, <)^I/\mathcal{D}$. \square

Claim 10.22. *Let $\mathcal{D}, I, \lambda, M, M_1$ be as in 10.19 and let \mathbf{s} be a cofinality spectrum problem given by that Claim. For $\kappa = \text{cf}(\kappa) \leq \lambda$, then the following are equivalent:*

- (1) *\mathcal{D} has κ^+ -treetops in the sense of 10.13.*
- (2) *$\kappa^+ \leq \mathfrak{t}_{\mathbf{s}}$.*

Proof. Clearly (1) \rightarrow (2).

To show (2) \rightarrow (1), we prove the contrapositive. That is, we show that

Subclaim. If $(\mathcal{T}, \leq_{\mathcal{T}})$ is any tree definable in M^+ , not necessarily an element of $\text{Tr}(\mathbf{s})$, and there is in $(\mathcal{T}, \leq_{\mathcal{T}})^I/\mathcal{D} = (\mathcal{T}, \leq_{\mathcal{T}})^{M_1^+}$ an increasing sequence of length $\kappa = \text{cf}(\kappa) \leq \lambda$ with no upper bound, then there is $\mathbf{a} \in \text{Or}(\mathbf{s})$ such that in $\mathcal{T}_{\mathbf{a}}$ there is an increasing sequence of length κ with no upper bound.

(While \mathbf{a} nontrivial will be guaranteed by choosing \mathbf{a} from 10.19(2), it follows from $\mathcal{T}_{\mathbf{a}}$ having arbitrarily long paths.)

So let such a $(\mathcal{T}, \preceq_{\mathcal{T}})$ be given. Let $\bar{c} = \langle c_{\alpha} : \alpha < \kappa \rangle$ be an increasing sequence in \mathcal{T} with no upper bound. By regularity, as $\kappa \leq \lambda$ there is a map $d : \kappa \rightarrow \mathcal{D}$ whose image is a regularizing family. In other words, by Los' theorem, we may assume that there is a sequence of finite trees $(\mathcal{T}_t, \preceq_{\mathcal{T}}^t)$ for $t \in I$ such that

$$\prod_t (\mathcal{T}_t, \preceq_{\mathcal{T}}^t) / \mathcal{D}$$

is a subtree of $(\mathcal{T}, \preceq_{\mathcal{T}})^{M_1^+}$ which includes the sequence \bar{c} . Let $\mathbf{a} \in \text{Or}(\mathbf{s})$ be given by 10.19(2).

Analogously to 10.17, we may choose at every (or almost every) index $t \in I$ a function $f_t : (\mathcal{T}_t, \preceq_{\mathcal{T}}^t) \rightarrow (\mathcal{T}_{\mathbf{a}}^{M_1^+}, \preceq_{\mathbf{a}}^{M_1^+})$ such that f_t is injective and respects the partial ordering; i.e., for $x, y \in \text{dom}(f_t)$ we have that $x \preceq_{\mathcal{T}}^t y$ iff $f_t(x) \preceq_{\mathbf{a}}^{M_1^+} f_t(y)$. Now let $f = \prod_t f_t / \mathcal{D}$ and suppose for a contradiction that $\langle b_{\alpha} := f(c_{\alpha}) : \alpha < \kappa \rangle$ has an upper bound in $\mathcal{T}_{\mathbf{a}}$; call it b_* . Consider the map $d' : \kappa \rightarrow \mathcal{D}$ given by $\alpha \mapsto d(\alpha) \cap \{t \in I : b_{\alpha}[t] \preceq b_*[t]\} \cap \{t \in I : f_t \text{ is injective and respects the partial ordering}\}$. Now for each $t \in I$, the set $B_t := \{b_{\alpha}[t] : \alpha < \kappa \wedge t \in d'(\alpha)\}$ is linearly ordered by \preceq , by the choice of b_* . For each $t \in I$, let b_t be the maximal element of B_t under this linear ordering. Then by Los' theorem and the choice of the f_t 's, we have that the element $c_* := \prod_t f_t^{-1}(b_t) / \mathcal{D}$ is well defined. By Los' theorem (recalling that \mathcal{T} is definable) $c_* \in \mathcal{T}$, and again by Los' theorem c_* is an upper bound for the sequence \bar{c} in \mathcal{T} , contradiction. We have shown that $\langle b_{\alpha} : \alpha < \kappa \rangle$ is an increasing sequence in $\mathcal{T}_{\mathbf{a}}$ with no upper bound, which completes the proof. \square

Conclusion 10.23. *Regular ultrapowers extending the theory of linear order may be regarded as cofinality spectrum problems, for which the specialized definitions $\mathcal{C}(\mathcal{D})$, $|I|^+$ -treetops retain their intended meaning as stated in 10.21 and 10.22.*

Proof. By 10.19, 10.21, 10.22. \square

We may now give the analogue of Theorem 9.1 for regular ultrapowers.

Theorem 10.1. *Let \mathcal{D} be a regular ultrafilter on I , $|I| = \lambda \geq \aleph_0$. If \mathcal{D} has λ^+ -treetops, then \mathcal{D} is λ^+ -good.*

Proof. Let \mathbf{s} be a cofinality spectrum problem associated to \mathcal{D} , given by 10.19. By Theorem 9.1, $\mathcal{C}(\mathbf{s}, \mathfrak{t}_{\mathbf{s}}) = \emptyset$. By Claim 10.22 and the assumption of λ^+ -treetops, $\lambda^+ \leq \mathfrak{t}_{\mathbf{s}}$, so necessarily $\mathcal{C}(\mathbf{s}, |I|^+) = \emptyset$. Apply Claim 10.21 to conclude $\mathcal{C}(\mathcal{D}) = \emptyset$. Then by Corollary 10.16 (or Fact 1.3) \mathcal{D} is λ^+ -good, which completes the proof. \square

Moreover, the result from Section 6 may here be strengthened to a characterization.

Lemma 10.24. *Let \mathcal{D} be a regular ultrafilter on I , $|I| = \lambda$. Then the following are equivalent:*

- (1) $\kappa = \text{cf}(\kappa) \leq \lambda = |I|$ implies $(\kappa, \kappa) \notin \mathcal{C}(\mathcal{D})$.
- (2) \mathcal{D} has $|I|^+$ -treetops.

Proof. (2) \rightarrow (1): Lemma 2.2.

(1) \rightarrow (2): We prove the contrapositive. Suppose that for some regular $\kappa \leq |I|$ and some model M (in a countable signature), M interprets, or without loss of

generality, defines a tree $(\mathcal{T}, \leq_{\mathcal{T}})$ whose \mathcal{D} -ultrapower contains a path of length κ with no upper bound. Also without loss of generality, M expands $(\omega, <)$; since ultrapowers commute with reducts, there is no harm in adding this order under a disjoint unary predicate, or as a separate sort.

Let \mathbf{s} be a cofinality spectrum problem given by Claim 10.19. By Claim 10.22, there is $\mathcal{T}_{\mathbf{a}} \in \text{Tr}(\mathbf{s})$ which contains a path of length κ with no upper bound. By Lemma 6.2, there is a definable (i.e., definable in M_1^+) linear order which has a (κ, κ) -cut. Now apply Claim 10.21 to conclude that $(\kappa, \kappa) \in \mathcal{C}(\mathcal{D})$, which completes the proof. \square

Thus we obtain the following new characterization of Keisler’s notion of goodness.

Main Theorem 10.25. *Let \mathcal{D} be a regular ultrafilter on I , $|I| = \lambda$. Then the following are equivalent:*

- (1) \mathcal{D} is λ^+ -good.
- (2) \mathcal{D} has λ^+ -treetops.
- (3) $\mathcal{C}(\mathcal{D})$ contains no symmetric cuts.
- (4) $\mathcal{C}(\mathcal{D}) = \emptyset$.

Proof. (2) \leftrightarrow (3): Lemma 10.24.

(4) \rightarrow (3): Immediate.

(2) \rightarrow (1): Theorem 10.1.

(1) \leftrightarrow (4): Fact 1.3. \square

Remark 10.26. *By a theorem of Malliaris and Shelah, to the equivalent conditions of Theorem 10.25 we may add: \mathcal{D} is λ^+ -excellent; see [28] Theorem 5.2.*

11. SOP_2 IMPLIES MAXIMALITY IN KEISLER’S ORDER

In this section, we first show that for a regular ultrafilter \mathcal{D} on I , \mathcal{D} has λ^+ -treetops precisely when \mathcal{D} -ultrapowers realize SOP_2 -types, as defined in 11.2(3). We then prove Main Theorem 11.11, showing that any theory with SOP_2 is maximal in Keisler’s order, which solves one of the two open problems mentioned in the introduction.

Definition 11.1. (Shelah [39]) *The theory T has SOP_2 if there is a formula $\psi(x; \bar{y})$ such that in some model $M \models T$ there exist $\langle \bar{a}_\eta : \eta \in {}^{\kappa >} \mu \rangle$, called an SOP_2 tree for ψ , such that*

- (1) for $\eta, \rho \in {}^{\kappa >} \mu$ incomparable, i.e., $\neg(\eta \leq \rho) \wedge \neg(\rho \leq \eta)$, we have that $\{\psi(x; \bar{a}_\eta), \psi(x; \bar{a}_\rho)\}$ is inconsistent.
- (2) for $\eta \in {}^{\kappa >} \mu$, $\{\psi(x; \bar{a}_{\eta|_i}) : i < \kappa\}$ is a consistent partial ψ -type.

By compactness, clearly we can replace ${}^{\kappa >} \mu$ by ${}^{\omega >} 2$ in the definition.

Definition 11.2 (Definitions and conventions on SOP_2). *In this section,*

- (1) *The formula $\psi = \psi(x; y)$ will denote a formula with SOP_2 , and $\ell(x), \ell(y)$ need not be 1.*
- (2) *If M^I/\mathcal{D} is a regular ultrapower, by “ SOP_2 -type” or “ SOP_2 - κ -type” we will mean a partial type $p(x) = \{\psi(x; a_\ell) : \ell < \kappa\}$ almost all of whose projections to the index model come from an SOP_2 -tree for ψ .*
- (3) *Say that \mathcal{D} realizes all SOP_2 -types to mean that all SOP_2 - $|I|$ -types are realized in all ultrapowers M^I/\mathcal{D} .*

- (4) All SOP_2 -types considered will be SOP_2 - μ -types for $\mu \leq |I|$. (A regular ultrapower of a nonsimple theory will fail to be $|I|^{++}$ -saturated by prior work of the authors [26].)
- (5) As ultrapowers commute with reducts, we will freely assume that the theory in question has enough set theory for trees, Definition 2.3.

Note that saying \mathcal{D} realizes all SOP_2 -types in the sense of Definition 11.2 certainly need not imply (a priori) that for any φ with SOP_2 , all φ -types are realized in \mathcal{D} -ultrapowers. By the usual coding tricks one may take the disjoint union of a formula with SOP_2 and one with, e.g., SOP ; such a formula will necessarily be maximal. Rather, Definition 11.2 captures the essential structure in the sense that any \mathcal{D} which is able to realize all ψ -types for *some* formula ψ with SOP_2 will necessarily realize all SOP_2 -types in the sense of Definition 11.2(5). That said, it will follow a posteriori from Theorems 9.1 and 11.9 that realizing this minimal set of SOP_2 types is, indeed, strong enough to guarantee λ^+ -saturation in general.

Corollary 11.3. *The type p is an SOP_2 -type in M^λ/\mathcal{D} iff we may add a predicate P of arity $\ell(y)$ to the vocabulary τ , and for each $i \in I$ define M_i as the index model M expanded to a model of $\tau \cup \{P\}$ in which P names an SOP_2 -tree, so that in the ultraproduct $N = \prod_i M_i/\mathcal{D}$ we have that p is a type whose parameters come from P^N .*

Proof. By Los' theorem and Definition 11.2. □

Given instances $\psi(x; a_i), \psi(x; a_j)$ belonging to some consistent partial SOP_2 -type, and some index $s \in I$, we may write $a_i[s] \trianglelefteq a_j[s]$ to indicate comparability in the chosen SOP_2 -tree at index s , and likewise $a_i \trianglelefteq a_j$ to indicate comparability in the SOP_2 -tree induced on N by P . Since ultrapowers commute with reducts, we may consider \trianglelefteq as an additional relation in some expansion of the language.

We may thus consider SOP_2 -types as arising in the following canonical way.

Definition 11.4. (The canonical presentation)

- (1) Let T'_{SOP_2} be the universal first-order theory in the vocabulary $\{P, Q, \trianglelefteq, R\}$ such that $M \models T'_{SOP_2}$ if
 - (a) M is the disjoint union of P^M, Q^M ;
 - (b) $R^M \subseteq Q^M \times P^M$;
 - (c) $\trianglelefteq^M \subseteq P^M \times P^M$;
 - (d) (P^M, \trianglelefteq) is a tree;
 - (e) if $a_1 \neq a_2 \in P^M, \neg(a_1 \trianglelefteq^M a_2) \wedge \neg(a_2 \trianglelefteq^M a_1)$, then $M \models \neg(\exists x)(xRa_1 \wedge xRa_2)$.
- (2) Let T_{SOP_2} be the model completion of T'_{SOP_2} .
- (3) We say that the regular ultrafilter \mathcal{D} on I

$$(\lambda^+, Q) - \text{saturates } T_{SOP_2}$$

if whenever $M \models T_{SOP_2}$ we have that M^I/\mathcal{D} realizes all 1-types $q(x)$ such that $|q(x)| \leq \lambda$ and $Q(x) \in q(x)$.

Remark 11.5. *Since ultrapowers commute with reducts, Section 10 Theorem A, for any regular ultrafilter \mathcal{D} on I and any model M whose theory has SOP_2 , we clearly have that $N = M^I/\mathcal{D}$ realizes all SOP_2 - μ -types if and only if $\mathcal{D}(\mu^+, Q)$ -saturates T_{SOP_2} . In what follows, we will use these two presentations interchangeably.*

Note that the goal of Definition 11.4 is simply to standardize the presentation of SOP_2 -types, which are focused on the single formula ψ (in the case of that definition, xRy); in particular, it makes no claim to have constructed a minimally complex SOP_2 theory from any point of view other than that of capturing the necessary xRy -types. Recall the definition of distribution, 10.10 above.

Having set the stage, we state a simple criterion for a regular ultrafilter \mathcal{D} to realize SOP_2 , in terms of upper bounds for increasing sequences in trees.

Lemma 11.6. (*SOP₂-types and treetops*) *Let $|I| = \lambda$. Let*

$$\mathbf{P} = \{p : p = \{\psi(x; a_i) : i < \lambda\} \text{ is an } SOP_2\text{-type in } N = M^I/\mathcal{D}, |p| \leq |I|\}.$$

Then the following are equivalent:

- (1) *Each $p \in \mathbf{P}$ is realized in N .*
- (2) *Each $p \in \mathbf{P}$ has a distribution d such that for \mathcal{D} -almost all s , for all $i, j < \lambda$,*

$$s \in d(i) \cap d(j) \text{ implies } (a_i[s] \trianglelefteq a_j[s]) \vee (a_j[s] \trianglelefteq a_i[s])$$

- (3) *If $(\mathcal{T}, \trianglelefteq_{\mathcal{T}})$ is any tree and $\langle c_i : i < \lambda \rangle$ is a $\trianglelefteq_{\mathcal{T}}$ -increasing sequence in $N_{\mathcal{T}} := (\mathcal{T}, \trianglelefteq_{\mathcal{T}})^I/\mathcal{D}$, then $\langle c_i : i < \lambda \rangle$ has an upper bound in $N_{\mathcal{T}}$. That is, there exists $c \in N_{\mathcal{T}}$ such that $i < \lambda$ implies $c_i \trianglelefteq_{\mathcal{T}} c$.*

Proof. (1) \rightarrow (2) Let p be given, let $\alpha \in N$ be a realization of p , and let $\langle X_i : i < \lambda \rangle$ be a λ -regularizing family for \mathcal{D} . Then the distribution $d : \mathcal{P}_{\aleph_0}(\lambda) \rightarrow \mathcal{D}$ given by

- $\{i\} \mapsto \{s \in I : M \models \psi(\alpha[s], a_i(s))\} \cap X_i$,
- for $|u| > 1, u \mapsto \bigcap \{d(\{i\}) : i \in u\}$

satisfies the criterion (2) by definition of SOP_2 .

(2) \rightarrow (1) For any given $p \in \mathbf{P}$, if (2) holds, then it is easy to define a realization $\alpha[s]$ in each index model by definition of SOP_2 , and any $\alpha \in N$ with $\alpha = \prod_s \alpha[s] \text{ mod } \mathcal{D}$ will realize the type by Los' theorem.

(3) \rightarrow (2) Let (P, \trianglelefteq) be the (infinite) SOP_2 -tree in $M \models T_{SOP_2}$. Let $p \in \mathbf{P}$ be given, where $p = \{\psi(x; a_i) : i < \mu\}$. Then the sequence $\langle a_i : i < \lambda \rangle$ is \trianglelefteq -increasing and thus has an upper bound c in M^I/\mathcal{D} . Then the distribution d given by

$$\{i\} \mapsto \{s \in I : a_i[s] \trianglelefteq c[s]\}$$

and for $|u| > 1, u \mapsto \bigcap \{d(\{i\}) : i \in u\}$, satisfies (2) by definition of SOP_2 : any consistent set of instances must lie along a branch.

(2) \rightarrow (3) By compactness, we may suppose that (P, \trianglelefteq) contains an ω -branching tree of height ω . We would like to realize the type $\{x > c_i : i < \lambda\}$ in $(\mathcal{T}, \trianglelefteq_{\mathcal{T}})^I/\mathcal{D}$. Fix some distribution $d_{\mathcal{T}}$ of this type. Then $d_{\mathcal{T}}$ assigns finitely many formulas to each index model, and we may build a corresponding SOP_2 -type $p = \{xRa_i : i < \lambda\}$ by copying the patterns at each index: for each $s \in I$ let $a_i[s] \trianglelefteq a_j[s]$ if and only if $c_i[s] \trianglelefteq_{\mathcal{T}} c_j[s]$, and then choose each $a_i \in M^I$ so that $a_i = \prod_{s \in I} a_i[s] \text{ mod } \mathcal{D}$. Since $\{c_i : i < \lambda\}$ is $\trianglelefteq_{\mathcal{T}}$ -increasing, by Los' theorem p will be a consistent SOP_2 -type, so it will, by assumption, have a distribution d satisfying (2). Then d naturally refines $d_{\mathcal{T}}$ and gives a distribution in which for each $s \in I$, the set $C[s] := \{c_i[s] : i < \lambda, s \in d(\{i\})\}$ is finite and $\trianglelefteq_{\mathcal{T}}$ -linearly ordered in $(\mathcal{T}, \trianglelefteq_{\mathcal{T}})$. Choose $c \in (\mathcal{T}, \trianglelefteq_{\mathcal{T}})^I$ so that $c[s]$ is the $<$ -maximum element of $C[s]$ in each index model, and c/\mathcal{D} will be an upper bound by Los' theorem. □

Remark 11.7. *In Lemma 11.6(2) \rightarrow (3), it is SOP_2 rather than simply the tree property which is used.*

On the level of theories, tree-tops therefore give a necessary condition for saturation.

Corollary 11.8. *Let \mathcal{D} be a regular ultrafilter on λ and suppose that \mathcal{D} saturates some theory with SOP_2 . Then \mathcal{D} has λ^+ -tree-tops.*

Proof. This simply translates Lemma 11.6 via Remark 11.5. □

Conclusion 11.9. *Let \mathcal{D} be a regular ultrafilter on I , $|I| = \lambda$. Then recalling Definition 11.2, the following are equivalent:*

- (1) \mathcal{D} has λ^+ -tree-tops.
- (2) $\mathcal{C}(\mathcal{D})$ contains no symmetric cuts.
- (3) \mathcal{D} realizes all SOP_2 -types over sets of size λ .

Proof. (1) \iff (2): Lemma 10.24.

(1) \iff (3): Lemma 11.6. □

Remark 11.10. *Clearly in 11.9, i.e., in its constituent claims, one could separate the role of λ from the size of the index set, using $\mathcal{C}^{ct}(\mathfrak{s})$ instead of $\mathcal{C}(\mathcal{D})$.*

This yields the following.

Main Theorem 11.11. *Every theory with SOP_2 is maximal in Keisler’s order.*

Proof. By Keisler’s characterization, Section 10 Theorem B, it suffices to show that if \mathcal{D} is a regular ultrafilter on I , $|I| = \lambda$ and $M \models T$ then M^I/\mathcal{D} is λ^+ -saturated only if \mathcal{D} is λ^+ -good.

By Conclusion 11.9, a necessary condition for any regular ultrafilter \mathcal{D} on λ to saturate T is that \mathcal{D} have λ^+ -tree-tops. By Theorem 10.1, any regular ultrafilter \mathcal{D} on λ with λ^+ -tree-tops must be λ^+ -good. So a necessary condition for \mathcal{D} to saturate T is that \mathcal{D} be good, which completes the proof. □

Discussion 11.12. To conclude this section, we review some evidence for Conjecture 1.2 from the Introduction, which says that SOP_2 characterizes maximality in Keisler’s order. Any nonsimple theory contains a minimally inconsistent tree, called TP_2 , a maximally inconsistent tree, called TP_1 , or both ([40] Theorem III.7.11). TP_1 may be identified with SOP_2 , the lowest level of the so-called SOP_n hierarchy of properties whose complexity, in some sense, approaches that of linear order as n grows. (Considered as a property of formulas, SOP_2 is much weaker than SOP_3 ; it is open whether they coincide for first order complete T .) Briefly, then, the move from SOP_3 to SOP_2 moves Keisler’s order out of the territory of the SOP_n hierarchy onto what appears to be a major dividing line for which there are strong general indications of a theory. $NSOP_2$ (= not SOP_2) is in some senses close to simplicity; we hope to develop this theory in light of our work here, leveraging the tool of Keisler’s order. $NSOP_2$ and Conjecture 1.2 also connect to work of Džamonja-Shelah [9] and Shelah-Usvyatsov [45] on a weaker, related ordering; there it was shown, for instance, that under GCH $NSOP_2$ is necessarily nonmaximal in that ordering ([45] 3.15(2)) thereby strengthening the case for Conjecture 1.2.

Finally, there is a key analogy in this case between the independence/strict order dichotomy for nonstable theories and the TP_2/SOP_2 dichotomy for nonsimple theories. There is a Keisler-minimum unstable theory, the random graph, and as already noted strict order implies maximality. By a theorem of Malliaris [24], quoted below in Section 13 as Theorem F, there is a Keisler-minimum theory among the

theories with TP_2 . In Section 13 below, we apply Theorem 11.11 to prove that this theory is indeed a minimum nonsimple theory in Keisler’s order.

12. IF \mathcal{D} IS GOOD FOR SOME NONSIMPLE THEORY, THEN \mathcal{D} IS FLEXIBLE

In this section, we connect treetops to several key model-theoretically meaningful properties of ultrafilters. The main result is that any ultrafilter which is good for some nonsimple theory must be flexible (defined below), Conclusion 12.16. Also, with an eye to Conjecture 1.2, we further develop the picture of internal maps between sequences from Corollary 3.7.

It is known that Keisler classes other than the maximal class may be characterized by properties of filters; for example, see the following.

Definition 12.1. *Let \mathcal{D} be a regular ultrafilter on I , $|I| = \lambda$. Say that \mathcal{D} has 2-separation if whenever $N = M^I/\mathcal{D}$ is infinite and $A, B \subseteq N$ are disjoint sets of size $\leq \lambda$, there is an internal predicate P which contains A and is disjoint from B .*

Fact 12.2. (The Keisler class of the random graph; see, e.g., [26]) *There is a minimum class in Keisler’s order among the unstable theories, which includes the theory of the random graph. It can be characterized set-theoretically as the class of countable complete theories which are saturated precisely by regular ultrafilters with 2-separation.*

Corollary 12.3. *If \mathcal{D} is a regular ultrafilter on λ which has λ^+ -treetops, then \mathcal{D} has 2-separation.*

Proof. By Conclusion 10.9 and Theorem 10.1. □

The following definition is standard for ultrapowers and coincides with the definition for CSPs when \mathcal{D} has treetops.

Definition 12.4. (Lower cofinality) *Let \mathcal{D} be a regular ultrafilter on I and let $\kappa \leq \lambda = |I|$ be a regular cardinal. Let $N = (\lambda, <)^I/\mathcal{D}$. The lower cofinality of κ modulo \mathcal{D} , written $\text{lcf}(\kappa, \mathcal{D})$, is the cofinality of the set $\{a \in N : \zeta \in \kappa \text{ implies } N \models a > \zeta\}$ considered with the reverse order type. In other words, it is the smallest regular cardinal ρ so that there is a (κ, ρ) -cut in N half of which is given by the diagonal embedding of κ . This is also called the coinitiality of κ with respect to \mathcal{D} .*

Theorem 3.1 (Uniqueness) need not hold for regular ultrapowers without the assumption of treetops: the following theorem gives a family of examples where it will fail.

Theorem D. (Shelah [41] Theorem VI.3.12 p. 357) *Suppose $\aleph_0 = \lambda_0 < \lambda_1 < \dots < \lambda_n = \lambda^+$, each λ_i is regular, and for $\ell < n$, μ_ℓ are regular such that $\lambda_{\ell+1} \leq \mu_\ell \leq 2^{\lambda_\ell}$. Then for some regular λ_1 -good ultrafilter \mathcal{D} on λ , $\text{lcf}(\kappa, \mathcal{D}) = \mu_\ell$ whenever $\lambda_\ell \leq \kappa < \lambda_{\ell+1}$.*

These results were generalized by Koppelberg [18].

Definition 12.5. (Good for equality, Malliaris [24]) *Let \mathcal{D} be a regular ultrafilter. Say that \mathcal{D} is good for equality if for any set $X \subseteq N = M^I/\mathcal{D}$, $|X| \leq |I|$, there is $d : X \rightarrow \mathcal{D}$ such that for any $a, b \in X$, $t \in \lambda, t \in d(a) \cap d(b)$ implies that $(M \models a[t] = b[t]) \iff (N \models a = b)$.*

In the language of homogeneity, Malliaris had shown that the minimum TP_2 -class is precisely the class of theories saturated by ultrafilters on λ whose ultrapowers admit an internal bijection between any two disjoint sets of size $\leq \lambda$. (That is, for any two enumerations $\langle a_i : i < \lambda \rangle, \langle b_i : i < \lambda \rangle$ of small sets in the ultrapower N , there is an internal bijection $f : N \rightarrow N$ such that $f(a_i) = (b_i)$ for $i < \lambda$. There is no harm in replacing “bijection” with partial injection whose domain contains the desired set.) The name *good for equality* reflects that these maps are not assumed to preserve any additional structure. So among the nonsimple theories, we have, on the one hand, a class of theories characterized by ultrapowers admitting internal partial injections which preserve equality (in the sense of Definition 12.5), and, on the other hand, a class of theories characterized by ultrapowers admitting internal partial injections which preserve order (in the sense of Corollary 3.7). Toward a possible separation between these classes, we investigate the relative strength of these hypotheses.

Corollary 12.6. *If \mathcal{D} is a regular ultrafilter on λ which has λ^+ -treetops, then \mathcal{D} is good for equality.*

Proof. Conclusion 10.9 and Theorem 10.1. □

Remark 12.7. *If Conjecture 1.2 is true, then the converse to Corollary 12.6 is false; i.e., if \mathcal{D} is a regular ultrafilter on λ which is good for equality, then it need not have λ^+ -treetops.*

Proof. Malliaris had shown that the Keisler class of the theory $T_{f_{eq}}$ of a parametrized family of independent equivalence relations may be characterized as the class of theories saturated precisely by those regular ultrafilters which are good for equality (quoted below as Section 13 Theorem F). This theory does not have SOP_2 . If Conjecture 1.2 is true, then $T_{f_{eq}}$ is not in the maximal class, so any ultrafilter able to saturate it will be good for equality but not good, so (by Theorem 10.1) will not have λ^+ -treetops. □

Corollary 12.8 (of 3.7). *Suppose that \mathcal{D} has λ^+ -treetops, and let $\kappa = cf(\kappa) \leq \lambda$. Let $M = (\lambda, <)$. Let $A = \langle a_i : i < \kappa \rangle$ be any strictly increasing, κ -indexed sequence of $N = M^\lambda/\mathcal{D}$. Then in N there is an internal partial injection f which takes κ to the diagonal embedding of κ .*

Corollary 12.9. (of 12.8) *Suppose that \mathcal{D} has λ^+ -treetops, and let $\kappa = cf(\kappa) \leq \lambda$. Then every strictly increasing (or strictly decreasing) κ -indexed sequence has a distribution, Definition 10.10, which is good for equality.*

Proof. The diagonal embedding of κ has such a distribution. □

Definition 12.10. (Near- κ -indexed) *Let M be an infinite model, \mathcal{D} a regular ultrafilter on λ , $N = M^\lambda/\mathcal{D}$ and κ a regular cardinal. Say that the set $A \subseteq N$, $|A| = \kappa$ is near- κ -indexed and there exists an internal linear order on N under which A is either monotonic increasing or monotonic decreasing of order-type κ .*

In the context of ultrapowers, what the proof of Corollary 3.7 actually shows is the following slightly stronger statement.

Observation 12.11. *If \mathcal{D} is a regular ultrafilter on $\lambda \geq \kappa$ with λ^+ -treetops, then any near- κ -indexed set X in any \mathcal{D} -ultrapower has a distribution which is good for equality.*

Let $N = M^I/\mathcal{D}$, $|I| = \lambda$ and suppose \mathcal{D} has λ^+ -treetops, $\kappa \leq \lambda$. Then Claim 3.1 shows that the cointiality of any two κ -indexed sequences in the ultrapower is the same. Thus if $\mathcal{C}(\mathcal{D})$ contains *some* (κ, θ) -cut, *every* monotonic κ -indexed sequence will represent half of a (κ, θ) -cut. This is a strong omission of types. It does not contradict the universality of regular ultrapowers since elementary embeddings need not preserve cuts.

How strong is the assumption that all sets are near- κ -indexed?

Conclusion 12.12. *When \mathcal{D} is a regular ultrafilter on λ , then (1) implies (2), where*

- (1) \mathcal{D} is good for equality.
- (2) For any infinite model M , $N = M^\lambda/\mathcal{D}$, we have that any $A \subseteq N$, $|A| = \kappa \leq \lambda$ is near- κ -indexed.

If in addition \mathcal{D} has λ^+ -treetops, then (2) implies (1).

Proof. The last line holds by Observation 12.11, so we prove (1) implies (2).

Without loss of generality, M is a two sorted structure, one side of which contains an infinite set (from which we choose A) and the other side of which contains $(\lambda, <)$. Fix an enumeration $\pi : \kappa \rightarrow A$ of A . Let $K = \langle k_i : i < \kappa \rangle$ be the image of the diagonal embedding of κ in N , so $k_i = {}^\lambda\{i\}$. Choose a distribution $d_A : A \rightarrow \mathcal{D}$ which is good for equality. Let $d_\kappa : K \rightarrow \mathcal{D}$ be the distribution given by $d_\kappa(k_i) = d_A(\pi(i))$, which will be good for equality by definition. Now simply expand each index model $M[t]$ by adding a linear order $<_*$ to the first sort in such a way that the existential $<_*$ -type of $\{a[t] : a \in A, t \in d_A(a)\}$ is the same as the existential $<$ -type of $\{i : t \in d_\kappa(k_i)\}$.

Then in N the order $<_*$ on A will agree with the order $<$ on the true copy of κ , as described by the given enumeration. □

Finally, the following property was introduced by Malliaris in [22] and studied by Malliaris and by Malliaris and Shelah in [24, 26]. It is equivalent to λ -OK; see [26] Section 6.

Definition 12.13. (Flexible filters, [22]) *We say that the filter \mathcal{D} is λ -flexible if for any $f \in {}^I\mathbb{N}$ with $n \in \mathbb{N}$ implies $n <_{\mathcal{D}} f$, we can find $X_\alpha \in \mathcal{D}$ for $\alpha < \lambda$ such that for all $t \in I$*

$$f(t) \geq |\{\alpha : t \in X_\alpha\}|.$$

Informally, we can find a λ -regularizing family below any nonstandard integer.

Fact 12.14. (Malliaris [24]) *Suppose that \mathcal{D} is regular and T is nonlow or has TP_2 . If \mathcal{D} saturates T , then \mathcal{D} must be flexible.*

We now complete “ T is nonlow or has TP_2 ” to all nonlow or nonsimple theories. Note that any regular ultrafilter on λ which saturates *some* unstable theory must satisfy $\text{lcf}(\aleph_0, \mathcal{D}) \geq \lambda^+$ by [40] VI.4. Alternately, one can derive this condition from Theorem 10.1. Indeed one can also derive Claim 12.15 from that theorem, but it is interesting to prove it directly.

Claim 12.15. *If \mathcal{D} is a regular ultrafilter on λ , $\text{lcf}(\aleph_0, \mathcal{D}) \geq \lambda^+$ and \mathcal{D} has λ^+ -treetops, then \mathcal{D} is flexible.*

Proof. Let $M = (\lambda, <)$ and let $N = (\lambda, <)^I/\mathcal{D}$. Let some \mathcal{D} -nonstandard integer n_* be given. We would like to show that there is a regularizing family below n_* .

By hypothesis, $\text{lcf}(\aleph_0, \mathcal{D}) \geq \lambda^+$ so there is $B \subseteq N \setminus \mathbb{N}$, $B = \langle b_i : i < \lambda \rangle$ such that $i < j < \lambda$, $m < \omega$ implies $m < b_j < b_i < n_*$. By Corollary 3.7 and Corollary 12.9, there is a distribution d of B which is good for equality; that is, for all $b, b' \in B$, and all $t \in I$,

$$t \in d(b) \cap d(b') \text{ implies } (M \models b[t] = b'[t] \iff N \models b = b').$$

Let $\{X_b : b \in B\} \subseteq \mathcal{D}$ be given by $X_b = d(b)$. By choice of B and goodness for equality, $\{X_b : b \in B\}$ is a regularizing family, and by choice of d , it is below n_* , which completes the proof. \square

Conclusion 12.16. *Suppose \mathcal{D} is a regular ultrafilter on λ and \mathcal{D} saturates some nonlow or nonsimple theory. Then \mathcal{D} is flexible.*

Proof. Suppose \mathcal{D} saturates some theory T which is not low or not simple. If T is not low, apply Fact 12.14. So we may assume T is not simple. We know from [40] Theorem III.7.11 that any nonsimple theory will have either TP_2 or SOP_2 . In the case where T has TP_2 , Fact 12.14 applies again. Suppose then that T has SOP_2 . By Corollary 11.8, \mathcal{D} has λ^+ -treetops, so by Theorem 10.25 we know that $\text{lcf}(\aleph_0, \mathcal{D}) \geq \lambda^+$. Then the hypotheses of Claim 12.15 are satisfied, and \mathcal{D} is flexible. \square

13. THERE IS A MINIMUM NONSIMPLE CLASS IN KEISLER'S ORDER

The main result of this section is the following theorem.

Theorem 13.1. *There is minimum class among the nonsimple theories in Keisler's order, which contains the theory T_{feq} of a parametrized family of independent equivalence relations.*

We will build on the following theorems. (The reader may take the property TP_2 , the tree property of the second kind, to be a black box in the following results.)

Theorem E. (Shelah [40] III.7.11, in our language) *Every nonsimple theory has either TP_2 or SOP_2 (equivalently, TP_1).*

Theorem F. (Malliaris [24] Theorems 6.9 and 6.10, and Malliaris [25] Theorem 5.21) *There is minimum class among the theories with TP_2 in Keisler's order, which contains the theory T_{feq} of a parametrized family of independent equivalence relations.*

Moreover, for a regular ultrafilter \mathcal{D} on λ , the following are equivalent:

- (1) \mathcal{D} saturates T_{feq} .
- (2) \mathcal{D} is good for equality, Definition 12.5 above.
- (3) For any $N = M^\lambda / \mathcal{D}$ and any two disjoint sets $\{a_i : i < \lambda\}$, $\{b_i : i < \lambda\}$ of elements of N , listed without repetition, there exists an internal partial injection f such that for all $i < \lambda$, $f(a_i) = b_i$.

The tools of the present paper will combine to give two distinct proofs for Theorem 13.1. First, in light of 11.11, i.e., the maximality of SOP_2 in Keisler's order, Theorem 13.1 follows from Theorem F and Theorem E by virtue of collapsing the SOP_2 case.

However, it is possible to give a direct and more illuminating proof of this theorem, as we do below. The direct proof is not specific to ultrapowers, but holds for any cofinality spectrum problem allowing an endless $X_{\mathbf{a}}$; i.e., one in which there is

no bound $d_{\mathbf{a}}$ (which extends the context of regular ultrapowers). The proof itself is deferred until after Claim 13.5. It supposes a reduction 13.1, an example of the phenomenon of reduction to few asymmetric cuts. Historically, this reduction was the turning point of our argument. Since the reduction is now trivially true by the umbrella Theorem 8.1, we do not give a separate proof of 13.1.

Reduction 13.1. *If \mathbf{s} is a cofinality spectrum problem, to show that $\mathcal{C}(\mathbf{s}, \mathbf{t}_{\mathbf{s}}) \neq \emptyset$, it suffices to rule out the case of a (κ, κ^+) -cut where $\kappa^+ = \mathfrak{p}_{\mathbf{s}}$. [See Discussion 8.1.]*

Corollary 13.2. (of Theorem F) *If we consider the regular ultrapower $M_1 = M^\lambda/\mathcal{D}$ as a cofinality spectrum problem (M, M_1, \dots) , then to show that \mathcal{D} saturates T_{feq} it suffices to verify Theorem F(3) in the case where the sets are taken from $X_{\mathbf{a}}$ for some pseudo-finite $\mathbf{a} \in \text{Or}(\mathbf{s})$.*

Proof. In Theorem F, by regularity of \mathcal{D} and the fact that ultrapowers commute with reducts there is no harm in assuming $M = (\mathbb{N}, <)$, so $N = M^\lambda/\mathcal{D}$, in which case this is precisely what is proved. \square

Recall also that by Fact 12.2, the theory T_{rg} of the random graph is minimum in Keisler’s order among the unstable theories, and \mathcal{D} saturates T_{rg} if and only if it has so-called 2-separation (disjoint sets of size $\leq \lambda$ in the ultrapower can be separated by an internal set; see 12.2). For an arbitrary cofinality spectrum problem this means the following definition.

Definition 13.3. (2-separation for \mathbf{s}) *Let \mathbf{s} be a cofinality spectrum problem. We say that \mathbf{s} has 2-separation if for any $\mathbf{a} \in \text{Or}(\mathbf{s})$ and any two disjoint sets $A, B \subseteq X_{\mathbf{a}}$ with $|A| + |B| \leq \mathfrak{p}_{\mathbf{s}}$, there is a definable $X \in M_1$ such that $A \subseteq X$ and $B \cap X = \emptyset$.*

Lemma 13.4. *Let \mathbf{s} be a cofinality spectrum problem and suppose that*

- (1) \mathbf{s} has 2-separation;
- (2) $\kappa^+ = \lambda = \mathfrak{p}_{\mathbf{s}} < \mathbf{t}_{\mathbf{s}}$;
- (3) $\mathbf{a} \in \text{Or}(\mathbf{s})$ is endless; i.e., there is no bound $d_{\mathbf{a}}$;
- (4) $\{d_\epsilon : \epsilon < \lambda\}, \{e_\epsilon : \epsilon < \lambda\}$ are disjoint subsets of $X_{\mathbf{a}}$, listed without repetition.

Then there is in M_1 a bijection $f : X_{\mathbf{a}} \rightarrow X_{\mathbf{a}}$ such that $f(d_\epsilon) = e_\epsilon$ for all $\epsilon < \lambda$.

Proof. We begin with $\mathbf{a} \in \text{Or}(\mathbf{s})$, $\{d_\epsilon : \epsilon < \lambda\}, \{e_\epsilon : \epsilon < \lambda\}$ is as given. We define $\mathcal{I} = \{S \subseteq \lambda : \text{there is } f \text{ such that } M_1 \models \text{“}f \text{ is a one-to-one function from } X_{\mathbf{a}} \text{ to } X_{\mathbf{a}}\text{” and } \epsilon \in S_1 \iff M_1 \models f(d_\epsilon) = e_\epsilon\}$.

First, we verify that if $M_1 \models \text{“}f \text{ is a one-to-one function with } \text{dom}(f) \subseteq X_{\mathbf{a}} \text{ and } \text{range}(f) \subseteq X_{\mathbf{a}}\text{”}$ then

$$\{\epsilon < \lambda : f(d_\epsilon) = e_\epsilon\} \in \mathcal{I}.$$

Equivalently, this holds if $f \in M_1$ is a one-to-one function on some M_1 -definable subset of $X_{\mathbf{a}}$.

We can define from f a total function g as follows. Necessarily the domain and range of f are definable sets. Let $g : X_{\mathbf{a}} \rightarrow X_{\mathbf{a}}$ be defined by

$$g(a) = \begin{cases} f(a) & \text{if } a \in \text{dom}(f) \\ f^{-1}(a) & \text{if } a \in \text{range}(f) \setminus \text{dom}(f) \\ a & \text{if } a \in X_{\mathbf{a}} \setminus \text{dom}(f) \setminus \text{range}(f) \end{cases}.$$

Second, we prove that \mathcal{I} is an ideal on λ , by checking the conditions for an ideal. In light of the above it suffices to specify bijections on definable subsets of $X_{\mathbf{a}}$.

(a) $\mathcal{I} \subseteq \mathcal{P}(\lambda)$ by definition.

(b) $\emptyset \in \mathcal{I}$ trivially.

(c) Suppose $S_1 \subseteq S_2 \subseteq \lambda$ and $S_2 \in \mathcal{I}$, exemplified by f_2 . By 2-separation, there is a set $X \in M_1$ such that $M_1 \models X \subseteq X_{\mathbf{a}}$ and for all $\epsilon < \lambda$,

$$(M_1 \models d_\epsilon \in X) \iff (\epsilon \in S).$$

Let $f_1 = f_2 \upharpoonright_Y$ where $Y = \text{dom}(f_2) \cap X$. Then f_1 witnesses that $S_1 \in \mathcal{I}$. So \mathcal{I} is downward closed.

(d) If $S_1, S_2 \in \mathcal{I}$, then by (c) the set $S'_2 = S_2 \setminus S_1$ belongs to \mathcal{I} . Let f_1, f'_2 witness that S_1, S'_2 belong to \mathcal{I} , respectively. Define $f_2 \in M_1$ by $\text{dom}(f) = \text{dom}(f_1) \cup \text{dom}(f'_2)$, and

$$f_2(a) = \begin{cases} f_1(a) & \text{if } a \in \text{dom}(f_1) \\ f'_2(a) & \text{if } a \in \text{dom}(f_2) \setminus \text{dom}(f_1) \end{cases}.$$

Iterating, we have that \mathcal{I} is closed under a finite union.

It follows from (a)–(d) that \mathcal{I} is an ideal on λ .

Third, we prove closure for $\text{cf}(\delta) < \kappa$. Let $\mathbf{b} \in \text{Or}(\mathbf{s})$ be such that $X_{\mathbf{b}} = X_{\mathbf{a}} \times X_{\mathbf{a}}$. Let $(\mathcal{T}, \trianglelefteq)$ be the subset of $\mathcal{T}_{\mathbf{b}}$ consisting of elements c whose range is the graph of a partial injection from $X_{\mathbf{a}}$ to $X_{\mathbf{a}}$, i.e.,

- for each $n < \text{lg}(c)$, $\mathbf{c}(n) \in X_{\mathbf{a}} \times X_{\mathbf{a}}$; as before, we denote these values by $\mathbf{c}(n, 0)$ and $\mathbf{c}(n, 1)$, respectively;
- the set $\{\mathbf{c}(n) : n \leq \max(\text{dom}(c))\}$ is the graph of a partial injection from $X_{\mathbf{a}}$ to $X_{\mathbf{a}}$;
- if $\mathbf{c}_1 \trianglelefteq \mathbf{c}_2$, then \mathbf{c}_2 extends \mathbf{c}_1 , considered as a function.

Note that for elements $\mathbf{c} \in \mathcal{T}$, we will refer extensively to the function whose graph is $\{(\mathbf{c}(n, 0), \mathbf{c}(n, 1)) : n \leq \max(\text{dom}(\mathbf{c}))\}$. For purposes of clarity, we denote this function by $\text{fun}(\mathbf{c})$ and will say, e.g., that $d_\epsilon \in \text{fun}(\mathbf{c})$ or that $\text{fun}(\mathbf{c})(d_\epsilon) = e_\epsilon$.

Subclaim. Suppose that δ is an ordinal with $\text{cf}(\delta) < \kappa$ and we are given $\mathbf{c}_\alpha, n_\alpha, S_\alpha$ ($\alpha < \delta$) and a set S , such that

- for each $\alpha < \delta$, $M_1 \models \mathbf{c}_\alpha \in \mathcal{T}$;
- $M_1 \models \mathbf{c}_\beta \trianglelefteq \mathbf{c}_\alpha$ for each $\beta < \alpha < \delta$;
- for each $\alpha < \delta$, $n_\alpha = \text{lg}(\mathbf{c}_\alpha) - 1$;
- for each $\alpha < \delta$, $S_\alpha = \{\epsilon < \lambda : d_\epsilon \in \text{dom}(\text{fun}(\mathbf{c}_\alpha))\}$, i.e., $= \{\epsilon < \lambda : (\exists n \leq n_\alpha)(\mathbf{c}_\alpha(n, 0) = d_\epsilon)\}$;
- for each $\alpha < \delta$, $\epsilon \in S_\alpha$ implies $\text{fun}(\mathbf{c}_\alpha)(d_\epsilon) = e_\epsilon$;
- $S = \bigcup \{S_\alpha : \alpha < \delta\}$.

Then there is \mathbf{c}_* such that

- (1) $M_1 \models \mathbf{c}_* \in \mathcal{T}$;
- (2) $M_1 \models \mathbf{c}_\alpha \trianglelefteq \mathbf{c}_*$ for $\alpha < \delta$;
- (3) for $\epsilon < \lambda$,

$$d_\epsilon \in \text{dom}(\text{fun}(\mathbf{c}_*)) \iff \epsilon \in S \iff \text{fun}(\mathbf{c}_*)(d_\epsilon) = e_\epsilon.$$

In particular, $\text{fun}(\mathbf{c}_*)$ witnesses that $S \in \mathcal{I}$.

Proof. Since $\text{cf}(\delta) < \kappa < \mathbf{t}_s$, by treetops there is $\mathbf{c} \in \mathcal{T}$ satisfying (1) and (2). Now the sequence $\langle n_\alpha : \alpha < \delta \rangle$ represents the left half of some cut (C_1, C_2) of $\text{dom}(\mathbf{c})$.

Let $\langle m_\beta : \beta < \theta \rangle$ represent the right half of this cut.

By Reduction 13.1, $\text{cf}(\delta) < \kappa < \kappa^+ = \lambda$ implies $\text{lcf}(\delta) > \lambda$, as otherwise there would be a corresponding cut in $\mathbf{c}(\mathcal{D})$. Thus, $\theta > \lambda$.

Now for each $\epsilon \in \lambda \setminus S$ there is $\beta(\epsilon) < \theta$ such that $d_\epsilon \notin \text{dom}(\mathbf{c}(m_\beta))$. Let $\beta = \sup\{\beta(\epsilon) : \epsilon \in \lambda \setminus S\}$. Since $\theta > \lambda$, $\beta > \alpha$ for all $\alpha < \delta$. Then $\mathbf{c} \upharpoonright m_\beta$ is the desired \mathbf{c}_* . \square

This completes the proof of closure.

Fourth, we prove that the ideal \mathcal{I} is λ -complete and contains each $\{\alpha\}$ for $\alpha \in \lambda$. The claim about containing the singletons is trivial since we know it is an ideal on λ . So let us show that \mathcal{I} is λ -complete.

Let $\langle S_\alpha : \alpha < \delta \rangle$ be an increasing sequence of elements of \mathcal{I} , with $\delta < \lambda$. Without loss of generality $\delta = \text{cf}(\delta)$; call it θ . So $\theta \leq \kappa$. By induction on $\alpha < \kappa$ we choose a sequence $\langle \mathbf{c}_\alpha : \alpha < \kappa \rangle$ just as before. The subclaim proved above says precisely that we can continue the induction for all $\alpha < \kappa$, and we now address the case of κ , i.e., θ .

That is, having chosen $\langle \mathbf{c}_\alpha : \alpha < \theta \rangle$, as $\kappa < \lambda \leq \mathfrak{t}_s$ we have κ^+ -treetops so we may choose an upper bound \mathbf{c} . Let $n = \max(\text{dom}(\mathbf{c}))$, and by definition $\text{fun}(\mathbf{c})$ is a one-to-one function.

Thus by definition $\text{fun}(\mathbf{c})$ witnesses that $S \in \mathcal{I}$ where $S = \{\epsilon : \text{fun}(\mathbf{c})(d_\epsilon) = e_\epsilon\}$. By choice of \mathbf{c} as an upper bound, $\bigcup\{S_\alpha : \alpha < \theta\} \subseteq S \in \mathcal{I}$, and thus necessarily $\bigcup\{S_\alpha : \alpha < \theta\} \in \mathcal{I}$ by (c) above. This finishes the proof that the ideal is λ -complete.

Fifth, we prove that $\kappa^+ = \lambda$ is the union of κ sets from I . By induction on $\alpha < \lambda$ we choose $\langle \mathbf{c}_\alpha, n_\alpha \rangle$ such that

- $\mathbf{c}_\alpha \in \mathcal{T}$, i.e., it represents an increasing pseudo-finite sequence of one-to-one functions from $X_{\mathbf{a}}$ to $X_{\mathbf{a}}$;
- $n_\alpha = \text{lg}(\mathbf{c}) - 1$;
- $\beta < \alpha$ implies $\mathbf{c}_\beta \trianglelefteq \mathbf{c}_\alpha$;
- if $\epsilon < \beta < \alpha$ and $n \in (n_\epsilon, n_\beta]$, then $d_\epsilon \in \text{dom}(\text{fun}(\mathbf{c}_\alpha \upharpoonright_n))$ and $\text{fun}(\mathbf{c}_\alpha \upharpoonright_n)(d_\epsilon) = e_\epsilon$.

The induction is straightforward. For $\alpha = 0$, let $\mathbf{c}_\alpha = \emptyset$.

For $\alpha = \beta + 1$, let $n_{\alpha+1} = n_\beta + 1$, and \mathbf{c}_α is determined by asking that $\text{dom}(\text{fun}(\mathbf{c}_\alpha)) = \text{dom}(\text{fun}(\mathbf{c}_\beta)) \cup \{d_\beta\}$, that $d \in \text{dom}(\text{fun}(\mathbf{c}_\beta))$ implies $\text{fun}(\mathbf{c}_\alpha)(d) = \text{fun}(\mathbf{c}_\beta)(d)$, and that $\text{fun}(\mathbf{c}_\alpha)(d_\beta) = e_\beta$.

For the $\alpha = \delta < \lambda$ limit, let \mathbf{c} be a \trianglelefteq -upper bound given by treetops. Now for each $\epsilon < \alpha$ we define

$$k_\epsilon^\alpha = \max\{n : n_{\epsilon+1} \leq n \leq \text{dom}(\mathbf{c}) \text{ and } \text{fun}(\mathbf{c} \upharpoonright_n)(d_\epsilon) = e_\epsilon\}.$$

Thus for all $\epsilon < \alpha$ and all $\beta < \alpha$, $k_\epsilon^\alpha > n_\beta$, while for fixed α and increasing $\epsilon < \alpha$ the k_ϵ^α form a descending sequence. In other words, $(\langle n_\beta : \beta < \alpha \rangle, \langle k_\epsilon^\alpha : \epsilon < \alpha \rangle)$ represent a pre-cut in $X_{\mathbf{b}}$. However, $\text{lcf}(\text{cf}(\alpha), \mathfrak{s}) \geq \lambda$ so necessarily it is a pre-cut and not a cut; we may find k_ϵ such that for all $\epsilon < \alpha$ and all $\beta < \alpha$, $n_\beta < k_\epsilon < k_\epsilon^\alpha$. Let $\mathbf{c}_\alpha = \mathbf{c} \upharpoonright_{k_\epsilon}$. This completes the inductive construction of the sequence.

Thus $\langle (\mathbf{c}_\alpha, n_\alpha) : \alpha < \lambda \rangle$ is well defined. As we assumed $\mathfrak{t}_s > \lambda$, we have λ^+ -treetops so we may choose c to be an upper bound in \mathcal{T} for $\langle \mathbf{c}_\alpha : \alpha < \lambda \rangle$. For each $\epsilon < \lambda$, let k_ϵ be as given in the previous paragraph.

By definition $\langle n_\alpha : \alpha < \lambda \rangle$ is an increasing sequence in $X_{\mathbf{b}}$. By assumption in the statement of the Claim, we are in the case where $\mathfrak{p}_s < \mathfrak{t}_s$, and thus by Reduction 13.1 $\text{lcf}(\lambda) = \kappa = \lambda^-$. Thus for some $\langle n_i^* : i < \kappa \rangle$ we have that

$(\langle n_\alpha : \alpha < \lambda \rangle, \langle n_i^* : i < \kappa \rangle)$ represents a cut in $X_{\mathbf{b}}$. For each $i < \kappa$, let

$$Y_i = \{\epsilon < \lambda : n_i^* < k_\epsilon\}.$$

Now, each $Y_i \in \mathcal{I}$, since this is witnessed by $\mathbf{c} \upharpoonright_{n_i^*}$ by what we have shown. Moreover, $\{Y_i : i < \kappa\}$ is an increasing sequence of subsets of λ whose union is λ . So we have presented λ as the union of κ elements of \mathcal{I} .

Having proved each of these properties, we may therefore assume λ is the union of κ sets from I , and that the ideal \mathcal{I} is λ -complete. Thus $\lambda \in \mathcal{I}$. As $\lambda \in \mathcal{I}$ shows the existence of the desired bijection, this completes the proof. \square

Corollary 13.5. *Let \mathcal{D} be a regular ultrafilter on λ which λ^+ -saturates the theory of the random graph. If \mathcal{D} has λ^+ -treetops, then \mathcal{D} λ^+ -saturates T_{feq} .*

Proof. As \mathcal{D} is a regular ultrafilter, we may choose any M with enough set theory for trees and consider the ultrapower as a cofinality spectrum problem $(M, M^\lambda/\mathcal{D}, \dots)$. By assumption, \mathcal{D} has λ^+ -treetops, and thus $\mathfrak{t}_{\mathbf{s}} > \lambda$. There are two cases. If $\mathfrak{t}_{\mathbf{s}} = \mathfrak{p}_{\mathbf{s}} > \lambda$, then by definition of $\mathfrak{p}_{\mathbf{s}}$ (2.9 above) and Corollary 10.16, \mathcal{D} is λ^+ -good. Thus necessarily \mathcal{D} saturates T_{feq} .

Otherwise, $\mathfrak{t}_{\mathbf{s}} > \mathfrak{p}_{\mathbf{s}}$. It will suffice by Corollary 13.2 to show that bijections exist. By Reduction 13.1, the case $\mathfrak{t}_{\mathbf{s}} > \mathfrak{p}_{\mathbf{s}}$ necessarily entails that λ is the successor of a regular cardinal κ and that $\mathfrak{t}_{\mathbf{s}} > \mathfrak{p}_{\mathbf{s}} = \lambda = \kappa^+$. Thus, for any $\mathbf{a} \in \text{Or}(\mathfrak{s})$ and any suitably chosen sequences $\{d_\epsilon : \epsilon < \lambda\}$, $\{e_\epsilon : \epsilon < \lambda\}$ of elements of $X_{\mathbf{a}}$, we may apply Lemma 13.4 to obtain a suitable bijection. We conclude by Theorem F or just Corollary 13.2 that \mathcal{D} saturates T_{feq} , as desired. \square

We now have the ingredients to prove Theorem 13.1.

Proof of Theorem 13.1. Let $\lambda \geq \aleph_0$ and let \mathcal{D} be a regular ultrafilter on λ . Suppose that \mathcal{D} saturates some nonsimple theory T_* . It suffices to show that \mathcal{D} necessarily also saturates T_{feq} . There are two cases which, by Fact E, cover all possibilities.

Case 1. T_* has TP_2 . Then \mathcal{D} saturates T_{feq} by Theorem F.

Case 2. T_* has SOP_2 . By Fact 12.2, \mathcal{D} saturates T_{rg} . By Conclusion 11.9, \mathcal{D} has λ^+ -treetops. So we may apply Claim 13.5 to conclude \mathcal{D} saturates T_{feq} .

This completes the proof of Theorem 13.1. \square

14. $\mathfrak{p} = \mathfrak{t}$

In this section we apply the main theorem of Section 9 to prove Theorem 14.1.

Definition 14.1. (See, e.g., van Douwen [8], Vaughan [49], Blass [5]) *We define several properties which may hold of a family $D \subseteq [\mathbb{N}]^{\aleph_0}$. Let $A \subseteq^* B$ mean that $\{x : x \in A, x \notin B\}$ is finite.*

- *D has a pseudo-intersection if there is an infinite $A \subseteq \mathbb{N}$ such that for all $B \in D$, $A \subseteq^* B$.*
- *D has the s.f.i.p. if every nonempty finite subfamily has infinite intersection.*
- *D is called a tower if it is well ordered by \supseteq^* and has no infinite pseudo-intersection.*
- *D is called open if it is closed under almost subsets, and dense if every $A \in [\mathbb{N}]^{\aleph_0}$ has a subset in D .*

We then define

- $\mathfrak{p} = \min\{|\mathcal{F}| : \mathcal{F} \subseteq [\mathbb{N}]^{\aleph_0} \text{ has the s.f.i.p. but has no infinite pseudo-intersection}\},$
- $\mathfrak{t} = \min\{|\mathcal{T}| : \mathcal{T} \subseteq [\mathbb{N}]^{\aleph_0} \text{ is a tower}\},$
- $\mathfrak{b} = \min\{|B| : B \subseteq {}^\omega\omega \text{ is unbounded in } ({}^\omega\omega, \leq^*)\},$
- $\mathfrak{h} = \text{the smallest number of dense open families with empty intersection.}$

Fact 14.2. \mathfrak{p} and \mathfrak{t} are regular. It is known that $\mathfrak{p} \leq \mathfrak{t} \leq \mathfrak{h} \leq \mathfrak{b}$.

Proof. Regularity of \mathfrak{t} and the first inequality are clear from the definitions. For regularity of \mathfrak{p} , due to Szymański, see, e.g., van Douwen [8] Theorem 3.1(e). The result $\mathfrak{t} \leq \mathfrak{b}$ is due to Rothberger [37], attributed in [8] p. 123, and the result $\mathfrak{t} \leq \mathfrak{h} \leq \mathfrak{b}$ is due to Balcar, Pelant, and Simon [3], attributed in [49] p. 200. For completeness: for the second inequality, see, e.g., Blass [5] Proposition 6.8; for the third, [5] Theorem 6.9. □

To begin, we look for a relevant cofinality spectrum problem.

Definition 14.3. We fix the following for the remainder of this section:

- (1) Let $M = (\mathcal{H}(\aleph_1), \in)$.
- (2) Let $\mathbf{Q} = ([\mathbb{N}]^{\aleph_0}, \supseteq^*)$ be our forcing notion, and \mathbf{V} a transitive model of ZFC.
- (3) Let \mathbf{G} be the canonical name of a generic subset of \mathbf{Q} (which is forced to be an ultrafilter on the Boolean algebra $\mathcal{P}(\mathbb{N})^{\mathbf{V}}$).
- (4) Let \mathbf{G} be a generic subset of \mathbf{Q} over \mathbf{V} , which we fix for this section. (Often, however, we will simply work in \mathbf{V} using the name \mathbf{G} .)
- (5) For $f \in \mathbf{V}$, let \check{f} denote the \mathbf{Q} -name for f .

Define the generic ultrapower in the forcing extension $\mathbf{V}[\mathbf{G}]$ as follows:

- (6) Given M, \mathbf{Q} , and \mathbf{G} , by the generic ultrapower M^ω/\mathbf{G} in $\mathbf{V}[\mathbf{G}]$ we will mean the model $\mathcal{N} \in \mathbf{V}[\mathbf{G}]$ with universe $\{f/\mathbf{G} : f \in ({}^\omega M)^{\mathbf{V}}\}$, such that
 - $\mathcal{N} \models “f_1/\mathbf{G} = f_2/\mathbf{G}”$ iff $\{n : f_1(n) = f_2(n)\} \in \mathbf{G}$ (this set is necessarily from \mathbf{V});
 - $\mathcal{N} \models “(f_1/\mathbf{G}) \in (f_2/\mathbf{G})”$ iff $\{n : f_1(n) \in f_2(n)\} \in \mathbf{G}$.

We denote by $\mathbf{j} = \mathbf{j}_{\mathbf{G}} : M \rightarrow \mathcal{N}$ the map given by $\mathbf{j}(a) = \langle \dots a \dots \rangle / \mathbf{G}$.

It will also be useful to refer to these objects in \mathbf{V} .

- (7) In \mathbf{V} , let $\check{\mathcal{N}}$ be the \mathbf{Q} -name of the generic ultrapower $M^\omega/\check{\mathbf{G}}$, i.e., the model with
 - (a) universe $\{f/\check{\mathbf{G}} : f \in ({}^\omega M)^{\mathbf{V}}\}$ such that
 - (b) $\Vdash_{\mathbf{Q}} “(\check{\mathcal{N}} \models f_1/\check{\mathbf{G}} = f_2/\check{\mathbf{G}})$ iff $\{n : f_1(n) = f_2(n)\} \in \check{\mathbf{G}}”$
(as noted, this set is necessarily from \mathbf{V});
 - (c) $\Vdash_{\mathbf{Q}} “(\check{\mathcal{N}} \models (f_1/\check{\mathbf{G}}) \in (f_2/\check{\mathbf{G}}))$ iff $\{n : f_1(n) \in f_2(n)\} \in \check{\mathbf{G}}.”$
- (8) Note that $\mathcal{N} = \check{\mathcal{N}}[\mathbf{G}]$.

By definition of \mathfrak{t} , \mathbf{Q} is \mathfrak{t} -complete. Thus, forcing with \mathbf{Q} adds no new bounded subsets of \mathfrak{t} (where new means “ $\notin \mathbf{V}$ ”) and no new sequences of length $< \mathfrak{t}$ of members of \mathbf{V} . Moreover, by the \mathfrak{t} -completeness of \mathbf{Q} , and since $\mathfrak{p} \leq \mathfrak{t}$, moving from \mathbf{V} to $\mathbf{V}[\mathbf{G}]$ will not affect whether $\mathfrak{p} < \mathfrak{t}$.

We now build a cofinality spectrum problem. We will let $M = M^+ = (\mathcal{H}(\aleph_1), \in)$ as above and $M_1 = M_1^+ = \mathcal{N}$. For generic ultrapowers, the parallel of Los’ theorem holds, so \mathbf{j} is an elementary embedding of M into \mathcal{N} .

Definition 14.4. Working in $\mathbf{V}[\mathbf{G}]$, let M, \mathcal{N} be as in 14.3. Let Δ_{psf} be the set of all first-order formulas $\varphi(x, y, \bar{z})$ in the vocabulary of M , i.e., $\{\in, =\}$, such that if $\bar{c} \in {}^{\ell(\bar{z})}M$, then $\varphi(x, y, \bar{c})$ is a linear order on the finite set $A_{\varphi, \bar{c}} = \{a : M \models \varphi(a, a, \bar{c})\}$, denoted by $\leq_{\varphi, \bar{c}}$. We require $\ell(x) = \ell(y)$ but do not require $\ell(x) = 1$.

Observation 14.5. If M, \mathcal{N} are as above and $\varphi \in \Delta_{\text{psf}}$, then

(a) for each $\bar{c} \in {}^{\ell(\bar{v})}\mathcal{N}$, $\varphi(x, y; \bar{c})$ is a discrete linear order on the set

$$\{a : \mathcal{N} \models \varphi(a, a; \bar{c})\};$$

(b) each nonempty \mathcal{N} -definable subset of $A_{\varphi, \bar{c}}$ has a first and a last element;

(c) we may in \mathcal{N} identify $(A_{\varphi, \bar{c}}, \leq_{\varphi, \bar{c}})$ with a definable subset of some

$$\langle (X_n, \leq_n) : n < \omega \rangle / \mathbf{G},$$

where each X_n is finite and linearly ordered by \leq_n .

Proof. Los' theorem. □

Claim 14.6. Working in $\mathbf{V}[\mathbf{G}]$, $(M, \mathcal{N}, Th(M), \Delta_{\text{psf}})$ is a cofinality spectrum problem which, for the remainder of this section, we call \mathbf{s} .

Proof. We check Definition 2.3. Conditions (1) and (2) are immediate. For (3), let $\Delta = \Delta_{\text{psf}}$, so the only other data we need to specify are in each instance to set $d_{\mathbf{a}}$ to be the maximum element. Condition (4) is by construction; see 14.5. Suppose we are given \mathbf{a}, \mathbf{b} with $X_{\mathbf{a}} = \{a : \mathcal{N} \models \varphi_1(a, a, \bar{c}_1)\}$ and $X_{\mathbf{b}} = \{a : \mathcal{N} \models \varphi_2(a, a, \bar{c}_2)\}$, where φ_1, φ_2 are from 14.4. Let $\theta(x_1y_1x_2y_2, \bar{c}_1, \bar{c}_2)$ be the formula which implies $(x_1, y_1), (x_2, y_2) \in X_{\mathbf{a}} \times X_{\mathbf{b}}$ and gives the Gödel pairing function; i.e., we order $(x_1, y_1), (x_2, y_2)$ first by maximum, then by first coordinate, and then by second coordinate. Clearly $\theta(x_1y_1, x_2y_2, z_1z_2) \in \Delta_{\text{psf}}$. This gives (5) and (6). It remains to check we have trees. Let \mathbf{a} be given, and suppose $\varphi_{\mathbf{a}} = \varphi(x, y, z)$. Let $\psi(w, z)$ be such that for each $\bar{c} \in {}^{\ell(\bar{z})}M$ we have that $\mathcal{T}_{\varphi, \bar{c}} = \{\eta : M \models \psi(\eta, \bar{c})\}$ is the set of finite sequences of members of $A_{\varphi, \bar{c}} = \{a : M \models \varphi(a, a, \bar{c})\}$ of length $< \max A_{\varphi, \bar{c}}$. Let $\trianglelefteq = \{(\eta, \nu) : \eta, \nu \in \mathcal{T}_{\varphi, \bar{c}} \text{ and } \eta \text{ is an initial segment of } \nu\}$. We can likewise define the functions lg and val . Clearly by Los' theorem these objects will behave as desired in \mathcal{N} . This completes the proof of (7), and (8) is immediate, so we are done. □

It follows from the definitions that $\mathbf{p} \leq \mathbf{t}$. If $\mathbf{p} = \mathbf{t}$, then Theorem 14.1 is immediately true. So we shall assume, toward a contradiction, that $\mathbf{p} < \mathbf{t}$ in \mathbf{V} .

Claim 14.7. Working in $\mathbf{V}[\mathbf{G}]$, let \mathbf{s} be the cofinality spectrum problem from 14.6. Then $\mathbf{t} \leq \mathbf{t}_{\mathbf{s}}$.

That is, let $\mathbf{a} \in \text{Or}(\mathbf{s})$ be given, so $\mathcal{N} \models \text{“}(\mathcal{T}_{\mathbf{a}}, \trianglelefteq_{\mathbf{a}})$ is a tree of finite sequences of $(X_{\mathbf{a}}, \leq_{\mathbf{a}})$.” Then any increasing sequence of cofinality $\kappa < \mathbf{t}$ in $(\mathcal{T}_{\mathbf{a}}, \trianglelefteq_{\mathbf{a}})^{\mathcal{N}}$ has an upper bound.

Remark 14.8. In fact $\mathbf{t} = \mathbf{t}_{\mathbf{s}}$, but this will not be needed.

Proof of Claim 14.7. First, a reduction: Recalling the definition of \mathbf{s} in 14.6 and 14.5(c), we may assume $(X_{\mathbf{a}}, \leq_{\mathbf{a}}, \mathcal{T}_{\mathbf{a}}, \trianglelefteq_{\mathbf{a}}) = \langle (X_{\mathbf{a}_n}, \leq_{\mathbf{a}_n}, \mathcal{T}_n, \trianglelefteq_{\mathcal{T}_n}) : n < \omega \rangle / \mathbf{G}$. So without loss of generality for each $n < \omega$ $(X_{\mathbf{a}_n}, \leq_{\mathbf{a}_n}, \mathcal{T}_n, \trianglelefteq_{\mathcal{T}_n})$ is standard; i.e., $X_{\mathbf{a}_n}$ is finite and \mathcal{T}_n is the set of finite sequences of elements of $X_{\mathbf{a}_n}$, partially ordered by inclusion.

For each $n < \omega$, there is an isomorphism $h_n : (X_{\mathbf{a}_n}, <_{\mathbf{a}_n}) \rightarrow (k_n, <_{k_n})$ where $k_n \in \omega$, $<_k$ is the usual order on ω restricted to k , and $\lim_{\mathbf{G}} \langle k_n : n < \omega \rangle$ is infinite. Then in \mathcal{N} , $h = \langle h_n : n < \omega \rangle / \mathbf{G}$ gives an isomorphism between $(\mathcal{T}_{\mathbf{a}}, \trianglelefteq_{\mathbf{a}})$ and a definable downward closed subset of $({}^{\omega}>\omega, \trianglelefteq)^{\mathcal{N}}$. So for the remainder of the proof, without loss of generality, we work in the tree $({}^{\omega}>\omega, \trianglelefteq)^{\mathcal{N}}$. This completes the reduction.

We work now in \mathbf{V} . Let $\theta = \text{cf}(\theta) < \mathfrak{t}$ be given and let $B \in \mathbf{Q}$ ($B \in \mathbf{G}$) be such that

$$B \Vdash_{\mathbf{Q}} \text{“}\langle \underline{f}_\alpha / \underline{\mathbf{G}} : \alpha < \theta \rangle \text{ is } \trianglelefteq^{\mathcal{N}}\text{-increasing in } ({}^{\omega}>\omega, \trianglelefteq)^{\mathcal{N}}\text{”},$$

where without loss of generality, $B \Vdash_{\mathbf{Q}} \text{“}\underline{f}_\alpha = f_\alpha\text{”}$ for $\alpha < \theta$ since forcing with \mathbf{Q} adds no new sequences of length $< \mathfrak{t}$.

By assumption, $\theta < \mathfrak{t}$ and thus $\theta < \mathfrak{b}$, the bounding number. So we may choose some increasing function $g : \mathbb{N} \rightarrow \mathbb{N} \setminus \{0\}$ such that for each $\alpha < \theta$ there is n_α satisfying

$$\text{if } n \geq n_\alpha, \text{ then } g(n) > \text{lg}(f_\alpha(n)) + \Sigma\{f_\alpha(n)(j) : j < \text{lg}(f_\alpha(n))\}.$$

Informally, for each $\alpha < \theta$, for all but finitely many n , $g(n)$ dominates the sum of all values in the range of f_α when the domain is restricted to n .

Now let $\bar{s} = \langle s_n : n < \omega \rangle$ be given by

$$s_n = {}^{g(n)}\geq g(n) = \{\eta : \eta \text{ a sequence of length } \leq g(n) \text{ of numbers } < g(n)\}.$$

Then

- (1) each s_n is a finite nonempty subtree of ${}^{\omega}>\omega$;
- (2) if $\alpha < \theta$, then $(\forall^\infty n)(f_\alpha(n) \in s_n)$.

We use the following notation:

$$({}^{\omega}>\omega)^{[\nu]} = \{\eta \in {}^{\omega}>\omega : \nu \trianglelefteq \eta\}.$$

Then for each $\alpha < \theta$, we define Y_α as follows:

$$Y_\alpha = \bigcup \{ \{n\} \times (s_n \cap ({}^{\omega}>\omega)^{[f_\alpha(n)]}) : n \in B \}.$$

Finally, let

$$Y_* = \bigcup \{ \{n\} \times s_n : n \in B \}.$$

Then for each $\alpha < \theta$, we have

- (1) $Y_\alpha \subseteq B \times {}^{\omega}>\omega$;
- (2) $Y_\alpha \cap (\{n\} \times {}^{\omega}>\omega)$ is finite, and $\subseteq \{n\} \times s_n$ for every n ;
- (3) Y_α is infinite;
- (4) $Y_\alpha \subseteq Y_*$.

Moreover, if $\alpha < \beta$, then $Y_\beta \subseteq^* Y_\alpha$. Why? If $\alpha < \beta$, then $\{n \in B : f_\alpha(n) \not\trianglelefteq f_\beta(n)\}$ is finite, as otherwise there is $B' \geq_{\mathbf{Q}} B$ contradicting

$$B \Vdash_{\mathbf{Q}} (\mathcal{N} \models \text{“}f_\alpha / \mathbf{G} \trianglelefteq f_\beta / \mathbf{G}\text{”}).$$

So as $\mathfrak{t} = \lambda > \theta$, there is an infinite $Z \subseteq Y_*$ such that $\alpha < \theta$ implies $Z \subseteq^* Y_\alpha$. Let $B_1 = \{n \in B : Z \cap (\{n\} \times s_n) \neq \emptyset\}$. For each $n \in B_1$ choose $\nu_n \in s_n$ such that $(n, \nu_n) \in Z \cap (\{n\} \times s_n)$. Choose $\nu_n = \langle 0 \rangle$ for $n \in \mathbb{N} \setminus B$. Then

$$B_1 \Vdash_{\mathbf{Q}} \text{“}\langle \nu_n : n < \omega \rangle / \mathbf{G} \text{ is an upper bound for } \langle f_\alpha / \mathbf{G} : \alpha < \theta \rangle \text{ in } ({}^{\omega}>\omega, \trianglelefteq)^{\mathcal{N}}\text{”}.$$

This completes the proof. □

Conclusion 14.9. Working in $\mathbf{V}[\mathbf{G}]$, let \mathbf{s} be the cofinality spectrum problem from 14.6. Then $\mathcal{C}(\mathbf{s}, \mathbf{t}) = \emptyset$.

Proof. By Theorem 9.1, $\mathcal{C}(\mathbf{s}, \mathbf{t}_\mathbf{s}) = \emptyset$, and by Claim 14.7, $\mathbf{t} \leq \mathbf{t}_\mathbf{s}$. □

Discussion 14.10. These results do not contradict the existence of Hausdorff gaps; see, e.g., Definition 2.26 of [46]. This is because to obtain treetops and the transfer of the peculiar cut below, we restrict ourselves to some infinite subset of ω .

Aiming for a contradiction, we will leverage Conclusion 14.9 against a cut existence result from Shelah [44], which we prove can be translated to our context.

Definition 14.11. (Peculiar cut, [44] Definition 1.10) Let κ_1, κ_2 be infinite regular cardinals. A (κ_1, κ_2) -peculiar cut in ${}^\omega\omega$ is a pair $(\langle g_i : i < \kappa_2 \rangle, \langle f_i : i < \kappa_1 \rangle)$ of sequences of functions in ${}^\omega\omega$ such that

- (1) $(\forall i < j < \kappa_2)(g_i <^* g_j)$;
- (2) $(\forall i < j < \kappa_1)(f_j <^* f_i)$;
- (3) $(\forall i < \kappa_1)(\forall j < \kappa_2)(g_j <^* f_i)$;
- (4) if $f : \omega \rightarrow \omega$ is such that $(\forall i < \kappa_1)(f <^* f_i)$, then $f <^* g_j$ for some $j < \kappa_2$;
- (5) if $f : \omega \rightarrow \omega$ is such that $(\forall j < \kappa_2)(g_j <^* f)$, then $f_i <^* f$ for some $i < \kappa_1$.

Theorem G. (Shelah [44] Theorem 1.12) Assume $\mathfrak{p} < \mathfrak{t}$. Then for some regular cardinal κ there exists a (κ, \mathfrak{p}) -peculiar cut in ${}^\omega\omega$, where $\aleph_1 \leq \kappa < \mathfrak{p}$.

We include here a definition which we plan to investigate in a future paper (it is not used in the main line of our proof here).

Discussion 14.12. We note here that for a D a filter on \mathbb{N} (if D is the cofinite filter we may omit it), one may also define

- (1) for δ_1, δ_2 limit ordinals, normally regular cardinals, we say (\vec{f}^1, \vec{f}^2) is a D -weakly peculiar (δ_1, δ_2) -cut when
 - (a) $\vec{f}^1 = \langle f_\alpha^1 : \alpha < \delta_1 \rangle$ is $<_D$ -increasing;
 - (b) $\vec{f}^2 = \langle f_\beta^2 : \beta < \delta_2 \rangle$ is $<_D$ -decreasing;
 - (c) $f_\alpha^1 <_D f_\beta^2$ if $\alpha < \delta_1, \beta < \delta_2$;
 - (d) for no $A \in D$ and $f \in {}^\omega\omega$ do we have

$$\alpha < \delta_1 \wedge \beta < \delta_2 \text{ implies } f_\alpha^1 <_D f <_D f_\beta^2.$$

(2) we define

$$\mathcal{C}(D, {}^\omega\omega) = \{(\kappa_1, \kappa_2) : \text{there is a } D\text{-weakly peculiar } (\kappa_1, \kappa_2)\text{-cut, } \kappa_1, \kappa_2 \text{ regular}\}.$$

In a work in preparation we will show that the lower cofinality exists for $\kappa < \mathfrak{p}$ in, e.g., $\mathcal{C}({}^\omega\omega)$.

We now connect the cut existence from Theorem G with the existence of a cut in the generic ultrapower \mathcal{N} .

Claim 14.13. Working in $\mathbf{V}[\mathbf{G}]$, suppose $\mathfrak{p} < \mathfrak{t}$ and let \mathcal{N}, \mathbf{s} be as above. Then for some regular κ_1 with $\aleph_1 \leq \kappa_1 < \mathfrak{p}$, we have that $(\kappa_1, \mathfrak{p}) \in \mathcal{C}(\mathbf{s}, \mathfrak{t})$.

Remark 14.14. We use Claim 14.13 in the case where $\kappa_1 < \mathfrak{p} = \kappa_2$, but it holds generally.

Proof of Claim 14.13. Let \mathcal{N} be as above. We will prove that \mathcal{N} has a (κ_1, κ_2) -cut for some regular κ_1, κ_2 with $\aleph_1 \leq \kappa_1 < \kappa_2 = \mathfrak{p}$. The construction will show that this cut is in a pseudo-finite linear order of \mathcal{N} , in the sense of 14.4 above. As we have assumed that $\mathfrak{p} < \mathfrak{t}$, this will suffice to prove that $(\kappa_1, \mathfrak{p}) \in \mathcal{C}(\mathfrak{s}, \mathfrak{t})$.

Let $(\langle g_i : i < \kappa_2 \rangle, \langle f_i : i < \kappa_1 \rangle)$ be as in Theorem G, i.e., a (κ_1, κ_2) -peculiar cut in ${}^\omega\omega$. Working in $\mathbf{V}[\mathbf{G}]$, consider in \mathcal{N} the set

$$I = \prod_{n < \omega} [0, f_0(n)]/\mathbf{G}$$

with the usual linear order, i.e., the order \leq_I induced on the generic ultrapower by the factors. Note that the product is in \mathbf{V} , though \mathbf{G} is not. Clearly I is pseudo-finite in \mathcal{N} . (In other words, recalling 14.4 and 14.6, there is a nontrivial $\mathfrak{a} = (I, \leq_I, \dots) = (X_{\mathfrak{a}}, <_{\mathfrak{a}}, \dots) \in \text{Or}(\mathfrak{s})$.) Moreover,

- $i < j < \kappa_1$ implies $g_i/\mathbf{G}, g_j/\mathbf{G} \in I = X_{\mathfrak{a}}$ and $g_i/\mathbf{G} <_{\mathfrak{a}} g_j/\mathbf{G}$;
- $i < j < \kappa_2$ implies $f_i/\mathbf{G}, f_j/\mathbf{G} \in I = X_{\mathfrak{a}}$ and $f_i/\mathbf{G} <_{\mathfrak{a}} f_j/\mathbf{G}$;
- $i < \kappa_2, j < \kappa_1$ implies $g_i/\mathbf{G} <_{\mathfrak{a}} g_j/\mathbf{G}$.

So $(\langle g_i/\mathbf{G} : i < \kappa_2 \rangle, \langle f_i/\mathbf{G} : i < \kappa_1 \rangle)$ represents a pre-cut in $X_{\mathfrak{a}}$, and it will suffice to show that it represents a cut.

We carry out the remainder of the proof in \mathbf{V} . Assume for a contradiction that the conclusion fails; i.e., our pre-cut is not a cut. Then this failure is forced by some $B \in \mathbf{Q}, B \in \mathbf{G}$. That is, for some $h \in ({}^\omega\omega)^{\mathbf{V}}, B \Vdash_{\mathbf{Q}} "g_i/\mathbf{G} < h/\mathbf{G} < f_j/\mathbf{G}"$ for $i < \kappa_2, j < \kappa_1$. Then B is infinite, and $i < \kappa_2$ implies that $\{n : g_i(n) < h(n)\} \supseteq^* B$, as otherwise (recalling the definition of \mathbf{Q}) there is $B_1 \geq_{\mathbf{Q}} B, B_1 \Vdash g_i/\mathbf{G} \geq h/\mathbf{G}$. Likewise, $j < \kappa_1$ implies that $\{n : h(n) < f_j(n)\} \supseteq^* B$. This contradicts Definition 14.11 and so completes the proof. \square

Conclusion 14.15. *In Claim 14.13 we have shown that if $\mathfrak{p} < \mathfrak{t}$ (in \mathbf{V}), then*

- (1) $\Vdash_{\mathbf{Q}} "\mathcal{N}$ has a (κ_1, κ_2) -cut for some $\kappa_1 < \kappa_2 = \mathfrak{p}"$.
- (2) In $\mathbf{V}[\mathbf{G}]$ for some regular κ_1, κ_2 with $\aleph_1 \leq \kappa_1 < \kappa_2 = \mathfrak{p}, (\kappa_1, \kappa_2) \in \mathcal{C}(\mathfrak{s}, \mathfrak{t})$, thus $(\kappa_1, \kappa_2) \in \mathcal{C}(\mathfrak{s}, \mathfrak{t}_{\mathfrak{s}})$. In particular, $\mathcal{C}(\mathfrak{s}, \mathfrak{t}_{\mathfrak{s}}) \neq \emptyset$.

Proof. So there is no confusion about the assumption, recall by 14.6 that $\mathfrak{p}^{\mathbf{V}} < \mathfrak{t}^{\mathbf{V}}$ iff $\mathfrak{p}^{\mathbf{V}[\mathbf{G}]} < \mathfrak{t}^{\mathbf{V}[\mathbf{G}]}$. Then both (1) and (2) are immediate from 14.13, noting in the case of (2) that $\mathfrak{t} \leq \mathfrak{t}_{\mathfrak{s}}$ by 14.7. \square

We now prove Theorem 14.1.

Theorem 14.1. $\mathfrak{p} = \mathfrak{t}$.

Proof. It follows from the definitions that $\mathfrak{p} \leq \mathfrak{t}$. Suppose, in \mathbf{V} , that $\mathfrak{p} < \mathfrak{t}$. Working now in $\mathbf{V}[\mathbf{G}]$, let \mathfrak{s} be the cofinality spectrum problem from 14.6. By Conclusion 14.9, which does not assume $\mathfrak{p} < \mathfrak{t}, \mathcal{C}(\mathfrak{s}, \mathfrak{t}_{\mathfrak{s}}) = \emptyset$. By Conclusion 14.15(2), which does assume $\mathfrak{p} < \mathfrak{t}, \mathcal{C}(\mathfrak{s}, \mathfrak{t}_{\mathfrak{s}}) \neq \emptyset$, a contradiction. So necessarily $\mathfrak{p} = \mathfrak{t}$, which completes the proof. \square

ACKNOWLEDGMENTS

The authors would like to warmly thank the anonymous referee for thoughtful, extensive reports which significantly improved the presentation of this paper.

REFERENCES

- [1] James Ax, *The elementary theory of finite fields*, Ann. of Math. (2) **88** (1968), 239–271. MR0229613 (37 #5187)
- [2] James Ax and Simon Kochen, *Diophantine problems over local fields. I*, Amer. J. Math. **87** (1965), 605–630. MR0184930 (32 #2401)
- [3] Bohuslav Balcar, Jan Pelant, and Petr Simon, *The space of ultrafilters on \mathbf{N} covered by nowhere dense sets*, Fund. Math. **110** (1980), no. 1, 11–24. MR600576 (82c:54003)
- [4] Murray G. Bell, *On the combinatorial principle $P(c)$* , Fund. Math. **114** (1981), no. 2, 149–157. MR643555 (83e:03077)
- [5] Andreas Blass, *Combinatorial cardinal characteristics of the continuum*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 395–489, DOI 10.1007/978-1-4020-5764-9_7. MR2768685
- [6] G. Cantor, *Über eine Eigenschaft des Inbegriffes aller reellen algebraischen Zahlen*, Journal für die reine und angewandte Mathematik (Crelle's journal) **77** (1874), no. 2, 258–262.
- [7] Alan Dow, *Good and OK ultrafilters*, Trans. Amer. Math. Soc. **290** (1985), no. 1, 145–160, DOI 10.2307/1999788. MR787959 (86f:54044)
- [8] Eric K. van Douwen, *The integers and topology*, Handbook of set-theoretic topology, North-Holland, Amsterdam, 1984, pp. 111–167. MR776622 (87f:54008)
- [9] Mirna Džamonja and Saharon Shelah, *On \triangleleft^* -maximality*, Ann. Pure Appl. Logic **125** (2004), no. 1–3, 119–158, DOI 10.1016/j.apal.2003.11.001. MR2033421 (2004j:03044)
- [10] T. E. Frayne, D. S. Scott, and A. Tarski, *Reduced products*, Not. Amer. Math. Soc. **5** (1958), 673–674.
- [11] D. H. Fremlin, *Consequences of Martin's axiom*, Cambridge Tracts in Mathematics, vol. 84, Cambridge University Press, Cambridge, 1984. MR780933 (86i:03001)
- [12] Haim Gaifman and Constantine Dimitracopoulos, *Fragments of Peano's arithmetic and the MRDP theorem*, Logic and algorithmic (Zurich, 1980), Monograph. Enseign. Math., vol. 30, Univ. Genève, Geneva, 1982, pp. 187–206. MR648303 (83j:03095)
- [13] F. Hausdorff, *Summen von \aleph_1 Mengen*, Fund. Math. **26** (1936), 241–255.
- [14] H. Jerome Keisler, *Good ideals in fields of sets*, Ann. of Math. (2) **79** (1964), 338–359. MR0166105 (29 #3383)
- [15] H. Jerome Keisler, *Ultraproducts and saturated models*, Nederl. Akad. Wetensch. Proc. Ser. A 67 = Indag. Math. **26** (1964), 178–186. MR0168483 (29 #5745)
- [16] H. Jerome Keisler, *Ultraproducts which are not saturated*, J. Symbolic Logic **32** (1967), 23–46. MR0218224 (36 #1312)
- [17] Simon Kochen, *Ultraproducts in the theory of models*, Ann. of Math. (2) **74** (1961), 221–261. MR0138548 (25 #1992)
- [18] Sabine Koppelberg, *Cardinalities of ultraproducts of finite sets*, J. Symbolic Logic **45** (1980), no. 3, 574–584, DOI 10.2307/2273424. MR583375 (81i:04006)
- [19] Linus Kramer, Saharon Shelah, Katrin Tent, and Simon Thomas, *Asymptotic cones of finitely presented groups*, Adv. Math. **193** (2005), no. 1, 142–173, DOI 10.1016/j.aim.2004.04.012. MR2132762 (2006d:20075)
- [20] Kenneth Kunen, *Ultrafilters and independent sets*, Trans. Amer. Math. Soc. **172** (1972), 299–306. MR0314619 (47 #3170)
- [21] Jerzy Łoś, *Quelques remarques, théorèmes et problèmes sur les classes définissables d'algèbres* (French), Mathematical interpretation of formal systems, North-Holland Publishing Co., Amsterdam, 1955, pp. 98–113. MR0075156 (17,700d)
- [22] M. Malliaris, *Persistence and regularity in unstable model theory*, Ph. D. thesis, University of California, Berkeley, 2009.
- [23] M. E. Malliaris, *Realization of ϕ -types and Keisler's order*, Ann. Pure Appl. Logic **157** (2009), no. 2–3, 220–224, DOI 10.1016/j.apal.2008.09.008. MR2499710 (2010b:03044)
- [24] M. E. Malliaris, *Hypergraph sequences as a tool for saturation of ultrapowers*, J. Symbolic Logic **77** (2012), no. 1, 195–223, DOI 10.2178/jsl/1327068699. MR2951637

- [25] M. E. Malliaris, *Independence, order, and the interaction of ultrafilters and theories*, Ann. Pure Appl. Logic **163** (2012), no. 11, 1580–1595, DOI 10.1016/j.apal.2011.12.010. MR2959661
- [26] M. Malliaris and S. Shelah, *Constructing regular ultrafilters from a model-theoretic point of view*, Trans. Amer. Math. Soc. (2015).
- [27] M. Malliaris and S. Shelah, *Model-theoretic properties of ultrafilters built by independent families of functions*, J. Symbolic Logic **79** (2014), no. 1, 103–134, DOI 10.1017/jsl.2013.28. MR3226014
- [28] M. Malliaris and S. Shelah, *A dividing line within simple unstable theories*, Adv. Math. **249** (2013), 250–288, DOI 10.1016/j.aim.2013.08.027. MR3116572
- [29] M. Malliaris and S. Shelah, *Saturating the random graph with an independent family of small range*, Logic Without Borders (A. Hirvonen, J. Kontinen, R. Kossak, and A. Villaveces, eds.), DeGruyter, 2015, pp. 422.
- [30] M. Malliaris and S. Shelah, *An axiomatic approach to cofinality spectrum problems*. Manuscript F1245.
- [31] M. Malliaris and S. Shelah, *Existence of optimal ultrafilters and the fundamental complexity of simple theories*. arXiv:1404.2919.
- [32] M. Malliaris and S. Shelah, *Keisler’s order has infinitely many classes*. arXiv:1503.08341.
- [33] M. Malliaris and S. Shelah, *Model-theoretic applications of cofinality spectrum problems*. Manuscript 1051, arXiv:math.LO/1503.08338.
- [34] Zbigniew Piotrowski and Andrzej Szymański, *Some remarks on category in topological spaces*, Proc. Amer. Math. Soc. **101** (1987), no. 1, 156–160, DOI 10.2307/2046568. MR897088 (88g:54007)
- [35] A. J. Wilkie and J. B. Paris, *On the scheme of induction for bounded arithmetic formulas*, Ann. Pure Appl. Logic **35** (1987), no. 3, 261–302, DOI 10.1016/0168-0072(87)90066-2. MR904326 (89g:03087)
- [36] F. Rothberger and J. B. Paris, *Une remarque concerenant l’hypothèse du continu*, Fund. Math. **31** (1939), no. 3, 224–226.
- [37] Fritz Rothberger, *On some problems of Hausdorff and of Sierpiński*, Fund. Math. **35** (1948), 29–46. MR0029958 (10,689a)
- [38] Saharon Shelah, *Every two elementarily equivalent models have isomorphic ultrapowers*, Israel J. Math. **10** (1971), 224–233. MR0297554 (45 #6608)
- [39] Saharon Shelah, *Simple unstable theories*, Ann. Math. Logic **19** (1980), no. 3, 177–203, DOI 10.1016/0003-4843(80)90009-1. MR595012 (82g:03055)
- [40] Saharon Shelah, *Classification theory and the number of nonisomorphic models*, Studies in Logic and the Foundations of Mathematics, vol. 92, North-Holland Publishing Co., Amsterdam–New York, 1978. MR513226 (81a:03030)
- [41] S. Shelah, *Classification theory and the number of nonisomorphic models*, 2nd ed., Studies in Logic and the Foundations of Mathematics, vol. 92, North-Holland Publishing Co., Amsterdam, 1990. MR1083551 (91k:03085)
- [42] Saharon Shelah, *Toward classifying unstable theories*, Ann. Pure Appl. Logic **80** (1996), no. 3, 229–255, DOI 10.1016/0168-0072(95)00066-6. MR1402297 (97e:03052)
- [43] Saharon Shelah, *Two cardinal invariants of the continuum ($\mathfrak{d} < \mathfrak{a}$) and FS linearly ordered iterated forcing*, Acta Math. **192** (2004), no. 2, 187–223, DOI 10.1007/BF02392740. MR2096454 (2005f:03074)
- [44] Saharon Shelah, *A comment on “ $\mathfrak{p} < \mathfrak{t}$ ”*, Canad. Math. Bull. **52** (2009), no. 2, 303–314, DOI 10.4153/CMB-2009-033-4. MR2518968 (2010h:03078)
- [45] Saharon Shelah and Alexander Usvyatsov, *More on SOP_1 and SOP_2* , Ann. Pure Appl. Logic **155** (2008), no. 1, 16–31, DOI 10.1016/j.apal.2008.02.003. MR2454629 (2009k:03045)
- [46] Stevo Todorćević, *Coherent sequences*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 215–296, DOI 10.1007/978-1-4020-5764-9_4. MR2768682
- [47] S. Todorćević and B. Velićković, *Martin’s axiom and partitions*, Compos. Math. **63** (1987), 391–408.
- [48] B. Tsaban, *Selection principles and special sets of reals* (E. Pearl, ed.), Elsevier, New York, 2007.
- [49] Jerry E. Vaughan, *Small uncountable cardinals and topology*, Open problems in topology, North-Holland, Amsterdam, 1990, pp. 195–218. With an appendix by S. Shelah. MR1078647

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVENUE,
CHICAGO, ILLINOIS 60637

E-mail address: mem@math.uchicago.edu

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM, THE
HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL, AND DEPARTMENT OF
MATHEMATICS, HILL CENTER - BUSCH CAMPUS, RUTGERS, THE STATE UNIVERSITY OF NEW
JERSEY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NEW JERSEY 08854-8019

E-mail address: shelah@math.huji.ac.il