

CARSLAW, (a) *Introd. to the Theory of Fourier's Series and Integrals*, 3rd ed. rev. and enl., London, 1930, p. 293-296; (b) "A historical note on Gibbs' phenomenon in Fourier's series and integrals," *Amer. Math. So., Bull.*, v. 31, 1925, p. 420-424; and also *Encycl. d. math. Wissen.*, v. II.3.2, p. 1203f.

R.C.A.

¹D. H. L. writes as follows: This value has a last-figure error; in fact $K = 1.1789\ 79744\ 47216\ 72702\ 32029$. It is interesting to note that Zygmund's value is $\frac{1}{2}(K + 1)$. A value of $k = Si(\pi)$ is given to 16S in NYMTP, *Table of Sine, Cosine and Exponential Integrals*, v. 2, 1940, p. 206. From this it may be seen that Corey's value, referred to later, is also in error in the last figure, for 6 —, read 7. $k = 1.851\ 937051\ 982466$.

QUERY

19. THE INTEGRAL $\int_0^x e^{-A \cos \theta} d\theta$.—This integral arises in radium therapy discussion, and since tables of the function are so important for calculating the intensity of rod-shaped preparations, ROLF M. SIEVERT published such tables in his memoir, "Die v -Strahlungsintensität an der Oberfläche und in der nächsten Umgebung von Radiumnadeln," *Acta Radiologica*, Copenhagen, v. 11, 1930, p. 249-301. The tables on p. 271-280 are for $x = 30^\circ(1^\circ)90^\circ$, $A = [.1(.01).5; 3D]$. In the recent work, C. W. WILSON, *Radium Therapy, its Physical Aspects*, London, Chapman & Hall, 1945, p. 213-214, there is an abridgment of these tables for $x = 30^\circ(1^\circ)90^\circ$, $A = [0(.05).4; 3D]$. Current work connected with integrated radiation from a line source of radioactive material suggests the great desirability of extension of Sievert's table for $x < 30^\circ$, and for $A > .5$. Have other tables of the integral been published?

ROBLEY D. EVANS

Dept. of Physics
Massachusetts Institute of Technology

QUERIES—REPLIES

25. BRIGGS' ARITHMETICA LOGARITHMICA (Q7, v. 1, p. 170; QR21, v. 2, p. 94).—In the library of the University of Michigan is a copy of this volume with the extra 12 pages described in the query.

LOUIS C. KARPINSKI

Univ. of Michigan

26. SCARCE MATHEMATICAL TABLES (Q2, v. 1, p. 66; QR5, p. 100; 6, p. 132).—Four libraries have already been noted where HENRY GOODWYN, *A Table of the Circles*. . ., 1823, may be consulted. We may now add that copies are also available in the libraries of Brown University and of L. J. C.

CORRIGENDA

V. 1

- P. 215, B₆ 1, for $s = 1(1)50$, read $s = 1(1)150$.
 P. 220, A₁ 1, for 15D, read 15-20D; for 21.5, read 25.5. A₂ 1, delete δ .
 P. 221, A₃ 5, for $x/8$, read $1/(2x)$.
 P. 223, B₂ 5, for $.5(.1)1$, read $.5, .6, .8(.1)1$.
 P. 226, A₁ 4, delete Δ ; A₁ 8, for $0(.01)1$, read $0(.01)5.1$.
 P. 229, B₂ 10, for $(9 + x^2)$, read $(9 + x^2)^{\frac{1}{2}}$.