number of primes in the above expression for h_N. It should be borne in mind that the approximant vanishes at the endpoints of the interval $[0, \pi]$; consequently if the approximant does not have this property, we should modify it accordingly; this may involve subtracting a linear trend as suggested in similar circumstances by Lanczos [3, p. 236].

University of Kentucky
Lexington, Kentucky 40506

1. P. J. Davis, *Interpolation and Approximation*, Blaisdell, New York, 1963. MR **28** #393.
2. I. P. Natanson, *Konstrucktive Funktionentheorie*, Akademie-Verlag, Berlin, 1955. MR **16**, 1100.
3. C. Lanczos, *Applied Analysis*, Prentice-Hall, Englewood Cliffs, N. J., 1956. MR **18**, 823.

A Note on Best Approximation in E^n†

By J. T. Day

Let D be a closed convex set with positive volume V in Euclidean n-dimensional space. Let f be a nonnegative function of class C^2 on D (see [2]), and Q be a linear polynomial on D, i.e.

$$Q(x) = a_0 + a_1x_1 + a_2x_2 + \cdots + a_nx_n, \qquad x \in D.$$

We consider the problem of "best" one sided approximation of f by Q in the sense that among all linear functions $Q(x)$ satisfying

$$Q(x) \leqq f(x), \qquad x \in D, \tag{1}$$

we are looking for that one which maximizes $\int_D Q\,dx$.

THEOREM 1. *The problem under consideration has a unique solution given by the tangent plane through the centroid p of D, provided that the eigenvalues of the Hessian matrix $(f_{ij}(x))$, $x \in D$, are nonnegative.*

The proof is by construction. Let the centroid p of D have cartesian coordinates $(p_1, p_2, \cdots, p_n)$. Then

$$\int_D Q\,dx = V\cdot Q(p_1, p_2, \cdots, p_n) \tag{2}$$

for all linear polynomials Q (see [3]). Since $Q(p) \leqq f(p)$, we choose $Q^*(p) = f(p)$. Choose $Q_1^*(p) = f_1(p)$, $Q_2^*(p) = f_2(p), \cdots, Q_n^*(p) = f_n(p)$. Here $f_1(x) = (\partial f/\partial x_1)(x)$, etc. The above conditions determine $Q^*(x)$.

By Taylor's theorem we have $f(x) = Q^*(x) + R(x, p)$. The remainder $R(x, p)$ is nonnegative, since the eigenvalues of the Hessian matrix are nonnegative (see [2]). Thus $f(x) \geqq Q^*(x)$. We conclude that $Q^*(x)$ is a "best" approximate.

Suppose there were another "best" approximate $T(x)$. Then $T(p)$ must equal $f(p)$. Consider a point $x = (x_1, p_2, \cdots, p_n)$ where $x_1 > p_1$. By Taylor's theorem we have

Received October 11, 1965.
† This work was sponsored by the Mathematics Research Center under Contract No. DA-11-022-ORD-2059.

(3) $f(x) = f(p) + f_1(p)(x_1 - p_1) + f_{11}(p_1 + sh, p_2, \cdots, p_n)(x_1 - p_1)^2/2.$

Here $h = x_1 - p_1$, $0 < s < 1$.

(4) $$T(x) = f(p) + T_1(p)(x_1 - p_1).$$

Since $f(x) \geqq T(x)$, we find that

(5) $$f_1(p) - T_1(p) + f_{11}(p_1 + sh, p_2, \cdots, p_n)(x_1 - p_1)/2 \geqq 0.$$

The quantity $f_1(p) - T_1(p)$ must be nonnegative, for otherwise we could choose $(x_1 - p_1)$ so small that (5) could not hold. (We note here $f_{11}(x) \geqq 0$ for $x \in D$ by hypothesis.) A similar consideration in the case where $p_1 > x_1$ shows that $f_1(p) - T_1(p) \leqq 0$. Hence $f_1(p) = T_1(p)$. In the same manner one can show that $f_i(p) = T_i(p)$, $i = 2, \cdots, n$. Thus $Q^*(x)$ and $T(x)$ are identical.

The idea for this note occurred to the author after hearing a lecture by Prof. Ranko Bojanic [1] on "best" one sided approximation in the case of functions of one variable.

Mathematics Research Center, U. S. Army
University of Wisconsin
Madison, Wisconsin, 53706

1. R. Bojanic, "On polynomials of best one sided approximation." (To appear.)
2. W. H. Fleming, *Functions of Several Variables*, Addison-Wesley, Reading, Mass., 1965. MR **30** #4875.
3. A. H. Stroud, "Quadrature methods for functions of more than one variable," *Ann. New York Acad. Sci.*, v. 86, 1960, p. 784. MR **22** #10179.

A Close Approximation Related to the Error Function*

By Roger G. Hart

A function has been found that closely approximates the integral function

$$F(x) = \int_x^\infty \exp(-t^2/2)\, dt$$

for all real values of x.

Let

$$P(x) = \frac{\exp(-x^2/2)}{x}\left[1 - \frac{(1 + bx^2)^{1/2}/(1 + ax^2)}{P_0x + [P_0^2x^2 + \exp(-x^2/2)(1 + bx^2)^{1/2}/(1 + ax^2)]^{1/2}}\right]$$
$$\equiv P_0 + x^{-1}\{\exp(-x^2/2) - [P_0^2x^2 + \exp(-x^2/2)(1 + bx^2)^{1/2}/(1 + ax^2)]^{1/2}\},$$

where $P_0 = (\pi/2)^{1/2} \cong 1.253314137$,

$$a = \frac{1 + (1 - 2\pi^2 + 6\pi)^{1/2}}{2\pi} \cong .212023887,$$

Received January 5, 1966.

* This work was performed under the auspices of the U. S. Atomic Energy Commission.