

# On the Numerical Computation of Eigenvalues and Eigenvectors of Symmetric Integral Equations

By Peter Linz\*

**Abstract.** The well-known error estimates for the numerical computation of eigenvalues of symmetric integral equations are extended to the computation of the eigenvectors. The results are used to justify the application of an improvement method to obtain an efficient algorithm for solving the eigenvalue problem.

1. **Introduction.** The eigenvalues and eigenvectors of the integral equation

$$(1) \quad \lambda_n u_n(x) = K u_n(x) \equiv \int_0^1 K(x, t) u_n(t) dt,$$

with symmetric kernel  $K(x, t) = K(t, x)$ , may be approximated numerically by replacing the integral in Eq. (1) by a numerical quadrature. The corresponding equation is

$$(2) \quad \Lambda_n v_n(x) = \sum_{i=1}^N w_i K(x, x_i) v_n(x_i),$$

where  $\Lambda_n$  and  $v_n(x)$  are approximations to  $\lambda_n$  and  $u_n(x)$ , respectively. We will make the usual assumptions that  $w_i > 0$  and  $\sum_{i=1}^N w_i = 1$ . Satisfying Eq. (2) at the points  $x_i$ , we obtain the matrix eigenvalue problem

$$(3) \quad \Lambda_n \mathbf{v}_n = A \mathbf{v}_n,$$

where  $\mathbf{v}_n$  is a vector with components  $v_n(x_i)$ , and  $A$  a matrix with elements

$$a_{ij} = w_j K(x_i, x_j).$$

$A$ , while generally not symmetric, can be symmetrized by a simple similarity transformation, and we will assume that this has been done. For simplicity, we will also assume that the eigenvalues of Eq. (1) are nondegenerate.

Error estimates for the computed eigenvalues have been considered by a number of authors (Wielandt [5], Brakhage [1], Keller [3]). Their results show that the order of accuracy of the computed eigenvalues is the same as the order of accuracy of the chosen quadrature formula. In this paper we show that similar estimates hold for the computed eigenvectors. These estimates are then used as a justification for a method for improving the accuracy of the eigenvalues. In practice this implies that we can obtain highly accurate eigenvalues without having to solve the eigenvalue problem for large matrices.

---

Received December 8, 1969, revised April 14, 1970.

AMS 1969 subject classifications. Primary 6540, 6575; Secondary 4512.

Key words and phrases. Integral equations, eigenvalues, eigenvectors, numerical computation of eigenvalues, error estimates, Rayleigh quotient.

\* Current address: Department of Mathematics, University of California, Davis, California 95616.

Copyright © 1971, American Mathematical Society

2. **Error Estimates for the Eigenvalues.** Brakhage [1] showed that if  $\Lambda_k$  is an eigenvalue of Eq. (3), then there exists an eigenvalue  $\lambda_j$  of Eq. (1) such that

$$(4) \quad |\Lambda_k - \lambda_j| \leq \delta / (|\Lambda_k|^2 - \delta)^{1/2},$$

provided that  $|\Lambda_k|^2 > \delta$ , where

$$\delta = \max_{0 \leq x, s \leq 1} \left| \int_0^1 K(x, t) K(t, s) dt - \sum_{i=1}^N w_i K(x, x_i) K(x_i, s) \right|.$$

The largest eigenvalues are here associated with the smallest error bounds. Intuitively, one expects that the accuracy of the eigenvalue is determined by the smoothness of the associated eigenvector rather than by its magnitude. This is more apparent in Keller's formulation [3] which states that if  $\lambda_j$  is an eigenvalue of Eq. (1) then there exists an eigenvalue  $\Lambda_k$  of Eq. (3) such that

$$(5) \quad |\Lambda_k - \lambda_j| \leq c \max_{0 \leq x \leq 1} |\bar{e}_j(x)|,$$

where  $c$  is a constant (of order unity) and

$$\bar{e}_j(x) = \int_0^1 K(x, t) \bar{u}_j(t) dt - \sum_{i=1}^N w_i K(x, x_i) \bar{u}_j(x_i),$$

where  $\bar{u}_j$  is an eigenfunction normalized such that

$$\int_0^1 \bar{u}_j^2(x) dx = 1.$$

3. **Error Estimates for the Eigenvectors.** The results of the previous section can be extended to the computation of the eigenvectors. Let  $\mathbf{u}_k$  denote the vector with components  $u_k(x_i)$ . Assume that all the eigenvectors  $\mathbf{u}_k$  and  $\mathbf{v}_k$  ( $k = 1, 2, \dots, N$ ) are normalized such that

$$\|\mathbf{u}_k\| = \frac{1}{\sqrt{N}} \langle \mathbf{u}_k, \mathbf{u}_k \rangle^{1/2} \equiv \frac{1}{\sqrt{N}} \left\{ \sum_{i=1}^N u_k^2(x_i) \right\}^{1/2} = 1, \quad \|\mathbf{v}_k\| = 1.$$

Let  $\mathbf{e}_j$  denote the vector with components  $e_j(x_i)$ , where

$$e_j(x) = \int_0^1 K(x, t) u_j(t) dt - \sum_{i=1}^N w_i K(x, x_i) u_j(x_i).$$

**THEOREM 1.** *Let  $\lambda_j$  and  $u_j$  be a solution of Eq. (1), and let  $\Lambda_k$  be a solution of Eq. (3) such that  $|\lambda_j - \Lambda_k|$  satisfies the inequality (5). Then the eigenvector  $\mathbf{v}_k$  associated with  $\Lambda_k$  satisfies*

$$(6) \quad \|\mathbf{u}_j - \mathbf{v}_k\| \leq \frac{1}{\Delta_{jk}} \|\mathbf{e}_j\| + O(\|\mathbf{e}_j\|^2 / \Delta_{jk}^2),$$

where  $\Delta_{jk} = \min_{i \neq k} |\Lambda_k - \Lambda_i| - c \max |\bar{e}_j(x)|$ , provided that  $\Delta_{jk} > 0$ .

*Proof.* From Eq. (1) we have

$$(7) \quad \lambda_j \mathbf{u}_j = A \mathbf{u}_j + \mathbf{e}_j.$$

Since  $A$  is symmetric its eigenvectors are complete and orthogonal and we can write

$$\mathbf{u}_j = \mathbf{v}_k + \sum_{i=1}^N \alpha_{ji} \mathbf{v}_i, \quad \mathbf{e}_j = \sum_{i=1}^N \beta_{ji} \mathbf{v}_i.$$

Substituting into Eq. (7),

$$\lambda_j \mathbf{v}_k + \lambda_j \sum_{i=1}^N \alpha_{ji} \mathbf{v}_i = \Lambda_k \mathbf{v}_k + \sum_{i=1}^N \Lambda_i \alpha_{ji} \mathbf{v}_i + \sum_{i=1}^N \beta_{ji} \mathbf{v}_i$$

Because of the orthogonality of the  $\mathbf{v}_i$  we then have

$$\alpha_{jl} = \frac{\beta_{jl}}{\lambda_j - \Lambda_l}, \quad \text{for } l \neq k.$$

Now

$$\begin{aligned} |\lambda_j - \Lambda_l| &\geq ||\lambda_j - \Lambda_k| - |\Lambda_k - \Lambda_l|| \\ &\geq \Delta_{jk}, \quad \text{for } \Delta_{jk} > 0. \end{aligned}$$

Also,

$$||\mathbf{e}_j||^2 = \sum_{i=1}^N (\beta_{ji})^2,$$

so that

$$|\beta_{ji}| \leq ||\mathbf{e}_j||, \quad |\alpha_{jl}| \leq ||\mathbf{e}_j||/\Delta_{jk}, \quad l \neq k,$$

and since

$$||\mathbf{u}_j||^2 = (1 + \alpha_{jk})^2 + \sum_{i \neq k} (\alpha_{ji})^2 = 1, \quad \alpha_{jk} = O(||\mathbf{e}_j||^2/\Delta_{jk}^2).$$

Finally,

$$\begin{aligned} ||\mathbf{u}_j - \mathbf{v}_k|| &= \left\{ (\alpha_{jk})^2 + \sum_{i \neq k} (\alpha_{ji})^2 \right\}^{1/2} \\ &\leq \left\{ (\alpha_{jk})^2 + \frac{1}{\Delta_{jk}^2} \sum_{i \neq k} (\beta_{ji})^2 \right\}^{1/2} \leq \frac{1}{\Delta_{jk}} ||\mathbf{e}_j|| + O(||\mathbf{e}_j||^2/\Delta_{jk}^2). \end{aligned}$$

Thus, if  $\Lambda_k$  is well separated from the other computed eigenvalues the norm of the error is essentially proportional to the quadrature error. From (6) we have immediately that

$$(8) \quad |u_j(x_i) - v_k(x_i)| \leq \frac{\sqrt{N}}{\Delta_{jk}} ||\mathbf{e}_j|| + O(||\mathbf{e}_j||^2/\Delta_{jk}^2)$$

at the meshpoints  $x_i, i = 1, 2, \dots, N$ . If  $v_k(x)$  is computed by Eq. (2), then a similar estimate holds for all points of the range.

**THEOREM 2.** *If  $K(x, t)$  is bounded in the closed unit square,  $|K(x, t)| \leq L$ , then, for all  $x \in [0, 1]$ ,*

$$(9) \quad \begin{aligned} |u_j(x) - v_k(x)| &\leq \frac{1}{|\lambda_j|} \max |e_j(x)| + \frac{c}{|\Lambda_k|} \max |\bar{e}_i(x)| |u_j(x)| \\ &\quad + \frac{L\sqrt{N}}{|\Lambda_k| \Delta_{jk}} ||\mathbf{e}_j|| + O(||\mathbf{e}_j||^2/\Delta_{jk}^2), \end{aligned}$$

where  $c$  is the same constant as in Eq. (5).

*Proof.*

$$\begin{aligned}
 |u_i(x) - v_k(x)| &\leq \left| \frac{1}{\lambda_j} \int_0^1 K(x, t)u_i(t) dt - \frac{1}{\Lambda_k} \sum_{i=1}^N w_i K(x, x_i)v_k(x_i) \right| \\
 &\leq \left| \frac{1}{\lambda_j} \int_0^1 K(x, t)u_i(t) dt - \frac{1}{\lambda_j} \sum_{i=1}^N w_i K(x, x_i)u_i(x_i) \right| \\
 &\quad + \left| \frac{1}{\lambda_j} \sum_{i=1}^N w_i K(x, x_i)u_i(x_i) - \frac{1}{\Lambda_k} \sum_{i=1}^N w_i K(x, x_i)u_i(x_i) \right| \\
 &\quad + \left| \frac{1}{\Lambda_k} \sum_{i=1}^N w_i K(x, x_i)u_i(x_i) - \frac{1}{\Lambda_k} \sum_{i=1}^N w_i K(x, x_i)v_k(x_i) \right|
 \end{aligned}$$

and, applying inequalities (5) and (8), the estimate (9) follows.

**4. An Efficient Method for Computing the Eigenvalues.** We denote the inner product

$$\int_0^1 f(t)g(t) dt$$

by  $(f, g)$ . It is well known that if  $v_k(x)$  is an approximation to  $u_k(x)$  such that  $v_k(x) = u_k(x) + O(\Delta)$ , then an approximate eigenvalue may be computed by the Rayleigh quotient

$$(10) \quad \hat{\Lambda}_k = (v_k, Kv_k)/(v_k, v_k) = \lambda_k + O(\Delta^2).$$

Since a solution of Eq. (3) produces eigenvalues and eigenvectors with the same order of accuracy we can use the computed eigenvectors to improve the accuracy of the eigenvalues. Normally one will not be able to evaluate the required integrals in closed form, but Eq. (10) may be replaced by its discrete analogue

$$(11) \quad \bar{\Lambda}_k = \frac{\bar{v}_k^T A_M \bar{v}_k}{\bar{v}_k^T \bar{v}_k},$$

where  $A_M$  is the  $M \times M$  matrix with elements

$$(12) \quad a_{ij} = \bar{w}_i K(\bar{x}_i, \bar{x}_j),$$

and  $\bar{v}_k$  is the vector with components  $v_k(\bar{x}_i)$ ,  $\bar{x}_i$  being the quadrature points for the  $M$ -point quadrature formula, and  $\bar{w}_i$  the associated weights. The  $v_k(\bar{x}_i)$  are to be obtained from  $v_k(x_i)$  by means of Eq. (2). It can then be shown that

$$(13) \quad \bar{\Lambda}_k = \Lambda_k^M + O(\|e_j\|^2),$$

where  $\Lambda_k^M$  is the eigenvalue we would obtain by solving Eq. (3) with  $M$  quadrature points. If we want to find eigenvalues to a certain accuracy we can use the following procedure:

- (1) Choose a quadrature formula with  $N$  points and solve (3) to find the eigenvalues and eigenvectors. The  $N$  in this case can be kept considerably smaller than what we would have to choose if the results of (3) were to be used directly.
- (2) Use Eq. (11) with  $M > N$  to evaluate improved eigenvalues. This step involves matrix multiplications only. Provided our initial approximation to the eigenvalues is

TABLE 1

	Example 1	Example 2	Example 3	Example 4
$N = 2$	.42678	1.33222	.25000	.27192
$N = 10$	.40612	1.35214	.34343	.24407
$N = 20$	.40549	1.35281	.34641	.24324
$N = 30$	.40538	1.35293	.34697	.24309
$N = 50$	.40532	1.35299	.34725	.24301
$N = 2, M = 50$	.40507	1.35293	.34482	.24261
$N = 10, M = 50$	.40532	1.35299	.34725	.24300
exact $\lambda_0$	.40528	1.35303	.34741	.24296

sufficiently accurate, the final eigenvalues will be essentially as accurate as if we had solved (3) using  $M$  points.

Thus, we avoid what is frequently the most time consuming part of the computation, the solution of the eigenvalue problem (3) for large matrices.

**5. Numerical Examples.** In the following examples we computed the largest eigenvalue  $\lambda_0$  by the midpoint quadrature rule using values of  $N = 2, 10, 20, 30, 50$  with Eq. (3), and values of  $N = 2, 10, M = 50$  with the indicated improvement method.

*Example 1.*

$$K(x, t) = \min(x, t), \quad \lambda_0 = 4/\pi^2 = .40528.$$

*Example 2 (Brakhage [1]).*

$$K(x, t) = e^{xt}, \quad \lambda_0 = 1.35303.$$

*Example 3 (Bückner [2, p. 49]).*

$$K(x, t) = |x - t|, \quad \lambda_0 = .34741.$$

*Example 4 (Mikhlin [4, p. 86]).*

$$\begin{aligned} K(x, t) &= \frac{1}{2}x(2 - t), & x \leq t, \\ &= \frac{1}{2}t(2 - x), & x \geq t, \\ \lambda_0 &= .24296. \end{aligned}$$

Results are given in Table 1.

**Acknowledgments.** The work presented in this paper is supported by the AEC Computing and Applied Mathematics Center, Courant Institute of Mathematical Sciences, New York University, under Contract AT (30-1)-1480 with the U. S. Atomic Energy Commission.

Courant Institute of Mathematical Sciences  
 New York University  
 251 Mercer Street  
 New York, New York 10012

1. H. BRAKHAGE, "Zur Fehlerabschätzung für die numerische Eigenwertbestimmung bei Integralgleichungen," *Numer. Math.*, v. 3, 1961, pp. 174–179. MR 23 #2185.
2. H. BÜCKNER, *Die praktische Behandlung von Integralgleichungen*, Ergebnisse der angewandten Mathematik, Band 1, Springer-Verlag, Berlin, 1952. MR 14, 210.
3. H. B. KELLER, "On the accuracy of finite difference approximations to the eigenvalues of differential and integral operators," *Numer. Math.*, v. 7, 1965, pp. 412–419. MR 32 #6706.
4. S. G. MIHLIN, *Integral Equations and Their Applications to Some Problems of Mechanics, Mathematical Physics and Engineering*, GITTL, Moscow, 1949; English transl., Pergamon Press, New York, 1957. MR 12, 712; MR 19, 428.
5. H. WIELANDT, *Error Bounds for Eigenvalues of Symmetric Integral Equations*, Proc. Sympos. Appl. Math., vol. 6, Amer. Math. Soc., Providence, R. I., 1956, pp. 261–282. MR 19, 179.