

TABLE ERRATA

501.—K. Y. CHOONG, D. E. DAYKIN & C. R. RATHBONE, *Regular Continued Fractions for π and γ* , University of Malaya Computer Centre, September 1970, ms. deposited in UMT file. (See *Math. Comp.*, v. 25, 1971, p. 403, UMT 23.)

By a more extended calculation, the continued fraction herein for Euler's constant has been found to be correct to only the first 3251 partial quotients of the 3470 listed. This error affects the accompanying statistical table as well as Table 1 on p. 390 of the authors' related paper [1].

WILLIAM A. BEYER

Los Alamos Scientific Laboratory
Los Alamos, New Mexico 87544

1. K. Y. CHOONG, D. E. DAYKIN & C. R. RATHBONE, "Rational approximations to π ," *Math. Comp.*, v. 25, 1971, pp. 387–392.

502.—A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER & F. G. TRICOMI, *Tables of Integral Transforms*, Vol. I, McGraw-Hill Book Co., New York, 1954.

On p. 31, in Formula 17 the Fourier cosine transform of

$$[\cosh(\beta x) + \cos c]^{-1} \cosh(ax)$$

should read

$$\pi\beta^{-1} \csc c \{ \cos[a(\pi - c)/\beta] \cosh[y(\pi + c)/\beta] - \cos[a(\pi + c)/\beta] \cosh[y(\pi - c)/\beta] \} \\ \times [\cosh(2\pi y/\beta) - \cos(2\pi a/\beta)]^{-1}.$$

HENRY E. FETTIS

Aerospace Research Laboratories
Wright-Patterson Air Force Base, Ohio 54533

EDITORIAL NOTE: For notices of additional errata see *Math. Comp.*, v. 25, 1971, p. 199, MTE 472; v. 24, 1970, pp. 239–240, MTE 451; v. 23, 1969, p. 468, MTE 436 and the footnote thereto.

503.—I. S. GRADSHTEYN & I. M. RYZHIK, *Table of Integrals, Series, and Products*, 4th ed., Academic Press, New York, 1965.

On p. 527, formula 4.224(11) is incorrect for the cases $a^2 < 1$ and $a^2 > 1$. When $a > 0$, the common value of the integrals $\int_0^{\pi/2} \ln(1 + a \sin x)^2 dx$ and

$\int_0^{\pi/2} \ln(1 + a \cos x)^2 dx$ can be written as

$$\pi \ln(a/2) + 4G + 4S(b),$$

where G is Catalan's constant, $b = (1 - a)/(1 + a)$, and

$$S(b) = \sum_{k=1}^{\infty} \frac{b^k}{k} \sum_{n=1}^k \frac{(-1)^{n+1}}{2n-1}.$$

AKIN OJO
J. SADIKU

Departments of Physics and Mathematics
University of Ibadan
Ibadan, Nigeria

EDITORIAL NOTE: Alternatively, when $a^2 \leq 1$, the value of these integrals can be expressed as $\pi \ln(1 + (1 - a^2)^{1/2})/2 - 2 \sin^{-1} a \ln(1 + (1 - a^2)^{1/2})/a + 4 \text{Cl}_2(\sin^{-1} a) - \text{Cl}_2(2 \sin^{-1} a)$, where $\text{Cl}_2(x)$ is Clausen's integral. When $a \geq 1$, the value is $\pi \ln(a/2) + 4 \text{Cl}_2(\sin^{-1} 1/a) - \text{Cl}_2(2 \sin^{-1} 1/a)$.