

## On Truncatable Primes

By I. O. Angell and H. J. Godwin

**Abstract.** Truncatable primes are those that yield a sequence of primes when digits are removed always from the left or always from the right. The sizes of the largest truncatable primes in a given number base are estimated by a probabilistic argument and compared with computed values.

The number 357686312646216567629137 is a prime, and, if successive digits are removed from the left, a sequence of primes ending 137, 37, 7 is obtained. We call such a number a left-truncatable prime and this particular one is the largest in decimal notation. Similarly 73939133 is the largest right-truncatable prime to base 10: it yields a sequence of primes . . . 73, 7 if we truncate from the right. The number 1979339339 has been quoted [1] as the largest right-truncatable prime; but we adhere to the convention that 1 is not a prime number, and so exclude it.

We have computed  $L_a$ , the largest left-truncatable prime with base  $a$ , for  $3 \leq a \leq 11$ , and  $R_a$ , the largest right-truncatable prime with base  $a$ , for  $3 \leq a \leq 15$ . The results are as follows (in decimal form).

---

$a$	$R_a$	$L_a$
3	71	23
4	191	4091
5	2437	7817
6	108863	4836525320399
7	6841	817337
8	4497359	14005650767869
9	1355840309	1676456897
10	73939133	357686312646216567629137
11	6774006887	2276005673
12	18704078369	
13	38901772669	
14	6525460043032393259	
15	927920056668659	

---

To investigate the length of the largest truncatable primes we use the fact that the density of primes in the neighborhood of  $n$  is  $1/\log n$ . Consider first right-

---

Received June 2, 1976.

AMS (MOS) subject classifications (1970). Primary 10A25; Secondary 60–01, 60–04.

Key words and phrases. Primes, number bases, probability.

Copyright © 1977, American Mathematical Society

truncatable primes. Given such a prime  $p$ , with  $r$  digits to base  $a$ , we look for one with  $r + 1$  digits by testing which of  $ap, ap + 1, \dots, ap + a - 1$  are prime. (Actually we tested for pseudoprimality: for the longest pseudoprime found, true primality for it and its sequence of truncations was then checked, using tables or Lehmer's test.)

The expected number of primes among these is  $a/\log \mu$ , where  $ap \leq \mu \leq ap + a - 1$ , i.e.  $a^{r-1} < \mu < a^r$ . In practice, we should consider  $ap + t$  where  $t$  runs through numbers  $1, \dots, a - 1$  prime to  $a$ : if  $a$  has prime divisors  $q_1, q_2, \dots, q_n$ , this means that we have  $a \prod_{i=1}^n (1 - 1/q_i)$  choices; but now the probability of such a choice being prime is

$$\frac{1}{\log \mu} \cdot \prod_{i=1}^n \left(1 - \frac{1}{q_i}\right)^{-1},$$

and the same result is obtained.

If there are  $\pi(a)$  primes less than  $a$ , then the expected number of right-truncatable primes with  $s$  digits lies between

$$\pi(a) \prod_{r=1}^{s-1} \left(\frac{a}{r \log a}\right) \quad \text{and} \quad \pi(a) \prod_{r=1}^{s-1} \left(\frac{a}{(r+1) \log a}\right),$$

i.e. between

$$\frac{\pi(a)a^{s-1}}{(s-1)!(\log a)^{s-1}} \quad \text{and} \quad \frac{\pi(a)a^{s-1}}{s!(\log a)^{s-1}}.$$

Using Stirling's approximation to the factorial, these bounds become

$$\frac{s\pi(a) \log a}{\sqrt{2\pi s} a} \left(\frac{ae}{s \log a}\right)^s \quad \text{and} \quad \frac{\pi(a) \log a}{\sqrt{2\pi s} \cdot a} \left(\frac{ae}{s \log a}\right)^s.$$

These expressions are dominated by  $(ae/(s \log a))^s$ , for growth if  $s < ae/\log a$  and for decrease if  $s > ae/\log a$ . Hence a rough estimate of the maximum  $s$  is  $ae/\log a$ . For small values of  $s$  we can obtain a better estimate by replacing  $\log \mu$ , where  $a^{s-1} < \mu < a^s$ , by the average of  $\log x$  over  $(a^{s-1}, a^s)$ , i.e. by  $((sa - s + 1) \log a)/(a - 1) - 1$ : the estimate of the largest  $s$  is then the largest value for which the expected number of primes is not less than a half.

For left-truncatable primes,  $p$ , of length  $s$ , is followed by primes among  $a^s + p, 2 \cdot a^s + p, \dots, (a - 1)a^s + p$ . We now have  $(a - 1)$  choices; but all the numbers are prime to  $a$ , and so the expected number of primes is

$$\frac{(a-1)}{\log \mu} \prod_{i=1}^n \left(1 - \frac{1}{q_i}\right)^{-1}.$$

The crude estimate of  $s$  is

$$\frac{ae}{\log a} \prod_{i=1}^n \left(1 - \frac{1}{q_i}\right)^{-1};$$

and a better estimate can be obtained in a similar way to that used for right-truncatable primes, except that now  $q_1, \dots, q_n$ , give rise to no primes of length greater than one.

It will be noted that we have excluded 0 as a leading digit. If we do not do this, the length of the longest left-truncatable prime becomes indeterminate since, for example, there are, with probability one, infinitely many primes of the form  $10^k + 3$ .

The comparison between observed lengths and lengths given by the better estimates are as follows:

$a$	Length (in base $a$ ) of			
	$R_a$		$L_a$	
	observed	better estimate	observed	better estimate
	3	4	5	3
4	4	6	6	9
5	5	7	6	7
6	7	8	17	19
7	5	8	7	8
8	8	9	15	16
9	10	10	10	13
10	8	10	24	23
11	10	11	9	11
12	10	12		
13	10	12		
14	17	13		
15	13	14		

Department of Statistics and Computer Science  
 Royal Holloway College  
 Egham Hill, Egham  
 Surrey, TW20 0EX, England

1. H. J. CLARK, Letter in *Computer Weekly*, No. 360, September 27, 1973, p. 26.