

TABLE ERRATA

574.—A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER & F. G. TRICOMI, *Higher Transcendental Functions*, vol. 2, McGraw-Hill Book Co., New York, 1953.

On p. 103, the right side of formula 52 should read in part:

$$-\frac{1}{2} + \pi^{1/2} x^{-1} \frac{\Gamma(\nu + 1)}{\Gamma\left(\nu + \frac{1}{2}\right)} \left(1 - \frac{t^2}{x^2}\right)^{\nu-1/2}, \quad 0 < t < x < \pi.$$

The right side of formula 54 corresponding to $0 < x < t < \pi$ should read $-1/\pi^{1/2}$ instead of $-(\frac{1}{2} + \nu)/\pi^{1/2}$. The portion of this formula relating to the interval $0 < t < x < \pi$ is correct, but a simpler expression for this region is

$$-\frac{1}{\pi^{1/2}} + \frac{\pi^{1/2}(2\nu + 1)}{x} \int_0^{\cos^{-1}(t/x)} \sin^{2\nu} \theta \, d\theta,$$

which may be written in terms of the hypergeometric function as given, or as

$$-\frac{1}{\pi^{1/2}} + \frac{\pi\Gamma\left(\nu + \frac{3}{2}\right)}{x\Gamma(\nu + 1)} - \frac{\pi^{1/2}(2\nu + 1)t}{x^2} F\left(\frac{1}{2} - \nu, \frac{1}{2}, \frac{3}{2}; \frac{t^2}{x^2}\right).$$

These formulas have been reproduced as formulas (12) and (13) on p. 123 of [1], and accordingly the same corrections are applicable therein.

HENRY E. FETTIS

1885 California, Apt. 62
Mountain View, California 94041

1. V. MARGULIS, *Handbook of Series for Scientists and Engineers*, Academic Press, New York and London, 1965.

On p. 250, Eq. 11.5(17), which is Rodrigues' formula for the associated Legendre functions, should end with $(1 - x^2)^n$ instead of $(1 - x^2)^m$.

S. N. STUART

Division of Chemical Physics
CSIRO
P. O. Box 160
Clayton, Victoria, Australia 3168

EDITORIAL NOTE: For notices of additional errata in this volume see *Math. Comp.*, v. 30, 1976, pp. 675–676, MTE 524 and the editorial footnote thereto. Further errors in the book by Margulis are noted in *Math. Comp.*, v. 21, 1967, pp. 750–751, MTE 417.

575.—W. MAGNUS, F. OBERHETTINGER & R. P. SONI, *Formulas and Theorems for the Special Functions of Mathematical Physics*, third enlarged edition, Springer-Verlag, New York, 1966.

The following necessary typographical corrections have been noted.

| <i>page</i> | <i>line</i> | <i>for</i> | <i>read</i> |
|-------------|----------------------|------------------------------------|--------------------------------|
| 92 | 9 | a | α |
| 99 | 5 | b^ν | b^ν |
| 124 | -7 | 4.13.1 | 3.13.1 |
| 167 | -4 | :: | ; |
| 212 | 12 | $-\frac{2}{1+x}$ | $\frac{2}{1+x}$ |
| 213 | -8 |) ₃ |) _x |
| 214 | 6 | t | z |
| 217 | -6 | $\Sigma \Gamma(\underline{\quad})$ | $\Sigma (\underline{\quad})$ |
| 242 | 6, 7 | t | x |
| 250 | 12 | 4 | 12 |
| 252 | -7 | $e^{-x^2/2}$ | $e^{x^2/2}$ |
| 254 | 9 | $\Sigma_{m=0}^\infty$ | $\Sigma_{m=0}^n$ |
| 257 | -10 | U | U_n |
| 268 | -6 | $-az$ | $-aw$ |
| 285 | 13 | \int_z^∞ | $e^z \int_z^\infty$ |
| 327 | 3 | $e^{z^2/4}$ | $e^{-z^2/4}$ |
| 327 | 12 | $\sqrt{\pi}$ | $\sqrt{2\pi}$ |
| 332 | 2, 4 | e^{-imv} | e^{imv} |
| 332 | 9 | $\Sigma_{n=0}^\infty$ | $\Sigma_{n=0}^N$ |
| 339 | 6 | Erf | Erfc |
| 340 | 3 | e^x | e^{-x} |
| 342 | 2 | $\sqrt{\frac{\pi}{2}} a$ | $\sqrt{\frac{\pi}{2}} a^{1/2}$ |
| 342 | 12 (second integral) | e^{-t} | e^t |
| 347 | 2 | e^{-x} | e^x |
| 356 | -1 | a | n |

Furthermore, on p. 86, line 1 *delete a*, and on p. 229, line 9 *delete n*. On p. 93, line 7 in the right member of the equation *read* I_ν . Similarly, on p. 250, line 7 *read* $2^{n/2} He_n(x\sqrt{2})$. On p. 471, line -7 *read* $\Sigma_{l=0}^n$; in line -4 *read*

$$\Sigma_{l=0}^n ((-1)^l (n+l)! / (n-l)! (2l)! (2 \sin x)^{2l});$$

and on p. 493, line -10 *read* ϵ_n .

It should also be noted that the formula on p. 28, line -3 is incorrect.

N. M. TEMME

Mathematisch Centrum
2^{de} Boerhaavestraat 49
1091 AL Amsterdam
The Netherlands

On p. 132, the right side of the last formula should read (in part):

$$-\frac{1}{2} + \pi^{1/2} x^{-1} \frac{\Gamma(\nu+1)}{\Gamma\left(\nu + \frac{1}{2}\right)} \left(1 - \frac{t^2}{x^2}\right)^{\nu-1/2}, \quad 0 < t < x < \pi.$$

ROBERT L. PEXTON

Lawrence Livermore Laboratory
University of California
Livermore, California 94550

EDITORIAL NOTE: For notices of additional errors in this and earlier editions, see *Math. Comp.*, v. 34, 1980, p. 332, MTE 569 and the editorial footnote thereto.

576.—P. F. BYRD & M. D. FRIEDMAN, *Handbook of Elliptic Integrals for Engineers and Physicists*, 2nd rev. ed., Springer-Verlag, New York, 1971.

On p. 12 formula 115.01 gives expressions for $F(\theta \pm i\phi, k)$ and $E(\theta \pm i\phi, k)$ in terms of $F(\beta, k)$, $E(\beta, k)$, $F(A, k')$, and $E(A, k')$ with real arguments. However, the amplitudes A and β of the latter integrals are defined implicitly in terms of θ and ϕ . Explicit expressions for these quantities are:

$$\begin{aligned}\sin \beta &= 2 \sin \theta \left\{ \left[(1 + k \sin \theta \cosh \phi)^2 + (k \cos \theta \sinh \phi)^2 \right]^{1/2} \right. \\ &\quad \left. + \left[(1 - k \sin \theta \cosh \phi)^2 + (k \cos \theta \sinh \phi)^2 \right]^{1/2} \right\}^{-1}, \\ \sin A &= \tanh \phi / (1 - k^2 \sin^2 \beta)^{1/2}.\end{aligned}$$

On p. 39 the sections referenced in the footnote should be those numbered 813 and 814, and in Section 164.02 the left side of the last equation should read $\Pi(\phi, \alpha_1^2, k_1)$.

HENRY E. FETTIS

EDITORIAL NOTE: For previous notices of errata in this handbook see *Math. Comp.*, v. 26, 1972, p. 597, MTE 488 and the editorial footnote thereto.

577.—I. S. GRADSHTEYN & I. M. RYZHIK, *Table of Integrals, Series, and Products*, 4th ed., Academic Press, New York, 1965.

Further corrections required in this edition are the following:

P. xxxii: The lower limit of the integral defining $E_n(z)$ should be 1 instead of 0.

P. 34: Delete x from the second cosine term on the right side of formula 1.395(1).

P. 367: In formula 3.614 the conditions on the parameters should read $0 < b < a < 1, p = 1, 2, 3, \dots$ (In the source [1] of this formula, the relative size of a and b is not specified.) The alternative formula

$$\int_0^\pi \frac{\sin x \sin px \, dx}{(a^2 - 2ab \cos x + b^2)(1 - 2a^p \cos px + a^{2p})} = \frac{\pi a^{p-1}}{2b(b^p - a^{2p})}$$

holds in the less restrictive range $0 < a < 1, a^2 < b, p = 1, 2, 3, \dots$

P. 384: In the right member of formula 3.666(1), read $(\beta^2 - 1)^{\mu/2}$ in place of $(z^2 - 1)^{\mu/2}$.

P. 908: The right member of formula 8.128(3) should read $k[K(k) + iK'(k)]$, and all three formulas in 8.128 should carry the restriction $\text{Im}(k) < 0$.

P. 929: In formula 8.241(1), for $x > 1$ read $x < 1$.

P. 944: In formula 8.363(6) the second term on the right side should be $-\ln(2q)$.

P. 948: In formula 8.375(1) the summation symbol Σ' should be used in order to indicate that only one-half the last term is to be taken. An alternative form of this sum is

$$-2 \sum_{k=0}^{E(q/2)-1} \cos \frac{p(2k+1)\pi}{q} \ln \sin \frac{(2k+1)\pi}{q}.$$

The same error occurs in the source [2] of this formula. As noted in MTE 428 (*Math. Comp.*, v. 22, 1968, pp. 903–907), the range of p should be 1, 2, 3, . . . , $q - 1$. Also, the reference following this formula should be to 8.363(5)–(7).

P. 1020: In formula 8.835(3) the algebraic sign between the terms on the right side should be minus instead of plus.

P. 1067: In formula 9.254(2) a minus sign should be prefixed to the right side.

HENRY E. FETTIS

1. D. BIERENS DE HAAN, *Nouvelles Tables d'Intégrales Définies*, Hafner Publishing Co., New York, 1957.

2. N. NIELSEN, *Handbuch der Theorie der Gammafunktion*, Teubner, Leipzig, 1906.

EDITORIAL NOTE: For previous notices of errata in this edition see *Math. Comp.*, v. 33, 1979, p. 1377, MTE 565 and the editorial footnote thereto.

578.—HENRY E. FETTIS & JAMES C. CASLIN, *A Table of the Complete Elliptic Integral of the First Kind for Complex Values of the Modulus*, Part I, Report ARL 69-0172, Aerospace Research Laboratories, Office of Aerospace Research, United States Air Force, Wright-Patterson Air Force Base, Ohio, November, 1969. [See *Math. Comp.*, v. 24, 1970, pp. 993–994, RMT 76.]

Page 3: The second term on the right side of Eq. 7 should be

$$+ \frac{a \cdot b}{c \cdot 1!} z.$$

Page 16: In Eq. 49 the numerator of the right member should read $2\sqrt{\rho_n}$, and in Eq. 51 the numerator of the third term on the right side should be 2 instead of 3.

Page 20: In lines 2 and 6 replace k and k' by K and K' ; also in the formulas on lines 3 and 7.

An errata sheet distributed to the original recipients of this report gave the following typographical corrections:

Page 4, line 9: Read “relations” for “reactions.”

Page 10, line 11: Read, in part, “. . . $K(k)$ when k is outside the unit circle.”

Page 15, Eq. 46: Read ϕ_1 for ϕ .

Page 23, line -2: Read $w = \frac{1}{2} + iy$.

Page 24, line 5: Read τ -plane instead of -plane.

HENRY E. FETTIS

579.—E. T. WHITTAKER & G. N. WATSON, *A Course of Modern Analysis*, 4th ed., Cambridge Univ. Press, New York and London, 1927, and subsequent reprints.

On p. 289, the first equation should read:

$$\frac{\Gamma(a)\Gamma(b)}{\Gamma(c)} F(a, b; c; z) = \frac{\Gamma(a)\Gamma(b-a)}{\Gamma(c-a)} (-z)^{-a} F(a, 1-c+a; 1-b+a; z^{-1}) \\ + \frac{\Gamma(b)\Gamma(a-b)}{\Gamma(c-b)} (-z)^{-b} F(b, 1-c+b; 1-a+b; z^{-1}).$$

S. N. STUART

Division of Chemical Physics
CSIRO
P. O. Box 160
Clayton, Victoria, Australia 3168

EDITORIAL NOTE: For a previously announced error in this edition, see *Math. Comp.*, v. 33, 1979, p. 431, MTE 560.

580.—P. F. BYRD & M. D. FRIEDMAN, *Handbook of Elliptic Integrals for Engineers and Physicists*, Springer, New York and Berlin, 1953, 2nd rev. ed., 1971.

The correction noted in MTE 557 and MTE 559 (*Math. Comp.*, v. 33, 1979, pp. 430–431) should also be made in formula 911.01 of this handbook. The correct expansion for sn^2 can be obtained by differentiating formula 905.01, and, as noted by O. G. Ruehr (*SIAM Rev.*, v. 22, 1980, p. 234), is given in [1] (p. 25; formula 2.23) and in [2] (Section 22.735, Ex. 5).

HENRY E. FETTIS

1. F. OBERHETTINGER, *Fourier Expansions*, Academic Press, New York, 1973.
2. E. T. WHITTAKER & G. N. WATSON, *A Course of Modern Analysis*, Cambridge Univ. Press, fourth edition reprinted, 1973.

581.—H. H. GOLDSTINE, *A History of Numerical Analysis from the 16th through the 19th Century*, Springer, New York, 1977.

On page 304, lines 9 and 10, for

$$\frac{(x-b)^\beta(x-c)^\gamma \cdots (x-l)^\lambda}{(a-b)^\beta(a-c)^\gamma \cdots (a-l)^\lambda} \\ \cdot \left[f(a) + f'(a)(x-a) + \frac{1}{2!} f''(a)(x-a)^2 \right. \\ \left. + \cdots + \frac{1}{(\alpha-1)!} f^{(\alpha-1)}(a)(x-a)^{\alpha-1} \right],$$

read

$$(x-a)^\alpha(x-b)^\beta(x-c)^\gamma \cdots (x-l)^\lambda \\ \cdot \sum_{s=0}^{\alpha-1} \left\{ \sum_{j=s}^{\alpha-1} \frac{[(x-b)^\beta(x-c)^\gamma \cdots (x-l)^\lambda]^{(j-s)}}{s!(j-s)!(x-a)^{\alpha-j}} \right\} f^{(s)}(a).$$

Lines 7 and 8 are also incorrect, because the author overlooked the need for the expansions of $1/(z - b)^\beta$, $1/(z - c)^\gamma$, ..., $1/(z - l)^\lambda$ in powers of $z - a$ to be combined with the expansion of $f(z)/(x - z)$ in order to obtain the coefficient of $(z - a)^{\alpha-1}$. The original derivation [1] by Hermite of his osculatory interpolation formula does not have this error.

HERBERT E. SALZER

941 Washington Avenue
Brooklyn, New York 11225

1. *J. Reine Angew. Math.*, v. 84, 1878, pp. 70–79.

582.—I. S. GRADSHTEYN & I. M. RYZHIK, *Table of Integrals, Series, and Products*, 4th ed., Academic Press, New York, 1965.

On p. 371 formula 3.624(6) is valid only for integer values of a . This restriction is not stated.

For all real values of a the appropriate formula is

$$\int_0^{\pi/2} \left(\frac{\sin ax}{\sin x} \right)^2 dx = \frac{\pi a}{2} + \frac{1}{2} \sin a\pi \left\{ 1 + a \left[x \left(\frac{a}{2} \right) - x \left(\frac{1+a}{2} \right) \right] \right\},$$

where $\psi(z) = d \ln \Gamma(z) / dz$.

For a derivation of this result, see [1].

HENRY E. FETTIS

1. H. E. FETTIS, "On some trigonometric integrals," *Math. Comp.*, v. 34, 1980, pp. 1325–1329.