

## On the Bisection Method for Triangles

By Andrew Adler

**Abstract.** Let  $UVW$  be a triangle with vertices  $U$ ,  $V$ , and  $W$ . It is "bisected" as follows: choose a longest edge (say  $VW$ ) of  $UVW$ , and let  $A$  be the midpoint of  $VW$ . The  $UVW$  gives birth to two daughter triangles  $UVA$  and  $UWA$ . Continue this bisection process forever.

We prove that the infinite family of triangles so obtained falls into finitely many similarity classes, and we obtain sharp estimates for the longest  $j$ th generation edge.

**1. Introduction.** Let  $UVW$  be the triangle with vertices  $U$ ,  $V$ , and  $W$ . We "bisect" triangles as follows: choose a longest edge (say  $VW$ ) of  $UVW$ , and let  $A$  be the midpoint of  $VW$ . Then  $UVW$  gives birth to two daughter triangles  $UVA$  and  $UAW$ . So the generation 0 triangle  $UVW$  gives rise to two generation 1 triangles. "Bisect" these in turn, giving rise to four generation 2 triangles, and so on. So  $UVW$  through this process gives rise to an infinite family of triangles. This bisection process and a generalization to three dimensions have a number of numerical applications; see, e.g., [1], [3], [4].

Let  $m_j$  be the length of the longest  $j$ th generation edge. A bound for the rate of convergence of  $m_j$  has been obtained in [2]. Sharp estimates for certain classes of triangles have been given in [5]. In this paper we prove that  $m_j \leq \sqrt{3} 2^{-j/2} m_0$ , if  $j$  is even, and that  $m_j \leq \sqrt{2} 2^{-j/2} m_0$ , if  $j$  is odd, with equality for equilateral triangles. We prove, moreover, the following geometrically interesting fact: the (infinite) family of  $UVW$  contains only finitely many similarity types.

*Definition.* If  $\Delta$  is a triangle, then  $\phi(\Delta) = (\text{area of } \Delta)/l^2(\Delta)$ , where  $l(\Delta)$  is the length of the longest edge of  $\Delta$ .  $\mathcal{F}_0(\Delta)$  is the collection of even generation descendants of  $\Delta$ , and  $\mathcal{F}_1(\Delta)$  is the collection of odd generation descendants.

Since our bisection process in particular bisects areas, in order to find out about  $m_j$ , it is enough to know how the dimensionless quantity  $\phi(\Delta)$  behaves under bisection of triangles. Our results will be proved by an induction on  $\phi$ . It is necessary to deal first with acute angled triangles, then with obtuse triangles. The squares of side-lengths needed in this paper are all calculated by straightforward use of the law of cosines.

**2. Acute Triangles.** Let  $\Delta = UVW$  be an acute angled triangle, with  $VW$  a longest edge. Write  $\|UV\|^2 = p$ ,  $\|UW\|^2 = q$ . For convenience let  $\|VW\|^2 = 1$ , and assume  $p \leq q \leq 1$ . Bisect edge  $VW$  at  $A$ .  $UW$  is then the longest edge of  $UAW$ . Bisect it at  $B$ . Then  $UA$  is the longest edge of  $UBA$ . There are now three different possibilities to consider.

---

Received June 3, 1980.

1980 *Mathematics Subject Classification.* Primary 51M20; Secondary 65N50, 65L50.

©1983 American Mathematical Society  
0025-5718/82/0000-1065/\$02.00

*Possibility 1.*  $UV$  is a longest edge of  $UVA$ . Examination of Figure 1 shows that bisection of  $UAB$  and of  $UVA$  gives rise to triangles similar to already occurring triangles, and so (up to similarity)  $\mathfrak{F}_0(\Delta)$  contains only  $UVW$  and  $UAB$ , while  $\mathfrak{F}_1(\Delta)$  only contains  $UVA$  and  $UAW$ .




FIGURE 1

Since  $\|UA\|^2 = \frac{1}{4}(2p + 2q - 1)$ ,  $\phi(UAB) = \phi(\Delta)/2p + 2q - 1$ . But since  $p + q \geq 1$  (from the acuteness of  $\Delta$ ), using elementary linear programming, we find that  $\phi(UAB) \geq \frac{1}{3}\phi(\Delta)$ , with equality if  $\Delta$  is equilateral. It is easy to see that  $\phi(UVA)$  and  $\phi(UAW)$  are both  $\geq \frac{1}{2}\phi(\Delta)$ .

Since  $\Delta$  is acute,  $\|AV\| \leq \|AU\|$ , so if Possibility 1 does not hold,  $AU$  is a longest edge of  $UVA$ . Bisect  $AU$  at  $C$ . It is not hard to show that  $AV$  is a longest edge of  $CVA$ . Bisect  $AV$  at  $D$ . We have reached the position illustrated in Figure 2.




FIGURE 2

There are two possibilities now left.

*Possibility 2.*  $UV$  is a longest edge of  $UVC$ . Examination of Figure 2 will show that further bisection produces triangles similar to already occurring triangles. So (up to similarity),  $\mathfrak{F}_0(\Delta)$  consists of  $UVW$ ,  $UAB$ ,  $UVC$ , and  $CVA$ , and  $\mathfrak{F}_1(\Delta)$  consists of  $UVA$ ,  $UAW$ , and  $CVD$ . Since  $UA$  is a longest edge of  $UVA$ ,  $\frac{1}{4}(2p + 2q - 1) \geq p$ , so  $q \geq p + \frac{1}{2}$ . Elementary linear programming now gives  $\phi(UAB) \geq \frac{1}{2}\phi(\Delta)$ , and  $\phi(UVC) \geq \frac{1}{2}\phi(\Delta)$ . Of course  $\phi(CVA) = \phi(\Delta)$ . It turns out that  $\|CV\|^2 = \frac{1}{16}(6p - 2q + 3)$ . Linear programming now gives  $\phi(CVD) \geq \frac{1}{2}\phi(\Delta)$ . Similarly, we find that  $\phi(UVA) \geq \phi(\Delta)$ , and  $\phi(UAW) \geq \frac{1}{2}\phi(\Delta)$ .

If  $UV$  is not the longest edge of  $UVC$ , there remains only

*Possibility 3.*  $CV$  is a longest edge of  $UVC$ . So  $\frac{1}{16}(6p - 2q + 3) \geq p$ , that is  $q \leq 3/2 - 5p$ . Then (up to similarity),  $\mathfrak{F}_0(\Delta)$  consists of  $UVW$ ,  $UAB$ , and  $\mathfrak{F}_1(UVA)$ , while  $\mathfrak{F}_1(\Delta)$  consists of  $UAW$  and  $\mathfrak{F}_0(UVA)$ . As usual,  $\phi(UAW) \geq \frac{1}{2}\phi(\Delta)$ . Since  $q \leq 3/2 - 5p$ , linear programming gives  $2p + 2q - 1 \leq \frac{6}{5}$ . So  $\phi(UAB) \geq \frac{5}{6}\phi(\Delta)$ ,

while  $\phi(UVA) \geq \frac{5}{3}\phi(\Delta)$ . So  $UVA$  is much “fatter” than  $\Delta$ . This enables us to push through an induction.

**LEMMA 1.** *Let  $\Delta$  be an acute triangle. Then the family of  $\Delta$  contains only finitely many similarity types. If  $\Gamma$  is in  $\mathfrak{F}_0(\Delta)$ ,  $\phi(\Gamma) \geq \frac{1}{3}\phi(\Delta)$ . If  $\Gamma$  is in  $\mathfrak{F}_1(\Delta)$ ,  $\phi(\Gamma) \geq \frac{1}{2}\phi(\Delta)$ .*

*Proof.* We show that if our assertions hold whenever  $\phi(\Delta) \geq (\frac{3}{5})^n$ , they hold whenever  $\phi(\Delta) \geq (\frac{3}{5})^{n+1}$ . So suppose  $\Delta = UVW$  is acute and  $\phi(\Delta) \geq (\frac{3}{5})^{n+1}$ . If  $\Delta$  satisfies Possibility 1 or Possibility 2, then, by our earlier calculations,  $\Delta$  certainly satisfies our lemma. So suppose that  $\Delta$  falls under Possibility 3. The elements of  $\mathfrak{F}_1(\Delta)$  are, up to similarity,  $UAW$  (and  $\phi(UAW) \geq \frac{1}{2}\phi(\Delta)$ ) together with  $\mathfrak{F}_0(UVA)$ . But since  $\phi(UVA) \geq \frac{5}{3}\phi(\Delta)$ , by induction assumption  $UVA$  satisfies our lemma, so if  $\Gamma$  is in  $\mathfrak{F}_0(UVA)$ ,  $\phi(\Gamma) \geq \frac{1}{3}\phi(UVA) \geq \frac{5}{9}\phi(\Delta) > \frac{1}{2}\phi(\Delta)$ . The same sort of calculation shows that under Possibility 3, if  $\Gamma$  is in  $\mathfrak{F}_0(\Delta)$ ,  $\phi(\Gamma) \geq \frac{1}{3}\phi(\Delta)$ , indeed  $\phi(\Gamma) \geq \frac{5}{6}\phi(\Delta)$ . This completes the induction.

The inequalities for  $\phi$  are sharp, for if  $\Delta$  is equilateral, no improvement is possible. One cannot expect to make significant improvements on estimates for  $\mathfrak{F}_1(\Delta)$ . But our proof shows that for the “general” acute triangle (Possibility 3), if  $\Gamma$  is in  $\mathfrak{F}_0(\Delta)$ ,  $\phi(\Gamma) \geq \frac{5}{6}\phi(\Delta)$ .

**3. Obtuse Triangles.** Suppose now we are “bisecting” a triangle  $\Delta = UVW$ , where as usual  $\|VW\|^2 = 1$ ,  $\|UW\|^2 = q$ ,  $\|UV\|^2 = p$ ,  $p \leq q < 1$ , and where the angle  $VUW$  is  $\geq 90^\circ$ . Bisect  $VW$  at  $A$ . Then  $UW$  is the longest edge of  $\Delta UAW$ . Bisect it at  $B$  (see Figure 1).

**LEMMA 2.** *If  $\Delta$  is obtuse, the family of  $\Delta$  contains only finitely many similarity types. If  $\Gamma$  is in  $\mathfrak{F}_1(\Delta)$ ,  $\phi(\Gamma) \geq \frac{1}{2}\phi(\Delta)$ . If  $\Gamma$  is in  $\mathfrak{F}_0(\Delta)$ ,  $\phi(\Gamma) \geq \frac{1}{3}\phi(\Delta)$ .*

*Proof.* Let  $\frac{1}{2} \leq \lambda < 1$ . We prove that if our result holds for all obtuse triangles  $\Delta$  such that the smallest angle of  $\Delta$  has cosine  $\leq \sqrt{\lambda}$  and such that  $\phi(\Delta) \geq \lambda^n$ , then the result holds for all such  $\Delta$  with  $\phi(\Delta) \geq \lambda^{n+1}$ . So suppose that  $\phi(\Delta) \geq \lambda^{n+1}$ , and that  $\Delta$  has smallest angle  $\alpha$ , where  $\cos^2 \alpha \leq \lambda$ . If the angle  $BAU (= AUV)$  is  $\geq 90^\circ$ , there is no problem. For it is easy to see that all angles of triangles  $UAB$ ,  $UVA$  are  $\geq \alpha$ . But

$$\phi(UAB) = \frac{1}{q}\phi(\Delta) \geq \frac{1}{\cos^2 \alpha}\phi(\Delta) \geq \lambda^n.$$

Also  $\phi(UVA) = 2\phi(\Delta) \geq 2\lambda^{n+1} \geq \lambda^n$  since  $\lambda \geq \frac{1}{2}$ . Now  $\mathfrak{F}_1(\Delta)$  consists, up to similarity, of  $UAW$ ,  $\mathfrak{F}_1(UAB)$ , and  $\mathfrak{F}_0(UVA)$ . By induction assumption, if  $\Gamma$  is in  $\mathfrak{F}_1(UAB)$ , then  $\phi(\Gamma) \geq \phi(UAB) \geq \frac{1}{2}\phi(\Delta)$ , while if  $\Gamma$  is in  $\mathfrak{F}_0(UVA)$ ,  $\phi(\Gamma) \geq \frac{1}{3}\phi(UVA) = \frac{2}{3}\phi(\Delta) > \frac{1}{2}\phi(\Delta)$ . Elements of  $\mathfrak{F}_0(\Delta)$  are dealt with in the same way.

So it remains to see what happens if the angle  $BAU$  is  $\leq 90^\circ$ . If  $UV$  is the longest edge of  $UVA$ , the family of  $\Delta$  has at most four similarity types, and a quick computation yields the result. Otherwise,  $\phi(UVA) = 2\phi(\Delta)$ , and of course  $\phi(UAB) \geq \phi(\Delta)$ , and our result follows quickly from Lemma 1.

The estimate for  $\mathfrak{F}_1(\Delta)$  cannot be significantly improved. But by a closer analysis of the possibilities that arise when the angle  $UAB$  is acute, one can show that in fact if  $\Gamma$  is in  $\mathfrak{F}_0(\Delta)$ ,  $\phi(\Gamma) \geq \phi(\Delta)$ .

**4. Summary, Problems.** By combining Lemma 1, Lemma 2, and the fact that area goes down by a factor of 2 each generation we obtain:

**THEOREM.** *Under the bisection process, the family of a triangle falls into finitely many similarity classes. If  $j$  is even,  $m_j \leq \sqrt{3} 2^{-j/2} m_0$ . If  $j$  is odd,  $m_j \leq \sqrt{2} 2^{-j/2} m_0$ .*

Both estimates are sharp, for we have equality when the triangle is equilateral. If the starting triangle is far from being equilateral, the bounds for  $m_j$  when  $j$  is even can be improved. By examining the details of the proof, one can find an upper bound for the number of similarity types in the family of  $\Delta$ , say as a function of  $\phi(\Delta)$ . But there appears to be nothing very interesting left to do for triangles.

But one can raise similar problems in a much more general setting. Let  $A_1, A_2, \dots, A_n$  be a configuration of  $n + 1$  points in  $d$ -dimensional space. Suppose  $\|A_0 - A_1\| \geq \|A_i - A_j\|$  for all  $i, j$ . Then the configuration gives birth to two daughter configurations  $A_0, (A_0 + A_1)/2, A_2, \dots, A_n$  and  $(A_0 + A_1)/2, A_1, A_2, \dots, A_n$ . One can define  $m_j$  as for triangles and ask about the behavior of  $m_j$ . It seems reasonable to conjecture that  $m_j = O(2^{-j/n})$ . One can make the even stronger conjecture that up to similarity any configuration has a finite family.

Already for four points in general position in 3-dimensional space, the problems seem difficult. We have a proof of the "finite family" conjecture for certain classes of tetrahedra. For example, it turns out that if a tetrahedron is nearly equilateral and the second largest edge is opposite the longest edge, then the family of the tetrahedron falls into  $\leq 37$  similarity classes. (The condition "nearly equilateral" is a little complicated to describe briefly, but, for example, it is satisfied if all edge lengths are within 5% of each other.)

Department of Mathematics  
University of British Columbia  
Vancouver, B.C., Canada V6T 1W5

1. C. HARVEY & F. STENGER, "A two dimensional analogue to the method of bisections for solving nonlinear equations," *Quart. Appl. Math.*, v. 33, 1976, pp. 351-368.
2. R. B. KEARFOTT, "A proof of convergence and an error bound for the method of bisection in  $R^n$ ," *Math. Comp.*, v. 32, 1978, pp. 1147-1153.
3. K. SIKORSKI, "A three-dimensional analogue to the method of bisections for solving nonlinear equations," *Math. Comp.*, v. 33, 1979, pp. 722-738.
4. F. STENGER, "Computing the topological degree of a mapping in  $n$ -space," *Numer. Math.*, v. 25, 1975, pp. 23-38.
5. M. STYNES, "On faster convergence of the bisection method for certain triangles," *Math. Comp.*, v. 33, 1979, pp. 717-721.