

Superconvergence of a Collocation-Type Method for Simple Turning Points of Hammerstein Equations

By Sunil Kumar

Abstract. In this paper a simple turning point ($y = y^c$, $\lambda = \lambda^c$) of the parameter-dependent Hammerstein equation

$$y(t) = f(t) + \lambda \int_a^b k(t, s)g(s, y(s)) ds, \quad t \in [a, b],$$

is approximated numerically in the following way. A simple turning point ($z = z^c$, $\lambda = \lambda^c$) of an equivalent equation for $z(t) := \lambda g(t, y(t))$ is computed first. This is done by solving a discretized version of a certain system of equations which has (z^c, λ^c) as part of an isolated solution. The particular discretization used here is standard piecewise polynomial collocation. Finally, an approximation to y^c is obtained by use of the (exact) equation

$$y(t) = f(t) + \int_a^b k(t, s)z(s) ds, \quad t \in [a, b].$$

The main result of the paper is that, under suitable conditions, the approximations to y^c and λ^c are both superconvergent, that is, they both converge to their respective exact values at a faster rate than the collocation approximation (of z^c) does to z^c .

1. Introduction. We consider parameter-dependent Hammerstein equations of the form

$$(1.1) \quad y(t) = f(t) + \lambda \int_a^b k(t, s)g(s, y(s)) ds, \quad t \in [a, b],$$

where $-\infty < a < b < \infty$, $\lambda \in \mathbf{R}$ is the parameter, f , k , and g are known functions, with $g(s, v)$ nonlinear in v , and the pair $(y = y^c, \lambda = \lambda^c)$ is a simple (that is, quadratic) turning point of (1.1) which is to be determined numerically. (For the definition of a simple turning point see the first paragraph of Section 3.)

We do this by an approximation procedure which hinges on an equivalent equation (see (1.4)) for the function z defined by

$$(1.2) \quad z(t) := \lambda g(t, y(t)), \quad t \in [a, b].$$

On substituting (1.2) into (1.1) we have immediately

$$(1.3) \quad y(t) = f(t) + \int_a^b k(t, s)z(s) ds, \quad t \in [a, b],$$

and hence it follows from (1.2) that z satisfies the nonlinear integral equation

$$(1.4) \quad z(t) = \lambda g \left(t, f(t) + \int_a^b k(t, s)z(s) ds \right), \quad t \in [a, b].$$

Received December 15, 1986; revised June 29, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 65R20, 45G10.

©1988 American Mathematical Society
 0025-5718/88 \$1.00 + \$.25 per page

The equivalence of (1.4) and (1.1) (Kumar and Sloan [13]) is in the sense that there is a one-to-one correspondence between their solution sets (see Lemma 1 for details). Thus, corresponding to the simple turning point (y^c, λ^c) of (1.1) is a simple turning point $(z = z^c, \lambda = \lambda^c)$ of (1.4). It is the latter turning point that is actually computed in this paper. This computation is done via the enlarged system approach of Moore and Spence [14], with piecewise polynomial collocation (Atkinson, Graham and Sloan [4], Joe [7]) as the underlying discretization.

The desired approximation to y^c is then obtained by use of the (exact) equation (1.3). Thus, essentially, y^c is approximated by the collocation-type method of Kumar and Sloan [13] (see also Kumar [11]).

The rest of the paper is organized as follows. Section 2 provides necessary background material, while Section 3 gives details of our approximation procedure. Section 3 also establishes the convergence of the approximations to z^c , λ^c , and y^c , this being done there by a simple application of a theorem of Spence and Moore [16]. In Section 4 the theorem of Spence and Moore is used again, and hence it is shown that, under suitable conditions, the approximations to y^c and λ^c are both superconvergent, that is, they both converge to their respective exact values at a faster rate than the collocation approximation (of z^c) does to z^c . In Section 5 numerical results are presented.

2. Preliminaries. It is convenient to consider (1.1) in the Banach space $C = C[a, b]$ of continuous, real-valued functions on $[a, b]$. Recall that this space is equipped with the uniform norm

$$\|x\|_\infty = \sup_{a \leq t \leq b} |x(t)|, \quad x \in C.$$

On the other hand, it is convenient to consider (1.4) in the Banach space $R = R[a, b]$ of bounded Riemann-integrable real-valued functions on $[a, b]$, the reason being that R permits piecewise continuous approximations to z^c to be dealt with (Anselone [3, p. 22], Atkinson, Graham and Sloan [4]). Like C , the space R is also equipped with the uniform norm. Since a bounded function on $[a, b]$ is Riemann-integrable if and only if it is continuous almost everywhere on $[a, b]$, it follows that C is a closed subspace of R .

It is also convenient to make the following assumptions on the functions f , k , and g in (1.1):

- A1. $f \in C$;
- A2. the kernel k satisfies

$$\sup_{a \leq t \leq b} \int_a^b |k(t, s)| ds < \infty$$

and

$$\lim_{t \rightarrow t'} \int_a^b |k(t, s) - k(t', s)| ds = 0, \quad t' \in [a, b];$$

- A3. the function $g(t, v)$ is defined and continuous on $[a, b] \times \mathbf{R}$;
- A4. the partial derivative $g_v(t, v) := \partial g(t, v) / \partial v$ exists and is continuous on $[a, b] \times \mathbf{R}$;
- A5. the partial derivative $g_{vv}(t, v) := \partial g_v(t, v) / \partial v$ exists and is continuous on $[a, b] \times \mathbf{R}$;

A6. the function g_{vv} satisfies the Lipschitz condition

$$|g_{vv}(t, y_1(t)) - g_{vv}(t, y_2(t))| \leq \alpha |y_1(t) - y_2(t)|,$$

for some constant $\alpha > 0$, $t \in [a, b]$, and all $y_1, y_2 \in B_1(y^c, \rho)$, where

$$B_1(y^c, \rho) = \{y \in C : \|y - y^c\|_\infty \leq \rho\}, \quad \rho > 0.$$

Note that assumption A2 ensures via the Arzelà-Ascoli theorem (Kantorovich and Akilov [9, p. 27]) that the linear integral operator K , defined by

$$(Kw)(t) := \int_a^b k(t, s)w(s) ds, \quad t \in [a, b], \quad w \in R,$$

is a compact operator from R to C (Anselone [1], [2], [3, Proposition 2.10]), and hence also from C to C . Being both compact and linear, it is necessarily (Kantorovich and Akilov [9, p. 244]) completely continuous.

Defining another completely continuous operator $T: R \rightarrow C$ by

$$T(w)(t) := f(t) + (Kw)(t), \quad t \in [a, b], \quad w \in R,$$

and a continuous, bounded operator $G: C \rightarrow C$ by

$$G(x)(t) := g(t, x(t)), \quad t \in [a, b], \quad x \in C,$$

we rewrite the integral equations (1.1) and (1.4) in operator form as

$$(2.1) \quad y = T(\lambda G(y)), \quad y \in C,$$

and

$$(2.2) \quad z = \lambda GT(z), \quad z \in R,$$

respectively. The two equations are equivalent in the sense of the following lemma.

LEMMA 1 (Kumar and Sloan [13]). *For $\lambda \in \mathbf{R}$, the sets*

$$\Theta_{TG} := \{y \in C : T(\lambda G(y)) = y\} \quad \text{and} \quad \Theta_{GT} := \{z \in R : \lambda GT(z) = z\}$$

are in one-to-one correspondence. Specifically, λG is a one-to-one operator from Θ_{TG} onto Θ_{GT} , with inverse T .

Thus $z^c = \lambda^c G(y^c)$, and $y^c = T(z^c)$.

Note that since T is completely continuous from R to C , and G is continuous and bounded on C , it follows (Krasnosel'skii and Zabreiko [10, p. 74]) that $T(\lambda G)$ is completely continuous on C , and GT is completely continuous on R . The assumptions A3 and A4 ensure that G is continuously Fréchet differentiable on C ; its Fréchet derivative at $x \in C$ is the bounded linear operator $G'(x)$ given by

$$[G'(x)w](t) = g_v(t, x(t))w(t), \quad t \in [a, b], \quad w \in C.$$

If, in addition, A1 and A2 hold, then the operator GT is continuously Fréchet differentiable on R ; its Fréchet derivative at $z \in R$ is the completely continuous linear operator $(GT)'(z)$ given by

$$[(GT)'(z)w](t) = G'(T(z))(Kw)(t), \quad t \in [a, b], \quad w \in R.$$

Furthermore, if A5 also holds, then the second Fréchet derivative of GT at $z \in R$ is the symmetric bilinear operator $(GT)''(z)$, on $R \times R$ to C , given by

$$[(GT)''(z)wx](t) = g_{vv}(t, T(z)(t))(Kw)(t)(Kx)(t), \quad t \in [a, b], \quad w, x \in R.$$

Similarly, $T(\lambda G)$ is twice Fréchet differentiable; its first and second Fréchet derivatives at $y \in C$ are given by

$$[(T\lambda G)'(y)w](t) = \lambda K G'(y)w(t), \quad t \in [a, b], \quad w \in C,$$

and

$$[(T\lambda G)''(y)wx](t) = \lambda K g_{vv}(t, y(t))w(t)x(t), \quad t \in [a, b], \quad w, x \in C,$$

respectively.

The final assumption (A6) implies that for all $y_1, y_2 \in B_1(y^c, \rho)$,

$$\begin{aligned} & \| (T\lambda G)''(y_1) - (T\lambda G)''(y_2) \| \\ & \leq |\lambda| \sup_{a \leq t \leq b} \int_a^b |k(t, s)| |g_{vv}(s, y_1(s)) - g_{vv}(s, y_2(s))| ds \\ & \leq \alpha |\lambda| \sup_{a \leq t \leq b} \int_a^b |k(t, s)| |y_1(s) - y_2(s)| ds \\ & \leq \alpha |\lambda| \|K\| \|y_1 - y_2\|_\infty, \end{aligned}$$

where

$$\|K\| = \sup_{a \leq t \leq b} \int_a^b |k(t, s)| ds < \infty.$$

Likewise, for all $z_1, z_2 \in B_2(z^c, \rho)$, where

$$B_2(z^c, \rho) = \{z \in R: \|z - z^c\|_\infty \leq \rho / \|K\|\},$$

it is easily verified that $T(z_1), T(z_2) \in B_1(y^c, \rho)$, and therefore

$$\begin{aligned} & \| (GT)''(z_1) - (GT)''(z_2) \| \\ & \leq \|K\|^2 \sup_{a \leq t \leq b} |g_{vv}(t, T(z_1)(t)) - g_{vv}(t, T(z_2)(t))| \\ & \leq \alpha \|K\|^2 \sup_{a \leq t \leq b} |T(z_1)(t) - T(z_2)(t)| \\ & \leq \alpha \|K\|^3 \|z_1 - z_2\|_\infty. \end{aligned}$$

3. The Approximate Method and its Convergence. Since (z^c, λ^c) is a simple turning point of (2.2), the following hold by definition (see, for example, Moore and Spence [14], [15]):

- D1. $z^c - \lambda^c GT(z^c) = 0$;
- D2. $I - \lambda^c (GT)'(z^c)$ has a one-dimensional null space spanned by ϕ^c ;
- D3. $I - \lambda^c (GT)'(z^c)$ has closed range;
- D4. $[I - \lambda^c (GT)'(z^c)]^*$ has a one-dimensional null space spanned by the linear functional σ^c ;

(Here $*$ denotes the conjugate of a bounded linear operator on R .)

D5. $\sigma^c(GT(z^c)) = \sigma^c(z^c) / \lambda^c \neq 0$;

D6. $\sigma^c([(GT)''(z^c)]\phi^c\phi^c) \neq 0$.

It follows from properties D1 and D2 that (z^c, λ^c) is a nonisolated solution of (2.2). However, (z^c, λ^c) is contained as part of an isolated solution of the following system (Moore and Spence [14]):

$$(3.1a) \quad z - \lambda GT(z) = 0,$$

$$(3.1b) \quad \phi - \lambda[(GT)'(z)]\phi = 0,$$

$$(3.1c) \quad l(\phi) - 1 = 0,$$

where $\lambda \in \mathbf{R}$, $z, \phi \in R$ and l is a bounded linear functional which normalizes ϕ .

For our convergence analysis we rearrange (3.1) as

$$(3.1'a) \quad z - \lambda GT(z) = 0,$$

$$(3.1'b) \quad \phi - \lambda[(GT)'(\lambda GT(z))]\phi = 0,$$

$$(3.1'c) \quad \lambda - [\lambda - l(\phi) + 1] = 0.$$

Note that (3.1) and (3.1') are equivalent in the sense that any solution of (3.1) is a solution of (3.1') and *vice versa*.

We now write (3.1') in abbreviated form as

$$(3.2) \quad Z - F(Z) = 0,$$

where $Z = (z, \phi, \lambda)$, and F is an operator on the Banach space $R \times R \times \mathbf{R}$. We equip this product space with the product norm

$$\|Z\| = \max(\|z\|_\infty, \|\phi\|_\infty, |\lambda|), \quad Z \in R \times R \times \mathbf{R}.$$

LEMMA 2 (Moore and Spence [14]). *Suppose A1 to A5 hold, and $l(\phi^c) = 1$. Then $Z^c = (z^c, \phi^c, \lambda^c)$ is an isolated solution of (3.2).*

3.1. *Algorithmic Details of the Approximation Procedure.* In the present work we approximate (z^c, ϕ^c, λ^c) by (z_n, ϕ_n, λ_n) , where z_n and ϕ_n are of the form

$$z_n(t) = \sum_{j=1}^n a_{nj} u_{nj}(t), \quad t \in [a, b],$$

$$\phi_n(t) = \sum_{j=1}^n b_{nj} u_{nj}(t), \quad t \in [a, b],$$

and $\{u_{n1}, \dots, u_{nn}\}$ is a set of basis functions for some chosen approximating space. We determine the unknowns $a_{n1}, \dots, a_{nn}, b_{n1}, \dots, b_{nn}$, and λ_n by collocating (3.1) at n distinct points $\tau_{n1}, \dots, \tau_{nn}$ in $[a, b]$:

$$\left. \begin{aligned} z_n(\tau_{ni}) - \lambda_n[GT(z_n)](\tau_{ni}) &= 0 \\ \phi_n(\tau_{ni}) - \lambda_n[(GT)'(z_n)\phi_n](\tau_{ni}) &= 0 \end{aligned} \right\}, \quad i = 1, \dots, n,$$

$$l(\phi_n) - 1 = 0.$$

Thus we discretize (3.1) by the standard collocation method.

In practice, the computation of the $2n + 1$ unknowns $a_{n1}, \dots, a_{nn}, b_{n1}, \dots, b_{nn}$, and λ_n may be arranged in the following manner. Let

$$x_j = \begin{cases} a_{nj}, & j = 1, \dots, n, \\ b_{n,j-n}, & j = n + 1, \dots, 2n, \\ \lambda_n, & j = 2n + 1, \end{cases}$$

$$f_i = f(\tau_{ni}), \quad i = 1, \dots, n,$$

$$U_{ij} = u_{nj}(\tau_{ni}), \quad 1 \leq i, j \leq n,$$

and

$$K_{ij} = \int_a^b k(\tau_{ni}, s) u_{nj}(s) ds, \quad 1 \leq i, j \leq n.$$

Then the unknowns x_1, \dots, x_{2n+1} satisfy the system of equations

$$\left. \begin{aligned} \sum_{j=1}^n x_j U_{ij} - x_{2n+1} g \left(\tau_{ni}, f_i + \sum_{j=1}^n x_j K_{ij} \right) &= 0 \\ \sum_{j=1}^n x_{n+j} U_{ij} - x_{2n+1} g_v \left(\tau_{ni}, f_i + \sum_{j=1}^n x_j K_{ij} \right) \left[\sum_{j=1}^n x_{n+j} K_{ij} \right] &= 0 \end{aligned} \right\},$$

$$i = 1, \dots, n,$$

$$l \left(\sum_{j=1}^n x_{n+j} u_{nj} \right) - 1 = 0.$$

Since $y^c = T(z^c)$, we define our approximation to y^c to be

$$y_n := T(z_n).$$

It then follows that

$$y_n(t) = f(t) + \sum_{j=1}^n x_j \int_a^b k(t, s) u_{nj}(s) ds, \quad t \in [a, b].$$

3.2. Analysis of the Approximate Method. We consider the case where the collocation approximations z_n and ϕ_n are sought in piecewise polynomial function spaces.

For any natural number N , let

$$\Pi_N : a = s_1 < s_2 < \dots < s_N < s_{N+1} = b$$

be a partition of $[a, b]$, and let $h = h(N) = \max_{1 \leq i \leq N} (s_{i+1} - s_i)$. We assume that $h \rightarrow 0$ as $N \rightarrow \infty$ and that the partition Π_N is quasi-uniform, that is, there exists a constant ζ such that, for all N , $h \leq \zeta \min_{1 \leq i \leq N} (s_{i+1} - s_i)$.

With r a positive integer and ν an integer satisfying $0 \leq \nu < r$, let $S_{r,N}^\nu$ denote the space of piecewise polynomial functions of order r and continuity ν . This terminology may be better understood by noting that a function $\omega \in S_{r,N}^\nu$ if and only if it is a polynomial of degree $\leq r - 1$ on each subinterval (s_i, s_{i+1}) , $1 \leq i \leq N$, and has $\nu - 1$ continuous derivatives on (a, b) . If $\nu = 0$, there is no continuity requirement at the breakpoints s_i , $1 \leq i \leq N + 1$, in which case we arbitrarily take each $\omega \in S_{r,N}^0$ to be right-continuous at $s_1 = a$, and left-continuous at every other s_i , $2 \leq i \leq N + 1$.

For $n = \dim(S_{r,N}^\nu) = (N - 1)(r - \nu) + r$, let $\{u_{nj}\}_{j=1}^n$ be a set of basis functions for $S_{r,N}^\nu$, and choose a set $\{\tau_{ni}\}_{i=1}^n$ of distinct points in $[a, b]$ such that the $n \times n$ matrix $(u_{nj}(\tau_{ni}))$ is nonsingular. Furthermore, let P_n be the interpolatory projection operator from $C + S_{r,N}^\nu$ onto $S_{r,N}^\nu$, which satisfies the condition

$$(P_n w)(\tau_{ni}) = w(\tau_{ni}), \quad i = 1, \dots, n,$$

for all $w \in C + S_{r,N}^\nu$. Then our discretization of (3.1) is represented by the system of equations

$$(3.3a) \quad z_n - \lambda_n P_n G T(z_n) = 0,$$

$$(3.3b) \quad \phi_n - \lambda_n P_n G'(T(z_n)) K \phi_n = 0,$$

$$(3.3c) \quad l(\phi_n) - 1 = 0,$$

where $z_n = P_n z_n \in S_{r,N}^\nu$, $\phi_n = P_n \phi_n \in S_{r,N}^\nu$, and $\lambda_n \in \mathbf{R}$.

While we use (3.3) for computational purposes, to prove the existence and convergence of a turning point of (3.3a), we consider the following discretization of (3.1')

$$(3.3'a) \quad z_n - \lambda_n P_n G T(z_n) = 0,$$

$$(3.3'b) \quad \phi_n - \lambda_n [(P_n G T)'(\lambda_n P_n G T(z_n))] \phi_n = 0,$$

$$(3.3'c) \quad \lambda_n - [\lambda_n - l(\phi_n) + 1] = 0.$$

This system we write in abbreviated form as

$$(3.4) \quad Z_n - F_n(Z_n) = 0,$$

where $Z_n = (z_n, \phi_n, \lambda_n)$, and F_n is an operator from $R \times R \times \mathbf{R}$ to $S_{r,N}^\nu \times S_{r,N}^\nu \times \mathbf{R}$.

Compact operator equations of the form (3.2) and (3.4) have been analyzed by Spence and Moore [16]. Here we apply their third theorem to prove the convergence of a solution Z_n of (3.4) to the solution Z^c of (3.2). But before we can do this, we need to impose the additional constraints on the (collocation) points $\{\tau_{ni}\}_{i=1}^n$ that they be chosen to ensure that P_n is uniformly bounded as an operator from $C + S_{r,N}^\nu$ to $S_{r,N}^\nu$, that is

$$(3.5) \quad \|P_n\| \leq c_3,$$

where $c_3 > 0$ is independent of n , and

$$(3.6) \quad \lim_{n \rightarrow \infty} \|w - P_n w\|_\infty = 0 \quad \text{for all } w \in C.$$

Remark 1. Throughout this paper, c_1, \dots, c_5 denote positive generic constants which may take different values at their different occurrences but which are always independent of n .

THEOREM 1. *Let (y^c, λ^c) be a simple turning point of (2.1), and let (z^c, λ^c) be the corresponding simple turning point of (2.2). Suppose A1 to A6 hold, the interpolatory operator P_n satisfies (3.5) and (3.6), and $l(\phi^c) = 1$. Then, for sufficiently large n , $(z_n, \phi_n, \lambda_n) \in S_{r,N}^\nu \times S_{r,N}^\nu \times \mathbf{R}$ is an isolated solution of system (3.3'), and*

$$\begin{aligned} &\max(\|z^c - z_n\|_\infty, \|\phi^c - \phi_n\|_\infty, |\lambda^c - \lambda_n|) \\ &\leq c_1 \|z^c - P_n z^c\|_\infty + c_2 \|\phi^c - P_n \phi^c\|_\infty. \end{aligned}$$

Proof. An application of Theorem 3 of Spence and Moore [16] yields the result. Two of the less trivial assumptions to check are the collective compactness (Anselone [3, p. 94]) of $\{P_n GT, n \geq 1\}$ on R , and the pointwise convergence of $P_n GT$ to GT on R . The former condition follows from Weiss [19], and the latter from the mapping properties of GT , and (3.6).

From Theorem 3 of Spence and Moore [16] we have immediately that

$$\begin{aligned} &\max(\|z^c - z_n\|_\infty, \|\phi^c - \phi_n\|_\infty, |\lambda^c - \lambda_n|) \\ &\leq c_5 \max(\|z^c - P_n z^c\|_\infty, \|\phi^c - \lambda^c [(P_n GT)'(P_n z^c)] \phi^c\|_\infty). \end{aligned}$$

The result now follows from the argument

$$\begin{aligned} &\|\phi^c - \lambda^c [(P_n GT)'(P_n z^c)] \phi^c\|_\infty \\ &\leq \|\phi^c - \lambda^c [(P_n GT)'(z^c)] \phi^c\|_\infty \\ &\quad + \|\lambda^c [(P_n GT)'(z^c)] \phi^c - \lambda^c [(P_n GT)'(P_n z^c)] \phi^c\|_\infty \\ &= \|\phi^c - P_n \phi^c\|_\infty + \|\lambda^c P_n [(GT)'(z^c) - (GT)'(P_n z^c)] \phi^c\|_\infty \\ &\leq \|\phi^c - P_n \phi^c\|_\infty + c_4 \|(GT)'(z^c) - (GT)'(P_n z^c)\|_\infty \\ &\leq \|\phi^c - P_n \phi^c\|_\infty + c_3 \|z^c - P_n z^c\|_\infty, \end{aligned}$$

where in going from the second last step to the final step we have used a mean value theorem (Kantorovich and Akilov [9, p. 500]) since $(GT)''(z)$ exists and is bounded for all z in a neighborhood of z^c . \square

Remark 2. The fact that (3.3') has an isolated solution (z_n, ϕ_n, λ_n) implies that (z_n, λ_n) is a simple turning point of (3.3a).

Remark 3. Conditions (3.5) and (3.6) do not hold in the case of collocation with global polynomials, and thus a different approach needs to be adopted to prove the convergence of Z_n to Z^c . For a possible approach see Kumar [12].

Since our approximation to y^c is

$$(3.7) \quad y_n := T(z_n),$$

where $z_n \in S_{r,N}^\nu$ is part of the isolated solution (z_n, ϕ_n, λ_n) of system (3.3') (see Theorem 1), it follows from

$$\|y^c - y_n\|_\infty = \|K(z^c - z_n)\|_\infty \leq \|K\| \|z^c - z_n\|_\infty$$

that the rate of convergence of y_n to y^c is no worse than that of z_n to z^c . In the next section we show that the convergence rate of y_n to y^c (and also that of λ_n to λ^c) may be better than that of z_n to z^c .

4. Superconvergence Results. For $1 \leq p < \infty$, let $L_p = L_p(a, b)$ denote the Banach space of measurable, real-valued functions on (a, b) which have integrable p th power. The norm on this space is defined by

$$\|w\|_p := \left(\int_a^b |w(s)|^p ds \right)^{1/p}, \quad w \in L_p.$$

Let $L_\infty = L_\infty(a, b)$ denote the Banach space of essentially bounded real-valued functions on (a, b) . This space has the norm

$$\|w\|_\infty = \operatorname{ess\,sup}_{a < t < b} |w(t)|, \quad w \in L_\infty.$$

Finally, let $W_p^m = W_p^m(a, b)$, for m a nonnegative integer and $1 \leq p \leq \infty$, be the Sobolev space of functions w such that $w^{(i)} \in L_p$ for $i = 0, 1, \dots, m$, where $w^{(i)}$ is the i th distributional derivative of w . This space we equip with the norm

$$\|w\|_{m,p} = \sum_{i=0}^m \|w^{(i)}\|_p.$$

The following corollary to Theorem 1 holds for any set of collocation points for which the operator P_n satisfies (3.5) and (3.6), and may be obtained quite easily from an approximation-theoretic result in Graham, Joe and Sloan [5, Theorem 3.1].

COROLLARY 1. *If $z^c \in W_\infty^\mu$, $\mu \geq 1$, and $\phi^c \in W_\infty^\delta$, $\delta \geq 1$, then*

$$\max(\|z^c - z_n\|_\infty, \|\phi^c - \phi_n\|_\infty, |\lambda^c - \lambda_n|) = O(h^\xi),$$

where $\xi = \min(\mu, \delta, r)$.

Up to this point, the analysis predicts a convergence rate of $O(h^\xi)$ for both λ_n and y_n . In this section we carry the analysis further, and hence show that, under suitable conditions, both y_n and λ_n may converge at a rate $o(h^\xi)$.

We begin by noting that corresponding to the enlarged system (3.1) for (z^c, λ^c) is an analogous system for (y^c, λ^c) :

$$(4.1a) \quad y - T(\lambda G(y)) = 0,$$

$$(4.1b) \quad \psi - \lambda K G'(y) \psi = 0,$$

$$(4.1c) \quad L(\psi) - 1 = 0,$$

where $\lambda \in \mathbf{R}$, $y, \psi \in C$ and L is a bounded linear functional which normalizes ψ .

For our subsequent analysis we rearrange (4.1) as

$$(4.1'a) \quad y - T(\lambda G(y)) = 0,$$

$$(4.1'b) \quad \psi - [(T\lambda G)'(T\lambda G(y))] \psi = 0,$$

$$(4.1'c) \quad \lambda - [\lambda - L(\psi) + 1] = 0.$$

Note that (4.1) and (4.1') are equivalent in the sense that any solution of (4.1) is a solution of (4.1') and *vice versa*.

We now write (4.1') in abbreviated form as

$$(4.2) \quad Y - H(Y) = 0,$$

where $Y = (y, \psi, \lambda)$, and H is an operator on the Banach space $C \times C \times \mathbf{R}$. We equip this product space with the product norm

$$\|Y\| = \max(\|y\|_\infty, \|\psi\|_\infty, |\lambda|), \quad Y \in C \times C \times \mathbf{R}.$$

LEMMA 3 (Moore and Spence [14]). *Let $\{\psi^c\}$, $L(\psi^c) = 1$, be a basis of the null space of the operator $I - \lambda^c K G'(y^c)$, and suppose A1 to A5 hold. Then $Y^c = (y^c, \psi^c, \lambda^c)$ is an isolated solution of (4.2).*

From Lemmas 1 to 3 it follows, if $z = z^c$ and $\lambda = \lambda^c$, that

$$\phi = \lambda^c G'(y^c) \psi^c$$

is a solution of (3.1b). Similarly, if $y = y^c$ and $\lambda = \lambda^c$, then

$$(4.3) \quad \psi = K \phi^c$$

is a solution of (4.1b). Thus, just as we approximate $y^c = T(z^c)$ by $y_n = T(z_n)$ (see (3.7)), we may approximate ψ^c by

$$(4.4) \quad \psi_n = K \phi_n, \quad L(\psi_n) = 1,$$

where $\phi_n \in S_{r,N}^\nu$ is part of the isolated solution (z_n, ϕ_n, λ_n) of system (3.3') (see Theorem 1).

Using (3.3'), we write (3.7) and (4.4) as the system of equations

$$(4.5) \quad \begin{aligned} y_n - T(\lambda_n P_n G(y_n)) &= 0, \\ \psi_n - [(T \lambda_n P_n G)'(T \lambda_n P_n G(y_n))] \psi_n &= 0, \\ L(\psi_n) - 1 &= 0, \end{aligned}$$

and then abbreviate (4.5) as

$$(4.6) \quad Y_n - H_n(Y_n) = 0,$$

where $Y_n = (y_n, \psi_n, \lambda_n)$, and H_n is an operator on $C \times C \times \mathbf{R}$. An application of Theorem 3 of Spence and Moore [16] now gives the following result.

THEOREM 2. *Let (y^c, λ^c) be a simple turning point of (2.1), and let (z^c, λ^c) be the corresponding simple turning point of (2.2). Suppose A1 to A6 hold, the interpolatory operator P_n satisfies (3.5) and (3.6), and $l(\phi^c) = 1$. Then, for sufficiently large n , $(y_n, \psi_n, \lambda_n) \in C \times C \times \mathbf{R}$ is an isolated solution of system (4.5), and*

$$(4.7) \quad \begin{aligned} \max(\|y^c - y_n\|_\infty, \|\psi^c - \psi_n\|_\infty, |\lambda^c - \lambda_n|) \\ \leq c_3 \|K(z^c - P_n z^c)\|_\infty + c_4 \|K(\phi^c - P_n \phi^c)\|_\infty. \end{aligned}$$

Proof. To derive (4.7), note that Theorem 3 of Spence and Moore [16] gives immediately that

$$\begin{aligned} \max(\|y^c - y_n\|_\infty, \|\psi^c - \psi_n\|_\infty, |\lambda^c - \lambda_n|) \\ \leq c_5 \max(\|K(z^c - P_n z^c)\|_\infty, \|K \phi^c - [(T \lambda^c P_n G)'(T P_n z^c)] \psi^c\|_\infty), \end{aligned}$$

and an argument similar to that given in the proof of Theorem 1 yields

$$\begin{aligned} \|K \phi^c - [(T \lambda^c P_n G)'(T P_n z^c)] \psi^c\|_\infty \\ \leq \|K(\phi^c - P_n \phi^c)\|_\infty + c_2 \|K(z^c - P_n z^c)\|_\infty. \quad \square \end{aligned}$$

Inequality (4.7) is the key to the derivation of our superconvergence results for y_n and λ_n . The results given here are for the cases $\nu = 0$ and $\nu = 1$ only, and appear in Theorem 3. It should be noted that Theorem 3 holds only if the collocation points $\{\tau_{nj}\}_{j=1}^n$ are chosen, for each case $\nu = 0$ and $\nu = 1$, in the specific way indicated below.

Case $\nu = 0$. Recall that the dimension of $S_{r,N}^0$ is $n = Nr$, and hence Nr collocation points are needed. These should be selected as follows. Let ζ_1, \dots, ζ_r be the zeros of the r th degree Legendre polynomial $\Phi_r(s)$, $s \in [-1, 1]$. (Note that these zeros are known as the Gauss-Legendre points.) Take as the collocation points the points ζ_1, \dots, ζ_r shifted linearly to each subinterval (s_i, s_{i+1}) , $1 \leq i \leq N$:

$$\tau_{n,(i-1)r+j} = (s_i + s_{i+1} + (s_{i+1} - s_i)\zeta_j)/2, \quad 1 \leq j \leq r, \quad 1 \leq i \leq N.$$

Case $\nu = 1$. In this case the collocation approximations to z^c and ϕ^c are sought in $S_{r,N}^1$, the space of continuous piecewise polynomial functions, and hence r is necessarily ≥ 2 . The $n = Nr - N + 1$ collocation points should be taken to be the breakpoints s_i , $1 \leq i \leq N + 1$, plus the $r - 2$ Lobatto points (that is, the zeros of the first derivative of $\Phi_{r-1}(s)$, $s \in [-1, 1]$) shifted linearly to each subinterval (s_i, s_{i+1}) , $1 \leq i \leq N$. Thus, if we let $\zeta_{r-1} = 1$ and, for $r \geq 3$, we let $\zeta_1, \dots, \zeta_{r-2}$ be the $r - 2$ Lobatto points, then the collocation points are

$$\tau_{n,(i-1)(r-1)+j+1} = (s_i + s_{i+1} + (s_{i+1} - s_i)\zeta_j)/2, \quad 1 \leq j \leq r - 1, \quad 1 \leq i \leq N,$$

with $\tau_{n1} = s_1 = a$.

For the above sets of collocation points, the corresponding interpolatory operator P_n satisfies (3.5) and (3.6) (Vainikko and Uba [18]), and Y_n may exhibit the (super)convergence rate indicated in the next theorem. (There, $k_t(s) := k(t, s)$ for $t, s \in [a, b]$.)

THEOREM 3. *Let (y^c, λ^c) be a simple turning point of (2.1), and let (z^c, λ^c) be the corresponding simple turning point of (2.2). Suppose A1 to A6 hold, and $l(\phi^c) = 1$. If $z^c \in W_1^\mu$, $\mu \geq 1$, $\phi^c \in W_1^\delta$, $\delta \geq 1$, and $k_t \in W_1^m$, $1 \leq m \leq r$, with $\|k_t\|_{m,1}$ bounded independently of t , then for sufficiently large n ,*

$$\max(\|y^c - y_n\|_\infty, \|\psi^c - \psi_n\|_\infty, |\lambda^c - \lambda_n|) = O(h^{\gamma_\nu}),$$

where $\gamma_\nu = \min(m + r, \mu, \delta, 2r - 2\nu)$, $\nu = 0$ or 1 .

Proof. The result is derived from inequality (4.7) by the arguments of Graham, Joe and Sloan [5] for the case $\nu = 0$, and Joe [7] for the case $\nu = 1$. \square

Thus, even though z_n exhibits at most an $O(h^r)$ convergence rate, both y_n and λ_n may exhibit up to an $O(h^{2r})$ convergence rate in the case $\nu = 0$, and up to an $O(h^{2r-2})$ convergence rate in the case $\nu = 1$. Note that there is no superconvergence when continuous piecewise-linear functions ($\nu = 1$, $r = 2$) are used.

5. A Numerical Example. In this section we compute the simple turning point of an integral equation reformulation of the nonlinear two-point boundary value problem

$$y''(t) - \lambda \exp(y(t)) = 0, \quad t \in (0, 1); \quad y(0) = y(1) = 0, \quad \lambda \geq 0,$$

which arises in the theory of spontaneous combustion of an infinite slab of exothermically reacting material (Gray and Wake [6], Thomas [17]). This problem is symmetric about $t = 1/2$, and has solutions of the form

$$y(t) = \ln(\beta) - 2 \ln(\cosh((2t - 1)\sqrt{\lambda\beta/8})), \quad t \in [0, 1],$$

where β satisfies the equation

$$\beta - \cosh^2(\sqrt{\lambda\beta/8}) = 0.$$

A simple turning point exists at $\lambda^c = 3.513830719125\dots$, and there are two solutions for $0 < \lambda < \lambda^c$.

The problem may be reformulated as the integral equation

$$y(t) = -\lambda \int_0^1 k(t, s) \exp(y(s)) ds, \quad t \in [0, 1],$$

where

$$k(t, s) = \begin{cases} -s(1-t), & s \leq t, \\ -t(1-s), & s > t. \end{cases}$$

The enlarged system that we solve for (z^c, ϕ^c, λ^c) is

$$\begin{aligned} z(t) - \lambda \exp\left(-\int_0^1 k(t, s)z(s) ds\right) &= 0, \\ \phi(t) + \lambda \exp\left(-\int_0^1 k(t, s)z(s) ds\right) \int_0^1 k(t, s)\phi(s) ds &= 0, \\ \phi(1/2) - 1 &= 0. \end{aligned}$$

Results for the case of piecewise constant functions ($\nu = 0, r = 1$), equally spaced breakpoints, and a set of collocation points consisting of the midpoints of each subinterval, are displayed in Table 1. The observed rates of convergence may be deduced from the columns headed EPOH which contain estimates of the power of h . Note that the observed rates support the theoretical predictions of Corollary 1 ($\xi = 1$) and Theorem 3 ($\gamma_0 = 2$).

TABLE 1 ($\nu = 0, r = 1$)

N	$\ z^c - z_n\ _\infty$	EPOH	$\ y^c - y_n\ _\infty$	EPOH	$ \lambda^c - \lambda_n $	EPOH
4	2.60 E00	0.984	1.24 E-2	2.14	5.64 E-2	1.99
8	1.31 E00	0.994	2.83 E-3	2.03	1.42 E-2	2.00
12	8.79 E-1	0.997	1.24 E-3	2.03	6.34 E-3	2.00
16	6.60 E-1	0.998	6.91 E-4	2.05	3.57 E-3	2.00
20	5.28 E-1	0.999	4.38 E-4	2.01	2.28 E-3	2.00
24	4.40 E-1		3.03 E-4		1.59 E-3	

Table 2 shows the results for the case of discontinuous piecewise linear functions ($\nu = 0, r = 2$), equally spaced breakpoints, and a set of collocation points consisting of the two Gauss-Legendre points, $-1/\sqrt{3}$ and $1/\sqrt{3}$, shifted linearly to each subinterval. Note that the results suggest an $O(h^4)$ convergence rate for both y_n

and λ_n whereas, because $k_t(s) \in W_1^1(0, 1)$ but $k_t(s) \notin W_1^2(0, 1)$, Theorem 3 predicts a rate of only $O(h^3)$. This discrepancy occurs because the Sobolev spaces used in this paper are of integral order only: in such a setting the prediction of Theorem 3 is the best possible. If, however, the analysis were to be carried out in the fractional derivative space setting used by Joe [8] in his study of the superconvergence phenomenon for (linear) second-kind Fredholm integral equations then, for this case, an h^4 order of convergence would be predicted for both y_n and λ_n .

TABLE 2 ($\nu = 0$, $r = 2$)

N	$\ z^c - z_n\ _\infty$	EPOH	$\ y^c - y_n\ _\infty$	EPOH	$ \lambda^c - \lambda_n $	EPOH
4	5.68 E-1	1.79	7.20 E-4	3.80	9.25 E-5	7.27
8	1.64 E-1	1.93	5.16 E-5	3.89	5.99 E-7	1.98
12	7.50 E-2	1.97	1.06 E-5	3.94	2.69 E-7	3.38
16	4.26 E-2	1.98	3.42 E-6	3.97	1.02 E-7	3.67
20	2.74 E-2	1.99	1.41 E-6	3.99	4.48 E-8	3.79
24	1.91 E-2		6.81 E-7		2.25 E-8	

Acknowledgement. The author is grateful to Professor I. Sloan for assistance in the development of this work.

School of Mathematics
University of New South Wales
Sydney, NSW 2033, Australia

1. P. M. ANSELONE, "Uniform approximation theory for integral equations with discontinuous kernels," *SIAM J. Numer. Anal.*, v. 4, 1967, pp. 245-253.
2. P. M. ANSELONE, "Collectively compact approximations of integral operators with discontinuous kernels," *J. Math. Anal. Appl.*, v. 22, 1968, pp. 582-590.
3. P. M. ANSELONE, *Collectively Compact Operator Approximation Theory and Applications to Integral Equations*, Prentice-Hall, Englewood Cliffs, N. J., 1971.
4. K. ATKINSON, I. GRAHAM & I. SLOAN, "Piecewise continuous collocation for integral equations," *SIAM J. Numer. Anal.*, v. 20, 1983, pp. 172-186.
5. I. G. GRAHAM, S. JOE & I. H. SLOAN, "Iterated Galerkin versus iterated collocation for integral equations of the second kind," *IMA J. Numer. Anal.*, v. 5, 1985, pp. 355-369.
6. B. F. GRAY & G. C. WAKE, "Criticality in the infinite slab and cylinder with surface heat sources," *Combustion and Flame*, v. 55, 1984, pp. 23-30.
7. S. JOE, "Collocation methods using piecewise polynomials for second kind integral equations," *J. Comput. Appl. Math.*, v. 12 & 13, 1985, pp. 391-400.
8. S. JOE, *The Numerical Solution of Second Kind Fredholm Integral Equations*, Ph. D. Thesis, University of New South Wales, Sydney, 1985.
9. L. V. KANTOROVICH & G. P. AKILOV, *Functional Analysis*, Pergamon Press, Oxford, 1982.
10. M. A. KRASNOSEL'SKII & P. P. ZABREIKO, *Geometrical Methods of Nonlinear Analysis*, Springer-Verlag, Berlin, 1984.
11. S. KUMAR, "Superconvergence of a collocation-type method for Hammerstein equations," *IMA J. Numer. Anal.*, v. 7, 1987, pp. 313-225.
12. S. KUMAR, "The numerical solution of Hammerstein equations by a method based on polynomial collocation," 1987, submitted for publication.
13. S. KUMAR & I. H. SLOAN, "A new collocation-type method for Hammerstein integral equations," *Math. Comp.*, v. 48, 1987, pp. 585-593.
14. G. MOORE & A. SPENCE, "The calculation of turning points of nonlinear equations," *SIAM J. Numer. Anal.*, v. 17, 1980, pp. 567-576.
15. G. MOORE & A. SPENCE, "The convergence of operator approximations at turning points," *IMA J. Numer. Anal.*, v. 1, 1981, pp. 23-38.

16. A. SPENCE & G. MOORE, "A convergence analysis for turning points of nonlinear compact operator equations," in *Numerical Treatment of Integral Equations*, ISNM, vol. 53 (J. Albrecht and L. Collatz, eds.), Birkhäuser Verlag, Basel, 1980, pp. 203–212.
17. P. H. THOMAS, "On the thermal conduction equation for self-heating materials with surface cooling," *Trans. Faraday Soc.*, v. 54, 1958, pp. 60–65.
18. G. VAINIKKO & P. UBA, "A piecewise polynomial approximation to the solution of an integral equation with weakly singular kernel," *J. Austral. Math. Soc. Ser. B*, v. 22, 1981, pp. 431–438.
19. R. WEISS, "On the approximation of fixed points of nonlinear compact operators," *SIAM J. Numer. Anal.*, v. 11, 1974, pp. 550–553.