

**ON THE EQUATION $\sum_{p|N} \frac{1}{p} + \frac{1}{N} = 1$,
 PSEUDOPERFECT NUMBERS,
 AND PERFECTLY WEIGHTED GRAPHS**

WILLIAM BUTSKE, LYNDIA M. JAJE, AND DANIEL R. MAYERNIK

ABSTRACT. We present all solutions to the equation $\sum_{p|N} \frac{1}{p} + \frac{1}{N} = 1$ with at most eight primes, improve the bound on the nonsolvability of the Erdős-Moser equation $\sum_{j=1}^{m-1} j^n = m^n$, and discuss the computational search techniques used to generate examples of perfectly weighted graphs.

Recent study of the unit fraction equation

$$(1) \quad \sum_{i=1}^k \frac{1}{n_i} + \prod_{i=1}^k \frac{1}{n_i} = 1,$$


$n_1 < n_2 < \dots < n_k$, has sparked renewed interest in the relation

$$(2) \quad \sum_{p|N} \frac{1}{p} + \frac{1}{N} = 1,$$

where the sum is taken over all distinct prime divisors of N . One purpose of this paper is to present all solutions of equation (2) with $k \leq 8$ primes. There is exactly one solution for each k in this range, verifying conjectures of Ke and Sun [9], and Cao, Liu and Zhang [7]. In the second section, properties of solutions will be applied to the Erdős-Moser equation

$$(3) \quad \sum_{j=1}^{m-1} j^n = m^n.$$

We improve the bound on m to $10^{9.3 \times 10^6}$ for the conjecture that no nontrivial solution to (3) exists. Finally, we apply search techniques developed in connection with equations (1) and (2) to the topic of perfectly weighted graphs (see [4]). Specifically, for $n \geq 3$ we have found all perfectible graphs of the following form.



Received by the editor June 19, 1996 and, in revised form, March 17, 1998.
 1991 *Mathematics Subject Classification*. Primary 11D68; Secondary 11Y50, 05C50.

The authors would like to thank the referee for several important improvements to the original manuscript.

1. PRIMARY PSEUDOPERFECT NUMBERS

Recall that a positive integer is called *perfect* if it is the sum of all its proper divisors, and *pseudoperfect* if it is the sum of *some* of its proper divisors ([8, p. 46]). A positive integer $N = \prod_{i=1}^k n_i$ with factors n_i satisfying equation (1) is clearly pseudoperfect since

$$N = \sum_{i=1}^k \frac{N}{n_i} + 1.$$

All solutions n_1, \dots, n_k to equation (1) are known for $k \leq 7$ ([5], [3]). For $k = 8$, the list of known solutions continues to grow, with 89 solutions announced by Brenton and Bruner in 1994 ([2]). At present 112 solutions are known to the authors.

In the case where the divisors n_i are precisely the distinct prime divisors of N , we obtain equation (2). Conversely, since equation (2) implies that N is square-free, a solution to (2) is a special case of (1). We will call an integer $N = \prod_{i=1}^k p_i$ satisfying (2) a *primary pseudoperfect number*. Through search methods described in Section 4, all primary pseudoperfect numbers with $k \leq 8$ prime factors have been found.

Theorem 1. *Table 1 comprises the complete list of solutions to the equation*

$$\sum_{p|N} \frac{1}{p} + \frac{1}{N} = 1$$

with eight or fewer primes.

TABLE 1. Primary pseudoperfect numbers with $k \leq 8$ prime factors

k	N	Factors
1	2	2
2	6	2,3
3	42	2,3,7
4	1806	2,3,7,43
5	47058	2,3,11,23,31
6	2214502422	2,3,11,23,31,47059
7	52495396602	2,3,11,17,101,149,3109
8	8490421583559688410706771261086	2,3,11,23,31,47059,2217342227,1729101023519

No solutions to equation (2) are known of length greater than 8. We do not know whether there are infinitely many solutions. As in the case of perfect numbers, no odd primary pseudoperfect number is known.

If we allow prime *powers* among the divisors, we have two additional solutions.

k	N	Factors
7	144508961850	2,3,11,25,29,1097,2753
8	20882840055109264384350	2,3,11,25,29,1097,2753,144508961851

These, together with the 8 solutions of Table 1, constitute the list of known solutions in primes p_i to the equation

$$(4) \quad \sum_{i=1}^k \frac{1}{p_i^{\alpha_i}} + \frac{1}{\prod_{i=1}^k p_i^{\alpha_i}} = 1.$$

There is also independent interest in the companion equation

$$(5) \quad \sum_{p|N} \frac{1}{p} - \frac{1}{N} = 1.$$

Reference [1] discusses the history of this equation and presents the eleven solutions that were known as of 1996. Recently, two new solutions have been found:

$$\begin{aligned} &1910667181420507984555759916338506 \\ &= 2 * 3 * 7 * 43 * 1831 * 138683 * 2861051 * 1456230512169437 \end{aligned}$$

by M. Hogan and C. Mangilin, and

$$\begin{aligned} &4200017949707747062038711509670656632404195753751630609228764416142557211- \\ &582098432545190323474818 \\ &= 2 * 3 * 11 * 23 * 31 * 47059 * 2217342227 * 1729101023519 \\ &* 8491659218261819498490029296021 * 58254480569119734123541298976556403 \end{aligned}$$

by R. Girgensohn (both unpublished).

2. THE ERDÖS-MOSER EQUATION

More than four decades ago Paul Erdős conjectured that no solution exists to the equation

$$1^n + 2^n + \dots + (m - 1)^n = m^n$$

except the trivial solution $1^1 + 2^1 = 3^1$. Although the conjecture remains unproven (see [8, p. 153–154]), in 1953 Leo Moser [11] verified that no solution exists for $m < 10^{10^6}$. This bound has recently been used by Pieter Moree [10] to obtain similar results for the equation $\sum_{j=1}^{m-1} j^n = am^n$. Moser’s proof proceeds by using elementary number theoretic considerations to show that if (m, n) is a solution, then the following expressions involving the prime divisors of $m - 1$ and of $2m \pm 1$ must be integers:

$$(\alpha) \quad \sum_{p|(m-1)} \frac{1}{p} + \frac{1}{m-1} = t_1,$$

$$(\beta) \quad \sum_{p|(2m-1)} \frac{1}{p} + \frac{2}{2m-1} = t_2,$$

$$(\gamma) \quad \sum_{p|(2m+1)} \frac{1}{p} + \frac{4}{2m+1} = t_3.$$

Furthermore, if m is odd, then $m \equiv 3 \pmod{8}$ and

$$(\delta) \quad \sum_{p|\frac{(m+1)}{2}} \frac{1}{p} + \frac{1}{\frac{(m+1)}{2}} = t_4.$$

No solution to any of these is known for $t_i > 1$. For $t_1, t_4 = 1$, equations (α) and (δ) imply that $m - 1$ and $\frac{m+1}{2}$ are a pair of primary pseudoperfect integers. No nontrivial solution is known to either (β) or (γ) . All of this comprises strong support for the conjecture that no solution to (3) exists.

Considering first the case $m \equiv \pm 1 \pmod{6}$, Moser notes that except for the primes 2 and 3, no prime can divide any two of $m \pm 1$, $2m \pm 1$. Therefore, the prime divisors of the square-free integer $M = \frac{(4m^4 - 5m^2 + 1)}{12} = \frac{1}{12}(m-1)(m+1)(2m-1)(2m+1)$ satisfy

$$(6) \quad \sum_{p|M} \frac{1}{p} + \frac{1}{m-1} + \frac{2}{m+1} + \frac{2}{2m-1} + \frac{4}{2m+1} \\ = t_1 + t_2 + t_3 + t_4 - \frac{1}{2} - \frac{1}{3} \geq 3\frac{1}{6}.$$

In the remaining cases $m \equiv 3 \pmod{6}$ and m even, similar analysis applied to $M' = \frac{1}{4}(m-1)(m+1)(2m-1)(2m+1)$ and to $M'' = (m-1)(2m-1)(2m+1)$, respectively, lead to similar inequalities, which are greatly more restrictive than (6), since in these cases the small primes 3, respectively 2, do not appear in the sum. The bound $m > 10^{10^6}$ then follows from estimates on the rate of growth of $\sum \frac{1}{p}$ taken over all primes.

In 1953 “computation” was the unwanted stepchild of “pure” mathematics, in part because adequate computational tools were lacking. Moser himself is (justly) proud of having achieved the startlingly immense bound 10^{10^6} by techniques of analytic number theory “without laborious computations” ([11, p. 84]).

Times change: now we can actually calculate these large numbers that previously could only be roughly estimated. All calculations reported in this paper were done on a network of 20 Sun Sparc stations over the course of about 10 months.

Theorem 2. *Let (m, n) be a solution to the Erdős-Moser equation (3), with $n > 1$. Then $m > 1.485 \times 10^{9321155}$.*

Proof. As above, the critical case is $m \equiv \pm 1 \pmod{6}$. In this case put $M = \frac{4m^4 - 5m^2 + 1}{12}$. We claim that M has at least 4990906 prime factors. For if not, then

$$\sum_{p|M} \frac{1}{p} \leq \sum_{i=1}^{4990905} \frac{1}{p_i},$$

where p_i is the i th prime. But by direct computation

$$\sum_{i=1}^{4990905} \frac{1}{p_i} = 3.1666666588101728584 \dots < 3\frac{1}{6} - 10^{-9}.$$

Since, by Moser’s bound $m > 10^{10^6}$, this is less than $3\frac{1}{6} - \frac{1}{m-1} - \frac{2}{m+1} - \frac{2}{2m-1} - \frac{4}{2m+1}$, contradicting (6). Thus $M \geq \prod_{i=1}^{4990906} p_i$. Again, direct computation gives

$$\sum_{i=1}^{4990906} \log p_i = 8.5851010694053365252 \dots * 10^7.$$

Solving the resulting inequality

$$\frac{m^4}{3} > M > e^{8.5851010694053365252 * 10^7}$$

gives the required bound $m > 1.485 * 10^{9321155}$. The cases $m \equiv 0 \pmod{3}$ and m even are similar. \square

Remark. While this bound appears to be the best available by the method pioneered by Moser, the authors hope that new insights will eventually make it possible to reach the more natural benchmark 10^{10^7} .

3. PERFECTLY WEIGHTED GRAPHS

The concept of a perfectly weighted graph was introduced by Brenton and Drucker in [4] in connection with the problem of classifying isolated singular points of algebraic surfaces by properties of the local fundamental group.

Definitions. Let G be a tree (a connected graph with no circuits) on n vertices v_1, \dots, v_n , with an integer weight $w_i > 1$ assigned to each vertex v_i . Then the weighted graph $G = G(w_1, \dots, w_n)$ is called *perfectly weighted* if the corresponding matrix

$$M_G = \begin{bmatrix} w_1 & 0 & \dots & 0 \\ 0 & w_2 & 0 & \vdots \\ \vdots & 0 & \ddots & 0 \\ 0 & \dots & 0 & w_n \end{bmatrix} - [\text{the adjacency matrix of } G]$$

is positive definite with determinant 1. An unweighted tree G is *perfectible* if there exist weights w_i for its vertices such that the resulting weighted graph $G(w_1, \dots, w_n)$ is perfectly weighted.

An isolated singular point x of an m -dimensional complex analytic variety X is called *homologically trivial* if x admits a neighborhood U in X which is homeomorphic to the cone on a homology $(2m - 1)$ -sphere. That is, $H_i(\partial U, Z) = 0$ for $0 < i < 2m - 1$.

The main theorem from [4] gives the following relation between perfectly weighted graphs and homologically trivial singularities in complex dimension 2.

Lemma 3.1. *Let X be a complex surface with a singularity at the point $x \in X$ and with no other singular points. Let \tilde{X} be the minimal nonsingular model of X and let $\rho : \tilde{X} \rightarrow X$ be the minimal resolution of singularities. Denote by C the exceptional curve $\rho^{-1}(x)$, and write $C = \bigcup_{i=1}^n C_i$ with each C_i irreducible. Suppose that the resolution ρ is normal and that each component C_i is rational. Then $x \in X$ is homologically trivial if and only if the dual intersection graph of ρ is a perfectly weighted tree.*


Here the *dual intersection graph* (call it G_ρ) is the graph on vertices v_1, \dots, v_n , where v_i meets v_j if and only if C_i meets C_j in \tilde{X} and with weight w_i on v_i equal to the negative of the Chern class of the normal bundle of the embedding of C_i in \tilde{X} . The essential element of the proof is the following presentation, due to Mumford [12], of the local fundamental group. Under the hypotheses of Lemma 3.1, if G_ρ is a tree, then the first homotopy group of a tubular neighborhood $T = \rho^{-1}(\partial U)$ of the exceptional curve C in \tilde{X} is given by generators x_1, \dots, x_n with relations

$$\prod_{j=1}^n x_j^{C_i \cdot C_j} = 1$$

for all i , and $x_i x_j = x_j x_i$ if C_i meets C_j , where $C_i \cdot C_j$ is the intersection number (the negative of the i, j th entry of G_ρ). Since the intersection matrix $[C_i \cdot C_j]$ is always negative definite in the complex case, the corresponding first homology group

$H_1(T, Z) = \pi_1(T)/(xyx^{-1}y^{-1} = 1)$ is a finite group of order $D = (-1)^n \det[C_i \cdot C_j] = \det[M_{G_\rho}]$. Thus, if $x \in X$ is homologically trivial, then $\pi_1(T)$ is a *perfect group* (generated by commutators), and the converse follows from Poincaré duality.

In the special case in which $G = G_\rho$ is the weighted star



direct computation shows that

$$D = \det[M_G] = \prod_{i=0}^k n_i - \sum_{i=1}^k \prod_{j \neq i} n_j.$$


This is equivalent to

$$\sum_{i=1}^k \frac{1}{n_i} + \frac{D}{\prod_{i=1}^k n_i} = n_0,$$

which exhibits the connection between this topic and our equations (1) and (2). Explicitly (allowing $n_0 = 1$), the group generated by x_0, \dots, x_n with relations $x_0 = \prod_{i=1}^k x_i = x_i^{n_i}$ for all i is perfect if and only if the integer $N = \prod_{i=1}^k n_i$ is pseudoperfect with factors n_i satisfying equation (1). We find it interesting that although the terms “perfect number” and “perfect group” were coined independently, the results of this paper reveal a relation between these apparently disparate topics.

From the point of view of number theory, perhaps the most interesting graphs are the so-called “weighted flowers”, which are weighted graphs of the form

$$J_{k,m} = J_{k,m}(n_1, \dots, n_k; w_0, \dots, w_m)$$



Lemma 3.2. *For the weighted graph $G = J_{k,m}(n_1, \dots, n_k; w_0, \dots, w_m)$ pictured above, we have*

$$(7) \quad \sum_{i=1}^k \frac{1}{n_i} + \frac{D}{Q \prod_{i=1}^k n_i} = \frac{P}{Q},$$

where $\frac{P}{Q}$ has the continued fraction expansion

$$[w_0, w_1, \dots, w_m] = w_0 - \frac{1}{w_1 - \frac{1}{w_2 - \frac{1}{\ddots - \frac{1}{w_m}}}}$$

and where $D = \det[M_G]$.


A simple proof follows by induction on m (cf. [4, Lemma 4.3]).

In view of Lemma 3.2, we can use computational techniques similar to those employed in finding solutions of equation (1) to find perfect weights for graphs of this type. A perfectible graph is called *minimal* if it contains no proper perfectible subgraphs. Table 2 presents the complete list of minimal perfectible flowers $J_{k,m}$ with $m > 2$. Since a graph containing a perfectible subgraph is itself perfectible, we have the following result.

Theorem 3. *Let $G = J_{k,m}$ be a flower with $m > 2$. Then G is perfectible if and only if G contains one of the five graphs in Table 2.*

Verifying that each of these weighted graphs is perfectly weighted is a direct application of Lemma 3.2. The proof of Theorem 3 consists of verifying that each graph is minimal and that the list is complete. This was accomplished by exhaustive searches, as discussed in Section 4.

The graphs in Table 2 are pictured with one set of perfect weights. The perfect weights may not be unique. For instance,



is another set of perfect weights on $J_{6,5}$.

The graphs $J_{3,28}$, $J_{4,6}$, and $J_{6,5}$ were discussed in [4]. $J_{10,3}$ was also introduced in [4], but at that time it was not known whether it was minimal or not. $J_{7,4}$ was derived only in 1995 after an earlier discovery by K. Conway (unpublished) of a set of perfect weights for $J_{8,4}$ shown below.





TABLE 2. Minimal perfectible flowers



4. SEARCH TECHNIQUES

The computational results reported thus far stem from finding solutions to the equation

$$(8) \quad \sum_{i=1}^k \frac{1}{n_i} + \frac{1}{Q \prod_{i=1}^k n_i} = \frac{P}{Q}.$$

Our main computational tool is Lemma 4.1, which gives a criterion for extending a solution of equation (7) to a solution of equation (8) by the adjunction of two more terms. Lemmas 4.1 and 4.2 generalize results of [5, Proposition 12 and Lemma 17].

Lemma 4.1. *Given a positive integer P and relatively prime positive integers n_1, \dots, n_k, Q , write*

$$\sum_{i=1}^k \frac{1}{n_i} + \frac{D}{Q \prod_{i=1}^k n_i} = \frac{P}{Q},$$

where $D = (\frac{P}{Q} - \sum_{i=1}^k \frac{1}{n_i})Q \prod_{i=1}^k n_i$. Let F be a factor of $Y = Q^2 \prod_{i=1}^k n_i^2 + D$ and write $Y = FG$. Suppose that F (and hence also G) is congruent to $-Q \prod_{i=1}^k n_i \pmod{D}$ and put

$$n_{k+1} = \frac{Q \prod_{i=1}^k n_i + F}{D} \quad \text{and} \quad n_{k+2} = \frac{Q \prod_{i=1}^k n_i + G}{D}.$$

Then the integers n_1, \dots, n_{k+2} satisfy the equation

$$\sum_{i=1}^{k+2} \frac{1}{n_i} + \frac{1}{Q \prod_{i=1}^{k+2} n_i} = \frac{P}{Q}.$$

Proof.

$$\begin{aligned} & \sum_{i=1}^k \frac{1}{n_i} + \frac{1}{n_{k+1}} + \frac{1}{n_{k+2}} + \frac{1}{Q \prod_{i=1}^k n_i (n_{k+1})(n_{k+2})} \\ &= \frac{P}{Q} - \frac{D}{Q \prod_{i=1}^k n_i} + \frac{D}{Q \prod_{i=1}^k n_i + F} + \frac{D}{Q \prod_{i=1}^k n_i + G} \\ & \quad + \frac{D^2}{Q \prod_{i=1}^k n_i (Q \prod_{i=1}^k n_i + F)(Q \prod_{i=1}^k n_i + G)} \\ &= \frac{P}{Q} + \frac{D(Y - FG)}{Q \prod_{i=1}^k n_i (Q \prod_{i=1}^k n_i + F)(Q \prod_{i=1}^k n_i + G)} = \frac{P}{Q}. \quad \square \end{aligned}$$

In addition, given a partial solution n_1, n_2, \dots, n_i , we know the bounds on a search for n_{i+1} .

Lemma 4.2. *Let $n_1 < n_2 < \dots < n_k, k > 2$, satisfy equation (8). Then for each index $i \leq k - 2$, we have*

$$\left(\frac{P}{Q} - \sum_{j=1}^i \frac{1}{n_j} \right)^{-1} < n_{i+1} < (k - i) \left(\frac{P}{Q} - \sum_{j=1}^i \frac{1}{n_j} \right)^{-1}.$$

Proof.

$$\frac{1}{n_{i+1}} = \frac{P}{Q} - \sum_{j \neq i+1} \frac{1}{n_j} - \frac{1}{Q \prod_{j=1}^k n_j} < \frac{P}{Q} - \sum_{j=1}^i \frac{1}{n_j},$$

so $n_{i+1} > (\frac{P}{Q} - \sum_{j=1}^i \frac{1}{n_j})^{-1}$ as required.

On the other hand, since $n_1 < n_2 < \dots < n_k$, we have

$$\begin{aligned} (k-i) \frac{1}{n_{i+1}} &\geq \frac{1}{n_{i+1}} + \frac{1}{n_{i+2}} + \left(\frac{1}{n_{i+1}} - \frac{1}{n_{i+2}} \right) + \sum_{j=i+3}^k \frac{1}{n_j} \\ &= \sum_{j=i+1}^k \frac{1}{n_j} + \frac{n_{i+2} - n_{i+1}}{n_{i+1}n_{i+2}} \\ &> \sum_{j=i+1}^k \frac{1}{n_j} + \frac{1}{Q \prod_{j=1}^k n_j} = \frac{P}{Q} - \sum_{j=1}^i \frac{1}{n_j}, \end{aligned}$$

and thus $n_{i+1} < (k-i)(\frac{P}{Q} - \sum_{j=1}^i \frac{1}{n_j})^{-1}$ as claimed. \square


To implement these ideas in a search program for fixed $\frac{P}{Q}$ and k , we use Lemma 4.2 to determine all possibilities for n_1, \dots, n_{k-2} , then we determine n_{k-1} and n_k by the technique of Lemma 4.1. The advantage of this method over simply searching for all possibilities for n_{k-1} (as many as 10^{13} choices for $k = 8$ and $\frac{P}{Q} = 1$) is that we can reduce computation time by making use of advanced factoring techniques. In contrast, a complete tree search for n_{k-1} , using the bounds of Lemma 4.2, is equivalent to factoring the large integer Y of Lemma 4.1 by trial division up to the square root. Computation time could be further reduced by incorporating the required congruence relations $F, G \equiv -Q \prod_{i=1}^k n_i \pmod{D}$ into the factoring methods and by taking advantage of the special form $Y = (Q \prod n_i)^2 + D$ for a known small number D . This program has proven to be the most useful tool for finding solutions to equation (8) and its special cases, equations (1) and (2). It yields both *nonsporadic solutions* (solutions of length k resulting from extending known solutions of length $k-1$ or $k-2$) and *sporadic solutions* (solutions not generated from such solutions of smaller length).

These searches have produced the following results. With respect to equation (1), 68 nonsporadic and 44 sporadic solutions have been discovered for $k = 8$. The 68 nonsporadic solutions are easy to find and were discussed in [3]. The 44 sporadic solutions include all except those in the string 2,3,7,43. The search is also complete with respect to solutions in *prime* integers n_i , giving a proof of the completeness of the list of primary pseudoperfect numbers in Table 1.

Similar computational searches give results about particular perfectible graphs. $J_{8,5}$, for instance, admits at least 21 sets of perfect weights. Sixteen of these sets result from extending perfect weights on the minimal perfectible graph $J_{6,5}$, four of them result from extending $J_{7,4}$, and one from extending $J_{8,4}$. These are the nonsporadic solutions, and there are possibly sporadic solutions for perfect weights on $J_{8,5}$ which have not yet been explored.

A special case of Lemma 3.2 reveals further interesting properties of the graphs $J_{k,m}$ and a tighter relation between the topics of perfectible graphs and pseudoperfect numbers.

Proposition 4.1. *The weighted graph*



is perfectly weighted if and only if n_1, \dots, n_k satisfy equation (1), where $n_k = m^2(w - 1) + m$.

Proof. Direct computation verifies that the continued fraction

$$[2, 2, \dots, 2, w] = \frac{(m + 1)w - m}{mw - (m - 1)} = 1 + \frac{1}{m} - \frac{1}{n_k}$$

for n_k as above. Thus, by Lemma 3.2 we have

$$\sum_{i=1}^{k-1} \frac{1}{n_i} + \frac{1}{m} + \frac{D}{(mw - (m - 1))m \prod_{i=1}^{k-1} n_i} = 1 + \frac{1}{m} - \frac{1}{n_k},$$

or

$$\sum_{i=1}^k \frac{1}{n_i} + \frac{D}{\prod_{i=1}^k n_i} = 1,$$

where D is the determinant of the weighted graph. Hence $D = 1$ if and only if n_1, \dots, n_k satisfy equation (1). \square

To apply this result we need only find solutions n_1, \dots, n_k to equation (1) in which one of the n_i 's happens to be congruent to $m \pmod{m^2}$ for some m (but $n_i \neq m$ to ensure that $w = 1 + (n_i - m)/m^2$ is greater than 1). For $k \leq 8$ we found 24 distinct solution sets n_1, \dots, n_k which contain an n_i with this special property for some integer m . Three of these sets have two different n_i 's with this property, and two have an n_i which satisfies this congruence for two different m 's. This gives a total of 29 examples of perfectly weighted graphs of the type $J_{k,m}$ with $k \leq 8$ and with weights as pictured in Proposition 4.1. They are presented in Table 3.

For $m < 5$ there are no solutions of this type for $k < 10$. The most challenging case in this range was $k = 9$ and $m = 3$. In this instance we found that there are only 5 solutions to equation (1) with no $n_i = 3$:

- 2, 5, 7, 9, 31, 73, 13327, 63582361, 110273083859;
- 2, 5, 7, 9, 37, 61, 383, 3226871, 2344136699;
- 2, 5, 7, 11, 17, 149, 1431, 64911433, 1169526576259;
- 2, 5, 7, 11, 17, 157, 961, 4398619, 8687184244716671;
- 2, 5, 7, 11, 17, 167, 1257, 1919, 9373.

This leads immediately to the result that $J_{9,3}$ is not perfectible. First, it is easy to reduce the general case of perfect weights for $J_{9,3}$ to those pictured in Proposition 4.1. Then we check that none of the n_i 's appearing in the five solutions above is congruent to 3 mod 9. In a similar manner, other graphs of type $J_{k,m}$ can be shown not to be perfectible, resulting in a proof of Theorem 3.

TABLE 3. Special perfect weights on $J_{k,m}, k \leq 8$


n_1, n_2, \dots, n_{k-1}	m	w
2,3,7,179,24323	5	3
2,3,7,55,179,24323	67	2240437
2,3,7,179,24323,10057317271	5	3
2,3,11,23,31,211031	71	13
2,3,11,23,31,12017087	7	965
2,3,7,43,1807,3263443,134811739261383753719	5	426002311687
2,3,7,43,1807,3263479,243811701792623	5	11527311163
2,3,7,43,1823,193667,637617223459	5	1250940688133154818523
2,3,7,43,1823,193667,637617223459	31	32542681793266254385
2,3,7,43,1831,132347,231679879	17	4142701692187
2,3,7,47,395,277442411,1701723083	361	7
2,3,7,47,403,19403,15435516179	5	3387914913502507
2,3,7,55,179,24323,101149630679497570171	67	2240437
2,3,7,55,179,24323,513449911932648503	37	7346471
2,3,7,179,24323,10057317271,101149630679497570171	5	3
2,3,7,179,24323,10057317287,5949978284730273323	5	3
2,3,7,179,24323,10057317311,2467064172726591731	5	3
2,3,7,179,24323,10057317467,513449911932648503	5	3
2,3,7,179,24323,10057317967,145121431390804003	5	3
2,3,7,179,24323,10057320619,30202945461748519	5	3
2,3,7,179,24323,10057325347,12523178395739983	5	3
2,3,7,179,24323,10057454579,736667018400959	5	3
2,3,11,17,79,1049,3696653	7	7
2,3,11,23,31,47059,3375982667	5	257468755
2,3,11,23,31,47059,165128325167	5	89784175
2,3,11,23,31,47059,165128325167	7	45808253
2,3,11,23,31,47147,11061526082145911	17	86259
2,3,11,23,31,211031,601432790177275	71	13
2,3,11,23,31,12017087,26715920281613179	7	965

Although $J_{9,3}$ is not perfectible, each of the five solutions to equation (1) with no $n_i = 3$ results in a perfectible flower of type $J_{9,6}$. They are presented in Table 4.

Solutions to equation (5), or more generally to






$$(9) \quad \sum_{i=1}^k \frac{1}{n_i} - \prod_{i=1}^k \frac{1}{n_i} = 1$$

also lead to perfect weights for graphs of type $J_{k,m}$. Namely, if n_1, \dots, n_k satisfy equation (9), then the graph



is perfectly weighted for $m = \prod_{i=1}^k n_i - 2$. Again, Lemma 3.2 provides the proof. All solutions to equation (9) are known for $k \leq 7$ (there are 50 of them), and more than 400 are known for $k = 8$ ([6], [1]).

TABLE 4. Examples of perfect weights on $J_{9,6}$



REFERENCES

1. D. Borewein, J. M. Borwein, P. B. Borwein and R. Girgensohn, *Giuga's conjecture on primality*, Amer. Math. Monthly **103** (1996), 40–50. MR **97b**:11004
2. L. Brenton and R. Bruner, *On recursive solutions of a unit fraction equation*, J. Aust. Math. Soc. **57** (1994), 341–356. MR **95i**:11024
3. L. Brenton and D. Drucker, *On the number of solutions of $\sum_{j=1}^s (\frac{1}{x_j}) + \frac{1}{(x_1 \dots x_s)} = 1$* , J. Number Theory **44** No. 2 (1993), 25–29. MR **94b**:11029
4. ———, *Perfect graphs and complex surface singularities with perfect local fundamental group*, Tôhoku Math. J. **41** (1989), 507–525. MR **91g**:14025
5. L. Brenton and R. Hill, *On the Diophantine equation $1 = \sum (\frac{1}{n_i}) + \frac{1}{(\prod n_i)}$ and a class of homologically trivial complex surface singularities*, Pacific J. Math. **133** (1988), 41–67. MR **89d**:32023
6. L. Brenton and M. K. Joo, *On the system of congruences $\prod_{j \neq i} n_j \equiv 1 \pmod{n_i}$* , The Fibonacci Quarterly **33** No. 3 (June–July 1995), 258–266. MR **96k**:11039
7. Z. Cao, R. Liu and L. Zhang, *On the equation $\sum_{j=1}^s (\frac{1}{x_j}) + \frac{1}{(x_1 \dots x_s)} = 1$ and Zám's problem*, J. Number Theory **27** No. 2 (1987), 206–211. MR **89d**:11023
8. R. Guy, *Unsolved Problems in Number Theory (2nd ed.)*, vol. I, Springer-Verlag, New York, 1994. MR **96e**:11002
9. Z. Ke and Q. Sun, *On the representation of 1 by unit fractions*, Sichuan Daxue Xuebao **1** (1964), 13–29.
10. P. Moree, *Diophantine equations of Erdős-Moser type*, Bull. Austral. Math. Soc. **53** (1996), 281–292. MR **97b**:11048
11. L. Moser, *On the Diophantine equation $1^n + 2^n + 3^n + \dots + (m-1)^n = m^n$* , Scripta Math. **19** (1953), 84–88. MR **14**:950g
12. D. Mumford, *The topology of normal singularities of an algebraic surface and a criterion for simplicity*, Inst. Hautes Études Sci. Publ. Math. No. **9** (1961), 5–22. MR **27**:3643

DEPARTMENT OF MATHEMATICS, WAYNE STATE UNIVERSITY, 1150 FAB, DETROIT, MICHIGAN 48202

Current address: Department of Mathematics, Purdue University, West Lafayette, Indiana 47906

E-mail address: butske@math.purdue.edu

Current address: EDS Office Centre, Mailstop 2061, 300 E. Big Beaver Road, Troy, Michigan 48083

E-mail address: lynda.jaje@eds.com

E-mail address: mayernik@math.wayne.edu