

**CORRIGENDUM TO  
“SPHERICAL MARCINKIEWICZ-ZYGMUND INEQUALITIES  
AND POSITIVE QUADRATURE”**

H. N. MHASKAR, F. J. NARCOWICH, AND J. D. WARD

Professor Dr. Joachim Stöckler has kindly pointed out to us that our proofs [2] of the estimates (4.5) and (4.9) are valid only in the case when we do not require the quadrature weights  $a_\xi$  to be nonnegative, since the Krein-Rutman extensions of nonnegative functionals are not guaranteed to be norm-preserving. The purpose of this note is to point out that the existence of nonnegative weights  $a_\xi$  satisfying (4.4) necessarily implies the following analogue (0.1) of (4.5). The existence of such weights was proved in the paper. (Here, and in the sequel, we use all the notation as in the paper.)

$$(0.1) \quad \left\| \frac{a_\xi}{\mu_q(R_\xi)} \right\|_{C, p'} \leq c, \quad 1 \leq p' \leq \infty.$$

The estimate (4.9) (being a special case of (4.5)) should also be modified accordingly in the case of nonnegative weights. The estimates are correct as stated in the case when the quadrature weights are not required to be nonnegative.

First, we prove (0.1) in the case when  $p' = \infty$ . Let  $N$  be an even integer with  $\delta_{C_0} \sim 1/N$ , and such that the quadrature formula (4.4) is exact for  $\Pi_N^q$ . Now fix  $\xi$ , and let  $S$  be the spherical cap circumscribing  $R_\xi$ . Then the proof of Proposition 3.2 shows that  $\mu_q(R_\xi) \sim \mu_q(S)$ . Moreover, one can write  $S := \{\mathbf{x} : \mathbf{x} \cdot \zeta \geq \cos \theta_\xi\}$  with  $\theta_\xi \sim N^{-1}$ . Now, let  $y := \cos \theta_\xi$  and  $\chi$  be the univariate function defined by

$$\chi(t) = \begin{cases} 1 & \text{if } y < t \leq 1, \\ 0 & \text{if } -1 \leq t \leq y. \end{cases}$$

Then there exist ([1, Section I.5]) polynomials  $\phi$  and  $\Phi$  in  $\Pi_N^1$  such that

$$(0.2) \quad \phi(t) \leq \chi(t) \leq \Phi(t), \quad t \in [-1, 1],$$

and

$$(0.3) \quad \int_{-1}^1 (\Phi(t) - \phi(t)) w_q(t) dt \leq \lambda_{N/2}(y),$$

---

Received by the editor September 21, 2001.

2000 *Mathematics Subject Classification*. Primary 65D32; Secondary 41A17, 42C10.

The research of the first author was supported, in part, by grant DMS-9971846 from the National Science Foundation and grant DAAD19-01-1-0001 from the U.S. Army Research Office.

The research of the second and third authors was sponsored by the Air Force Office of Scientific Research, Air Force Materiel Command, USAF, under grant number F49620-98-1-0204.

The research of the third author was also supported by grant DMS-9971276 from the National Science Foundation.

where  $\lambda_{N/2}$  denotes the Christoffel function for the weight  $w_q$ . In view of the quadrature formula (4.4), one obtains from (0.2) that

$$\begin{aligned} \int_{\mathbb{S}^q} \phi(\mathbf{x} \cdot \zeta) d\mu_q(\mathbf{x}) &= \sum_{\mathbf{z} \in \mathcal{C}_0} a_{\mathbf{z}} \phi(\mathbf{z} \cdot \zeta) \leq \sum_{\mathbf{z} \in \mathcal{C}_0} a_{\mathbf{z}} \chi(\mathbf{z} \cdot \zeta) \\ &\leq \sum_{\mathbf{z} \in \mathcal{C}_0} a_{\mathbf{z}} \Phi(\mathbf{z} \cdot \zeta) = \int_{\mathbb{S}^q} \Phi(\mathbf{x} \cdot \zeta) d\mu_q(\mathbf{x}). \end{aligned}$$

Further, we obtain from (0.2) also that

$$\int_{\mathbb{S}^q} \phi(\mathbf{x} \cdot \zeta) d\mu_q(\mathbf{x}) \leq \int_{\mathbb{S}^q} \chi(\mathbf{x} \cdot \zeta) d\mu_q(\mathbf{x}) = \mu_q(S) \leq \int_{\mathbb{S}^q} \Phi(\mathbf{x} \cdot \zeta) d\mu_q(\mathbf{x}).$$

Therefore, in view of (0.3), we obtain

$$\begin{aligned} (0.4) \quad &\left| \sum_{\mathbf{z} \in \mathcal{C}_0} a_{\mathbf{z}} \chi(\mathbf{z} \cdot \zeta) - \mu_q(S) \right| \leq \int_{\mathbb{S}^q} (\Phi(\mathbf{x} \cdot \zeta) - \phi(\mathbf{x} \cdot \zeta)) d\mu_q(\mathbf{x}) \\ &= c \int_{-1}^1 (\Phi(t) - \phi(t)) w_q(t) dt \leq c \lambda_{N/2}(y). \end{aligned}$$

Now, using [3, Lemma 5, p. 108] with  $a = b = q/2 - 1$  and taking into account the fact that  $y = \cos \theta_\xi$  with  $\theta_\xi \sim 1/N$ , we deduce that

$$\lambda_{N/2}(y) \leq \frac{c}{N} \left( \sqrt{1-y} + 2/N \right)^{2a+1} \leq cN^{-q} \leq c\mu_q(S).$$

Therefore, from (0.4),

$$a_\xi \leq \sum_{\mathbf{z} \in \mathcal{C}_0} a_{\mathbf{z}} \chi(\mathbf{z} \cdot \zeta) \leq c\mu_q(S) \leq c\mu_q(R_\xi).$$

This proves the estimate (0.1) in the case when  $p' = \infty$ . The case when  $p' = 1$  follows by taking  $P$  to be the polynomial identically equal to 1 in (4.4). In the case when  $1 < p' < \infty$ , we use these cases to conclude that

$$\left\| \frac{a_\xi}{\mu_q(R_\xi)} \right\|_{C, p'}^{p'} = \sum_{\xi \in \mathcal{C}_0} a_\xi \left\{ \frac{a_\xi}{\mu_q(R_\xi)} \right\}^{p'-1} \leq c.$$

#### REFERENCES

1. G. Freud, *Orthogonal polynomials*, Pergamon Press, 1971.
2. H. N. Mhaskar, F. J. Narcowich, and J. D. Ward, *Spherical Marcinkiewicz-Zygmund inequalities and positive quadrature*, *Math. Comp.*, **70** (2001), 1113–1130.
3. P. Nevai, *Orthogonal polynomials*, *Mem. Amer. Math. Soc.*, **18** (1979) no. 213. MR **80k**:42025

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE UNIVERSITY, LOS ANGELES, CALIFORNIA 90032

*E-mail address*: hmhaska@calstatela.edu

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843-3368

*E-mail address*: fnarc@math.tamu.edu

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843-3368

*E-mail address*: jward@math.tamu.edu