

A FIXED POINT THEOREM FOR PSEUDO-ARCS AND CERTAIN OTHER METRIC CONTINUA

O. H. HAMILTON

E. E. Moise has defined a pseudo-arc [1]¹ and has shown that it has the property of being homeomorphic to each of its nondegenerate subcontinua. Bing has shown that a pseudo-arc is homogeneous [2]. F. B. Jones, in a letter to the author, has raised the question as to whether a pseudo-arc has the fixed point property with respect to continuous transformations. It is the purpose of this note to show that a pseudo-arc is a member of a more general class of metric continua which have the fixed point property.

We shall use Moise's definition of a chain, namely: a chain Y is a collection of mutually exclusive open sets (called links), $y_1, y_2, y_3, \dots, y_k$, such that y_i and y_j have a boundary point in common if and only if i and j are identical or consecutive integers. If Y designates a chain, $C(Y)$ will designate the closure of the set of points each of which is in a link of the chain Y , and ΔY will designate the maximum diameter of a link of the chain Y . For convenience the closure of a link, $C(y_i)$, of Y will be called a closed link of Y . A closed link $C(y_i)$ of Y will be said to precede $C(y_j)$ in Y if $i < j$ and to follow $C(y_j)$ if $i > j$.

THEOREM. *Let Y_1, Y_2, Y_3, \dots be a sequence of chains such that: (1) $C(Y_1)$ is a compact nonvacuous metric space, (2) $C(Y_{i+1})$ is a subset of $C(Y_i)$ for each i , (3) $\lim_{i \rightarrow \infty} \Delta Y_i = 0$. Let M designate the continuum which is the intersection of the sets $C(Y_i)$. Then if T is a continuous transformation of M into a subset of itself, there is a point p of M such that $T(p) = p$.*

PROOF. Let ϵ be a positive real number. Let m be a positive integer such that ΔY_m is less than ϵ . Let A be the subset of M consisting of all points p of M such that either each of the closed links of Y_m which contain $T(p)$ follows all the closed links of Y_m which contain p or such that $T(p)$ is contained in some closed link of Y_m which contains p . Let B be the subset of M consisting of all points p of M such that either each closed link of Y_m which contains $T(p)$ precedes all the closed links of Y_m which contain p or such that $T(p)$ is contained in some closed link of Y_m which contains p . The points of M in the first closed link of Y_m are in A and the points of M in the last

Presented to the International Congress of Mathematicians, September 6, 1950; received by the editors December 28, 1949.

¹ Numbers in brackets refer to the references cited at the end of the paper.

closed link of Y_m are in B , and hence these sets are nonvacuous. It is easily shown that A and B are closed. Then the compact continuum M is the sum of the two closed sets A and B , which therefore have a point q in common. Hence q and $T(q)$ lie together in some closed link z of Y_m , and the distance from q to $T(q)$ is less than ϵ . Since ϵ is an arbitrary positive number, and since T is continuous and M is closed, it follows that for some point p of M , $T(p) = p$.

COROLLARY. A pseudo-arc has the fixed point property for continuous transformations.

PROOF. A pseudo-arc is a point set satisfying the definition of the set M of the theorem.

It should be noted that the theorem applies to certain indecomposable continua which are not pseudo-arcs. An example is the one given by Brouwer [3] and cited by Urysohn [4].

REFERENCES

1. E. E. Moise, *An indecomposable plane continuum which is homeomorphic to each of its non-degenerate subcontinua*, Trans. Amer. Math. Soc. vol. 63 (1948) pp. 581-594.
2. R. H. Bing, *A homogeneous indecomposable plane continuum*, Duke Math. J. vol. 15 (1948) pp. 729-742.
3. L. E. J. Brouwer, *Zur Analysis Situs*, Math. Ann. vol. 68 (1910) pp. 422-434.
4. Paul Urysohn, *Mémoire sur les multiplicités cantorienes*, Fund. Math. vol. 8 (1926) pp. 225-251.

OKLAHOMA A. AND M. COLLEGE