

**THE FUNDAMENTAL GROUP OF THE PRINCIPAL  
COMPONENT OF A COMMUTATIVE  
BANACH ALGEBRA<sup>1</sup>**

EDWARD K. BLUM

We consider an arbitrary commutative Banach algebra over the complex numbers. Let  $B$  denote the algebra,  $\{a, b, c, x, y, z, \dots\}$  its elements, and  $\{\lambda, \mu, \nu, \dots\}$  complex numbers. We assume that  $B$  contains a unit element,  $e$ , with  $\|e\| = 1$ .

If  $a^{-1}$  exists ( $aa^{-1} = a^{-1}a = e$ ), the element  $a$  is called "regular." The set of regular elements will be denoted by  $G$ . It is well known that (1)  $G$  is a topological group relative to multiplication and (2)  $G$  is an open subset of  $B$ . Since  $G$  is open, it is a union of maximal open connected sets, its components. We call the component  $G_1$  containing the unit  $e$  the "principal component" [1]. It is easy to see that  $G_1$  is a subgroup of  $G$ .

The function  $\exp(x) \equiv e + \sum_1^\infty x^n/n!$  is defined for all  $x$  in  $B$  and has the usual properties of the classical exponential function. If we let  $\pi_1(G_1)$  denote the fundamental group of  $G_1$ , we may state our main result as follows:

**THEOREM 1.** *Let  $P = \{x | \exp(x) = e\}$ .  $P$  is an additive group which is isomorphic to  $\pi_1(G_1)$ .*

We shall give a complete proof based on Schreier's theory of the universal covering group and then we shall outline a second proof which depends only on results from the theory of Banach algebras.<sup>2</sup> The result of Schreier which we shall use may be stated as follows [2]:

**THEOREM.** *Let  $B$  be a simply-connected, locally-connected and locally simply-connected topological group. If  $P$  is a discrete normal subgroup of  $B$ , then the fundamental group of the topological space  $B/P$  is isomorphic to the group  $P$ .*

The algebra  $B$ , regarded as an additive group with the metric topology of the norm, clearly satisfies the hypotheses of Schreier's

---

Presented to the Society, February 28, 1953; received by the editors September 10, 1952.

<sup>1</sup> This result is contained in the author's doctoral dissertation, *The theory of analytic functions in Banach algebras*, completed in June 1952 under E. R. Lorch at Columbia University.

<sup>2</sup> The second proof is the one used in the author's dissertation. It was later pointed out by S. Eilenberg that a different proof is possible if one brings the Schreier theory to bear.

theorem. The set  $P = \{x \mid \exp(x) = e\}$  is obviously a normal subgroup of  $B$  since  $\exp(x - y) = \exp(x) \exp(-y) = \exp(x) [\exp(y)]^{-1}$ . Further,  $\exp(x)$  maps  $B$  onto the principal component,  $G_1$  [3]. It is clearly a continuous map. It is also an open mapping since its inverse,  $\log y$ , is continuous. Thus,  $\exp(x)$  is an open homomorphism of the additive topological group  $B$  onto the multiplicative topological group  $G_1$ . The kernel of this homomorphism is  $P$ , so that  $B/P$  is isomorphic to  $G_1$  as topological groups. Hence  $\pi_1(B/P)$  is isomorphic to  $\pi_1(G_1)$ . If it can be shown that  $P$  is discrete, the Schreier theorem is applicable and Theorem 1 follows immediately.

To prove  $P$  discrete it suffices to show that 0 is an isolated element of  $P$ . Suppose there is a sequence of elements  $\{z_n\}$  such that  $z_n \in P$ ,  $z_n \neq 0$  and  $\lim z_n = 0$ . By a theorem of Lorch [1], this implies that  $z_n = 2\pi i \sum_{j=1}^k n_j e_j$ , where the  $e_j$  are idempotent elements ( $e_j^2 = e_j$ ) and the  $n_j$  are rational integers. Furthermore, the spectrum of  $z_n$  consists of the points  $2\pi i n_1, \dots, 2\pi i n_k$ . However,  $\lim \|z_n\| = 0$  and, as is well known, the spectrum of  $z_n$  contains no points exterior to the circle of radius  $\|z_n\|$ , center at the origin. For sufficiently large  $n$ , this means that  $n_j = 0$  for  $j = 1, \dots, k$ ; i.e.  $z_n = 0$ . This contradiction completes the proof.

Now we shall indicate a more elementary and constructive proof in which no recourse is had to the Schreier theory.

First we establish a lemma concerning the function  $\exp(x)$ .

LEMMA: Let  $w_0 = \exp(x_0)$  and  $0 < \epsilon < 1/\|w_0^{-1}\|$ . Let

$$\delta = \sum_1^\infty \frac{(\|w_0^{-1}\|\epsilon)^n}{n}$$

If  $E$  is the set  $\{w \mid \|w - w_0\| < \epsilon\}$ , then for every  $w$  in  $E$  there is an  $x$  such that  $\|x - x_0\| < \delta$  and  $\exp(x) = w$ .

PROOF. Choosing  $w = w_0 + b$  where  $\|b\| < \epsilon$ , we have  $\|w_0^{-1}w - e\| = \|w_0^{-1}b\| \leq \|w_0^{-1}\|\|b\| < 1$ . Hence  $w_0^{-1}w$  is in  $G_1$  and there is an element  $c$  in  $B$  such that  $\exp(c) = w_0^{-1}w$ . In fact, we may take

$$c = \sum_1^\infty - (1/n)(e - w_0^{-1}w)^n$$

so that  $\|c\| < \sum_1^\infty (1/n)(\|w_0^{-1}\|\epsilon)^n = \delta$ . The element  $x = x_0 + c$  satisfies the conclusion of the lemma. It is important to note that  $\delta$  approaches 0 as  $\epsilon$  approaches 0.

Using this lemma, we are able to prove

THEOREM 2. Let  $K: \{f(s), 0 \leq s \leq 1\}$  be a curve in  $G_1$  joining  $e = f(0)$

to  $w=f(1)$ . There exists an element  $z$  in  $B$  such that  $\exp(z)=w$  and the curve  $K(z): \{\exp(tz), 0 \leq t \leq 1\}$  is homotopic to  $K$  in  $G_1$ .

PROOF. If  $z_s$  is such that  $\exp(z_s)=f(s)$ , let  $K(z_s)$  denote the curve  $\{\exp(tz_s), 0 \leq t \leq 1\}$ .  $K(z_s)$  is a compact set in  $G_1$ . Hence, there is a number  $\rho > 0$  such that every sphere with center on  $K(z_s)$  and radius  $\rho$  is contained in  $G_1$ . Choose  $\epsilon$  such that  $0 < \epsilon < \min\{\rho, 1\}$ . There is a number  $\gamma(\epsilon) > 0$  such that  $\|f(r)-f(s)\| < \epsilon$  whenever  $|r-s| < \gamma(\epsilon)$ . By the lemma, there is an element  $z_r$  such that  $\|z_r-z_s\| < \delta$  and  $\exp(z_r)=f(r)$ , where  $\delta = \sum_1^\infty (1/n)(\|[f(s)]^{-1}\|\epsilon)^n$ , that is,  $z_r=z_s+b$  where  $\|b\| < \delta$ .

For all  $t, 0 \leq t \leq 1, \|\exp(tz_r) - \exp(tz_s)\| \leq \|\exp(tz_s)\| \cdot \|\exp(tb) - e\| \leq \sum_0^\infty (1/n!) \|z_s\|^n \cdot \sum_1^\infty (1/n!) \|b\|^n \leq \exp\|z_s\| \sum_1^\infty \delta^n/n!$ . By choosing  $\epsilon$  sufficiently small, thereby making  $\delta$  small, we have  $\|\exp(tz_r) - \exp(tz_s)\| < \rho$ . It follows that  $K(z_s)$  is homotopic in  $G_1$  to the curve  $K(z_r) \cup K_r^s$  consisting of  $K(z_r)$  followed by the arc  $K_r^s: \{f(u), r \leq u \leq s\}$ .

In particular, since  $\exp(0)=f(0)=e$ , there is an  $r > 0$  and an element  $z_r$  such that  $K(z_r)$  is homotopic to the arc  $K_0^r: \{f(s) | 0 \leq s \leq r\}$ . The set of all real numbers  $r$  for which this holds has a least upper bound,  $\mu \leq 1$ . Suppose  $\mu < 1$ . We obtain a contradiction.

Let  $\exp(z'_\mu)=f(\mu)$ . There is an element  $z'_s$  such that  $K(z'_\mu)$  is homotopic to  $K(z'_s) \cup K_s^\mu, s < \mu$ . But there is a  $z_s$  such that  $K(z_s)$  is homotopic to  $K_0^s$ . Since  $\exp(z'_s)=\exp(z_s)=f(s)$ , we have  $z_s=z'_s+c$  where  $\exp(c)=e$ . Let  $z_\mu=z'_\mu+c$ . It is simple to show that  $K(z_\mu)$  is homotopic to  $K(z_s) \cup K_s^\mu$ , which is, in turn, homotopic to  $K_0^s \cup K_s^\mu = K_0^\mu$ . Hence  $K(z_\mu)$  is homotopic to  $K_0^\mu$ . By the same reasoning, there is a number  $s_1 > \mu$  such that  $K_0^{s_1}$  is homotopic to  $K(z_{s_1})$ , contradicting the assumption on  $\mu$ . Therefore,  $\mu = 1$ .

The next theorem then follows easily.

**THEOREM 3.** Every closed curve in  $G_1$  with  $e$  as initial and end point is homotopic in  $G_1$  to a curve  $K(b)$  of the form  $\{\exp(tb) | 0 \leq t \leq 1, \exp(b)=e\}$ . If  $\exp(b')=e$  and  $b \neq b'$ , then  $K(b')$  is not homotopic to  $K(b)$ . Thus each homotopy class contains precisely one curve of the form  $\{\exp(tb), 0 \leq t \leq 1\}$  where  $\exp(b)=e$ .

PROOF. The first part of the theorem is obtained by choosing any point  $w=f(s)$  on  $K$ . By Theorem 2, there is a  $z$  such that  $K(z)$  is homotopic to  $K_0^s$  and a  $z'$  such that  $K(z')$  is homotopic to  $K_s^1$ . The element  $b=z'-z$  gives the desired result.

Noting that  $\int_{K(b)} x^{-1} dx = b$  and  $\int_{K(b')} x^{-1} dx = b'$ , we see that  $K(b)$  cannot be homotopic to  $K(b')$ , for in that event, the integrals would

be equal by the Cauchy integral theorem (by [1] and a result in the author's dissertation not yet published).

Theorem 1 follows directly from Theorem 3.

#### BIBLIOGRAPHY

1. E. R. Lorch, *The theory of analytic functions in normed abelian vector rings*, Trans. Amer. Math. Soc. vol. 54 (1943) pp. 414-425.
2. L. Pontrjagin, *Topological groups*, Princeton University Press, 1946, p. 231.
3. M. Nagumo, *Einige analytische Untersuchungen in linearen metrischen Ringen*, Jap J. Math. (1936).

COLUMBIA UNIVERSITY AND  
UNIVERSITY OF MARYLAND

#### REMARK ON A FORMULA FOR THE BERNOULLI NUMBERS

L. CARLITZ

Some years ago Garabedian [1] proved the following formula:

$$(1) \quad B_{k+1} = \frac{(-1)^{k+1}(k+1)}{2^{k+1}-1} \sum_{r=0}^k (-1)^r \frac{\Delta^r 1^k}{2^{r+1}} \quad (k \geq 0),$$

where the even suffix notation is employed for the Bernoulli numbers. The proof of (1) made use of the sum of a certain divergent series.

We wish to point out that (1) is not new. It can be found (in somewhat different notation) in [3, p. 224, formula (68)].

It may be of interest to give a short proof of (1). We use the formula [2, p. 28]

$$(2) \quad C_k = 2^{k+1}(1 - 2^{k+1}) \frac{B_{k+1}}{k+1},$$

where the  $C_k$  are the coefficients in the Euler polynomial:

$$(3) \quad E_k(x) = \left(x + \frac{C}{2}\right)^k = \sum_{s=0}^k \binom{k}{s} 2^{-s} C_s x^{k-s}.$$

Then in view of

$$(4) \quad E_k(x+1) + E_k(x) = 2x^k,$$

we have

$$(5) \quad E_k(x) = \left(1 + \frac{1}{2} \Delta\right)^{-1} x^k = \sum_{s=0}^k (-1)^s 2^{-s} \Delta^s x^k.$$

Received by the editors September 5, 1952.