

A CONNECTED COUNTABLE HAUSDORFF SPACE

R. H. BING

In Math. Ann. vol. 94 (1925) pp. 262–295, Urysohn gave an example of a connected Hausdorff space with only countably many points. Here is another.

EXAMPLE 1. The points of the space are the rational points in the plane on or above the x -axis. If (a, b) is such a point and $\epsilon > 0$, $(a, b) + \{(r, 0) \mid \text{either } |r - (a + b/3^{1/2})| < \epsilon \text{ or } |r - (a - b/3^{1/2})| < \epsilon\}$ is a neighborhood.

To construct geometrically a neighborhood with center at (a, b) , consider an equilateral triangle with base on the x -axis and apex at (a, b) . (If $b = 0$, regard (a, b) as the triangle.) Then (a, b) plus all rational points on the x -axis whose distances from a base vertex of the triangle are less than ϵ is an ϵ -neighborhood with center at (a, b) .

This space satisfies the Hausdorff axioms and has the property that for each pair of neighborhoods, there is a point common to their closures. Hence, the space is connected.

Although this space has a countable basis, it is not regular and hence not metric. Its dimension depends on the definition of dimension used. In the Menger-Urysohn sense (the dimension is defined inductively in terms of boundaries of open sets) the space is one-dimensional, and in the Lebesgue sense (the dimension is defined in terms of orders of coverings) it is infinite-dimensional.

EXAMPLE 2. We may enlarge our description to get a connected countable Hausdorff space of any positive dimension (even infinite) in the Menger-Urysohn sense. We give one of dimension two.

The points of the space are the rational points of Euclidean 3-space whose second and third coordinates are non-negative. If $(a, b, 0)$ is a point, an ϵ -neighborhood with center at $(a, b, 0)$ is $(a, b, 0) + \{(r, 0, 0) \mid \text{either } |r - (a + b/3^{1/2})| < \epsilon \text{ or } |r - (a - b/3^{1/2})| < \epsilon\}$. If (a, b, c) , $c \neq 0$, is a point, an ϵ -neighborhood with center at (a, b, c) is the sum of (a, b, c) and all ϵ -neighborhoods with centers at points $(a, r, 0)$ where either $|r - (b + c/2^{1/2})| < \epsilon$ or $|r - (b - c/2^{1/2})| < \epsilon$.

THE UNIVERSITY OF WISCONSIN

Received by the editors August 15, 1952.